

22-002
no. 11
D 1957
May 29
c. 2

Historical File Copy

N BUREAU OF STATISTICS

OTTAWA - CANADA

Authority of the Rt. Hon. C. D. Howe, Minister of Trade and Commerce



DOMINION BUREAU
OF STATISTICS

For Release 1957

May 29, 1957, 3 p.m.

PROPERTY OF THE
LIBRARY

C.R. No. 7, 1957

Price: \$2.00 for series of 23 Crop Reports.

TELEGRAPHIC CROP REPORT — PRAIRIE PROVINCES

This is the fourth of the 1957 series of thirteen telegraphic reports, issued by the Dominion Bureau of Statistics, covering crop conditions in the Prairie Provinces. A selected list of crop correspondents chosen from the Federal and Provincial Departments of Agriculture, private crop observers and grain men supply the information on which these reports are based. The weather data included in this release are furnished by the Meteorological Division, Department of Transport.

S U M M A R Y

Wheat seeding is rapidly drawing to a close in the Prairie Provinces and most of the coarse grains have also been sown. A somewhat smaller proportion of rape and flax seedings has been completed. Moisture supplies have been sufficient to ensure good germination of early-seeded crops. However, soils are becoming dry over fairly wide areas, particularly in Saskatchewan and Alberta and general rains would be welcome in such areas.

Wet weather held up seeding in Manitoba for a day or two last week. Most of the wheat and a large proportion of the coarse grains have been seeded but much of the flaxseed and rapeseed remains to be done. In a number of instances farmers are awaiting wild oat growth before completing seeding. Moisture conditions are generally satisfactory and germination of early-sown grains has been good. Pastures and meadows are also showing good growth. Rain would be welcome soon, however, in some areas as stubble lands are dry. Grasshoppers are starting to hatch in significant numbers in the Carberry area.

Wheat seeding is nearing completion in most districts of Saskatchewan and farmers are making favourable progress in the remaining areas. About 90 per cent of the wheat and 75 per cent of the oats and barley are seeded. Less than 5 per cent of the wheat seeding remains to be done in southern, central, and west-central districts while in the Yorkton, Prince Albert and Lloydminster districts 15 to 20 per cent remains. Some 65 per cent of the intended flax acreage has been planted. Growth in early-seeded fields is reported as good but germination in late-seeded fields is slow due to cold weather and lack of adequate surface moisture. Moisture conditions are sufficient to start seeded fields but early rains are needed to promote proper development. Frost killed some early-seeded fields of flax and rape seed and some re-seeding is necessary but damage to other grains was not extensive. Farmers had some success in the control of wild oats this year as early growth enabled many to obtain a kill before seeding grains. Seeding trends continued to indicate a reduction in wheat acreage and a significant increase in barley, flax, and rapeseed acreage. Grasshopper activity is reported in some southern areas, with control measures under way.

Prepared in the Crops Section, Agriculture Division

Wheat seeding is all but completed in Alberta and sowing of coarse grains should be practically finished by June 1. Conditions have been generally ideal this spring for destroying weeds and for early seeding. Fair rains have fallen in western areas but only showers have been received in other regions. Early crops have germinated well, but will soon need rain, especially on stubble. Late crops will need rain to ensure even germination. Dry soil may reduce forage crop seedings to below earlier intentions. Recent cool weather is retarding growth but frosts last week did little apparent damage. Some slight crop injury by cutworms is reported in southern localities. The earliest hatch of grasshoppers in at least seven years is now occurring in southern Alberta. However, no serious general damage is expected. The pea aphid infestation on alfalfa is still severe in localized areas.

MANITOBA In southeastern Manitoba seeding is practically completed except in the Dominion City area where approximately 70 per cent has been done. Intermittent rain delayed field work last week. Germination has been good. Grasshoppers are slow in hatching. Since our last report only one-half day's seeding has been completed in the Altona area but activity is expected to be renewed by Wednesday morning. All crops are coming along well. About 20 per cent of the crop area, mostly intended for flax remains to be sown. These late seedings will benefit from a good weed kill. Around Morden most wheat is showing above ground, with late rains improving germination and pastures. Seeding of coarse grains is mostly complete, row crops are nearing completion, while some flax remains to be sown. Crop prospects are improving. Moisture is sufficient for the present but warm weather is needed.

In south-central Manitoba at Pilot Mound seeding of wheat is completed in most sections with coarse grains and flax 60 to 75 per cent done. Up to one-half inch of rain last week has improved stands of forage crops. Moisture supplies are ample at Melita, with 1.2 inches of rain received in the past week. Ninety-five per cent of the wheat, 80 per cent of the oats and barley and 70 per cent of the flax have been seeded. Farmers are waiting for a good growth of weeds before seeding the remainder of the crop. Pastures and hay lands are in good condition. The crop outlook is generally good but the rye crop is only fair to poor.

At Beausejour east of Winnipeg spring seeding progressed favourably in spite of last week's rain. Wheat seeding is almost completed with barley and oats 75 per cent seeded. Early-sown wheat is already up. Pastures and hay fields are in good growing condition. Wild oat growth on second-crop fields has been prolific. Some grasshopper hatching has been reported. At Portage la Prairie seeding ranges from completed on some farms to 75 per cent done on others. Moisture supplies are good. Some early-sown land will be re-seeded because of wild oats. Seed germination is good and pastures are in excellent condition. Some reports of grasshoppers have been received.

At Brandon in western Manitoba early-sown crops show uniform emergence and strong growth. Seeding is nearing completion in most districts. Some stubble fields are too dry for optimum germination of late-sown crops. Growth on pasture and hay fields is only fair and rain would be beneficial. Prospects for grain crops remain good. At Virden seeding is progressing somewhat faster than normal with quite a few farmers finished. Early fields are up and look quite clean. Recent rains have greatly improved the moisture situation.

In the Interlake area at Teulon seeding operations are in full swing. A few farmers have finished and most of the wheat has been sown. Some farmers are waiting for wild oat growth and will sow barley. Sugar beet planting has been completed but very little rape has been sown as yet. Up to 95 per cent of the wheat, 50 per cent of the

coarse grains and 25 per cent of the rape has been sown in the Arborg district. The moisture situation is very good. Pastures are much improved and the outlook is for a good hay crop. There has been some delay in seeding due to rain.

Seeding is 85 per cent completed in the Neepawa area. A good control has been obtained on weeds. Pastures are in fair condition, summerfallows are in good shape and the moisture situation is fair. Grasshoppers are commencing to hatch. Seeding was halted for two days last week in the Minnedosa area on account of rain. Wheat seeding is 90 per cent completed while oats and barley are 80 per cent sown. Some flax has been seeded. Most early-sown crops are in good condition but the odd field has had to be re-seeded because of heavy wild oat growth. The sweet clover weevil is doing much damage. Pastures are good in this area. At Shoal Lake cold, rainy weather during the early part of last week has delayed seeding. A light rain helped pasture and forage crops somewhat but a heavier rain is needed to bring them along more rapidly. One report was received of light frost damage on early-sown potatoes.

In northern Manitoba at Swan River the cool weather continues. Approximately 90 per cent of the wheat, 50 per cent of the barley, and 35 per cent of the oats have been sown. Forage growth is slow because of cold weather. Many farmers are waiting for wild oats to germinate before continuing with spring cultivation.

The Entomological Laboratory at Brandon reports that grasshoppers are hatching in economic numbers in the Carberry area. Control measures have not yet commenced. Average precipitation for the province since April 1 has been 3 per cent below normal as compared with 22 per cent below normal a week ago, 11 per cent below normal two weeks ago, and 42 per cent below normal a year ago. Mean temperature for the week ending May 27 was 2.7 degrees below normal as compared with 2.0 degrees above normal a week ago, 3.8 degrees above normal two weeks ago and 2.3 degrees above normal for the week ending May 28, 1956.

SASKATCHEWAN Our correspondent from Fillmore in southeast Saskatchewan reports that seeding is almost completed and summerfallowing is under way. There is no evidence of frost damage and growth generally is even and rapid. The Experimental Farm at Indian Head reports that dry weather during the past week has enabled farmers to proceed with seeding without interruption and wheat seeding is now completed. Early-seeded crops have emerged and growth in some fields has reached three to four inches. Seeding of wheat and barley is from 80 to 90 per cent completed. Flax seeding is getting under way. Wild oat infestation is severe in many early-seeded fields and this will have an adverse effect on crops later on.

The Experimental Farm at Swift Current indicates that seeding is practically completed and a start has been made on summerfallow operations. The weather has been cool and dry and, because of this, growth has been rather slow. Dry weather has also reduced seedings from original intentions. Rye is in shot blade. Livestock are in good condition but pastures are only fair and stands are deteriorating rapidly due to lack of moisture. Around Val Marie rains are badly needed. Crops are generally above ground but growth is slow due to cool weather. In the Leader district seeding of all crops has been completed. Sub-soil moisture is adequate but good rains are required to ensure uniform germination. Summerfallow operations are well under way. Some slight frost damage is reported from local areas.

Around Togo crop conditions remain favourable and given another week of good weather seeding will be completed. Many farmers are still holding up operations while waiting to obtain a kill of wild oats. Some local showers have been received but a general soaking rain will be needed soon. From Sturgis our correspondent reports that

wheat seeding is completed and some 65 per cent of the coarse grains are in the ground. Rain is needed to germinate late-seeded crops, while the wild oat infestation in early-seeded crops is heavy. In the Willowbrook district wheat seeding is completed and about 60 per cent of the oats and barley has been sown. Moisture reserves are generally adequate and the weather is cool, and cloudy. Pastures and gardens are very good.

The Entomological Laboratory at Saskatoon reports that crop development is relatively slow due to cold weather. Sub-soil moisture is satisfactory but surface soil is dry and late fields are emerging unevenly. Grasshoppers are beginning to hatch but no insect damage to cereal crops has been observed or reported. A severe infestation of pea aphid is reported on alfalfa in the irrigated area around Maple Creek. Around Rosetown seeding is completed and conditions are generally fair. Wheat is above ground but germination in the late fields is somewhat uneven because of dry surface soil. Only odd fields of flax and rapeseed are showing above ground. Slight frost damage is reported from local areas. The Experimental Farm at Scott reports that there is need for effective rains in most districts of west-central Saskatchewan as evidenced by parched hay and pasture fields. Growth of early-seeded grain is slow and germination in late-seeded fields is uneven. Frost during the past week, with 12 degrees in some areas, damaged barley and killed a small percentage of newly emerged flax and rapeseed.

In the Melfort district of northeast Saskatchewan warm weather during the past week has permitted good progress to be made with seeding operations. Some 30 to 50 per cent of the seeding is completed, and, although wild oat growth is slow, many farmers are waiting to obtain a kill before seeding. Hay and pasture crops are making fair growth although not as good as last year. In the Meota district of northwest Saskatchewan seeding is completed but moisture conditions are only fair and pastures are poor. Rain since April 1 to date has totalled only three-quarters of an inch.

Average precipitation for the province since April 1 has been 12 per cent below normal, as compared with 20 per cent above normal a week ago, 9 per cent above normal two weeks ago and 35 per cent below normal a year ago. Mean temperature for the week ending May 27 was 3.3 degrees below normal in contrast to 4.6 degrees above normal a week ago, 3.4 degrees above normal two weeks ago, and 6.4 degrees above normal for the week ending May 28, 1956.

ALBERTA Seeding is now completed in the Manyberries area in the southeast and summer-fallow land has been worked. Wheat is up three inches and coarse grains one and one-half inches. Moisture reserves are poor and rain is needed. The range is in only fair condition and is dry in some localities. Light frosts have occurred in low areas. Seeding is also completed at Medicine Hat where early grain is up four inches. Rain is needed. Some frost damage occurred on May 20, particularly to flax.

Grain seeding is almost completed in the area south of Calgary. Germination has been good but moisture is needed, particularly east of Lethbridge. Seeding of canning corn, peas and beans is completed, and about 30 per cent of the sugar beet crop has been thinned. Irrigating is in full swing in all districts. The weather has been dry and warm and growth good in the Cardston district. Rain is needed. At Claresholm seeding will be completed this week. Early-sown grains are well above ground, with fall rye heading. Winter wheat is a good crop. Wild oats are a problem in some localities. Moisture supplies are adequate and general prospects good.

Seeding is practically completed in the Vulcan and Brooks districts of south-central Alberta. However, a heavy frost last week did some damage to potatoes, flax, barley and gardens. Light cutworm damage is reported from some areas. A good general rain is needed in these districts.

East of Calgary at Hanna wheat seeding will be completed this week and only 20 per cent of the coarse grains remains to be sown. Early-sown grain is coming up nicely but the topsoil is very dry and later seedings are somewhat patchy. Seeding is completed at Calgary, with early-sown grain up two to three inches. An eight degree frost has damaged a small percentage of flaxseed. Moisture supplies are generally satisfactory but rain would benefit late-sown grains. Farther north at Olds wheat seeding is completed, with three-quarters of the oats and barley in the ground. Rain is required in the next ten days.

All of the wheat and 80 per cent of the coarse grains have been seeded in the Stettler district of central Alberta. Moisture conditions are fair but rain will be needed in a week or ten days to bring crops along favourably. Farther north, at Donalda, seeding is practically completed, apart from a little barley and greenfeed. Crops are growing well with wheat covering the ground in some fields. Moisture reserves are fair and hay and pasture crops are in good condition. At Sedgewick, 95 per cent of the wheat and 60 per cent of the coarse grains are seeded. Frost has damaged some early rape seedings. Pastures are good but rain will be needed soon.

Excellent growing conditions have prevailed in the Red Deer area of west-central Alberta, with crops showing good germination and growth. Seeding is completed except for late barley and greenfeed. Light frosts have been reported but no damage is anticipated. Pastures are good and a heavy hay crop is in prospect. The presence of sweet clover weevil has been noted in the district. Farther north at Lacombe seeding has been completed except for about 20 per cent of the barley acreage which will not go in the ground until after June 1. With continued dry weather, moisture reserves are becoming depleted and hay and pastures particularly would benefit from immediate heavy rains. Wheat in some fields is averaging six inches in height.

At Vermilion east of Edmonton wheat seeding is completed and germination of all grains is good. Growth is well under way and pastures and livestock are in good condition. Rain is needed soon. All of the wheat and 80 per cent of the coarse grains have been seeded in the Edmonton district. Moisture is needed for later seedings.

Seeding operations are nearing completion in the Bonnyville area northeast of Edmonton. However, germination and growth has been slow due to abnormally cool weather. First-seeded grain is just emerging and forage crops are at a standstill. Many farmers are still dry feeding cattle to save pastures. Rain would be welcome. Rain totalling .92 inches has provided excellent soil moisture conditions in the Athabasca area. Crops are germinating very well except for the occasional early field which baked following earlier rains. Seeding should be completed by the end of this week. Pastures are in excellent condition. Nearby at Colinton, and also at Athabasca, crops are making excellent recovery from earlier frosts. Moisture supplies are adequate.

Around Falher in the eastern part of the Peace River district seeding has been completed except for purposely delayed late barley. Early-seeded flax has been slightly damaged by frost. Early-sown grain is up but moisture conditions are only fair to poor, with only light showers received so far this season. Prospects are good for grasses and legumes but pasture growth is slow. The Experimental Farm at Beaverlodge in the western part of the Peace River area reports seeding of spring cereals virtually completed except for some barley which is being held back for wild oat control. Germination has been slow, but is uniform and vigorous with first-seeded cereals now emerging. At Fort Vermilion in the far north emergence of wheat and early-seeded flax is general. Seeding of oats, barley and flax is still in progress. Soil moisture

conditions are very good, with snow and rain during the past week temporarily delaying field operations. Adapted varieties of winter cereals show satisfactory winter survival.

Average precipitation for the province since April 1 has been 23 per cent below normal as compared with 13 per cent below normal a week ago, 16 per cent below normal two weeks ago and 47 per cent below normal a year ago. Mean temperature for the week ending May 27 was 1.6 degrees below normal as compared with 2.9 degrees above normal a week ago, 0.1 degrees below normal two weeks ago and 7.2 degrees above normal for the week ending May 28, 1956.

Precipitation and Temperature Data, Prairie Provinces^{1/}

Province and Crop District	Station	Precipitation			Mean Temperature	
		Week ending	Total	Normal	Week ending 8 a.m.	
		8 a.m. May 27, 1957	since April 1	since April 1	May 27 1957	Normal
		- inches -			- degrees F. -	
MANITOBA						
1	Pierson	1.82	4.12	2.92	49	53
	Melita	1.46	2.72	3.07	51	53
	Waskada	N.R.	1.16 ^{2/}	2.33	N.R.	54
	Deloraine	.77	2.31	2.69	47	54
2	Boissevain	.75	3.18	3.02	51	53
	Deerwood	1.39	3.08	2.69	52	54
	Ninette	.75	1.90	3.20	51	53
	Portage la Prairie	1.67	3.31	2.73	51	54
3	Macdonald	.64	2.24	2.73	51	54
	Graysville	N.R.	1.82 ^{2/}	2.70	N.R.	53
	Morden	1.33	2.73	2.87	52	54
	Altona	1.35	2.32	2.91	52	54
	Morris	1.07	2.80	2.56	52	54
	Roland	1.66	2.90	2.54	52	54
	Gretna	1.04	2.25	2.23	52	54
	Emerson	1.60	3.13	2.23	51	54
4	Winnipeg	1.44	3.19	3.08	48	55
6	Sprague	1.80	3.65	3.04	51	52
	Seven Sisters Falls	.45	.88	1.97	51	52
	Steinbach	1.39	4.04	2.56	51	54
7	Virden	.73	1.79	2.03	50	52
	Reston	N.R.	1.04 ^{2/}	2.58	N.R.	53
	Rivers	.41	1.69	2.59	51	53
8	Brandon	.54	1.78	2.56	51	53
	Cypress River	.42	2.47	2.67	51	53
9	Neepawa	.27	2.41	2.59	50	53
10	Russell	.20	.86	2.23	N.R.	51
	Rosburn	.40	2.42	2.23	48	51
	Birtle	.55	1.85	2.28	49	51
11	Dauphin	.41	2.44	2.01	49	52
12	Gimli	.85	2.22	3.15	51	51
13	Swan River	.47	2.33	1.95	47	50
	The Pas	.04	1.84	1.75	43	51
MANITOBA AVERAGE		.92	2.50	2.57	50.2	52.9
SASKATCHEWAN						
1A	Estevan	.67	3.47	2.53	49	52
	Carlyle	.49	1.74	2.84	47	52
	Oxbow	N.R.	.67 ^{2/}	2.75	N.R.	52
	Willmar	.65	1.89	2.60	N.R.	N.R.
1B	Broadview	.38	1.19	2.46	47	52
	Moosomin	.59	1.93	2.13	51	52
2A	Yellow Grass	.57	2.08	2.41	48	52
	Creelman	.76	3.05	2.22	48	52
	Weyburn	.64	2.44	2.65	50	52
	Midale	.82	3.02	2.88	50	52

Precipitation and Temperature Data, Prairie Provinces^{1/}

Province and Crop District Station		Precipitation			Mean Temperature	
		Week ending	Total	Normal	Week ending 8 a.m.	
		8 a.m. May 27, 1957	since April 1	since April 1	May 27	
		- inches -			- degrees F. -	
SASKATCHEWAN (continued)						
2B	Moose Jaw	.54	2.71	2.30	50	54
	Regina	.41	2.37	2.11	49	52
	Rowatt	.52	1.73 ^{2/}	2.16	N.R.	52
	Francis	.43	1.89	1.59	48	52
	Qu'Appelle	N.R.	.57 ^{2/}	2.83	N.R.	52
	Indian Head	.41	3.18	2.42	49	52
	Wilcox	.59	1.98	2.25	49	52
3AS	Assiniboia	.78	2.52	1.82	47	54
	Orniston	.46	2.08	2.06	49	54
	Readlyn	.60	2.50 ^{2/}	1.82	50	54
	Viceroy	.83	2.74 ^{2/}	1.82	49	52
	Minton	.65	3.18	2.93	47	53
	Cardross	.46	2.10	2.06	49	54
	Ceylon	.73	2.59	3.31	48	53
3AN	Chaplin	.03	1.12	2.69	49	54
	Gravelbourg	.47	2.43	1.86	47	52
	Coderre	.60	2.83	2.01	48	52
3BS	Shaunavon	.29	1.59	2.00	48	50
	Cadillac	.45	1.65 ^{2/}	3.21	46	50
	Pambrun	.20	2.70	2.06	48	53
	Aneroid	.50	1.74	2.21	48	51
3BS	Instow	.14	1.21	2.02	48	52
3BN	Pennant	.03	1.45	2.53	49	54
	Swift Current	.16	1.65	2.32	48	55
	Hodgeville	.02	1.76	2.11	49	54
	Hughton	.02	2.43	2.76	50	53
	Maple Creek	NIL	.54	2.31	52	53
4A	Consul	NIL	.41	2.35	N.R.	51
	Roadene	.07	.99	2.78	49	53
5A	Cupar	.33	2.04	2.02	50	52
	Leross	.29	2.78	2.21	47	50
	Melville	.36	1.80	2.35	49	51
	Yorkton	.31	1.28	2.27	49	52
	Bangor	.30	1.22	2.35	50	52
	Dafoe	.15	1.45	1.83	49	49
5B	Foam Lake	.21	1.41 ^{2/}	2.19	47	49
	Lintlaw	.15	1.24	2.55	46	49
	Kamsack	.26	1.88	1.67	49	50
	Buchanan	.21	1.74	2.19	48	50
	Pelly	.28	2.28	1.85	46	50
	Davidson	.16	2.86	2.07	48	51
	Dilke	.20	2.62 ^{2/}	2.26	N.R.	N.R.
6A	Imperial	N.R.	1.56 ^{2/}	1.87	N.R.	51
	Semans	.30	1.40	1.66	50	50
	Strasbourg	.05	1.44	2.33	48	52
	Watrous	.20	2.16	1.67	48	52

Precipitation and Temperature Data, Prairie Provinces^{1/}

Province and Crop District	Station	Precipitation			Mean Temperature	
		Week ending 8 a.m. May 27, 1957	Total since April 1	Normal since April 1	Week ending 8 a.m. May 27 1957	Normal
		- inches -			- degrees F. -	
SASKATCHEWAN (concluded)						
6B	Harris	NIL	2.05	1.59	48	52
	Outlook	.08	1.48	1.72	51	52
	Saskatoon	.11	1.78	1.86	49	52
	Elbow	.12	1.91	1.93	51	53
	Tugaske	.30	2.44	1.90	49	52
	Dundurn	.16	1.57	2.02	49	52
	Northern	NIL	1.91	2.20	49	52
7A	Kindersley	Trace	.63	1.89	50	50
	Rosetown	.02	2.10	2.26	47	50
7B	Macklin	.09	.58	2.87	49	49
	Scott	.04	.77	2.07	49	52
	Biggar	.01	1.13	1.87	50	51
	Ruthilda	.04	.58 ^{2/}	2.09	N.R.	N.R.
8A	Hudson Bay	.18	1.98	2.06	46	52
	Porcupine Plain	.24	1.62 ^{2/}	2.07	46	51
8B	Humboldt	.06	2.08	1.88	47	52
	Melfort	.79	3.07	2.27	48	52
9A	North Battleford	Trace	.78	1.86	51	55
	Rabbit Lake	.03	1.50	1.78	47	51
	Leask	Trace	2.61	2.03	49	52
	Prince Albert	.09	3.13	2.06	49	53
	Island Falls	.24	3.60	2.05	42	49
9B	Waseca	.06	.83	2.05	48	51
SASKATCHEWAN AVERAGE		.29	1.94	2.20	48.5	51.8
ALBERTA						
1	Bindloss	NIL	2.14	2.30	52	54
	Empress	Trace	.60	2.22	51	56
	Excel	NIL	1.21	1.63	50	53
	Foremost	.03	1.12	3.68	52	58
	Hanna	Trace	1.62	2.65	50	54
	Manyberries	.06	1.07	2.57	51	55
	Medicine Hat	Trace	1.80	2.03	52	55
	Naco	Trace	1.21 ^{2/}	2.91	49	51
	Oyen	NIL	1.70	1.63	N.R.	N.R.
	Winnifred	NIL	1.69 ^{2/}	2.71	N.R.	N.R.
	Brooks	.05	1.16	2.27	52	55
	Drumheller	.09	1.87	2.31	53	52
	Gleichen	Trace	2.31	2.31	53	52
	Hays	.02	.94	2.06	52	53
2	Hussar	.01	2.54	2.66	N.R.	N.R.
	Lethbridge	.44	2.45	2.56	53	54
	Strathmore	.05	1.18	2.40	49	50
	Taber	.09	1.06 ^{2/}	2.46	52	53
	Three Hills	N.R.	1.93 ^{2/}	1.95	N.R.	50
	Trochu	.01	1.64	2.24	50	50
	Vauxhall	.06	.79	2.15	51	54
	Vulcan	.09	2.09	2.45	51	51



1010505541

- 10 -

Precipitation and Temperature Data, Prairie Provinces^{1/}

Province and Crop District	Station	Precipitation			Mean Temperature	
		Week ending 8 a.m.	Total since	Normal since	Week ending 8 a.m.	May 27
		May 27, 1957	April 1	April 1	1957	Normal
		- inches -			- degrees F. -	
<u>ALBERTA</u> (concluded)						
3	Calgary	.02	1.86	2.67	49	52
	Cardston	.49	3.52	4.27	51	54
	Cowley	.46	4.09	3.07	50	51
	Fort McLeod	.28	1.67	2.25	53	55
	High River	.15	3.27	3.31	48	51
	Magrath	.17	2.00	3.32	52	54
	Olds	.05	2.39	2.94	50	51
4	Alliance	.68	1.98	1.99	50	52
	Camrose	.24	1.81	2.71	51	50
	Coronation	.06	1.01	2.22	50	52
	Hardisty	.18	1.87 ^{2/}	1.77	N.R.	N.R.
	Hughenden	.05	1.17	2.28	N.R.	54
	Lloydminster	.05	1.29	1.82	50	49
	Stettler	.26	2.05	3.31	51	51
	Vegreville	.10	.45	2.65	51	51
	Vermilion	.09	1.85	2.23	49	50
	5	Edmonton	.15	1.21	2.27	48
Lacombe		.17	2.25	2.39	50	51
Red Deer		.06	2.30	3.06	48	50
Rocky Mountain House		Trace	1.66	3.10	50	49
Wetaskiwin		.33	1.54	2.03	50	51
6	Athabasca	1.12	2.50	2.31	48	51
	Campsie	.28	1.25 ^{2/}	2.17	50	51
	Edson	.11	2.04	2.02	48	51
	Elk Point	.15	.71	1.98	47	49
	Embarras	NIL	.13	1.34	46	47
	Lac la Biche	.18	.83	2.22	48	51
	Whitecourt	.20	1.51	2.64	49	51
7	Beaverlodge	.04	1.75	1.92	51	51
	Berwyn	N.R.	.30 ^{2/}	1.89	N.R.	51
	Fairview	.02	.74	1.41	52	52
	Grande Prairie	.14	1.56	2.12	52	51
	High Prairie	.08	.70	1.83	50	53
	Keg River	.28	1.38	2.32	N.R.	51
	Rycroft	.04	.70	1.35	51	51
	Wagner	.23	1.54	2.34	49	50
	ALBERTA AVERAGE		.15	1.82	2.37	50.3

N.R. - No report.

^{1/} - Source: Meteorological Service of Canada.

^{2/} - Incomplete; not included in average.