

Acute Toxicity to Juvenile Pacific Salmonids of Garlon 3A™, Garlon 4™, Triclopyr, Triclopyr Ester, and Their Transformation Products: 3,5,6-Trichloro-2-pyridinol and 2-Methoxy-3,5,6-trichloropyridine

M. T. Wan, D. J. Moul, and R. G. Watts

Environment Canada, Conservation and Protection, Environmental Protection,
Pacific Region, Kapilano 100, Park Royal, West Vancouver, B.C., Canada V7T 1A2

Garlon 3A™ and Garlon 4™ are formulated triclopyr herbicides manufactured by The Dow Chemical Company. These products are selective post-emergent herbicides used for the control of broadleaf vegetation. The use of Garlon herbicides for conifer release in coastal British Columbia may result in the unintentional introduction of triclopyr and its degradation products into waterbodies inhabited by salmonids. The objective of this study was to evaluate the acute toxicity of formulated and technical triclopyr and its transformation products to juvenile salmonids of the Pacific Northwest.

MATERIALS AND METHODS

The common names of the test chemicals are summarized in Table 1. All test materials were supplied by Dow Chemical U.S.A., Midland, Michigan.

A series of 96-h static acute toxicity tests was conducted from March to June 1986 and March to April 1987 in accordance with the protocols of Environment Canada (1980) and Buchanan (1982). Under yearling juvenile Pacific salmonids were obtained as fry in fresh water from various British Columbia hatcheries and used as test fish, (Table 2).

Before testing, all fish were acclimated to laboratory conditions for a minimum period of 2 wk. They were reared in fiberglass tanks at an average ambient temperature of 6°C (range 5-8°C) and were fed a maintenance diet of Oregon moist pellets. Fish under test were not fed. The average lengths of fish, weights, loading densities and test volumes were recorded, (Table 3). The loading density and test volume were adjusted according to the size of the fish.

Ten fish per concentration were placed in each test vessel. Observations of fish mortality were recorded at 24

Send reprint requests to Michael T. Wan.

Table 1. Nomenclature of test chemicals

Common name	Chemical name, active ingredient, purity
Garlon 3A TM	triethylamine of triclopyr; 36% acid equivalent of triclopyr
Garlon 4 TM	butoxyethyl ester of triclopyr; 48% acid equivalent of triclopyr
Triclopyr ester	butoxyethyl ester of triclopyr; 99.7%
Triclopyr	triclopyr[[(3,5,6-trichloro-2-pyridinyl)oxy]acetic acid]; 99.2%
Pyridinol	3,5,6-trichloro-2-pyridinol; 99.7%
Pyridine	2-methoxy-3,5,6-trichloropyridine; 99.7%

Table 2. Test fish species, ages, and suppliers

Fish Species	Date Obtained	Age (mo)	Hatchery
Coho salmon (<i>Oncorhynchus kisutch</i>)	05/15/86	1	Capilano Hatchery Vancouver, B.C.
Chinook salmon (<i>O. tshawytscha</i>)	05/07/86	4-5	Capilano Hatchery Vancouver, B.C.
Chum salmon (<i>O. keta</i>)	03/04/87	2-3	Little Qualicum Spawning Channel, Qualicum Beach, B.C.
Pink salmon (<i>O. gorbuscha</i>)	05/01/86	1	Quinsam Hatchery Campbell River, B.C.
Sockeye salmon (<i>O. nerka</i>)	04/02/86	0.75	Fisheries/Ocean Canada Research Laboratory, Cultus Lake, B.C.
	03/03/87	1.75	
Rainbow trout (<i>Salmo gairdneri</i>)	05/08/86	1	Fraser Valley Trout Hatchery, Abbotsford, B.C.
	05/12/86	2	
	25/02/86	1	

48, 72, and 96 h after introduction. At each observation period, the temperature, pH, and dissolved oxygen of each test vessel were recorded. A fish was considered dead when movement ceased and no further responses could be obtained by prodding or moving the animal. Fish found dead during the test were removed at each observation. Cumulative fish mortality was recorded and the results used for the calculation of LC₅₀ values based on the "Lethal" computer program developed by Stephan (1983).

The tests were conducted in 35-L glass aquaria. A maximum volume of either 20 or 30 L of dechlorinated Vancouver City tap water was used in each vessel, depending

Table 3. Average sizes and densities, and

Fish	Length cm (range)	Density
Coho	4.0 (3.5-4.5)	0
Chum	4.5 (3.9-5.0)	0
Chinook	6.8 (5.8-7.5)	2
Pink	3.5 (3.4-3.7)	0
Sockeye	3.9 (3.5-4.3)	0
Rainbow	4.1 (3.7-4.5)	0

on the size of the test aerated to prevent possible oil free compressed air column at the rate of 7.5

The starting temperature it stabilised at 14°C after 30-min aeration. The test was designed to minimize the shock to fish by gradually raising the ambient water temperature.

Chemical quality of dilute water samples was checked throughout the study period from March to April, 1987. Heavy metal concentrations were 3.25 to 3.81 mg/L during the test. Heavy metals were less than the background levels for most elements, i.e., (< 0.002) for Cd, Mo, Sr; (< 0.002) for Cd, for Ni, Sn. The average for other elements was: K (0.07); Mg (0.10); Na (0.07). Dissolved oxygen averaged 10.3 ± 0.1 mg/L to 6.0.

An appropriate amount of Garlon 4TM was dissolved in 25-mL beakers with distilled water to the exposure concentration. The solution which readily mixed and a volume of acetone was added to the solution. There were six nominal concentrations of 100 mg/L to 1000 mg/L and a control.

Duplicate 10 mL water samples were collected from one concentration of each treatment after 30-min aeration just before the test. The samples were analyzed to determine the concentration of chemicals in the test vessels.

The water samples were pre-

icals

ingredient, purity

olopyr; 36% acid
pyr
triclopyr; 48% acid
pyr
triclopyr; 99.7%
ichloro-2-pyridinyl)
).2%
/ridinol; 99.7%
chloropyridine; 99.7%

, and suppliers

Hatchery

Capilano Hatchery
Vancouver, B.C.

Capilano Hatchery
Vancouver, B.C.
Little Qualicum
Spawning Channel,
Qualicum Beach,
B.C.

Quinsam Hatchery
Campbell River,
B.C.

Fisheries/Ocean
Canada Research
Laboratory,
Cultus Lake, B.C.
Fraser Valley
Trout Hatchery,
Abbotsford, B.C.

ction. At each observa-
pH, and dissolved oxygen
ed. A fish was consider-
and no further responses
g or moving the animal.
est were removed at each
mortality was recorded and
ulation of LC50 values
er program developed by

5. glass aquaria. A max-
0 of dechlorinated Van-
in each vessel, depending

Table 3. Average sizes of fish, weights, loading densities, and test volumes

Fish	Length	Weight	Loading	Test
	cm (range)	g (range)	Density (g/L)	Volume (L)
Coho	4.0 (3.5-4.5)	0.5 (0.3-0.9)	0.28	20
Chum	4.5 (3.9-5.0)	0.5 (0.3-0.8)	0.27	20
Chinook	6.8 (5.8-7.5)	2.7 (1.4-3.8)	1.00	30
Pink	3.5 (3.4-3.7)	0.2 (0.2-0.2)	0.10	20
Sockeye	3.9 (3.5-4.3)	0.5 (0.3-0.6)	0.24	20
Rainbow	4.1 (3.7-4.5)	0.7 (0.4-0.9)	0.33	20

on the size of the test fish. The test solutions were aerated to prevent possible premature fish mortality. Oil free compressed air was bubbled through the water column at the rate of 7.5 mL/min/L during the test.

The starting temperature of the test water was 8°C, and it stabilised at 14°C after 24 h in the test chamber. The test was designed to start at a lower temperature to minimize the shock to fish from the holding tank where the ambient water temperature ranged from 6 to 8°C.

Chemical quality of diluent water was generally similar throughout the study periods of March to June, 1986 and March to April, 1987. Hardness as CaCO₃ ranged from 3.25 to 3.81 mg/L during the study periods. Inorganic metals were less than the detection limits (mg/L) for most elements, i.e., (< 0.005) for Pb, As, Cr, Cu, Mn, Mo, Sr; (< 0.002) for Cd, B, Be, Ti, Zn; and (< 0.02) for Ni, Sn. The average concentration (mg/L) for the other elements was: K (0.13); Al (0.12); Ca (1.2); Fe (0.07); Mg (0.10); Na (0.50), and Si (1.10). Dissolved oxygen averaged 10.3 ± 0.8 mg/L, and pH varied from 5.6 to 6.0.

An appropriate amount of each test chemical was dissolved in 25-mL beakers with 10 mL of acetone to obtain the exposure concentrations, except for Garlon 3A™ and Garlon 4™ as both materials contained an emulsifier which readily mixed and dispersed in water. An equal volume of acetone was added to the appropriate controls. There were six nominal concentrations ranging from 0.30 mg/L to 1000 mg/L and a control for each test.

Duplicate 10 mL water samples were collected at random from one concentration of each test chemical after the 30-min aeration just before fish introduction. They were analyzed to determine the actual amount of chemicals in the test vessels.

The water samples were prepared and extracted according

to the method outlined by Lee et al. (1986). Ethyl acetate was used as the extraction solvent instead of dichloromethane. The ethylated extracts were analyzed with a Hewlett Packard Model 5890 gas chromatograph equipped with a Ni-63 electron capture detector. The residue recovery rates of this method of analysis according to Lee et al. (1986) were greater than 95 %.

RESULTS AND DISCUSSION

The analytical results show that the measured concentration of each candidate material in the water of the test vessels immediately prior to fish introduction was less than the corresponding nominal concentration (Table 4).

Table 4. Concentrations of Test Chemicals in Water

Test chemical	Concentration of Test chemical (mg/L)		Recovery (%)
	Nominal	Measured*	
Garlon 3A TM	37.0	21.8	59
Garlon 4 TM	0.75	0.70	93
Triclopyr ester	1.17	0.98	84
Triclopyr	1.50	1.00	67
Pyridine	2.50	1.16	46
Pyridinol	1.56	1.03	66

* mean of duplicate samples

The data suggest that chemical loss occurred and the extent of loss appeared to vary for the different test chemicals. A number of factors may have contributed to this loss, viz., volatilization during the initial aeration process (Doudoroff et al. 1951); chemical degradation via photolysis and hydrolysis (McCall and Gavit 1986); reduced solubility of test chemicals (Dow Chemical 1981); and glass adsorption of test chemicals (Meehan et al. 1974).

Because of this chemical loss, the calculation of the 24, 48, 72, and 96-h LC₅₀ values for the test chemicals was adjusted to the measured concentration (Table 5). The results show that the LC₅₀ values changed only slightly after 96 h of exposure. The order of increasing toxicity to salmonids is as follows: Garlon 3ATM, triclopyr, pyridine, pyridinol, Garlon 4TM, and triclopyr ester, (Table 6). Triclopyr ester was the most toxic test chemical. Both Garlon 4TM and one of its transformation products, pyridinol were equally toxic to the salmonids tested.

Table 5. Acute toxicity to salmonids of triclopyr and triclopyr-2-trichloropyr

Test Chemical	Test Fish Species
Garlon 3A TM	coho chum sockeye rainbow chinook
Garlon 4 TM	coho chum sockeye rainbow chinook pink
Triclopyr ester	coho chum sockeye rainbow chinook pink
Triclopyr	coho chum sockeye rainbow chinook pink
Pyridine	coho chum sockeye rainbow chinook pink
Pyridinol	coho chum sockeye rainbow chinook pink

(1986). Ethyl vent instead of s were analyzed s chromatograph detector. The od of analysis is ter than 95 %.

sured concentra- ater of the test duction was less tion (Table 4).

micals in Water

Recovery (%)

59
93
84
67
46
66

occurred and the e different test ve contributed to ing the initial (1951); chemical ysis (McCall and st chemicals (Dow of test chemicals

luculation of the he test chemicals ation (Table 5). es changed only order of increas- ws: Garlon 3ATM, n 4TM, and tri- ster was the most M and one of its e equally toxic to

Table 5. Acute toxicities to juvenile Pacific salmonids of Garlon 3ATM, Garlon 4TM, triclopyr acid, triclopyr ester, 3,5,6-triclopyr-2-pyridinol, and 2-methoxy-3,5,6-trichloropyridine

Test Chemical	Test Fish Species	LC50 (mg/L)			
		24 h	48 h	72 h	96 h
Garlon 3A TM	coho	498	476	476	463
	chum	316	290	275	267
	sockeye	353	311	311	311
	rainbow	457	435	420	420
	chinook	472	312	283	275
Garlon 4 TM	coho	2.1	2.1	2.1	2.1
	chum	2.1	1.8	1.7	1.7
	sockeye	2.5	1.5	1.4	1.4
	rainbow	4.1	2.9	2.7	2.7
	chinook	4.2	2.7	2.7	2.7
Triclopyr ester	coho	1.0	1.0	1.0	1.0
	chum	0.4	0.3	0.3	0.3
	sockeye	0.5	0.4	0.4	0.4
	rainbow	1.1	1.1	1.1	1.1
	chinook	1.6	1.1	1.1	1.1
Triclopyr	coho	9.9	9.6	9.6	9.6
	chum	7.9	7.5	7.5	7.5
	sockeye	7.8	7.5	7.5	7.5
	rainbow	8.4	7.8	7.6	7.5
	chinook	9.7	9.7	9.7	9.7
Pyridine	coho	4.3	4.0	3.8	3.8
	chum	4.0	4.0	3.9	3.7
	sockeye	6.9	6.9	6.9	6.3
	rainbow	4.6	4.6	4.6	4.6
	chinook	3.2	2.9	2.9	2.9
Pyridinol	coho	1.8	1.8	1.8	1.8
	chum	1.8	1.8	1.8	1.8
	sockeye	2.5	2.5	2.5	2.5
	rainbow	1.5	1.5	1.5	1.5
	chinook	2.1	2.1	2.1	2.1
Triclopyr-2-pyridinol	coho	1.8	1.8	1.8	1.8
	chum	1.8	1.8	1.8	1.8
	sockeye	2.5	2.5	2.5	2.5
	rainbow	1.5	1.5	1.5	1.5
	chinook	2.1	2.1	2.1	2.1
2-methoxy-3,5,6-trichloropyridine	coho	1.8	1.8	1.8	1.8
	chum	1.8	1.8	1.8	1.8
	sockeye	2.5	2.5	2.5	2.5
	rainbow	1.5	1.5	1.5	1.5
	chinook	2.1	2.1	2.1	2.1

Table 6. Order of increasing toxicity to Pacific salmonids of test chemicals

Test Chemicals	Toxicity 96 h LC50 (Mean±S.E.) ^b	n	Student "t" test	"t" value	Level of Significance
1. Garlon 3A TM	347 + 44	5*	1 vs. 5	7.84	< 0.01
2. Triclopyr	7.9 ± 0.7	6	1 vs. 2	7.71	< 0.01
3. Pyridine	3.7 ± 0.8	6	5 vs. 6	4.60	< 0.01
4. Pyridinol	2.1 ± 0.2	6	2 vs. 6	9.89	< 0.001
5. Garlon 4 TM	2.0 ± 0.2	6			
6. Triclopyr ester	0.7 ± 0.2	6	(n = no. of fish species; * pink salmon not tested)		

Garlon 3ATM, the amine product of triclopyr, is significantly less toxic ($p < 0.01$) to salmonids than Garlon 4TM, ester formulation (Table 6). The results show that Garlon 3ATM was about 170 times less toxic to salmonids than Garlon 4TM. Within the various salmonid species, Garlon 3ATM is more toxic to chum, chinook and sockeye salmon than to coho salmon and rainbow trout (Table 5). Garlon 4TM is slightly more toxic to pink and sockeye salmon than to chum, coho, chinook salmon, and rainbow trout.

Moreover, Garlon 3ATM and Garlon 4TM are significantly less toxic ($p < 0.01$) to salmonids than their respective active ingredients triclopyr and triclopyr ester (Table 6). This observation suggests that the emulsifiers used in formulating these products do not increase fish toxicity.

Triclopyr is significantly less toxic ($p < 0.001$) to salmonids than triclopyr ester, (Table 6). This acid is about 11 times less toxic to the fish than triclopyr ester. Within species, however, triclopyr is more toxic to pink, chum, sockeye salmon and rainbow trout than to coho and chinook.

Triclopyr ester is more toxic to juvenile chum, pink and sockeye salmon than to coho and chinook salmon and rainbow trout (Table 5). The 96-h LC50 of coho salmon for triclopyr ester is 1.0 mg/L, confirming a recent study by Mayes et al. (1986) who reported that this material to be highly toxic to both the juvenile and alevin stages of coho salmon. The respective 96-h LC50 values were 1.3 and 0.3 mg/L.

Test results of the two principal transformation products of triclopyr suggest that the pyridinol metabolite is somewhat more toxic to juvenile salmon than the pyridine (Table 6). The pyridinol degradation product is also as toxic to young salmon as the formulated tri-

clopyr ester. This toxic to all species however, is more to five salmon species rainbow trout).

Under field conditions in a stream univalent operation would not 10 mg/L in 15 cm water rate of application, except perhaps during this product causing it is used under use of the lower rate 2.4 kg active ingredient generate toxic concentration (15 cm water) if the flushed out of the a

It is doubtful that pyridinol and pyridine in contaminated streams the occurrence of cases of the pyridine microbial activity may be potential for its transformation contaminated by Garlon. Studies have shown occurred in streams et al. 1976 and water toxicity of triclopyr products to young salmon presently not known.

To sum up, this bi Garlon 4TM, the ester formulation to salmonids of the Garlon 3ATM, the amine toxic to fish. Of products, the pyridinol 4TM to juvenile salmon more toxic to young species tested. Moreover that the emulsifier 3ATM and Garlon

Acknowledgments.

Wilson, R.H. Kussat, S. Szeto (Agriculture) S. Samis (Fisheries and (2) Agriculture

to Pacific

"t" value	Level of Significance
7.84	< 0.01
7.71	< 0.01
4.60	< 0.01
9.89	< 0.001

of fish species;
salmon not tested)

lopyr, is significantly more toxic to salmonids than Garlon 4TM. Results show that Garlon 4TM is more toxic to salmonid species, chinook and sockeye salmon and rainbow trout (Table 5). Garlon 4TM is more toxic to pink and sockeye salmon, and rainbow trout than to

are significantly more toxic than their respective parent compounds. Triclopyr ester is more toxic than the emulsifiable concentrate products do not

c (p < 0.001) to rainbow trout (Table 6). This acid is more toxic to fish than triclopyr ester. Triclopyr is more toxic to rainbow trout than to

file chum, pink and sockeye salmon and rainbow trout. In a recent study of coho salmon for rainbow trout, it was found that this material is more toxic to juvenile and alevin coho salmon than the emulsifiable concentrate LC₅₀ values

transformation product of triclopyr ester is more toxic to juvenile salmon than the parent compound. The degradation product of the emulsifiable concentrate formulated tri-

clopyr ester. This transformation product is equally toxic to all species of juvenile salmon. The pyridine, however, is more toxic to pink salmon than the other five salmon species (coho, chum, chinook, sockeye and rainbow trout).

Under field conditions, the concentration of Garlon 3ATM in a stream unintentionally oversprayed during an aerial operation would not likely exceed a level greater than 10 mg/L in 15 cm water even at the highest recommended rate of application, (i.e., 10 kg active ingredient/ha), except perhaps during an accident. The potential of this product causing fish kills is therefore small when it is used under prescribed conditions. However, the use of the lower recommended rate of Garlon 4TM, (i.e., 2.4 kg active ingredient/ha), has some potential to generate toxic concentrations (approximately 4 mg/L in 15 cm water) if the residues are not rapidly diluted and flushed out of the aquatic system.

It is doubtful that acutely toxic levels of both the pyridinol and pyridine degradation products would occur in contaminated streams, as their appearance requires the occurrence of certain chemical processes and, in the case of the pyridine, an incubation period involving microbial activity (Lee et al. 1986). However, there may be potential for sub-lethal effects of triclopyr and its transformation products in salmon waters contaminated by Garlon 3ATM and Garlon 4TM overspray. Studies have shown that low levels of these materials occurred in streams following aerial operations (Norris et al. 1976 and Wan 1986). The potential sub-lethal toxicity of triclopyr and its two transformation products to young salmon and other aquatic organisms is presently not known.

To sum up, this bioassay study indicates that Garlon 4TM, the ester formulation of triclopyr, is highly toxic to salmonids of the Pacific Northwest. In contrast, Garlon 3ATM, the amine formulation is considerably less toxic to fish. Of the two principal degradation products, the pyridinol metabolite is as toxic as Garlon 4TM to juvenile salmonids, while pyridine seems to be more toxic to young pink salmon than the other salmon species tested. Moreover, this study suggests indirectly that the emulsifiers used for the formulation of Garlon 3ATM and Garlon 4TM do not increase fish toxicity.

Acknowledgments. We wish to thank (1) Messrs. D.M. Wilson, R.H. Kussat, G. Rawn (Environmental Protection), S. Szeto (Agriculture Canada), J. Servizi, R.W. Gordon, S. Samis (Fisheries & Oceans Canada) for their comments; and (2) Agriculture Canada Research Laboratory,

Environment Canada Laboratory Services, Vancouver, B.C., and Dow Chemical U.S.A., Midland, Michigan, for technical support.

REFERENCES

- Buchanan RJ (1982) Provincial guidelines and laboratory procedures for measuring acute lethal toxicity of liquid effluents to fish. British Columbia Bioassay Task Force. British Columbia Ministry of Environment, Victoria. ISBN 0-7719-9174-6
- Doudoroff P, Anderson BG, Burdick GE, Galtsoff PS, Hart WB, Patrick R, Strong ER, Surber EW, VanHorn WM (1951) Bioassay methods for evaluation of acute toxicity of industrial wastes to fish. *Sewage Ind Wastes* 23:1380-1397
- Dow Chemical Company Inc (1981) Technical information on triclopyr - a new experimental herbicide. Sarnia, Ontario
- Environment Canada (1980) Standard procedure for testing the acute lethality of liquid effluents. Environmental Protection, Ottawa, Ontario. EPS 1-WP-80-1
- Lee CH, Oloffs PC, Szeto SY (1986) Persistence, degradation, and movement of triclopyr and its ethylene glycol butyl ether ester in a forest soil. *J Agric Food Chem* 34: 1075-1079
- Mayes MA, Murphy PG, Hopkins DL, Gersich FM, Blanchard A. (1986) The toxicity and metabolism of triclopyr butoxyethyl ester: coho salmon. *The Toxicologist. Society of Toxicology. Abstracts of the 25th Anniversary Meeting, New Orleans, Louisiana.* 6(1): p 26
- McCall PJ, Gavit PD (1986) Aqueous photolysis of triclopyr and its butoxyethyl ester and calculated environmental photodecomposition rates. *Environ Tox and Chem* 5: 879-885
- Meehan WR, Norris LA, Sears HS (1974) Toxicity of various formulations of 2,4-D to salmonids in southeast Alaska. *J Fish Res Board Canada* 33: 480-485
- Norris LA, Montgomery ML, Warren, LA (1987) Triclopyr persistence in western Oregon hill pastures. *Bull Environ Contam Toxicol* 39: (in press)
- Stephan CE (1983) Lethal programme for computer analyses of LC50. Environmental Research Laboratory, U.S. Environmental Protection Agency, Duluth, Minnesota
- Wan MT (1986) The persistence of triclopyr and its transformation product pyridinol in a coastal British Columbia stream. Environmental Protection, Vancouver, British Columbia. Regional Program Report: 86-24

Received January 26, 1987; accepted May 22, 1987