



Canadian National Vegetation Classification (CNVC) Classification nationale de la végétation du Canada

<http://cnvc-cnvc.ca>

North American Boreal Conifer Poor Swamp

Macrogroup M299

Marécages pauvres à conifères boréaux de l'Amérique du Nord

Boreal Flooded Swamp & Forest

D016 North American Boreal Flooded & Swamp Forest

M300 North American Boreal Flooded & Rich Swamp Forest

M299 North American Boreal Conifer Poor Swamp

CG0016 Atlantic Boreal Black Spruce - Balsam Fir Poor - Intermediate Treed Wetland

CG0019 Ontario-Quebec Boreal Black Spruce Poor - Intermediate Treed Wetland

CG0022 West-Central Boreal Black Spruce - Tamarack Poor - Intermediate Treed Wetland

CG0025 Subarctic Black Spruce Poor - Intermediate Treed Wetland



Concept

M299 describes nutrient-poor to medium wetland forests and woodlands throughout the boreal region of North America. These include bog and fen woodlands as well as poor to intermediate swamps, usually developed on peat deposits. This vegetation is characterized by black spruce (*Picea mariana*) and/or tamarack (*Larix laricina*) in the tree layer, overwhelming dominance of ericaceous species in the understory and ground cover of *Sphagnum* mosses. Typical understory species include common Labrador tea (*Rhododendron groenlandicum*), leatherleaf (*Chamaedaphne calyculata*), blueberries (*Vaccinium* spp.), shrub willows (*Salix* spp.), shrub birches (*Betula* spp.), bog laurel (*Kalmia polifolia*), three-leaved false Solomon's seal (*Maianthemum trifolium*), cloudberry (*Rubus chamaemorus*), small cranberry (*V. oxycoccus*) and bog rosemary (*Andromeda polifolia*). Sheep laurel (*K. angustifolia*) and rhodora (*Rhododendron canadense*) often replace *R. groenlandicum* in Atlantic Canada. Peat mosses (*Sphagnum* spp.) dominate the ground cover, but feathermosses (esp. red-stemmed feathermoss [*Pleurozium schreberi*] and staircase moss [*Hylocomium splendens*]) are common on dry microsites (e.g., peat hummocks).

These are generally stable ecosystems that are maintained by persistently high water tables within a cold climate. M299 occurs in boreal and subarctic climates, characterized by long, cold winters and short, cool to moderately warm summers. Mean annual temperatures in the Canadian range vary from approximately -10°C in Inuvik, Northwest Territories to >3.5°C in parts of insular Newfoundland. Mean annual precipitation generally follows a west to east gradient, increasing from <300 mm in the western subarctic to as high as 1800 mm in parts of Nova Scotia and insular Newfoundland. Substrates are usually *Sphagnum*-derived peats with nutrient regimes ranging from poor to medium, depending on local site-scale hydrology. Within a stand, hummocky microtopography associated with the growth of certain *Sphagnum* spp. provides a range of micro-scale moisture and nutrient gradients. Permafrost is a feature of subarctic and some northern boreal peatlands.



Bog woodland dominated by black spruce (*Picea mariana*) and common Labrador tea (*Rhododendron groenlandicum*) in northern Alberta.
Source: L. Allen



Black spruce (*Picea mariana*) poor swamp forest with understory dominated by common Labrador tea (*Rhododendron groenlandicum*) in northwestern Ontario.
Source: K. Baldwin



Canadian National Vegetation Classification (CNVC) Classification nationale de la végétation du Canada

<http://cnvc-cnvc.ca>

North American Boreal Conifer Poor Swamp

Macrogroup M299

Marécages pauvres à conifères boréaux de l'Amérique du Nord

Vegetation

Physiognomy and Structure

M299 describes boreal wetland forests and woodlands (i.e., >10% cover by trees >5 m height) that are dominated by *Picea mariana* and/or *Larix laricina*. These include bog and fen woodlands as well as poor to intermediate swamps, typically developed on peatlands. Physiognomically, the tree layer is usually evergreen or deciduous coniferous, although small amounts of broad-leaved cold-deciduous species (e.g., *Betula papyrifera*, *Alnus incana* or *Salix* spp.) occasionally occur. Stand density is highly variable, ranging from closed moderately productive forests to open stunted woodlands. Stands often contain trees of multiple ages that vary in height from 2 to >10 m; on poor sites small trees can be >100 yr. old. Although the main tree species are *P. mariana* and *L. laricina*, *Abies balsamea* can be a significant component in the Atlantic portion of the range. Understories usually have a well-developed low or dwarf shrub layer dominated by ericaceous species. The moss layer typically consists of continuous cover of *Sphagnum* mosses, feathermosses and lichens over shallow to deep peat substrates.

Floristics

Picea mariana characterizes M299 conditions across the entire (i.e., poor to medium) nutrient spectrum of these wetland forests and woodlands. *Larix laricina* occurs where the nutrient status is enhanced, even marginally, by groundwater inputs. In eastern Canada, *Abies balsamea* is consistently present on all but the poorest sites and *A. balsamea*-dominated intermediate swamps are encountered in Atlantic Canada.

Overall species diversity is low and the understory is dominated by ericaceous shrubs. *Rhododendron groenlandicum* is ubiquitous, except in Atlantic Canada where *Kalmia angustifolia* and *R. canadense* often replace it. *Vaccinium* spp. are also common: *V. myrtilloides* throughout the Canadian range except insular Newfoundland, *V. angustifolium* in eastern Canada and *V. vitis-idaea* in the northern and western portions of the range. *V. oxycoccus*, *K. polifolia* and *Chamaedaphne calyculata* are characteristic of bog and poor fen woodlands throughout the range, as are *Rubus chamaemorus* and *Maianthemum trifolium*. On sites with slightly enhanced nutrient status (intermediate fens and poor swamps), shrub birches (*Betula* spp.) and shrub willows (*Salix* spp.), as well as *Alnus incana* (mainly in eastern Canada; see Comments), are more common. These nutrient-medium sites also support greater abundance of graminoids (especially *Carex* spp.) and, primarily in western Canada, *Equisetum* spp.

The majority of these plant communities are characterized by a moss layer that is dominated by *Sphagnum* spp. However, some of the common *Sphagnum* mosses on the poorest sites (e.g., *S. fuscum*) form dense hummocks with dry exposed tops that are covered by feathermosses (typically *Pleurozium schreberi*, *Hylocomium splendens* and *Ptilium crista-castrensis*) and lichens (especially *Cladina* spp.). On sites with slightly enhanced nutrient status (usually intermediate fens), the moss layer is sometimes dominated by *Aulacomnium palustre* and other brown mosses (e.g., *Tomentypnum nitens*, *Sanionia uncinata*).

In northern parts of the range, *Vaccinium uliginosum*, *Empetrum nigrum* and *Rhododendron tomentosum* are common species in bog and poor fen woodlands. In eastern Canada, *Gaultheria hispidula* is ubiquitous. In Atlantic Canada, *A. balsamea*, *Ilex mucronata*, *Viburnum nudum* (see Comments) and *Osmundastrum cinnamomeum* are important species on sites with slightly enhanced nutrient status. Some species that are characteristic of open bogs and fens (e.g., *Sarracenia purpurea*, *Drosera* spp. and *Eriophorum* spp.; see M876 [North American Boreal & Sub-boreal Acidic Bog & Fen]) are uncommon in their M299 counterparts.

Dynamics

These are generally stable ecosystems that are maintained by persistently high water tables. Local hydrology is the main ecological determinant of vegetation characteristics. Any changes to water chemistry or level of the long-term water table will affect nutrient status, degree of aeration and soil temperature in the rooting zone, and thus influence overall species composition and tree productivity.

Enrichment of the rooting layer, typically by groundwater flow, results in the development of fen or swamp conditions because of increased supply of oxygen and macronutrients, and reduced acidity. If decomposition rates are very slow, bogs often develop from fen conditions as peat accumulates, elevating the rooting zone above contact with groundwater influences. In these cases, unless the local hydrology changes, depth of peat tends to increase over time, further reducing nutrient availability and promoting succession to open bog conditions.

A rise or drop in the water table (either by anthropogenic activities or natural causes [e.g., beaver dams]) can result in changes to the vegetation community. A rise in the water table can result in tree mortality and transition to open wetland vegetation. Paludification can also accompany elevated water tables in cold or humid climates, resulting in the expansion of peatlands onto neighbouring upland sites. A drop in the water table can reverse some of these effects and sometimes promote the development of more productive treed vegetation (e.g., as described in M495 & M496).

Cryoturbation associated with permafrost in the northern portions of the range can also disrupt plant growth on a local scale. "Drunken forests" occur when melting permafrost creates subsidence of surface substrates supporting M299 stands.

Although fires occur on peatlands, they are infrequent and of limited extent because these sites are so wet. Consequently, stands of M299 tend to be long lived and multi-aged, with trees up to or exceeding 200 years. *P. mariana* can establish from seed under favourable conditions (e.g., suitable seedbed) but typically self-replaces on these sites by vegetative layering. *L. laricina* usually establishes by seed.



Canadian National Vegetation Classification (CNVC) Classification nationale de la végétation du Canada

<http://cnvc-cnvc.ca>

North American Boreal Conifer Poor Swamp Marécages pauvres à conifères boréaux de l'Amérique du Nord

Macrogroup M299

Environment

M299 forests and woodlands typically develop on nutrient-poor to medium peatlands. Wetlands develop where the interaction between climate and landscape results in permanent or semi-permanent water tables at or near the soil surface. Peatlands form when the rate of plant production exceeds the rate of decomposition in a wetland environment. Local peat development is influenced by abiotic factors such as geological and hydrological conditions but, since this process is ultimately dependent on high levels of soil moisture and slow rates of plant decomposition, climate is a crucial factor affecting peatland formation. Cooler temperatures reduce rates of evapotranspiration and decomposition; as a consequence, the frequency and extent of peatlands on the landscape generally increase northward wherever level terrain and surficial conditions retain water near the land surface. When water, oxygen, nutrient and temperature regimes provide the necessary conditions for tree establishment and growth, forests and woodlands are able to develop on peatlands.

Climate

M299 occurs across the boreal zone of North America. Although bogs and fens are the dominant wetland classes (see the Canadian Wetland Classification System [CWCS]) throughout the boreal, regional differences in prevalent forms (see CWCS) are a response to climatic gradients of temperature and precipitation (as well as to variations in underlying geology and geochemistry). M299 develops within boreal and subarctic climates, characterized by long, cold winters and short, cool to moderately warm summers. Mean annual temperatures in the Canadian range vary from approximately -10°C in Inuvik, Northwest Territories to >3.5°C in parts of insular Newfoundland. Growing degree days above 5°C (GDD) vary from <600 in the subarctic to >1600 in southern Manitoba. Although some high elevation Cordilleran locations in northwestern British Columbia receive >1000 mm, mean annual precipitation generally follows a west to east gradient, increasing from <300 mm in the western subarctic to as high as 1800 mm in parts of Nova Scotia (Cape Breton Island) and insular Newfoundland.

Physiography, Geology, Topography and Soils

M299 occurs in all Canadian physiographic regions except those in the arctic and on the continental shelves. Terrain and geology (in addition to regional climate) control the distribution of wetlands on the landscape. M299 stands often occur within large wetland complexes in a mosaic with other treed and non-treed bog, fen, marsh and swamp conditions that reflect local hydrological gradients.

In the subhumid climate of west-central Canada, boreal peatlands occur only in water-collecting landscape depressions or where an external water source creates wetland conditions. Fens constitute the prevalent peatland class due to groundwater inputs and the nutrient-rich geochemistry of bedrock and glacial sediments of the Interior Plains and Cordilleran physiographic regions. In the humid climate of eastern Canada, bogs constitute the most frequent peatland class in the boreal zone. Greater precipitation and the predominantly acidic, nutrient-poor bedrock of the Precambrian Shield and Shield-derived glacial soils contribute to growth of *Sphagnum* mosses that create peat surfaces that are often raised above the surrounding water table (raised bogs). The generally humid environment of eastern Canada also contributes to paludification, giving rise to expansion of peatlands onto neighbouring upland sites. Blanket bogs occasionally develop on sloping terrain in parts of the Atlantic boreal region, where maritime climatic influences result in high levels of precipitation and atmospheric humidity.

Local to regional topography and geochemistry influence hydrological characteristics such as groundwater flow and nutrient content. For example, in the Rocky Mountain foothills of Alberta, generally calcareous tills and long gentle slopes supply nutrient-rich groundwater that supports the development of locally extensive fen woodlands and swamps. Conversely, Shield landscapes tend to comprise rolling, bedrock-controlled terrain with short drainage slopes and low relief; bedrock and glacial soils are typically acidic and nutrient-poor. Under these conditions, bogs and poor fens develop in topographic depressions; M299 forests and woodlands are often found around the margins of these basins, with non-treed conditions in the center.

Substrates are usually shallow to deep *Sphagnum*-derived peats, dominated by soils in the Organic order (according to the Canadian System of Soil Classification). *Sphagnum* mosses contribute to soil acidity by sequestering cations and releasing hydrogen ions into the immediate environment. Except in the spring, standing water is less common in M299 stands than in the mineral swamps described in M300 [North American Boreal Flooded & Rich Swamp Forest]. Nutrient regimes for forests and woodlands of M299 range from poor to medium, depending on local hydrology. Within a stand, hummocky microtopography associated with growth of certain *Sphagnum* spp. provides a range of micro-scale moisture and nutrient gradients. Where thick moss accumulations elevate the peat surface above the water table, minerotrophic groundwater may be brought to the rooting zone by the capillary action of the moss, but hummock tops often remain drier. In some cases, nutrient inputs are supplied largely by precipitation and dustfall (ombrotrophy); the poorest sites may not be able to support treed vegetation.

Permafrost is a characteristic of subarctic and some northern boreal peatlands. In the northern subarctic of central Canada, permafrost features often develop (e.g., palsas, peat plateaux and ice wedge polygons); southwards, these features occur only sporadically. Depth to permafrost affects the temperature of the active peat layer and thus, the effective growing season for peatland vegetation. Elevated permafrost features such as palsas and peat plateaux sometimes raise the peat surface above the surrounding water table, resulting in bog forms (often woodlands) surrounded by open fens.



Canadian National Vegetation Classification (CNVC) Classification nationale de la végétation du Canada


<http://cnvc-cnvc.ca>

North American Boreal Conifer Poor Swamp Marécages pauvres à conifères boréaux de l'Amérique du Nord

Macrogroup M299

Distribution and Geographic Range

M299 wetland forests and woodlands occur throughout the entire boreal region of North America, including Alaska and the subarctic. This vegetation occurs occasionally in the northern temperate region (esp. in eastern North America) where local climate, substrate and hydrological conditions create wetlands that do not support the growth of temperate species.



Related Concepts

M299 includes nutrient-poor to medium wetland forests and woodlands that have been described in provincial/territorial publications for the Boreal Low, Boreal High and Subarctic Woodland bioclimatic zones in Yukon; the Taiga Shield, Taiga Plains, Taiga Cordillera and Boreal Cordillera Level II ecoregions in Northwest Territories; the Boreal White & Black Spruce and Spruce – Willow – Birch biogeoclimatic zones in British Columbia; the Boreal Forest, Canadian Shield and Foothills natural regions in Alberta; the Boreal Shield, Taiga Shield and Boreal Plains ecozones in Saskatchewan and Manitoba plus the Hudson Plains ecozone in Manitoba; ecoregions OE [Hudson Bay Coast], 1E [Northern Taiga Ecoregion], 2E [James Bay], 2W [Big Trout Lake], 3E [Lake Abitibi], 3W [Lake Nipigon], 3S [Lake St. Joseph] and parts of 4E [Lake Temagami], 4W [Pigeon River], 4S [Lake Wabigoon], 5E [Georgian Bay] and 5S [Agassiz Clay Plain] in Ontario; the Balsam Fir – White Birch, Spruce – Moss, Spruce – Lichen and Forest Tundra bioclimatic domains in Quebec; the Highlands ecoregion of New Brunswick; the Northern Plateau and Cape Breton Highlands ecodistricts in Nova Scotia; the Taiga and Boreal Shield ecozones in Labrador; and all of insular Newfoundland.

Boreal upland forests and woodlands are described by M495 [Eastern North American Boreal Forest], M496 [West-Central North American Boreal Forest & Woodland], M156 [Alaskan-Yukon North American Boreal Forest] and M179 [North American Subarctic Woodland]. Temperate wetland forests and woodlands are described by M504 [Laurentian – Acadian – North Atlantic Coastal Flooded & Swamp Forest], M028 [Great Plains Floodplain Forest], M034 [Rocky Mountain - Great Basin Montane Riparian & Swamp Forest] and M035 [Vancouverian Flooded & Swamp Forest].

Comments

M299 describes nutrient-poor to medium wetland forests and woodlands, mostly occurring on shallow to deep peat deposits, in boreal North America. These communities are characterized by *Picea mariana* and/or *Larix laricina* in the tree layer, overwhelming dominance of ericaceous species in the understory and ground cover of *Sphagnum* mosses. Nutrient-rich boreal swamp and floodplain forests, described by M300 [North American Boreal Flooded & Rich Swamp Forest], include *Populus balsamifera* and, in eastern Canada, *Fraxinus nigra* and *Thuja occidentalis* among the dominant tree species. M300 vegetation also includes a diverse suite of understory species that are characteristic of moist to wet, nutrient-rich sites (e.g., *Alnus incana*, *Ribes* spp., *Cornus stolonifera* and *Rubus pubescens*). M300 soils can be either organic or mineral, but peat is usually well-decomposed and typically not as deep as in M299 stands. Although overall floristics are very similar to those of M299, open boreal bogs and poor fens (i.e., >10% cover by trees >5 m height) are described separately by M876 [North American Boreal & Sub-boreal Acidic Bog & Fen].

Within M299, CNVC Groups break out four geographic subsets: CG0016 [Atlantic Boreal Black Spruce - Balsam Fir Poor - Intermediate Treed Wetland] describes these conditions in Atlantic Canada, CG0019 [Ontario-Quebec Boreal Black Spruce Poor - Intermediate Treed Wetland] in Ontario and west-central Quebec, CG0022 [West-Central Boreal Black Spruce - Tamarack Poor - Intermediate Treed Wetland] in western Canada and CG0025 [Subarctic Black Spruce Poor - Intermediate Treed Wetland] in the northern part of the range. No Group has been defined for the Alaskan-Yukon boreal region yet. Within each of these Groups, CNVC Alliances aggregate Associations that share floristics and dominance characteristics within similar nutrient regimes.

Alnus incana (grey alder) here refers to ssp. *rugosa* (speckled alder) in eastern Canada and ssp. *tenuifolia* (mountain alder) in western Canada.

Viburnum nudum here refers to *V. nudum* var. *cassinoides* (wild raisin).



Canadian National Vegetation Classification (CNVC) Classification nationale de la végétation du Canada

<http://cnvc-cnvc.ca>

North American Boreal Conifer Poor Swamp Marécages pauvres à conifères boréaux de l'Amérique du Nord

Macrogroup M299

Source Information

Number of Source Plots for M299: 2742 (Canadian National Vegetation Classification. 2015. CNVC Master Database [VPro13/MSAccess 2010 format]. Natural Resources Canada, Sault Ste. Marie, ON.)

Information Sources (data):

Alberta Environment and Parks. 2014. Ecological Site Information System (ESIS). Govt. AB, Edmonton, AB. (250 plots)

Basquill, S.P. (compiler). 2015. Maritime provinces of Canada regional forest ecosystem plot database. Standardized forest ecosystem plot data compilation and classification from N.B. Dept. Nat. Resour.; P.E.I. For., Fish, & Wildlife Div., Dept. Commun., Land, & Environ.; N.S. Dept. Nat. Resour.; N.S. Environ.; Parks Can.; the Atlantic Can. Conserv. Data Centre; and other sources. Atlantic Can. Conserv. Data Centre, Sackville, NB. (10 plots)

Biogeoclimatic Ecosystem Classification Program of British Columbia. 2011. BECMaster ecosystem plot database [VPro13/MSAccess 2010 format]. W.H. MacKenzie, (ed.) B.C. Min. For., Lands, and Nat. Res. Ops., Smithers, BC. Available:

www.for.gov.bc.ca/hre/becweb/resources/informationrequests (accessed: June 2015). (128 plots)

McLaughlan, M.S.; Wright, R.A.; Jiricka, R.D. 2010. Saskatchewan forest ecosystem classification [data set]. Sask. Min. Environ. For. Serv., Prince Albert, SK. (151 plots)

McMurray, S.C., Johnson, J.A., Zhou, K., Uhlig, P.W.C. 2015. Ontario ecological land classification program - Ecological Data Repository (EDR). Ont. Min. Nat. Resour. & For., Sci. & Info. Branch, Sault Ste. Marie, ON. (428 plots)

Ministère des Ressources naturelles, de la Faune et des Parcs, Forêt Québec. 2003. Base de données des points d'observation écologique (version 2003). Gouv. du Qué., Min. des Res. nat., de la Faune et des Parcs, Forêt Qué., Dir. des inv. for., QC. (1720 plots)

Natural Resources Canada, Canadian Forest Service, Atlantic Region. 2006. Forest vegetation plot descriptions from the following publications; Damman, A.W.H. (1963, 1964, 1967); Meades, W.J. (1976, 1986). Nat. Res. Canada, Corner Brook, NL. (55 plots)

Concept Authors: L. Allen, K. Baldwin, S. Basquill, K. Chapman, W. MacKenzie, M. Major, B. Meades, D. Meidinger, C. Morneau, P. Uhlig, USNVC

Description Authors: K. Baldwin, D. Downing

Date of Concept: July, 2015

Date of Description: November, 2016

References

Archibald, J.H.; Klappstein, G.D.; Corns, I.G.W. 1996. Field guide to ecosites of southwestern Alberta. Nat. Resour. Can., Can. For. Ser., North. For. Cent., Edmonton, AB. Spec. Rep. 8.

Arnup, R.W.; LeBlanc, P.A.; Becker, G. 2006. Field guide to ecosites of the Mid-Boreal Upland ecoregion of Manitoba. Louisiana-Pacific Canada Ltd, For. Resour. Div. and Man. Conserv., For. Branch, Swan River and Winnipeg, MB.

Basquill, S.; Beaudette, D.; Cameron, R.; Curley, R.; Fenton, N.; Glen, W.; Gordon, S.; Hutchinson, J.; Kelly, G.; Loo, J.; Lynds, A.; MacAskill, D.; MacKinnon, D.; MacQuarrie, K.; Makepeace, S.; Matson, B.; Neily, P.; Quigley, E.; Zelazny, V. 2009 (updated 2015). Forest communities of the Maritime provinces of Canada. Atlantic Canada Conservation Data Centre, Sackville, NB.

Beckingham, J.D.; Archibald, J.H. 1996. Field guide to ecosites of northern Alberta. Nat. Resour. Can., Can. For. Serv., North. For. Cent., Edmonton, AB. Spec. Rep. 5.

Beckingham, J.D.; Corns, I.G.W.; Archibald, J.H. 1996. Field guide to ecosites of west-central Alberta. Nat. Resour. Can., Can. For. Serv., North. For. Cent., Edmonton, AB. Spec. Rep. 9.

Bostock, H.S. 1970. Physiographic subdivisions of Canada. Geol. Surv. Can. Econ. Geol. Rep. No. 1. Pages 10-30 in: R.J.W. Douglas (ed.) Geology and economic minerals of Canada. Geol. Surv. Can., Ottawa, ON.

Boulanger, Y.; Gauthier, S.; Burton, P.J. 2014. A refinement of models projecting future Canadian fire regimes using homogeneous fire regime zones. Can. J. For. Res. 44(4):365-376.

Brandt, J.P. 2009. The extent of the North American boreal zone. Environ. Rev. 17:101-161.



Canadian National Vegetation Classification (CNVC) Classification nationale de la végétation du Canada

<http://cnvc-cnvc.ca>

North American Boreal Conifer Poor Swamp

Macrogroup M299

Marécages pauvres à conifères boréaux de l'Amérique du Nord

References (cont'd)

- Brouillet, L.; Coursol, F.; Meades, S.J.; Favreau, M.; Anions, M.; Bélisle, P.; Desmet, P. 2010+. VASCAN, the database of vascular plants of Canada. Available: <http://data.canadensys.net/vascan/search> (accessed: September 2015).
- Crins, W.J.; Gray, P.A.; Uhlig, P.W.C.; Wester, M. 2009. The ecosystems of Ontario, Part 1: Ecozones and ecoregions. Ont. Min. Nat. Resour., Inven. Monit. Assess., Peterborough, ON. SIB TER IMA TR-01.
- Crum, H.A.; Planisek, S. 1988. A focus on peatlands and peat mosses. Univ. of Michigan Press, MI, US.
- Damman, A.W.H. 1963. A reconnaissance survey of the ecological conditions in the forests of the Roddickton area, Newfoundland. For. Res. Branch, Can. Dept. For., NL. Mimeo 63-N-1.
- Damman, A.W.H. 1964. Some forest types of central Newfoundland and their relation to environmental factors. The Society of American Foresters, US. Monograph 8.
- Damman, A.W.H. 1967. The forest vegetation of western Newfoundland and site degradation associated with vegetation change. PhD thesis, Univ. of Michigan, Ann Arbor, MI, US.
- Damman, A.W.H. 1983. An ecological subdivision of the island of Newfoundland. Pages 163-206 in: G.R. South (ed.) Biogeography and ecology of the island of Newfoundland. Dr W. Junk Publishers, The Hague, The Netherlands.
- DeLong, C.; Banner, A.; MacKenzie, W.H.; Rogers, B.J.; Kaytor, B. 2011. A field guide to ecosystem identification for the Boreal White and Black Spruce zone of British Columbia. B.C. Min. For. Range, For. Sci. Prog., Victoria, BC. Land Manage. Handb. No. 65.
- Ducks Unlimited. 2014. Field guide: boreal wetland classes in the Boreal Plains ecozone of Canada. Ducks Unlimited Canada, Canada.
- Ecological and Landscape Classification Program. 2015. Bioclimate zones of Yukon - Version 6 (November 2015). Report and map at 1:250,000 scale. Department of Environment, Whitehorse, YT.
- Ecological Stratification Working Group. 1995. A national ecological framework for Canada. Agric. and Agri-Food Can., Res. Branch, Centre Land and Biol. Resour. Res., and Environ. Can., State of Environ. Direct., Ecozone Analysis Branch, Ottawa/Hull, ON/QC.
- Ecoregions Working Group. 1989. Eoclimatic regions of Canada. W. Strong and S.C. Zoltai (compilers). Sustain. Dev. Branch, Can. Wildlife Serv., Conserv. and Prot., Environ. Can., Ottawa, ON. ELC Series No. 23.
- Ecosystem Classification Group. 2007 (revised 2009). Ecological regions of the Northwest Territories - Taiga Plains. Govt. N.W.T., Dept. Environ. & Nat. Resour., Yellowknife, NT.
- Ecosystem Classification Group. 2008. Ecological regions of the Northwest Territories – Taiga Shield. Gov. N.T., Dept. Env. & Nat. Resour., Yellowknife, NT.
- Ecosystem Classification Group. 2010. Ecological regions of the Northwest Territories – Cordillera. Gov. N.T., Dept. Env. & Nat. Resour., Yellowknife, NT.
- Environment Canada. 2015. Canadian climate normals, 1961-1990. Gov. Canada, Available: http://climate.weather.gc.ca/climate_normals/index_e.html (accessed: January 29, 2015).
- Flora of North America Editorial Committee. 2007+. Flora of North America north of Mexico, vols 27, 28, 29. Oxford University Press, New York and Oxford. Available: <http://www.mobot.org/plantscience/bfna/BFNAMenu.htm> (accessed: November, 2015).
- Fryer, J.L. 2014. *Picea mariana*. In: Fire Effects Information System. U.S. Dept. Agric., For. Serv., Rocky Mt. Res. Stn., Fire Sci. Lab., Missoula, MT, US. Available: <http://www.fs.fed.us/database/feis/plants/tree/picmar/all.html> (accessed: May 26, 2015).
- Greene, D.F.; Zasada, J.C.; Sirois, L.; Kneeshaw, D.; Morin, H.; Charron, I.; Simard, M.J. 1999. A review of the regeneration dynamics of North American boreal forest tree species. Can. J. For. Res. 29:824-839.



Canadian National Vegetation Classification (CNVC) Classification nationale de la végétation du Canada

<http://cnvc-cnvc.ca>

North American Boreal Conifer Poor Swamp

Macrogroup M299

Marécages pauvres à conifères boréaux de l'Amérique du Nord

References (cont'd)

Grondin, P.; Blouin, J.; Racine, P. 1998. Rapport de classification écologique du sous-domaine bioclimatique de la sapinière à bouleau blanc de l'ouest. Min. des Res. nat. du Qué., Dir. des inv. for., QC.

Grondin, P.; Blouin, J.; Racine, P.; D'Avignon, H.; Tremblay, S. 2000. Rapport de classification écologique du sous-domaine bioclimatique de la sapinière à bouleau blanc de l'est. Forêt Qué., Dir. des inv. for., Min. des Res. nat. du Qué., QC.

Hare, F.K. 1950. Climate and zonal divisions of the boreal forest formation in eastern Canada. *Geogr. Rev.* 40:615-635.

Hare, F.K.; Hay, J.E. 1974. The climate of Canada and Alaska. Vol. 11, pages 49-192 in: R.A. Bryson and F.K. Hare (eds.) *World survey of climatology*. Elsevier Scientific Publishing Company, Amsterdam, The Netherlands.

Harris, A.G., McMurray, S.C., Uhlig, P.W.C., Jeglum, J.K., Foster, R.F., Racey, G.D. 1996. Field guide to the wetland ecosystem classification for northwestern Ontario. *Ont. Min. Nat. Resour., Northwest Sci. Tech., Thunder Bay, ON. NWST Field Guide FG-01*.

Li, T.; Hélie, R. (compilers). 2014. Ecozones of Canada / Écozones du Canada [map]. Canadian Council on Ecological Areas / Conseil Canadien des Aires Écologiques, CA. Scale 1:25,000,000.

MacKenzie, W.H.; Moran, J.R. 2004. Wetlands of British Columbia: a guide to identification. B.C. Min. For., Res. Branch, Victoria, BC. *Land Manage. Handb. No. 52*.

McLaughlan, M.S.; Wright, R.A.; Jiricka, R.D. 2010. Field guide to the ecosites of Saskatchewan's provincial forests. *Sask. Min. Environ., For. Serv., Prince Albert, SK*.

Meades, W.J. 1976. Vegetation of Terra Nova National Park. In: *Biophysical classification of Terra Nova National Park: environmental component 3c*. Can. Dept. Indian & North. Aff., Parks Branch, CA.

Meades, W.J. 1983. Heathlands. Pages 267-318 in: G.R. South (ed.) *Biogeography and ecology of the island of Newfoundland*. Dr W. Junk Publishers, The Hague, The Netherlands.

Meades, W.J. 1986. Successional status of ericaceous dwarf-shrub heath in eastern Newfoundland. PhD thesis, Univ. of Connecticut, Storrs, CT.

Meades, W.J.; Moores, L. 1994. Forest site classification manual: a field guide to the Damman forest types of Newfoundland. 2nd ed. Corner Brook, Western Newfoundland Model Forest, Inc., NL. FRDA Rep. 003.

Meidinger, D.; Pojar, J. (eds.). 1991. *Ecosystems of British Columbia*. B.C. Min. For., Res. Branch, BC. Spec. Rep. Series No. 6.

Meidinger, D.; Faber-Langendoen, D.; Baldwin, K. 2016. Macrogroup Detail Report: M299 North American Boreal Conifer Poor Swamp Macrogroup [15 Jan 2016]. United States National Vegetation Classification. Fed. Geogr. Data Comm., Washington DC, US.

Ministère des Ressources naturelles du Québec, Forêt Québec. 2002+. Les guides de reconnaissance des types écologiques. Gouv. du Québec, Québec, QC. Available: <http://www.mffp.gouv.qc.ca/forets/inventaire/guide-types-ecologiques-carte.jsp> (accessed: May 2015).

Morneau, C. In prep. Rapport de classification écologique du sous-domaine bioclimatique de la pessière à mousses de l'est. Min. des forêts, de la Faune et des Parcs, Dir. des inv. for., QC.

National Wetlands Working Group. 1988. *Wetlands of Canada*. Sustain. Dev. Branch, Environ. Can., Ottawa, ON and Polyscience Publications Inc., Montreal, QC. ELC Series No. 24.

National Wetlands Working Group. 1997. *The Canadian wetland classification system*. B.G. Warner, and C.D.A. Rubec, (eds.) Wetlands Res. Centre, Univ. of Waterloo, Waterloo, ON.

Natural Regions Committee. 2006. Natural regions and subregions of Alberta. D.J. Downing and W.W. Pettapiece (compilers). Gov. AB, Min. Environ., AB. Pub. No. T/852.



Canadian National Vegetation Classification (CNVC) Classification nationale de la végétation du Canada

<http://cnvc-cnvc.ca>

North American Boreal Conifer Poor Swamp

Macrogroup M299

Marécages pauvres à conifères boréaux de l'Amérique du Nord

References (cont'd)

- Neily, P.; Basquill, S.; Quigley, E.; Stewart, B.; Keys, K. 2011. Forest ecosystem classification for Nova Scotia, Part I: Vegetation types. N.S. Dept. Nat. Resour., Renew. Resour. Branch, NS.
- Neily, P.; Quigley, E.; Basquill, S.; Keys, K. 2015. Ecological land classification for Nova Scotia. N.S. Dept. Nat. Resour., Renew. Resour. Branch, NS. Rep. FOR 2016-*
- Ontario Ministry of Natural Resources. 2009. Ecological land classification ecosites field manual – operational draft, April 20th, 2009 – boreal. Ecol. Land Class. Working Grp, Ont. Min. Nat. Resour., Sci. & Info Branch, Inven. Monit. Assess. Sect., Sault Ste. Marie, ON.
- Riley, J.L.; Michaud, L. 1989. Peat and peatland resources of northwestern Ontario. Min. North. Dev. & Mines, Ont. Geol. Surv., ON. Misc. Paper 144.
- Riley, J.L. 2011. Wetlands of the Ontario Hudson Bay Lowland: an Ontario overview. Nature Conservancy of Canada. Toronto, ON.
- Rydin, H.; Jeglum, J.K. 2006. The biology of peatlands. Oxford Univ. Press, Oxford, UK.
- Sanchez-Mata, D.; Rivas-Martinez, S. 2010. Bioclimatic dossier for the 'Circumboreal Vegetation Mapping Project' (CBVM). Pages 42-52 in: S.S. Talbot (ed.) Proc. 7th Intl. Conserv. Arctic Flora and Fauna (CAFF) Flora Gp. Workshop. January 28-February 3, 2011. Akureyri, Iceland. CAFF International Secretariat, CAFF Flora Expert Group (CFG), CAFF Proc. Series Rep. No. 8.
- Saucier, J.-P.; Baldwin, K.; Krestov, P.; Jorgenson, T. 2015. Boreal forests. Chapter 2 in: K.S.-H. Peh, R.T. Corlett and Y. Bergeron (eds.) Routledge handbook of forest ecology. Routledge, Abingdon, UK.
- Soil Classification Working Group. 1998. The Canadian system of soil classification. NRC Research Press, Ottawa, ON. Agric. and Agri-Food Can. Pub. 1646.
- Stockdale, C. 2014. Fire regimes of western boreal Canada and the foothills of Alberta. A discussion document and literature review for the LANDWEB Project.
- Uchtyl, R.J. 1991. *Abies balsamea*. In: Fire Effects Information System. U.S. Dept. Agric. For. Serv. Rocky Mt. Res. Stn., Fire Sci. Lab., Missoula, MT, US. Available: <http://www.fs.fed.us/database/feis/plants/tree/abibal/all.html> (accessed: May 26, 2015).
- Uchtyl, R.J. 1991. *Larix laricina*. In: Fire Effects Information System. U.S. Dept. Agric. For. Serv. Rocky Mt. Res. Stn., Fire Sci. Lab., Missoula, MT, US. Available: <http://www.fs.fed.us/database/feis/plants/tree/larlar/all.html> (accessed: October 06, 2008).
- Uhlig, P.W.C., Chapman, K., Baldwin, K., Wester, M., Yanni, S. 2016. Draft boreal treed vegetation type factsheets. Ecol. Land Class. Prog., Ont. Min. Nat. Resour. & For., Sci. & Info Branch, Sault Ste. Marie, ON.
- USNVC [United States National Vegetation Classification]. 2016. United States National Vegetation Classification Database ver. 2.0. Fed. Geogr. Data Comm., Veg. Subcomm., Washington DC, US. Available: <http://usnvc.org> (accessed: March 10, 2016).
- Vitt, D.H.; Halsey, L.A. 1994. The bog landforms of continental western Canada in relation to climate and permafrost patterns. *Arct. Alp. Res.* 26:1-13.
- Wells, E.D. 1996. Classification of peatland vegetation in Atlantic Canada. *J. Veg. Sci.* 7:847-878.
- Zoladeski, C.A.; Wickware, G.M.; Delorme, R.J.; Sims, R.A.; Corns, I.G.W. 1995. Forest ecosystem classification for Manitoba: field guide. Nat. Res. Can., Can. For. Serv., North. For. Centre, Edmonton, AB. Special Rep. 2.

The information contained in this factsheet is based on data and expert knowledge that is current to the date of description. As new information becomes available, the factsheet will be updated.

Suggested Citation: Baldwin, K.; Downing, D. North American Boreal Conifer Poor Swamp [online]. Sault Ste. Marie, Ontario, Canada: Canadian National Vegetation Classification. November, 2016; generated 30-November-2016; cited **ENTER DATE ACCESSED**. 10 p. Canadian National Vegetation Classification Macrogroup: M299. Available from <http://cnvc-cnvc.ca>. System Requirements: Adobe Acrobat Reader v. 7.0 or higher. ISSN 1916-3266.



Canadian National Vegetation Classification (CNVC) Classification nationale de la végétation du Canada

<http://cnvc-cnvc.ca>

North American Boreal Conifer Poor Swamp

Macrogroup M299

Marécages pauvres à conifères boréaux de l'Amérique du Nord

Comparison of Vegetation Characteristics for Boreal Wetland Forest and Woodland Macrogroups

Layer	Spp	n=2742	n=238	Common Name
		M299 Poor Swamp	M300 Flooded and Rich Swamp	
Tree	<i>Larix laricina</i>	***		tamarack
	<i>Picea mariana</i>	■■■■■	■■■■■	black spruce
	<i>Abies balsamea</i>	■■■	■■■■■	balsam fir
	<i>Betula papyrifera</i>	*	■■■■■	paper birch
	<i>Picea glauca</i>		■■■	white spruce
	<i>Populus balsamifera</i>		***	balsam poplar
	<i>Fraxinus nigra</i>		*	black ash
	<i>Thuja occidentalis</i>		*	eastern white cedar
Shrub	<i>Betula spp. (shrub)</i>	***		shrub birches
	<i>Kalmia polifolia</i>	■■		pale bog laurel
	<i>Chamaedaphne calyculata</i>	■■■■		leatherleaf
	<i>Salix spp. (shrub)</i>	■■■	***	shrub willows
	<i>Vaccinium myrtilloides</i>	■■■	***	velvet-leaved blueberry
	<i>Kalmia angustifolia</i>	■■■■	***	sheep laurel
	<i>Rhododendron groenlandicum</i>	■■■■■	■■■	common Labrador tea
	<i>Alnus incana</i>	■■■■	■■■■■	grey alder
	<i>Ribes glandulosum</i>		■■■	skunk currant
	<i>Viburnum edule</i>		■■■	squashberry
	<i>Cornus stolonifera</i>		****	red-osier dogwood
	<i>Acer spicatum</i>		***	mountain maple
Herb/ Dwarf Shrub	<i>Vaccinium vitis-idaea</i>	***		lingonberry
	<i>Vaccinium oxycoccos</i>	■■		small cranberry
	<i>Rubus chamaemorus</i>	■■■		cloudberry
	<i>Maianthemum trifolium</i>	■■■	***	three-leaved false Solomon's seal
	<i>Gaultheria hispidula</i>	■■■	■■■	creeping snowberry
	<i>Carex spp.</i>	■■■	■■■■■	sedges
	<i>Rubus pubescens</i>		■■■	dwarf raspberry
	<i>Maianthemum canadense</i>		■■■	wild lily-of-the-valley
	<i>Aralia nudicaulis</i>		■■	wild sarsaparilla
	<i>Gymnocarpium dryopteris</i>		**	common oak fern
Moss/Lichen	<i>Aulacomnium palustre</i>	***		ribbed bog moss
	<i>Cladina spp.</i>	■■■	■■	reindeer lichens
	<i>Sphagnum spp.</i>	■■■■■	■■■■■	peat mosses
	<i>Pleurozium schreberi</i>	■■■■	■■■	red-stemmed feathermoss
	<i>Hylocomium splendens</i>	■■■	■■■	stairstep moss

Legend

Constancy:	Black bar >= 50%	Cover:	5 bars >= 25%	2 bars >= 1%
	Grey bar >= 30%		4 bars >= 10%	1 bar <= 1%
	Asterisk >= 20%		3 bars >= 3%	



Canadian National Vegetation Classification (CNVC) Classification nationale de la végétation du Canada

<http://cnvc-cnvc.ca>

North American Boreal Conifer Poor Swamp

Macrogroup M299

Marécages pauvres à conifères boréaux de l'Amérique du Nord

Comparison of Vegetation Characteristics for Regional Groups in M299

Layer	Spp	n=574	n=2102	n=56	Common Name
		CG0022 West-Central	CG0019 Ontario-Quebec	CG0016 Atlantic	
Tree	<i>Larix laricina</i>	■■■■	****		tamarack
	<i>Picea mariana</i>	■■■■■	■■■■■	■■■■■	black spruce
	<i>Abies balsamea</i>		■■■	■■■■■	balsam fir
	<i>Betula papyrifera</i>		***	***	paper birch
Shrub	<i>Betula</i> spp. (shrub)	■■■■			shrub birches
	<i>Salix</i> spp. (shrub)	■■■	■■■		shrub willows
	<i>Chamaedaphne calyculata</i>	***	■■■■		leatherleaf
	<i>Rhododendron groenlandicum</i>	■■■■■	■■■■■	■■■	common Labrador tea
	<i>Alnus incana</i>		■■■■		grey alder
	<i>Kalmia polifolia</i>		■■		pale bog laurel
	<i>Vaccinium angustifolium</i>		■■■	■■■	early lowbush blueberry
	<i>Kalmia angustifolia</i>		■■■■	■■■■	sheep laurel
	<i>Ilex mucronata</i>			■■■■	mountain holly
	<i>Viburnum nudum</i>			■■■	wild raisin
	<i>Rhododendron canadense</i>			■■■■	rhodora
Herb/ Dwarf Shrub	<i>Calamagrostis canadensis</i>	***			bluejoint reedgrass
	<i>Petasites frigidus</i>	**			arctic sweet coltsfoot
	<i>Mitella nuda</i>	■■			naked mitrewort
	<i>Vaccinium vitis-idaea</i>	■■■			lingonberry
	<i>Vaccinium oxycoccos</i>	■■	■■		small cranberry
	<i>Maianthemum trifolium</i>	■■■	■■■	***	three-leaved false Solomon's seal
	<i>Rubus chamaemorus</i>	■■■	■■■	■■■	cloudberry
	<i>Carex</i> spp.	■■■■	■■■	■■■■	sedges
	<i>Clintonia borealis</i>		■■	■■■■	yellow clintonia
	<i>Maianthemum canadense</i>		**	■■	wild lily-of-the-valley
	<i>Osmundastrum cinnamomeum</i>			■■■■	cinnamon fern
	<i>Solidago macrophylla</i>			***	large-leaved goldenrod
	<i>Epigaea repens</i>			**	trailing arbutus
Moss/Lichen	<i>Tomentypnum nitens</i>	■■■■			golden fuzzy fen moss
	<i>Aulacomnium palustre</i>	■■■			ribbed bog moss
	<i>Cladina</i> spp.	■■■■	■■■	****	reindeer lichens
	<i>Sphagnum</i> spp.	■■■■■	■■■■■	■■■■■	peat mosses
	<i>Pleurozium schreberi</i>	■■■■	■■■■	■■■■	red-stemmed feathermoss
	<i>Bazzania trilobata</i>			■■■	three-lobed whipwort

Legend

Constancy:	Black bar >= 50%	Cover:	5 bars >= 25%	2 bars >= 1%
	Grey bar >= 30%		4 bars >= 10%	1 bar <= 1%
	Asterisk >= 20%		3 bars >= 3%	