



Agriculture
Canada

Research
Branch

Direction générale
de la recherche

Technical Bulletin 1986-4E

The T.M.E. system of feed evaluation: methodology, feed composition data and bibliography



The map on the cover has dots representing Agriculture Canada research establishments.

ONE HUNDRED YEARS OF PROGRESS

The year 1986 is the centennial of the Research Branch, Agriculture Canada.

On 2 June 1886, *The Experimental Farm Station Act* received Royal Assent. The passage of this legislation marked the creation of the first five experimental farms located at Nappan, Nova Scotia; Ottawa, Ontario; Brandon, Manitoba; Indian Head, Saskatchewan (then called the North-West Territories); and Agassiz, British Columbia. From this beginning has grown the current system of over forty research establishments that stretch from St. John's West, Newfoundland, to Saanichton, British Columbia.

The original experimental farms were established to serve the farming community and assist the Canadian agricultural industry during its early development. Today, the Research Branch continues to search for new technology that will ensure the development and maintenance of a competitive agri-food industry.

Research programs focus on soil management, crop and animal productivity, protection and resource utilization, biotechnology, and food processing and quality.

The T.M.E. system of feed evaluation: methodology, feed composition data and bibliography

I. R. SIBBALD
Animal Research Centre
Ottawa, Ontario

Animal Research Centre Contribution 85-19

Research Branch
Agriculture Canada
1986

Copies of this publication are available from:

Director

Animal Research Centre

Research Branch, Agriculture Canada

Ottawa, Ontario

K1A 0C6

Produced by Research Program Service

© Minister of Supply and Services Canada 1986

Cat. No. A54-8/1986-4E

ISBN 0-662-14628-X

SUMMARY

The true metabolizable energy (TME) system of feed evaluation comprises assays for bioavailable energy, amino acids, lipids and minerals. In this Bulletin the principles of the system are explained in the 'Introduction' which includes a diagram showing the partition of energy within the bird. A section entitled 'Bioassay Procedures' describes the basic methodology while sub-sections contain: a list of precautions to be observed when using the assays; a detailed description of the precision-feeding technique; details of the design and construction of cages for adult cockerels; and a discussion of excreta collection techniques. The section on 'Feedingstuff Composition' contains data describing the nitrogen, ether extract, crude fibre, neutral detergent fibre, ash, calcium, phosphorus, gross energy, TME, TME corrected to zero nitrogen balance (TME_n), total amino acids, and true available amino acids (TAAA) of an array of feed ingredients; the data are expressed per unit of dry matter. An important component of the Bulletin is the 'Bibliography' which lists references to more than 500 publications concerned, directly or indirectly, with the derivation and application of TME-type data.

RÉSUMÉ

La méthode de l'énergie métabolisable vraie (EMV) pour évaluer les aliments des animaux comporte des dosages biologiques de l'énergie, des acides aminés, des lipides et des minéraux biodisponibles. Dans ce bulletin, on explique d'abord les principes du système (Introduction) à l'aide d'un diagramme montrant la répartition de l'énergie dans le corps de la volaille. Une section intitulée "Méthodes de dosage biologique" décrit les techniques de base et comprend des sous-sections couvrant diverses questions comme: les précautions à prendre dans la conduite des dosages, une description détaillée de la technique d'alimentation de précision, les détails de la conception et de la construction de cages pour les coqs adultes et une analyse de la technique de collecte des excréments. La partie sur la "Composition des aliments" contient des données sur les teneurs en azote, extrait éthéré,


cellulose brute, fibre au détergent neutre, cendres, calcium, phosphore, énergie brute, EMV, EMV ramené au bilan azoté zéro (EMV_n), acides aminés totaux et acides aminés disponibles vrais (AADV), de tout un assortiment de matières alimentaires. Les données sont calculées sur la matière sèche. Complément important du bulletin, la "Bibliographie" contient plus de 500 titres de publications touchant de près ou de loin à l'obtention et à l'utilisation des données basées sur l'énergie métabolisable vraie.

PREFACE

In 1981 the Animal Research Centre published Technical Bulletin No. 3 entitled: Bioassays Based on Precision Feeding of Poultry. The French version was entitled: Dosage biologique en cage métabolique chez la volaille. The Bulletin contained a brief description of the precision feeding technique and of the assays based thereon. The primary purpose of the Bulletin was to be a bibliography of relevant literature and 162 references were listed.

An enlarged and updated version entitled: The T.M.E. System of Feed Evaluation was published in 1983 as Contribution 1983-20E of the Research Branch of Agriculture Canada. The French edition was Contribution 1983-20F and was entitled: La méthode d'évaluation des aliments par le calcul de l'énergie métabolisable vraie (E.M.V.). Features of the 1983 version were an enlarged section on assay methodology, inclusion of a section describing feedingstuff composition, and extension of the bibliography to 314 references. Demand for the revised Bulletin exceeded all expectations and approximately 1200 copies were distributed in less than two years.

The present Bulletin is an updated version of the 1983 edition. The introduction and methods sections have been revised in the light of recent research. The feedingstuff composition section is greatly expanded by data describing additional samples and by the inclusion of total and true available amino acid values. More than two hundred references are added to the bibliography.



Digitized by the Internet Archive
in 2013

<http://archive.org/details/tmesystemoffeede19864sibb>

TABLE OF CONTENTS

	<u>Page</u>
Summary/Résumé	1
Preface	3
Table of Contents	5
Acknowledgements	6
Introduction	7
Bioassay Procedures	12
General	12
Precautions	16
Precision Feeding	17
Cage Design and Construction	22
Excreta Collection	28
Feedingstuff Composition	34
Table 1. Nitrogen, fat, fibre, mineral (g/kg) and energy (MJ/kg) concentrations in the dry matter of several feedingstuffs	37
Table 2. Total amino acids in the dry matter of several feedingstuffs (g/kg)	52
Table 3. True available amino acids in the dry matter of several feedingstuffs (g/kg)	59
Table 4. Bioavailability of amino acids in the dry matter of several feedingstuffs (%)	66
Bibliography	73
1917	73
1924	73
1926	73
1966	73
1970	73
1975	73
1976	74
1977	74
1978	76
1979	78
1980	80
1981	86
1982	92
1983	99
1984	105
1985	112

ACKNOWLEDGEMENTS

The author wishes to thank all those who helped with the preparation of this Bulletin. Several people provided references to TME-related publications while others provided samples of feedingstuffs for analysis; their cooperation is appreciated. The preliminary drawings for the figures illustrating cage design and construction were prepared by F. Nachbaur; final drawings and all photographs were prepared by Research Program Service, Agriculture Canada. The feed composition tables could not have been compiled without the technical help of S. Winter, R. Cochrane and L. Hope; the proximate and amino acid analyses were made, in part, by the Chemistry and Biology Research Institute, Agriculture Canada. The staff of the poultry unit of the Animal Research Centre, Research Farm provided valuable help by caring for birds and participating in bioassay work. Finally, my thanks to D. Leger for editorial assistance throughout the preparation of the Bulletin.

DISCLAIMER

Mention of a trade name, proprietary product, or specific equipment does not imply its approval to the exclusion of other products which may be suitable.

INTRODUCTION

The true metabolizable energy (TME) system of feed evaluation, which is gaining wide acceptance (Figure 1), comprises bioassays for TME, TME corrected to zero nitrogen balance (TME_n), true available amino acids (TAAA), true available lipids (TAL) and true available minerals (TAM). Each assay involves corrections for metabolic plus endogenous losses. The desirability of making such corrections was recognized by Armsby (1)* who wrote: "since the digestion experiment as ordinarily conducted ignores the presence in the faeces of excretory products, the results obtained by its use will necessarily be too low." Goodwin's translation of Kellner (3) states: "the assumption is made that dung consists only of undigested food. The assumption, however, is not quite correct, for the faeces are always mixed with small quantities of substances which come from the digestive organs and which are termed the products of metabolism." Despite this early recognition of the problems associated with metabolic and endogenous losses, little attention was given to the introduction of appropriate corrections. The assay for the 'biological value' of protein included corrections for metabolic fecal and endogenous urinary nitrogen losses (2) and nitrogen-free diets have been used in some amino acid availability studies. However, the need for corrections has generally been ignored and the teachings of Kellner and Armsby seem to have been forgotten.

By making corrections for metabolic and endogenous losses, the assays of the TME system should provide data which are independent of the species, strain, sex, age and physiological state of the assay bird. Ideally, the data should be independent of species in general but this state is unattainable because of the wide variation in microbial activity within the alimentary tracts of diverse species such as the cow, pig and chicken. Such microbial activity has profound effects on the composition of ingesta offered for digestion and absorption by the host animal.

* Numbers in brackets refer to publications listed in the bibliography which starts on page 73.




Figure 1. Adoption of the TME system.

COUNTRIES WHERE THERE ARE KNOWN USERS (BASED ON PUBLICATIONS
IN THE SCIENTIFIC PRESS).

OTHER PROBABLE USERS.

The term TME was defined by Harris (4). The effect of metabolic fecal and endogenous urinary energy losses on apparent metabolizable energy (AME) values was described, with theoretical data, by Guillaume and Summers (5). The relationships between AME and TME are described in detail in two recent publications (526,556). Corrections for metabolic and endogenous losses are equally important and have similar effects in TAAA, TAL and TAM assays.

It is now common practice to correct TME values to zero nitrogen balance. The reasons are explained in detail by Wolynetz and Sibbald (526) who conclude that the correction causes estimates of bioavailable energy "to be based on the assumption that the available energy is being used for a single common purpose, the production of heat, and that the composition of the bird is unchanged." There are no comparable corrections in the TAAA, TAL or TAM bioassays. However, the TAM assay differs from other assays of the TME system because an intake of a mineral in excess of the requirement and storage capacity is likely to be excreted. Therefore in the TAM assay the mineral intake must be kept below the requirement. Intakes in excess of the requirement have no effect on the TAAA bioassay because the surpluses are voided as uric acid not amino acid. Similarly, in the ⁿTME and TAL bioassays excesses are stored as fat or lost as heat and do not affect the results.

The assays usually involve precision-feeding a fasted bird with a known quantity of the test feedingstuff and collecting the resulting excreta. Each feedingstuff is fed at two, or more, levels of input to establish the relationship between nutrient input and output. For simplicity and convenience one input level is usually zero. Voluntary feeding has been used successfully as a substitute for precision feeding but it is subject to errors associated with feed spillage and consequent inaccurate estimates of intake. Furthermore, since birds eat at different rates the estimates of metabolic and endogenous losses may be subject to greater error.

Fasted birds tend to catabolize more body protein than fed birds thus increasing metabolic energy excretion and introducing a bias into the TME assay. Similarly, the loss of body protein as energy-containing excretory

products is affected by the amount and composition of the test feedingstuff, a secondary cause of bias. The problem is controlled by correcting the excreta energy outputs of both fed and fasted birds to zero nitrogen balance. The method for making the correction is explained later.

A schematic representation of the partition of ingested feed energy by the bird is presented as Figure 2. The terminology and abbreviations are similar to those published by the National Research Council of the United States (256). The figure differs in certain respects from that published in the second edition of this Bulletin. The feces are assumed to comprise three energy containing fractions: the fecal energy of feed origin ($F_i E$); the endogenous fecal energy ($F_e E$); and the metabolic fecal energy ($F_m E$). The $F_e E$ is contained in microflora and microbial debris and, for most practical purposes, may be considered part of the $F_i E$. The urine also comprises three energy containing fractions: the urinary energy of feed origin ($U_i E$); the metabolic urinary energy ($U_m E$); and the endogenous urinary energy ($U_e E$). The $U_m E$ is contained in products of tissue catabolism, it being assumed that anything absorbed across the gut wall immediately becomes part of a tissue. The $U_e E$ is contained in the excretory products resulting from protein turnover.

The partition shown in Figure 2 differs from that of the National Research Council (256) inasmuch as the $F_m E$ and $U_e E$ are included as part of the net energy for maintenance. Further, the $U_m E$ is placed under the net energy for production because it results from tissue catabolism which may occur at the same time as tissue synthesis as well as in the absence thereof. The figure helps to explain the differences between AME and TME and shows that feces and urine contain energy components which are derived from the body rather than the feed.




Figure 2. The partition of ingested feed energy in the bird (the forerunner of this scheme is described in reference 336).

BIOASSAY PROCEDURES

General

The assays of the TME system have several procedures in common and may be performed simultaneously providing that there are sufficient excreta available for the physical and chemical analyses. Birds are fasted, to empty their alimentary canals of feed residues, and then precision-fed a known quantity of the material to be assayed. Each bird is placed in a wire cage, where water is available ad libitum. The time is recorded and the excreta are collected quantitatively for a predetermined period of time. One bird in each replication remains fasted and serves as a negative control for the estimation of metabolic plus endogenous losses. The excreta, together with samples of the test materials, are assayed for gross energy, amino acids, lipid or mineral elements as appropriate. The basic calculation is as follows:

$$TX = IX - (FX + UX) + (F_m X + U_m X + U_e X)$$

where: TX is the true available nutrient X;

IX is the amount of X placed in the fed bird;

(FX + UX) is the amount of X excreted by the fed bird; and

(F_m X + U_m X + U_e X) is the amount of X excreted by the fasted bird.

The preferred bird is a dubbed adult cockerel of an egg-laying strain, which has never had access to grit. Other birds may be used but chicks have limited feed capacity, while fasted laying hens often produce shell-less eggs which tend to break and contaminate the excreta. Laying birds may be useful in assays for TAM where a high mineral requirement is desirable (338), but this is an atypical situation. Grit is avoided because it may be retained in the gizzard and voided on an irregular basis. Grit in excreta damages grinding equipment and introduces major errors into short-term mineral balances.

All birds used in an assay must have been maintained on the same diet. The composition of the diet is not of critical importance providing that it allows the birds to satisfy their nutrient requirements. Many laboratories

feed a laying hen diet, containing 15% of protein, during the maintenance period between assays. The preliminary fast is usually for 24 hours but a longer period may be required if the maintenance diet contains substantial quantities of indigestible materials. Where uncertainty exists it is advisable to measure the gut-clearance time of the maintenance diet before embarking on a series of assays.

As the input of test material increases, the effects of experimental errors are reduced but the possibility of regurgitation increases; 30 to 40 g of air-dry feed is usually satisfactory. Large inputs, particularly of bulky feedingstuffs can lead to crop impaction. Impacted birds have extended feed residue retention times and consequently may yield misleading data. An exception to the foregoing is the TAM bioassay in which the input of the test mineral should be no greater than the bird's requirement because excess minerals are voided as minerals in the excreta.

The use of fasted birds to measure metabolic plus endogenous losses is simple and convenient. The data obtained from a single group of such birds can be used to correct the excreta outputs of several other groups, in the same assay, fed a variety of feedingstuffs. The optimal number of fasted birds in an assay depends on several factors as discussed by Wolynetz and Sibbald (526). However, the use of fasted birds is not essential. The objective is to measure the relationship between the nutrient excreted and the nutrient input and this can be achieved by using two or more levels of input, none of which need be zero.

Initially, it was recommended that the test material be pelleted but this is not necessary if the stem of the feeding funnel has an internal diameter of about 1.0 cm; however, care must be taken to avoid loss of feed by adherence to the funnel. Very dusty or hygroscopic materials are best fed in conjunction with a carrier, such as 90% ground maize plus 10% of vegetable oil; this makes it necessary to assay the carrier. The use of precision feeding is not essential although it is preferred. Assay data have been obtained successfully from birds allowed to consume feed voluntarily.

Test materials are weighed prior to an assay and held in containers until used. Clear polypropylene containers (130 ml) with close fitting lids such as are used for urine samples are satisfactory. Sub-samples of the test materials must be weighed for dry matter determination at the same time as the containers are being prepared. The timing is important because it avoids errors associated with water uptake or loss by the test materials. All analytical data describing the test materials should be expressed on the basis of dry matter.

In this laboratory, birds are housed and maintained in individual wire cages on the lower level of a two-tier system. Cage design and construction are described later (page 22). Water is provided by a nipple system and feed is available in a trough running the full length of each block of cages. At the start of an assay, fasting is initiated by removal of the feed trough. Under other management systems, such as when water is provided in troughs, it is important that feed in the water system or adhering to the cages be removed.

Fasted birds are taken from the lower tier of cages, given the appropriate treatment, and then housed in alternate cages of the upper tier. The technique of precision feeding is described later (page 17). The upper cages are used only during excreta collection periods and are scrupulously cleaned prior to each assay. Excreta collection trays are placed under each bird. The trays, preferably made of smooth plastic, are larger than the bottom of each cage thus reducing the chance of excreta loss. An alternative excreta collection technique is described later (page 28).

Handling of birds causes losses of scale and feathers making quantitative excreta collection difficult. Blowing scale off the excreta collection trays an hour after housing reduces the problem. Excreta are collected at about 24 and again at exactly 48 hours after housing. A single 48 hour collection may be satisfactory but the double collection is favoured, when trays are used, because it reduces excreta deterioration and contamination. When the assays were first described a collection period of 24 hours was suggested but subsequent work showed this to be insufficient for clearance of the residues

of some test materials. Excreta adhering to the cage must be collected and excreta must be washed from any feathers trapped on the excreta collection tray. The trays must be checked for regurgitated feed which, if present, invalidates the use of the bird in the assay. The two excreta samples from each bird are frozen, dried, equilibrated with atmospheric moisture, weighed, pooled, ground, mixed and assayed. Freeze-drying is preferred because it leaves the excreta in a sponge-like, easy-to-grind form; however, oven drying has been shown to be satisfactory for TME (97) and TAAA (370) bioassays. In some laboratories the excreta from several birds are pooled to reduce the analytical work. The procedure should not alter the estimated TME, TAAA, TAL or TAM values but it does restrict the ability to assess variation and to make comparisons between samples (343).

Recent work has demonstrated the advisability of correcting TME values to zero nitrogen balance (TME_n). As a first step in the calculations the excreta energy output ($FE + UE$) is corrected to zero nitrogen balance ($(FE_n + UE_n)$), as follows:

$$(FE_n + UE_n) = (FE + UE) + k(IN - FN - UN)$$

where: k is a constant which estimates the gross energy content of the excretory products resulting from the catabolism of a unit weight of tissue nitrogen;

IN is the nitrogen input as test material; and

FN and UN are the fecal and urinary nitrogen outputs.

For fasted birds, IN is zero. In most assays the term $k(IN - FN - UN)$ is negative; consequently, $(FE_n + UE_n)$ is usually smaller than $(FE + UE)$.

The nitrogen-corrected energy excretion of a fasted bird is better described as $(FE_{mn} + UE_{mn} + UE_n)$. TME_n values are calculated as follows:

$$TME_n = IE - (FE_n + UE_n) + (FE_{mn} + UE_{mn} + UE_n)$$

where: IE is the amount of energy, as test material, placed in the bird.

There is no evidence that similar corrections are required in any of the other bioassays of the TME system.

Precautions

The following is a list of precautions to be observed if high quality assay data are to be obtained. The list concludes with the most common causes of abnormally large and small values.

1. Assay birds must be in good health.
2. All assay birds must be fed the same maintenance diet between assays.
3. Assay birds should be grit-free.
4. Test materials must be assayed for dry matter at the time that they are weighed into containers in preparation for feeding to the birds.
5. Dusty and hygroscopic test materials should be fed in conjunction with a carrier; the carrier must also be assayed.
6. Assay birds must be fasted for sufficient time to allow all feed residues to be voided.
7. Feed removal for fasting must be total. Feed adhering to the cage and usually ignored by the bird will be eaten if no other is available.
8. Birds must have continuous access to fresh water.
9. Excreta trays must be checked for regurgitated feed which, when found, eliminates the bird from the assay.
10. The excreta collection period must be exactly the same for all birds in an assay.
11. When using adult cockerels, and feed inputs of 30 to 40 g, an excreta collection period of 48 hours should be sufficient. For other birds and inputs a preliminary experiment may be necessary to establish the length of the collection period.
12. Excreta collection must be quantitative. Feathers must not be included and scale is to be avoided as much as possible.
13. Dried excreta should be equilibrated with atmospheric moisture or be held in such a manner that its moisture content remains constant between weighing and sub-sampling for analysis.

Abnormally large values may result from:

- a. incomplete clearance of the residues of the test material, check for crop impaction;
- b. incomplete excreta collection, excreta may have been voided beyond the collection tray;

- c. unobserved regurgitation beyond the collection tray causing excreta output to be too small;
- d. errors in weighing, preparing or administering the test material; and
- e. analytical errors.

Abnormally small values are obtained when:

- a. the preliminary fast is inadequate and residues of the maintenance diet are assumed to be derived from the test material;
- b. the bird has access to feed during the fast;
- c. regurgitated feed is mixed with the excreta;
- d. scale or feathers are included with the excreta; and
- e. preparatory and analytical errors.

The assays are simple and relatively fast. However, like all assays they require care and attention to detail if high quality data are to be obtained.

Precision Feeding

The purpose of precision feeding is to ensure that a known quantity of a feedingstuff enters the alimentary canal of a bird at a known time. The procedure avoids the need to recover waste feed, prevents feed selection and eliminates variation of intake among birds. All of these problems are encountered when birds ingest feed voluntarily.

Precision feeding involves insertion of a tube from the beak, via the esophagus, into the crop, causing feed to move through the tube, and removal of the tube. The initial method developed at the Animal Research Centre used a simple glass tube into which pelleted feed was placed and then pushed into the crop with a glass rod. From this was developed a glass funnel and metal plunger. With experience came the ability to insert tubes with larger diameters and an effective device was prepared by taping a glass powder funnel to a piece of 1.2 cm diameter copper water pipe. A metal rod was used as a plunger. Today, a stainless steel funnel is used with a stem 40 cm long, an external diameter of 1.2 cm and an internal diameter of 1.1 cm. The plunger consists of a 1.0 cm diameter aluminum rod to which a 3.0 cm diameter spherical knob is attached. A plastic sleeve is riveted to the rod to

prevent the plunger from projecting more than 0.5 cm beyond the end of the funnel. Ease of operation demands that the funnel be light in weight and well balanced. Heavy funnels, particularly those with uneven weight distribution, are difficult to control when in the bird. Plastic funnels have the advantage of light weight but many possess electrostatic properties which make quantitative feed delivery impossible. Some of the feeding devices which have been used in this laboratory are shown in Figures 3 and 4.

Successful precision feeding depends upon control of the bird. A simple, rapid technique used at the Animal Research Centre is as follows. The operator sits on a stool, or chair without sides, and crosses his left leg over his right. A bird is taken with both hands and placed with the keel (breast bone) pushed into the groove formed by the left thigh and abdomen of the operator. The body of the bird is at 45 degrees from the vertical. The legs of the bird project to the left and are unable to gain leverage on anything. The bird is held firmly in place with the elbow of the left arm, and the beak is grasped from above the head with the left thumb and forefinger. The neck of the bird is slightly extended, the beak is opened and the stem of the funnel is inserted. The funnel should move down the esophagus and into the crop with a minimum of effort. If a blockage is encountered, wait a few seconds for the bird to swallow or relax. If this is unsuccessful, remove the funnel and try again. Feed is poured into the funnel from a container held in the right hand and pushed into the crop by a second operator. If the funnel tends to move out of the bird as feed is pushed down, the end of the funnel is in the esophagus not the crop. Deposition of feed in the esophagus can lead to regurgitation. The funnel is withdrawn using the right hand. The left hand is moved to the neck of the bird and gentle pressure is applied to the esophagus. This, coupled with rotary withdrawal of the funnel, removes feed particles which may have adhered to the outside of the stem of the funnel. The complete operation usually takes less than one minute per bird.

The precision feeding technique is illustrated in Figures 5 to 12. A six-minute, 16 mm, colour, sound film entitled "Poultry Force Feeding - An Experimental Technique" is available by writing to: Communications Branch, Agriculture Canada, Ottawa, Ontario, Canada, K1A 0C7.




Figure 3.

Some early feeding devices: (A) glass funnel and rod, (B) copper water pipe attached to a glass powder funnel and an aluminum plunger, (C) simple glass tube and rod.




Figure 4.

Stainless steel funnel (A) with aluminum plunger (B). Note plastic sleeve (C) on plunger which controls penetration (D) beyond the end of the funnel.



Figure 5. Bird is placed at a 45° angle with keel in groove formed by the operator's thigh and abdomen.



Figure 6. The legs project and cannot gain leverage on anything. The bird is held in place by the left elbow.



Figure 7. The beak is opened from above using the thumb and forefinger of the left hand.



Figure 8. The neck is slightly extended and the stem of the funnel inserted.



Figure 9. The inserted funnel is held in place by the thumb and forefinger.



Figure 10. Feed is poured into the funnel.



Figure 11. Feed is pushed down the funnel by a second operator. Note that the first operator helps to hold the funnel in place.



Figure 12. The funnel is removed with a rotary action, pressure is applied to the esophagus with the left hand to dislodge feed particles adhering to the stem of the funnel.

Cage Design and Construction

The cages used to house the adult cockerels have the following dimensions: depth (front to back) 40.6 cm, height 50.8 cm and width 30.5 cm. Figures 13 to 21 describe the cages, their assembly and use; all dimensions are in centimetres. The back, top and floor of a row of 10 cages is fabricated from a single piece of wire mesh approximately 1.3 m wide x 3.05 m long. The wire is 2.03 mm diameter (14 gauge). The mesh is a 5.1 x 2.5 cm grid welded at each corner. The internal partitions and ends are made of the same material. The cage fronts, also 3.05 m in length, are made from 4.11 mm diameter (8 gauge) wire. The door opening of 23 x 28 cm provides adequate access to remove the birds. The cages are set up in a double-deck arrangement and the droppings board is placed horizontally to hold the excreta collection trays. Water is available from a nipple system on the top of the cages.




Figure 13. End view of the single piece of wire mesh which is folded to create the body (top, back and bottom) of a row of 10 cages.




Figure 14. A typical cage partition or end. Partitions are located at 30.5 cm intervals and are welded in place.




Figure 15. A. The front for a row of cages.
B. Side view of cage front.




Figure 16. A. The front with cage doors installed.
B. Side view of cage door.
C. Detail of cage door.




Figure 17. The relationship between the cage body with partitions and the front. For simplicity, only 2 doors are shown on the front.




Figure 18. An assembled cage unit with feed trough.




Figure 19 Side view of a 4-row double deck battery of cages showing, (A) position of droppings board to hold excreta collection trays and (B) normal position of droppings board.




Figure 20. Cages being used to maintain birds between assays.




Figure 21. Birds housed in cages of upper tier being used in an assay.

Excreta Collection

Initially, excreta were collected on smooth plastic trays which were larger than the bottoms of the cages. The procedure continues to be used in many laboratories but has the disadvantage that scale and feathers may be trapped in the excreta. There is also the possibility of excreta falling beyond the edges of the trays.

A procedure whereby human colostomy bags are attached to birds to collect excreta was described in the previous edition of this Bulletin. After working with the procedure in a number of assays it was abandoned because: a) there was a high incidence of adhesive failure and samples were lost due to bags falling off the birds; and b) removal of those bags which remained attached caused feathers to be pulled out so that the area around the cloaca was denuded.

A harness designed to hold an excreta collection bag has been developed in Portugal (444). Modifications have been made to the design and the harness is now used routinely at the Animal Research Centre. The major modification is the addition of two short tapes which are tied around the base of the tail to prevent vertical slippage. When using the harness four holes are cut in the collection bag and the tapes are threaded through them. The neck of the bag is then folded back. With this procedure the use of adhesive tape to hold the bag to the harness is avoided. When the harness is on the bird, the ends of the two long tapes are tied together as a precaution against lateral slippage. The harness, its method of attachment, and its use are illustrated in Figures 22 to 35.

At the Animal Research Centre, the collection bags are labelled, holes for the tapes are cut, the weight of each bag is recorded, and the bags are attached to the harnesses. The harnesses, with bags, are attached to the birds immediately after precision feeding. At the conclusion of an excreta collection, the harnesses are removed from the birds and the bags plus excreta are separated from the harnesses. The excreta are frozen, freeze-dried, and equilibrated with atmospheric moisture while in the bags. The procedure is efficient and provides excreta free of scale and feathers.

Care is required when attaching a harness to a bird. If the tension of the tapes is too great the bird has difficulty maintaining its balance and bruising appears in the wings. Even when the minimum required tension is used there is a temporary disturbance of balance for one or two minutes.

Birds do peck at the bags and can cause additional damage with their spurs. Debeaking and spur removal reduces the problem and has the additional benefit of making the birds easier to handle. Detection of leaking bags, and of regurgitation, is made easier if excreta collection trays are placed below harnessed birds. Clipping the feathers around the cloaca is recommended to prevent excreta adherance.




Figure 22. The excreta collection harness comprises a plastic ring (OD 7.5 cm, ID 5.75 cm) (A), 2 long cotton tapes (40 cm) (B), 2 short cotton tapes (20 cm) (C) and 2 elastic tapes (10 cm) (D) each attached to two small plastic rings (OD 1.9 cm) (E) which serve as buckles. Each D is sewn to the corresponding C at F. All the tapes are 1.25 cm wide and the ends of the cotton tapes are varnished to prevent fraying.



Figure 23. The tapes of the harness are threaded through holes in a labelled and weighed plastic bag which measures 15.2 x 25.4 cm. x .10 mm thick (see also Figure 34).



Figure 24. Harness and bag about to be attached to the bird. The first step is to attach the short tapes around the base of the tail. Note that the neck of the bag has been everted over the large ring.



Figure 25. Tying the short tapes.



Figure 26. A long tape is placed between the thigh and the abdomen and is then brought around the shoulder.



Figure 27. Each long tape is buckled to the elastic tape on the opposite side of the back using the pair of small rings. When in place these tapes form a cross on top of the back.



Figure 28. The ends of the long tapes are tied together to reduce the chance of lateral slippage.



Figure 29. The location of the plastic ring when the harness is in place. The collection bag has been removed. Note that the feathers around the cloaca have been clipped.



Figure 30. The location of the collection bag when the harness is attached.



Figure 31. Bird immediately after attaching bag and harness.



Figure 32. Bird with plastic bag containing excreta.



Figure 33. Rear view of bird showing collected excreta.



Figure 34. Bag and harness removed from bird.



Figure 35. Bag with excreta ready for further processing.

FEEDINGSTUFF COMPOSITION

Changes in feedingstuffs over time and the introduction of new analytical procedures combine to create a continuing demand for data describing feedingstuff composition. The second edition of this Bulletin contained a simple table listing the proximate composition, gross energy, TME and TME_n values of a variety of feed ingredients. The information has now been expanded in two directions. The original table has been revised and now includes data describing more than 200 additional samples; for many of the new samples calcium, total phosphorus and neutral detergent fibre values are presented. Three new tables present total and true available amino acid values for many of the new samples. Data are expressed as g/kg or MJ/kg of dry matter except in Table 4 where percentages are used.

Samples of feedingstuffs were obtained from several sources including commercial feed manufacturers, the Canadian Wheat Board, plant breeders and the Animal Research Centre Research Farm. On receipt, each sample was given a code number and all available information, descriptive of the sample, was entered in a catalogue. The samples were stored in sealed containers in a freezer until required for analysis.

At the start of a bioassay the required number of samples was withdrawn from storage, in the order of their receipt, and allowed to warm to room temperature. Coarse samples, such as whole grains, were ground. Each sample was thoroughly mixed and sub-samples were taken for the bioassay and for dry-matter measurement. If a sample was finely divided, additional sub-samples were taken for physical and chemical analysis. Alternatively, less fine material was reground prior to sub-sampling for the physical and chemical analyses; when this step was introduced new dry-matter measurements were made to minimize errors associated with variation in water content.

The feedingstuffs were assayed for dry matter, ether extract, nitrogen, crude fibre, ash and calcium using the procedures of the A.O.A.C. (Official Methods of Analysis, 1980). Phosphorus was measured by Hambleton's method

(Advances in Automated Analysis, 7th Technicon Int. Conf., 1976) and neutral detergent fibre by the procedure of Van Soest (J.A.O.A.C. 50:50-55, 1967). Gross energy was measured with a Parr adiabatic oxygen bomb calorimeter fitted with a 'master control' unit. Amino acids were measured with a Beckman 121M automatic analyzer following hydrolysis with 6N HCl (J. Agr. Food Chem. 13:266-288, 1965). A performic acid digestion (J. Biol. Chem. 238:235-237, 1963) was used in the analysis for sulphur amino acids.

The TME and TME_n values were measured as described herein. The preliminary fast was 24 hours, the feed input was usually 30 g of air-dry material per bird, and the excreta collection period was 48 hours. Each TME and TME_n value is the mean of 6 to 8 replicated observations. The TAAA assays were made using the excreta samples collected during the TME assays; however, excreta from birds given a particular feedingstuff were pooled, in proportion to the amounts voided by individual birds, to yield two composite samples each containing excreta from three or four birds. Amino acid profiles were obtained for each pooled sample and used to compute two sets of TAAA data from which the means of Tables 3 and 4 were calculated. Amino acid analyses of feedingstuffs were made in duplicate.

In Table 4 there are 18 values in excess of 100%. Most describe the bioavailabilities of arginine and lysine in ground yellow corn. Such values are erroneous and probably arose because the estimates of metabolic and endogenous amino acid losses, measured with fasted birds, differed slightly from those of the fed birds. Examination of the relevant duplicate values obtained with fasted birds showed them to be more variable and to have larger means than average. In other experiments (57, 95) the bioavailabilities of corn amino acids have approached 100% so it is reasonable that even small errors could result in substantial over-estimates. Additional factors possibly contributing to error are the low amino acid concentrations in corn and the relatively high lysine concentrations in the excreta of fasted birds.

Each feedingstuff is described in Table 1. Only abbreviated descriptions appear in Tables 2, 3 and 4 but reference to Table 1 descriptions is possible

by use of the code numbers which appear in the left hand column of each table. The code numbers also serve to allow readers to identify information added since the second edition (621-862). Those who have included the earlier table in their data banks will wish to avoid double entry of information and the consequent bias.

The nitrogen data are recorded as g/kg of dry matter. No attempt was made to estimate protein concentrations because the validity of some conversion factors is debatable. Recently, the assay for crude fibre was replaced by that for neutral detergent fibre; the latter assay does not apply to animal products and their fibre values are no longer estimated. Compositional means for feedingstuffs are not presented because they may be misleading if an anomalous sample is included.

The term 'whole ground' used to describe some grains means that whole, intact kernels were ground and assayed; there was no intentional classification or separation of the ground product. The term 'hull-less' denotes a genotype which produces a grain with little or no hull and should not be confused with 'dehulled' which implies physical removal of the hull. Oat groats, for example, are dehulled oats.

There are many sources of additional data which could be of interest to those constructing data banks. Many references will be found in the bibliography. The data recorded in Tables 1, 2, 3 and 4 were obtained within the facilities of Agriculture Canada subsequent to publication of a table appearing in reference 32.

TABLE I

Nitrogen, fat, fibre, mineral (g/kg) and energy (MJ/kg) concentrations in the dry matter of several feedingstuffs^{1,2}

Code	Description of Feedingstuff ³	Nitrogen			Ether Extract	Crude Fibre	Neutral Detergent Fibre	Total Phosphorus	Gross Energy	TME	TME _n
		Ether	Nitrogen	Extract							
476	Alfalfa. dehydrated, meal.	28.4	23.0	250	-	-	91	-	18.9	6.5	6.1
359	Alfalfa. dehydrated, pellets.	30.2	26.1	290	-	-	92	-	18.8	6.6	-
372	Alfalfa. dehydrated, pellets.	25.4	34.0	291	-	-	91	-	18.6	4.9	-
440	Alfalfa. dehydrated, pellets.	23.3	30.7	276	-	-	93	-	18.9	4.4	4.4
MS	Alfalfa. dehydrated, pellets.	27.6	23.0	250	-	-	90	-	18.9	5.7	-
655	Alfalfa. dehydrated, pellets, second cut, Minnesota.	26.2	19.4	-	443	93	15.5	2.2	18.6	4.8	4.8
656	Alfalfa. dehydrated, pellets, first cut, Minnesota.	30.8	15.4	-	383	103	19.5	2.6	18.0	6.3	5.5
341	Alfalfa. dehydrated, pellets, 15% protein.	25.7	20.6	315	-	-	76	-	18.8	6.6	-
653	Alfalfa. dehydrated, pellets, (16%), 2nd cut, Minn.	26.2	19.0	-	429	92	16.1	2.2	18.6	4.8	4.5
839	Alfalfa. dehydrated, pellets, 17% protein, Nebraska.	28.2	18.5	-	385	92	8.8	2.4	18.8	4.2	4.0
378	Alfalfa. dehydrated, pellets, 18% protein.	30.1	51.8	261	-	-	78	-	18.7	5.6	-
657	Alfalfa. dehydrated, pellets, 18% protein, Nebraska.	30.1	18.5	-	382	94	12.0	2.6	18.7	4.2	4.0
652	Alfalfa. dehydrated, pellets, 19% protein, Nebraska.	29.8	23.3	-	394	95	14.0	2.8	18.7	4.8	4.6
338	Alfalfa. dehydrated, pellets, 20% protein.	36.2	35.0	213	-	-	95	-	19.1	7.0	-
654	Alfalfa. dehydrated, pellets, 20% protein.	29.8	17.5	-	409	104	10.4	2.4	18.6	3.8	3.7
348	Alfalfa. dehydrated, reground pellets, 15% protein.	26.8	20.7	265	-	-	111	-	18.1	4.8	-
349	Alfalfa. dehydrated, reground pellets, 17% protein.	29.4	24.7	238	-	-	116	-	18.2	5.6	-
350	Alfalfa. dehydrated, reground pellets, 20% protein.	37.2	32.9	224	-	-	123	-	18.2	6.1	-
351	Alfalfa. dehydrated, reground pellets, 22% protein.	37.2	30.5	199	-	-	128	-	18.5	6.6	-
339	Alfalfa. dehydrated, pellets, (17%) + vegetable fat.	28.2	21.8	301	-	-	90	-	18.8	4.9	-
340	Alfalfa. dehydrated, pellets, (22%) + ethoxyquin.	38.0	32.7	203	-	-	92	-	18.9	7.4	-
649	Algae. dehydrated.	60.6	19.2	-	6	69	4.9	13.9	22.2	9.2	8.5
650	Algae. dehydrated.	59.0	27.0	-	24	142	43.0	7.4	21.0	10.8	9.8
607	Alpha-protein. Nutritional Biochemicals.	143.9	-	-	-	20	-	-	23.0	15.4	14.3
609	Assay protein. Ralston Purina.	121.7	-	-	-	49	-	-	23.1	18.5	17.8
577	Babassu meal.	30.1	7.1	346	-	-	75	-	-	18.4	4.6
578	Babassu meal.	29.6	13.9	425	-	-	66	-	-	18.6	4.8
534	Barley. whole ground.	20.1	10.0	45	-	-	23	-	-	18.3	4.0
435	Barley. whole ground.	20.8	15.8	39	-	-	22	-	-	18.6	4.3
689	Barley. whole ground.	23.4	24.3	-	154	23	-	.5	3.0	13.5	13.0
690	Barley. whole ground.	21.2	15.0	-	183	25	-	-	18.7	14.1	13.6

TABLE 1 cont.

Code	Description of Feedingstuff	Nitrogen	Ether Extract	Crude Fibre	Neutral Detergent Fibre	Ash	Calcium	Gross Energy	TME	TME _n
691	Barley. whole ground.	15.8	14.8	-	181	25	.5	3.9	13.7	13.4
692	Barley. whole ground.	21.1	16.4	-	139	25	.5	3.8	13.9	13.5
693	Barley. whole ground.	18.4	16.0	-	178	26	.4	3.7	13.8	13.3
694	Barley. whole ground.	19.5	14.1	-	175	29	.4	6.7	18.2	14.3
695	Barley. whole ground.	22.2	15.3	-	214	35	.6	3.9	18.5	13.0
696	Barley. whole ground.	20.2	20.3	-	164	27	.5	6.4	18.3	13.4
697	Barley. whole ground.	20.7	14.4	-	134	26	.5	6.2	18.3	13.4
698	Barley. whole ground.	23.4	15.7	-	140	27	.5	6.0	18.3	13.3
392	Barley. whole ground. Ontario.	22.1	18.0	47	-	24	-	-	18.7	14.5
473	Barley. whole ground. Quebec.	18.1	14.5	66	-	31	-	-	18.5	14.1
531	Barley. whole ground. (cv. Bruce).	20.0	16.5	57	-	29	-	-	18.2	13.5
367	Barley. whole ground. (cv. Conquest).	20.1	16.9	58	-	32	-	-	18.2	14.7
532	Barley. whole ground. (cv. Leger).	20.4	14.7	58	-	27	-	-	18.4	13.4
592	Barley. whole ground. (cv. Massey).	20.6	15.6	61	-	30	-	-	18.4	12.8
530	Barley. whole ground. (cv. Vanier).	18.1	15.2	64	-	29	-	-	18.3	14.0
769	Barley. whl. grd., No. 2 CW, 2-row, immat., 60kg/hL.	22.5	17.2	-	236	28	.7	7.0	18.5	13.6
765	Barley. whl. grd., No. 2 CW, 2-row, immat., 64kg/hL.	20.9	16.3	-	238	27	.6	5.3	18.4	14.3
762	Barley. whl. grd., No. 1 feed, green.	22.8	16.6	-	254	30	.5	5.3	18.5	13.6
775	Barley. whl. grd., No. 1 feed, green, 60 kg/hL.	17.9	15.0	-	213	32	.6	5.5	18.2	13.6
777	Barley. whl. grd., No. 1 feed, inf. var., 60 kg/hL.	26.4	15.7	-	261	25	.7	3.1	18.6	13.5
767	Barley. whl. grd., No. 1 feed, standard.	16.5	15.2	-	195	25	.6	3.3	18.2	14.1
768	Barley. whl. grd., No. 1 feed, standard, 72kg/hL.	18.9	16.0	-	186	27	.5	3.1	18.3	14.6
761	Barley. whl. grd., No. 1 feed, 46% plump, 6% thin.	20.8	16.5	-	272	27	.6	5.2	18.4	13.7
763	Barley. whl. grd., No. 1 feed, 60% plump, 8% thin.	19.0	11.1	-	239	27	.5	5.3	18.3	14.2
766	Barley. whl. grd., No. 1 feed, 38% plump, 6% thin,	26.9	17.5	-	256	20	.6	2.8	18.7	14.8
				(68 kg/hL.)						
403	Barley. hull-less, ground.	26.8	14.0	34	-	29	-	-	18.8	13.5
404	Barley. hull-less, ground.	29.6	15.4	22	-	27	-	-	18.7	15.4
405	Barley. hull-less, ground.	28.2	13.2	31	-	27	-	-	18.8	14.1
406	Barley. hull-less, ground.	27.8	16.9	37	-	28	-	-	18.7	13.3
408	Barley. hull-less, ground.	28.3	14.8	30	-	29	-	-	18.8	13.9
409	Barley. hull-less, ground.	24.3	12.2	50	-	28	-	-	18.5	14.6
410	Barley. hull-less, ground.	27.8	12.7	29	-	27	-	-	18.6	14.0
411	Barley. hull-less, ground.	-	-	-	-	-	-	-	18.8	14.6
449	Barley. hull-less, ground.	27.8	13.5	22	-	17	-	-	19.0	14.9
450	Barley. hull-less, ground.	24.6	12.5	22	-	16	-	-	18.9	14.6
451	Barley. hull-less, ground.	26.7	13.2	21	-	17	-	-	18.3	15.1
452	Barley. hull-less, ground.	24.3	12.2	21	-	15	-	-	18.9	15.6

TABLE I cont.

Code	Description of Feedingstuff	Nitrogen	Ether Extract	Crude Fibre	Neutral Detergent Fibre	Ash	Calcium	Total Phosphorus	Gross Energy	TME	TME _n
453	Barley. hull-less, ground.	27.4	15.6	23	-	18	-	-	18.9	15.3	14.8
454	Barley. hull-less, ground.	25.5	10.2	22	-	18	-	-	18.8	15.3	14.6
455	Barley. hull-less, ground.	27.3	13.1	22	-	17	-	-	18.9	15.4	14.9
456	Barley. hull-less, ground.	27.5	11.4	23	-	17	-	-	19.0	15.1	14.7
457	Barley. hull-less, ground.	26.8	14.4	21	-	17	-	-	18.9	15.2	14.8
458	Barley. hull-less, ground.	26.4	13.1	21	-	16	-	-	18.8	14.8	14.4
495	Barley. hull-less, ground.	26.0	17.4	33	-	20	-	-	18.7	13.3	13.1
496	Barley. hull-less, ground.	22.8	14.6	26	-	17	-	-	18.6	13.6	13.2
497	Barley. hull-less, ground.	20.4	12.1	24	-	18	-	-	18.4	14.2	13.8
498	Barley. hull-less, ground, (cv. Bichita).	24.5	14.9	21	-	18	-	-	18.5	14.1	13.6
407	Barley. hull-less, ground, (cv. Kun Lun).	26.3	15.5	27	-	24	-	-	18.6	14.0	13.3
549	Beans. black, ground, (cv. Loop).	39.2	12.3	46	-	49	-	-	18.4	6.9	7.0
546	Beans. faba, ground, (cv. Aladin).	46.5	10.9	92	-	30	-	-	18.6	11.6	11.3
545	Beans. faba, ground, (cv. Herz Freya).	43.8	12.4	91	-	34	-	-	18.4	12.0	11.2
547	Beans. mung, ground.	43.2	8.2	55	-	44	-	-	18.5	12.6	12.0
548	Beans. mung, ground, (cv. Morden 39).	39.9	8.7	58	-	85	-	-	17.7	11.9	11.4
544	Beans. white, ground, (cv. Seafarer).	37.7	16.8	47	-	47	-	-	18.5	8.6	8.3
483	Blood meal.	143.7	20.5	23	-	28	-	-	24.6	16.5	15.4
484	Blood meal.	151.6	9.0	14	-	22	-	-	24.6	16.4	15.1
617	Blood meal.	153.3	3.5	4	-	15	-	-	24.8	17.1	15.7
618	Blood meal.	154.0	3.7	3	-	14	-	-	24.8	17.3	16.0
666	Blood meal.	152.2	4.0	-	32	18	1.1	1.6	24.8	16.3	15.4
667	Blood meal.	158.8	5.1	-	12	17	.9	1.3	24.6	17.0	15.6
717	Blood meal.	157.0	1.1	-	-	15	.7	1.0	24.6	15.5	14.1
718	Blood meal.	154.8	5.3	-	-	16	1.4	1.4	24.7	17.7	15.5
719	Blood meal.	154.2	5.1	-	-	23	2.6	2.1	24.5	18.0	15.9
739	Blood meal.	158.0	0.9	-	-	16	.6	1.1	24.5	15.8	14.4
740	Blood meal.	154.9	12.6	-	-	14	1.0	1.5	24.7	17.8	15.8
741	Blood meal.	153.1	8.8	-	-	28	2.3	2.2	24.2	17.0	15.2
742	Blood meal.	158.7	1.9	-	-	15	2.4	2.4	24.3	16.8	15.1
792	Blood meal.	159.0	3.2	-	-	18	1.2	1.8	24.3	16.6	15.5
793	Blood meal.	153.6	31.6	-	-	16	.9	1.8	25.2	18.1	16.2
794	Blood meal.	149.6	2.3	-	-	22	1.0	1.5	23.5	15.3	14.0

TABLE 1 cont.

Code	Description of Feedingstuff	Nitrogen	Ether Extract	Crude Fibre	Neutral Detergent Fibre	Ash	Calcium	Gross Energy	TME	TME _n
795	Blood meal.	157.4	1.7	—	—	18	.5	1.4	24.5	15.3
817	Blood meal.	158.0	11.3	—	—	14	.3	1.4	24.8	17.1
599	Bone meal.	18.7	65.1	—	—	778	—	—	6.4	5.1
537	Brewer's dried grains.	39.0	63.4	198	—	47	—	—	20.6	7.9
562	Brewer's wet grains. freeze-dried.	46.6	75.8	142	—	45	—	—	21.4	10.3
	Brewer's yeast. See Yeast.									9.8
519	Buckwheat. whole ground.	19.7	26.2	140	—	23	—	—	18.5	14.1
556	Buckwheat. whole ground, (cv. Mancan).	23.4	26.1	125	—	27	—	—	18.7	13.7
	Canola meal. See Rapeseed meal.									12.9
606	Casein. vitamin-free, General Biochemicals.	145.4	—	—	—	57	—	—	24.2	20.5
604	Cellulose. Alphacel, Nutritional Biochemicals.	1.6	—	833	—	3	—	—	17.3	ND
543	Cereal offal.	28.7	22.5	25	—	32	—	—	18.5	15.8
	Cerelose. See Glucose.									15.0
643	Coconut meal.	37.2	28.8	—	597	72	1.6	10.8	18.8	11.4
363	Corn. ground, yellow.	22.1	48.7	21	—	18	—	—	19.0	17.4
380	Corn. ground, yellow.	15.7	39.8	23	—	14	—	—	18.3	16.8
387	Corn. ground, yellow.	17.6	43.0	21	—	17	—	—	18.9	17.3
436	Corn. ground, yellow.	13.5	40.6	27	—	15	—	—	18.8	16.6
563	Corn. ground, yellow.	13.2	36.6	23	—	13	—	—	18.4	16.4
838	Corn. ground, yellow.	16.2	39.8	—	13	14	.2	3.2	18.8	16.9
852	Corn. ground, yellow.	15.6	—	—	—	—	—	—	18.6	17.1
853	Corn. ground, yellow.	15.0	38.1	—	110	15	.1	3.4	18.9	17.0
855	Corn. ground, yellow.	16.8	29.2	—	111	15	.1	3.0	18.9	16.8
856	Corn. ground, yellow.	15.6	33.8	—	99	14	ND	3.4	18.8	17.4
857	Corn. ground, yellow.	16.5	20.4	—	98	12	.1	3.0	18.9	17.8
858	Corn. ground, yellow.	16.7	11.0	—	114	15	.1	3.6	18.9	17.3

TABLE 1 cont.

Code	Description of Feedingstuff	Nitrogen	Ether Extract	Crude Fibre	Neutral Detergent Fibre	Ash	Calcium	Total Phosphorus	Gross Energy	TME	TME _n
859	Corn. ground, yellow.	17.2	43.8	-	125	16	.1	3.7	18.9	17.3	16.4
860	Corn. ground, yellow.	17.5	43.7	-	111	14	.1	3.5	19.1	18.0	17.0
861	Corn. ground, yellow.	17.3	32.8	-	130	15	.1	3.3	18.9	17.0	16.2
862	Corn. ground, yellow.	16.5	35.0	-	110	15	.1	3.6	18.9	16.7	16.1
470	Corn. ground, yellow, Quebec.	13.6	44.8	21	-	12	-	-	18.9	17.4	16.6
467	Corn. ground, yellow, No. 3, Ontario.	13.3	41.1	21	-	14	-	-	18.7	17.4	16.4
468	Corn. ground, yellow, No. 3, U.S.	14.1	42.9	22	-	14	-	-	18.8	17.6	16.6
469	Corn. ground, yellow, No. 3, U.S.	15.0	44.7	23	-	13	-	-	18.8	17.1	16.1
646	Corn germ meal.	38.1	67.8	-	308	21	.2	7.2	20.6	9.0	8.8
507	Corn gluten feed.	40.5	55.1	88	-	49	-	-	19.9	11.7	11.0
536	Corn gluten feed.	37.1	15.7	104	-	77	-	-	18.5	8.3	8.0
651	Corn gluten feed.	44.1	62.9	-	287	56	.6	9.8	20.5	12.4	11.6
674	Corn gluten feed.	33.4	38.9	-	310	49	.3	11.1	19.2	11.1	10.0
726	Corn gluten feed.	37.0	42.7	-	325	50	.7	10.4	19.4	10.1	9.1
727	Corn gluten feed.	39.0	42.0	-	286	49	.7	10.3	19.5	10.8	10.0
754	Corn gluten feed.	41.2	43.7	-	293	53	.2	11.5	19.7	10.3	9.8
755	Corn gluten feed.	40.1	42.0	-	333	55	.2	10.8	19.4	9.7	9.2
805	Corn gluten feed.	41.8	29.9	-	323	59	.3	11.2	19.4	11.0	10.6
806	Corn gluten feed.	40.3	39.0	-	355	54	.3	10.5	19.7	10.5	10.0
827	Corn gluten feed.	37.6	36.2	-	318	48	.3	10.5	19.4	10.2	9.6
828	Corn gluten feed.	34.4	7.4	-	257	58	.2	10.3	18.4	10.8	9.9
445	Corn gluten feed. flakes.	36.4	33.1	92	-	54	-	-	19.5	10.2	9.6
493	Corn gluten meal.	101.8	11.6	16	-	41	-	-	22.8	17.9	17.2
616	Corn gluten meal.	107.0	10.6	8	-	37	-	-	23.2	17.9	17.3
675	Corn gluten meal.	87.0	10.0	-	38	16	ND	5.3	22.0	18.5	17.3
725	Corn gluten meal.	112.7	7.9	-	12	18	.4	6.7	23.2	18.4	17.3
752	Corn gluten meal.	106.3	9.8	-	14	26	-	5.8	23.0	18.7	17.3
753	Corn gluten meal.	106.4	9.3	-	12	29	.2	5.9	22.8	18.6	17.3
807	Corn gluten meal.	112.5	11.7	-	ND	34	ND	5.2	23.3	18.1	17.2
808	Corn gluten meal.	118.6	12.3	-	ND	26	ND	5.3	23.7	18.2	17.2
829	Corn gluten meal.	110.6	15.1	-	5	15	ND	5.6	23.5	18.4	17.4
830	Corn gluten meal.	109.6	15.4	-	13	16	ND	4.8	24.0	19.1	17.7
831	Corn gluten meal.	107.6	3.6	-	7	15	ND	5.1	23.3	18.5	17.5
521	Corn starch. pearlized.	1.0	1.8	4	-	ND	-	-	17.0	17.6	16.2
600	Corn starch. pearlized.	.9	-	-	-	-	-	-	17.2	17.0	16.9

TABLE 1 cont.

Code	Description of Feedingstuff	Nitrogen	Ether Extract	Crude Fibre	Detergent Fibre	Ash	Calcium	Total Phosphorus	Gross Energy	TME	TME _n
373	Cottonseed meal.	82.2	16.2	113	-	-	-	19.7	9.8	-	-
374	Cottonseed meal.	73.8	20.4	148	-	-	-	19.5	10.4	-	-
504	Crab meal. New Brunswick.	54.4	25.4	225	-	418	-	11.5	4.0	3.6	-
567	Distiller's dried grains.	48.7	138.4	102	-	52	-	23.2	14.8	13.8	-
510	Distiller's dried grains. corn.	47.7	31.1	100	-	44	-	22.5	13.4	13.3	-
501	Distiller's dried grains. with solubles.	45.2	92.1	101	-	45	-	22.2	12.8	12.4	-
540	Distiller's dried grains. with solubles.	43.8	78.6	105	-	48	-	21.8	12.7	12.2	-
561	Distiller's wet grains. freeze-dried.	66.8	69.6	214	-	11	-	-	22.6	11.8	10.7
601	Egg albumen solids. spray-dried, pasteurized.	132.6	-	-	-	70	-	-	22.0	16.3	15.7
596	Egg white. raw, spray-dried.	134.1	-	-	-	66	-	-	22.0	16.3	14.8
394	Feather meal.	129.2	53.0	-	-	34	-	-	23.3	16.1	-
477	Feather meal.	137.6	91.7	13	-	25	-	-	24.8	17.1	15.3
478	Feather meal.	143.1	56.0	13	-	18	-	-	24.2	15.7	14.0
500	Feather meal.	142.8	34.5	12	-	19	-	-	24.0	14.6	13.6
663	Feather meal.	131.6	111.4	-	-	136	32	8.5	5.7	24.8	17.4
664	Feather meal.	139.8	61.0	-	-	159	22	4.7	3.0	24.0	15.2
665	Feather meal.	143.0	80.9	-	-	292	23	3.4	5.2	24.5	15.6
714	Feather meal.	142.5	80.1	-	-	18	3.4	2.3	24.8	15.5	13.6
715	Feather meal.	141.7	84.3	-	-	16	3.1	1.9	24.7	17.0	14.7
716	Feather meal.	140.7	99.3	-	-	20	4.4	2.9	24.8	17.0	14.6
737	Feather meal.	142.4	77.6	-	-	22	3.7	2.4	24.4	15.2	13.6
738	Feather meal.	142.6	53.2	-	-	19	3.8	2.1	23.8	15.4	13.2
790	Feather meal.	143.2	85.5	-	-	18	3.8	2.7	24.7	15.8	14.2
791	Feather meal.	144.6	71.7	-	-	21	5.0	3.2	24.3	17.5	15.1
815	Feather meal.	141.1	90.9	-	-	28	3.9	2.9	24.8	15.6	14.0
816	Feather meal.	141.7	86.2	-	-	14	2.2	2.4	24.6	16.9	15.4
360	Fish meal.	96.9	72.8	5	-	246	-	-	18.3	13.4	-
389	Fish meal.	117.0	25.0	-	-	229	-	-	18.3	13.3	-
398	Fish meal.	102.6	-	-	-	-	-	-	17.2	12.3	11.5
662	Fish meal.	107.0	96.3	-	-	30	198	61.8	35.8	19.9	14.3
712	Fish meal.	106.7	72.2	-	-	248	77.7	60.4	18.8	14.0	12.2
713	Fish meal.	110.5	45.9	-	-	-	235	71.6	59.6	18.5	14.3

TABLE 1 cont.

Code	Description of Feedingstuff	Nitrogen	Ether Extract	Crude Fibre	Neutral Detergent Fibre	Ash	Calcium	Total Phosphorus	Gross Energy	TME	TME _n
734	Fish meal.	117.6	119.4	-	-	130	34.6	22.8	22.3	17.0	15.6
735	Fish meal.	112.0	74.3	-	-	206	59.6	30.9	19.8	14.2	12.3
736	Fish meal.	108.5	56.5	-	-	241	74.6	58.6	18.9	14.0	12.3
788	Fish meal.	106.5	44.4	-	-	263	90.4	45.5	17.8	12.7	12.1
789	Fish meal.	109.0	106.7	-	-	203	65.8	38.4	20.7	16.6	15.0
814	Fish meal.	108.8	109.4	-	-	91	58.2	35.0	21.0	15.6	14.4
836	Fish meal.	117.6	50.6	-	23	171	54.1	31.9	20.4	14.0	12.8
527	Fish meal. 60% protein.	105.1	56.3	10	-	257	-	-	17.9	12.3	11.0
528	Fish meal. 66% protein.	100.3	20.3	17	-	319	-	-	15.8	10.4	9.2
441	Fish meal. Herring.	108.0	105.1	10	-	184	-	-	21.2	15.8	14.3
353	Fish meal. Menhaden.	107.2	79.2	4	-	198	-	-	20.0	13.8	-
354	Fish meal. Menhaden.	103.8	110.0	5	-	214	-	-	19.8	14.5	-
620	Fish meal. Menhaden.	111.0	119.0	9	-	210	58.2	35.4	20.6	16.1	14.1
621	Fish meal. Menhaden.	111.6	106.6	9	-	211	59.5	35.0	20.2	15.3	13.6
622	Fish meal. Menhaden.	117.0	138.2	6	-	190	52.9	31.1	21.2	16.5	14.6
623	Fish meal. Menhaden.	107.8	122.6	7	-	214	53.8	36.4	20.7	16.1	14.4
624	Fish meal. Menhaden.	112.6	111.6	7	-	209	58.5	32.0	20.6	15.7	13.9
625	Fish meal. Menhaden.	107.8	109.0	6	-	214	58.9	32.7	20.6	15.9	14.1
626	Fish meal. Menhaden.	115.8	109.2	5	-	198	48.1	30.4	20.2	15.5	13.4
627	Fish meal. Menhaden.	119.0	113.4	6	-	187	45.0	29.2	20.5	15.7	13.8
628	Fish meal. Menhaden.	117.4	113.1	6	-	189	47.5	30.6	20.3	15.6	13.6
629	Fish meal. Menhaden.	113.0	99.1	7	-	197	59.7	35.0	20.5	15.9	14.1
630	Fish meal. Menhaden.	109.8	106.4	7	-	210	53.6	30.3	20.3	15.5	13.7
631	Fish meal. Menhaden.	112.0	91.6	10	-	215	58.8	34.9	20.0	15.0	13.2
632	Fish meal. Menhaden.	106.3	166.1	9	-	163	38.7	25.5	21.6	16.9	15.0
633	Fish meal. Menhaden.	113.8	136.5	11	-	172	36.6	24.7	21.0	16.5	14.5
634	Fish meal. Menhaden.	110.7	113.9	10	-	176	31.3	23.1	20.5	15.6	13.9
635	Fish protein concentrate.	150.3	1.4	6	-	91	31.2	16.1	21.8	17.6	14.4
636	Fish protein concentrate.	143.0	1.1	6	-	99	34.0	16.8	21.5	17.5	14.4
581	Fish solubles.	102.6	123.0	ND	-	162	-	-	19.7	14.8	13.4
597	Gelatin. purified protein. Nutritional Biochemicals.	166.2	-	-	8	-	-	-	21.2	16.1	14.4
328	Glucose. monohydrate.	-	-	-	-	-	-	-	16.0	15.7	-
MS	Glucose. monohydrate.	-	-	-	-	-	-	-	15.4	15.9	-
605	Glucose. D-dextrose, anhydrous.	-	-	-	-	-	-	-	15.6	15.6	15.5

TABLE 1 cont.

Code	Description of Feedingstuff	Nitrogen	Ether Extract	Crude Fibre	Neutral Detergent Fibre	Ash	Calcium	Total Phosphorus	Gross Energy	TME	TME _n
558	Lathyrus cicera, ground.	42.2	6.3	62	-	64	-	-	11.7	11.7	11.3
557	Lathyrus sativus, ground.	45.4	6.0	77	-	34	-	-	18.4	11.8	11.3
560	Lentils, ground, (cv. Eston).	41.8	8.4	55	-	35	-	-	18.5	12.8	12.2
559	Lentils, ground, (cv. Laird).	39.7	7.4	54	-	93	-	-	17.3	12.3	11.8
499	Linseed meal.	58.4	66.7	114	-	60	-	-	20.4	9.2	8.6
506	Linseed meal.	59.9	43.6	128	-	79	-	-	19.8	9.1	8.0
512	Linseed meal.	64.8	28.0	124	-	63	-	-	19.6	8.4	7.9
432	Lupin seed, ground, New Zealand.	49.3	57.2	171	-	37	-	-	19.7	9.4	8.9
362	Meat meal.	84.4	98.6	43	-	259	-	-	17.7	11.4	-
397	Meat meal.	86.4	126.0	-	-	248	-	-	19.5	13.1	-
444	Meat meal.	74.4	118.2	29	-	342	-	-	16.4	10.7	9.7
479	Meat meal.	89.5	99.6	28	-	268	-	-	17.8	12.8	11.2
480	Meat meal.	85.5	107.3	28	-	255	-	-	18.6	12.8	11.6
481	Meat meal.	85.5	128.8	30	-	253	-	-	19.0	12.5	11.7
482	Meat meal.	84.5	130.1	26	-	230	-	-	19.1	12.9	11.6
658	Meat meal.	83.0	133.2	-	159	254	89.8	50.8	19.1	13.1	12.1
659	Meat meal.	84.4	95.4	-	78	340	128.2	76.2	16.1	7.6	7.2
660	Meat meal.	80.5	108.1	-	39	328	108.5	69.7	16.9	11.2	10.2
661	Meat meal.	82.4	89.9	-	152	334	102.6	67.0	16.4	11.4	10.3
709	Meat meal.	79.0	163.1	-	-	279	95.6	58.8	19.2	12.5	11.3
710	Meat meal.	84.8	96.5	-	-	283	104.0	66.3	17.6	12.0	10.5
711	Meat meal.	81.1	113.5	-	-	332	110.9	68.4	17.0	12.1	10.7
731	Meat meal.	84.4	115.0	-	-	280	91.4	59.2	17.9	12.4	11.0
732	Meat meal.	84.0	112.0	-	-	292	100.9	60.5	17.3	11.4	10.2
733	Meat meal.	85.9	98.5	-	-	318	107.6	66.6	17.2	12.2	10.8
785	Meat meal.	82.3	128.8	-	-	281	91.1	53.7	18.5	12.3	11.3
786	Meat meal.	87.9	111.7	-	-	249	73.4	50.2	19.1	13.6	12.4
787	Meat meal.	79.2	96.9	-	-	322	74.4	65.1	16.3	10.9	9.9
812	Meat meal.	86.3	112.8	-	-	284	90.0	43.6	18.0	12.3	11.3
813	Meat meal.	82.5	90.6	-	-	327	107.3	55.0	16.5	11.2	10.0
598	Milk, skim, spray-dried powder.	56.2	-	-	-	84	-	-	18.0	11.9	11.2
570	Millet, ground, Indiana.	19.1	38.3	98	-	33	-	-	18.8	15.5	14.6

TABLE 1. cont.

Code	Description of Feedingstuff	Nitrogen			Ether Extract			Crude Fibre			Neutral Detergent Fibre			Total Phos-phorus			Gross Energy	TME	TME _n
		18.9	32.4	29	-	16	-	111	-	-	-	-	-	18.6	16.7	16.1			
580	Milo. ground.	7.7	4.4	127	-	111	-	-	-	-	-	-	-	16.2	6.8	6.9			
442	Molasses. dried.																		
382	Oats. whole ground.	25.1	42.0	95	-	30	-	-	-	-	-	-	-	18.5	14.3	-			
388	Oats. whole ground.	21.0	41.0	105	-	30	-	-	-	-	-	-	-	19.3	14.3	-			
420	Oats. whole ground.	-	-	-	-	-	-	-	-	-	-	-	-	19.8	13.8	-			
421	Oats. whole ground.	-	-	-	-	-	-	-	-	-	-	-	-	19.8	14.7	-			
437	Oats. whole ground.	16.1	47.6	145	-	29	-	-	-	-	-	-	-	19.8	12.8	12.9			
533	Oats. whole ground.	18.9	53.0	117	-	28	-	-	-	-	-	-	-	19.5	13.3	12.9			
699	Oats. whole ground.	18.4	43.8	-	334	28	-	-	-	-	-	-	-	19.3	11.9	11.2			
700	Oats. whole ground.	18.2	45.6	-	374	34	-	-	-	-	-	-	-	19.3	11.9	11.2			
701	Oats. whole ground.	17.9	46.4	-	349	29	-	-	-	-	-	-	-	19.2	11.5	10.9			
702	Oats. whole ground.	19.1	46.2	-	279	37	-	-	-	-	-	-	-	19.1	12.9	12.2			
703	Oats. whole ground.	22.9	41.0	-	305	31	-	-	-	-	-	-	-	19.3	12.8	11.9			
595	Oats. whole ground, (cv. Donald).	18.5	34.0	146	-	26	-	-	-	-	-	-	-	19.2	13.5	13.0			
365	Oats. whole ground, (cv. Harmon).	19.7	52.2	108	-	33	-	-	-	-	-	-	-	19.2	13.1	-			
364	Oats. whole ground, (cv. Hinoat).	26.4	39.8	119	-	30	-	-	-	-	-	-	-	19.3	13.0	-			
594	Oats. whole ground, (cv. Woodstock).	23.4	38.8	145	-	36	-	-	-	-	-	-	-	19.3	13.4	12.7			
764	Oats. whole ground, No. 2 CW, green, 50 kg/hL.	20.7	53.7	-	281	32	-	-	-	-	-	-	-	19.7	13.4	12.9			
774	Oats. whole ground, No. 2 CW, green, 50 kg/hL.	19.6	45.8	-	296	36	-	-	-	-	-	-	-	19.2	13.1	12.6			
770	Oats. whole ground, No. 2 CW, green, 53 kg/hL.	16.2	64.6	-	287	31	-	-	-	-	-	-	-	19.6	13.5	13.3			
760	Oats. whole ground, No. 1 feed, green, 51 kg/hL.	17.1	61.1	-	296	35	-	-	-	-	-	-	-	19.5	13.4	12.7			
759	Oats. whole ground, No. 2 feed, 44 kg/hL.	18.9	49.5	-	365	28	-	-	-	-	-	-	-	19.4	12.3	11.9			
771	Oats. whl. grd., No. 2 feed, 18% wheat, 50 kg/hL.	21.9	42.6	-	310	30	-	-	-	-	-	-	-	19.4	13.3	12.8			
422	Oats. hull-less, ground.	28.3	53.7	23	-	25	-	-	-	-	-	-	-	19.9	16.9	16.1			
423	Oats. hull-less, ground.	31.2	50.1	23	-	25	-	-	-	-	-	-	-	19.8	16.5	15.9			
424	Oats. hull-less, ground.	26.2	57.0	20	-	23	-	-	-	-	-	-	-	19.9	17.2	16.4			
426	Oats. hull-less, ground.	26.2	55.4	23	-	21	-	-	-	-	-	-	-	19.8	17.0	16.4			
590	Oats. hull-less, ground.	30.9	59.9	29	-	23	-	-	-	-	-	-	-	19.8	16.6	16.0			
591	Oats. hull-less, ground.	29.8	55.1	27	-	23	-	-	-	-	-	-	-	19.8	16.5	15.9			
592	Oats. hull-less, ground.	30.0	54.0	27	-	23	-	-	-	-	-	-	-	19.7	16.8	16.2			
593	Oats. hull-less, ground.	30.6	49.9	29	-	24	-	-	-	-	-	-	-	19.7	16.6	16.0			
842	Oats. hull-less, ground.	31.8	63.7	-	-	-	-	-	-	-	-	-	-	19.8	17.6	16.8			
589	Oats. hull-less, ground, (cv. Terra).	29.1	48.2	28	-	24	-	-	-	-	-	-	-	19.5	16.3	15.8			
385	Oat groats. ground, Eastern Canada.	24.0	22.8	34	-	44	-	-	-	-	-	-	-	19.8	16.9	-			
384	Oat grouts. ground, Western Canada.	21.2	22.4	37	-	66	-	-	-	-	-	-	-	19.8	17.7	-			

TABLE 1. cont.

Code	Description of Feedingstuff	Nitrogen	Ether Extract	Crude Fibre	Neutral Detergent Fibre	Ash	Calcium	Total Phosphorus	Gross Energy	TME	TME _n
425	Oats, wild, dehulled, ground.	29.0	85.3	35	-	23	-	-	21.1	17.9	16.9
573	Peanut hulls. Ground.	13.9	18.5	539	-	91	-	-	18.8	4.6	4.5
352	Peanut hulls. Ground, skins.	24.6	164.0	116	-	26	-	-	22.0	13.7	-
575	Peanut meal.	86.0	6.8	74	-	50	-	-	19.6	12.2	11.2
517	Peas. whole ground.	38.8	10.5	74	-	31	-	-	18.3	12.7	12.1
553	Peas. whole ground, (cv. Century).	43.1	9.5	65	-	27	-	-	18.5	12.7	12.2
552	Peas. whole ground, (cv. Tara).	36.1	9.5	78	-	31	-	-	18.3	12.8	12.5
551	Peas. whole ground, (cv. Trapper).	41.7	10.8	73	-	30	-	-	18.6	12.7	12.1
550	Peas. whole ground, (cv. Triumph).	39.9	9.5	63	-	29	-	-	18.4	13.2	12.8
386	Potato waste. dried, ground.	12.1	70.2	40	-	40	-	-	18.8	15.7	-
494	Potato waste. dried, ground.	13.9	74.1	44	-	30	-	-	19.4	15.5	15.1
502	Poultry by-product meal.	104.7	163.6	17	-	135	-	-	22.9	16.9	15.5
518	Poultry by-product meal.	106.5	128.2	27	-	133	-	-	22.2	15.6	14.3
525	Rapeseed. small seed, screenings ⁴ .	31.7	142.7	178	-	52	-	-	22.5	14.6	13.8
524	Rapeseed. whole ground, (cv. Candle).	35.6	270.7	244	-	42	-	-	28.0	15.2	14.6
303	Rapeseed. whole ground, (cv. Tower).	-	-	-	-	-	-	-	27.9	19.7	-
523	Rapeseed. whole ground, (cv. Tower).	36.0	319.7	222	-	39	-	-	28.6	17.0	16.3
526	Rapeseed dockage ⁵ .	24.7	91.2	204	-	72	-	-	19.9	9.5	9.0
395	Rapeseed meal.	63.0	40.1	106	-	79	-	-	19.8	10.2	-
443	Rapeseed meal.	58.8	23.9	127	-	74	-	-	19.9	9.1	8.3
485	Rapeseed meal.	55.9	101.8	111	-	71	-	-	21.1	11.2	10.6
486	Rapeseed meal.	67.8	27.0	121	-	57	-	-	20.0	9.9	9.2
487	Rapeseed meal.	64.6	28.1	125	-	71	-	-	20.0	9.4	9.0
488	Rapeseed meal.	65.2	21.4	113	-	77	-	-	19.8	9.1	8.5
796	Rapeseed meal.	62.1	20.9	-	-	358	78	7.5	12.7	20.1	10.4
797	Rapeseed meal.	61.1	31.3	-	-	342	79	7.4	12.4	19.9	10.0
798	Rapeseed meal.	59.4	31.0	-	-	388	75	7.9	12.3	20.4	10.7
818	Rapeseed meal.	65.0	18.5	-	-	76	7.1	13.6	19.5	9.6	9.0
819	Rapeseed meal.	60.6	34.8	-	-	305	80	7.8	13.0	19.6	8.8

TABLE 1 cont.

Code	Description of Feedingstuff	Nitrogen	Ether Extract	Crude Fibre	Neutral Detergent Fibre	Ash	Calcium	Phos-phorus	Gross Energy	TME	TME _n
820	Rapeseed meal.	66.8	27.1	-	301	74	6.4	13.1	20.1	10.7	10.0
821	Rapeseed meal.	61.5	32.6	-	308	79	8.0	13.1	19.6	9.5	8.6
822	Rapeseed meal. (cv. Tower).	59.5	18.7	-	302	82	8.2	12.8	19.4	10.1	9.2
322	Rapeseed meal. Canola.	-	-	-	-	-	-	-	19.9	12.8	-
522	Rapeseed meal. Canola.	67.6	23.2	133	-	71	-	-	19.8	9.6	8.9
668	Rapeseed meal. Canola.	64.3	16.4	-	253	74	7.0	14.8	19.7	9.6	8.6
669	Rapeseed meal. Canola.	61.4	45.8	-	257	70	6.9	13.7	20.1	9.5	8.6
670	Rapeseed meal. Canola.	63.5	31.8	-	257	73	6.9	14.8	19.7	9.6	8.8
720	Rapeseed meal. Canola.	67.3	25.2	-	254	72	6.5	13.9	19.8	9.9	8.7
721	Rapeseed meal. Canola.	64.8	10.6	-	241	76	6.9	13.6	19.3	10.0	8.7
722	Rapeseed meal. Canola.	66.8	23.3	-	235	76	7.6	13.6	19.8	10.6	9.0
743	Rapeseed meal. Canola.	63.8	22.2	-	241	72	6.8	12.9	19.7	10.3	9.2
744	Rapeseed meal. Canola.	71.6	27.6	-	221	72	6.1	13.2	19.8	10.4	9.2
745	Rapeseed meal. Canola.	61.9	33.9	-	234	75	8.2	13.4	19.6	10.1	8.9
746	Rapeseed meal. Canola.	68.8	37.7	-	232	72	6.0	13.3	20.1	10.8	9.5
747	Rapeseed meal. Canola.	71.3	21.4	-	220	72	5.8	13.2	19.8	10.4	9.2
579	Rice bran.	23.2	161.8	122	-	104	-	-	20.9	14.2	13.6
641	Rice bran.	21.6	110.8	-	175	91	3.5	18.8	19.6	14.7	14.0
642	Rice bran.	23.3	167.0	-	275	107	1.3	23.7	22.6	14.0	14.0
647	Rice bran.	25.1	116.8	-	186	68	.6	13.5	20.4	15.0	14.6
648	Rice bran. infested with weevils.	25.1	140.0	-	222	118	.9	28.2	20.4	12.3	12.0
644	Rice bran. parboiled.	27.0	319.3	-	336	112	1.6	27.0	24.5	14.6	14.1
571	Rice bran. polish.	23.2	132.6	52	-	80	-	-	20.2	15.1	14.8
587	Rye. whole ground. (cv. Carmen).	20.2	10.3	28	-	16	-	-	18.4	14.5	14.0
554	Rye. whole ground. (cv. Cougar).	24.2	14.9	34	-	18	-	-	18.4	15.0	14.4
585	Rye. whole ground. (cv. Gazeile).	17.5	11.6	31	-	17	-	-	18.3	14.4	13.9
390	Rye. whole ground. (cv. Kodiak).	19.5	13.0	26	-	16	-	-	18.3	15.3	-
582	Rye. whole ground. (cv. Musketeer).	19.6	11.3	32	-	19	-	-	18.3	14.1	13.5
584	Rye. whole ground. (cv. Puma).	18.8	10.5	30	-	17	-	-	18.2	14.1	13.9
555	Rye. whole ground. (cv. Puma).	21.7	12.1	25	-	17	-	-	18.3	14.4	13.9
586	Rye. whole ground. (cv. Puma).	19.1	11.6	31	-	19	-	-	18.3	13.6	13.9
31.7	Screenings.	149.7	-	421	67	5.8	8.2	22.4	13.2	12.4	
25.5	Screenings.	140.6	-	401	67	5.2	6.6	21.2	11.2	11.2	
24.3	Screenings.	33.7	-	152	23	.8	6.1	18.6	15.1	14.2	
18.3	Screenings.	37.4	-	520	118	2.8	3.4	17.6	6.9	6.4	

TABLE 1 cont.

Code	Description of Feedingstuff	Nitrogen	Ether Extract	Crude Fibre	Neutral Detergent Fibre	Ash	Calcium	Total Phosphorus	Gross Energy	TME	TME _n
708	Screenings.	21.5	51.1	-	390	45	1.1	3.5	19.5	11.9	10.9
569	Screenings. feed.	17.9	16.0	148	-	63	-	-	18.1	8.9	8.6
568	Screenings. No. 1 feed.	24.6	30.9	50	-	22	-	-	19.0	14.5	13.7
576	Sesame seed meal.	75.0	47.1	78	-	129	-	-	18.7	10.7	9.5
639	Sesame seed meal.	75.3	12.7	-	198	161	29.5	16.8	17.1	8.6	8.2
645	Sesame seed meal.	73.7	30.8	-	200	174	31.4	16.5	17.3	9.6	9.0
678	Shrimp meal.	64.5	4.2	-	273	389	138.8	28.8	13.0	5.8	5.1
840	Single cell protein. Pruteen.	121.2	16.6	-	2	109	12.5	29.6	21.2	15.2	13.7
447	Soybean, whole seeds.	62.9	207.5	59	-	50	-	-	23.8	12.8	12.1
448	Soybean, whole seeds, rolled.	64.5	205.9	61	-	50	-	-	23.5	14.8	13.9
383	Soybean flakes. solvent extracted, raw.	86.5	10.6	44	-	64	-	-	19.8	9.9	-
446	Soybean hulls. ground.	26.6	25.0	357	-	48	-	-	18.2	4.6	4.2
327	Soybean meal.	-	-	-	-	-	-	-	-	19.4	13.4
368	Soybean meal.	-	-	43	-	67	-	-	-	19.5	13.0
393	Soybean meal.	91.0	6.0	-	-	-	-	-	-	19.8	12.8
399	Soybean meal.	88.5	-	-	-	-	-	-	-	19.9	12.8
489	Soybean meal.	78.9	61.3	35	-	65	-	-	-	20.7	13.6
490	Soybean meal.	78.7	14.8	55	-	65	-	-	-	19.7	11.4
491	Soybean meal.	81.3	39.0	43	-	66	-	-	-	20.0	12.2
492	Soybean meal.	82.8	6.6	40	-	67	-	-	-	19.7	11.2
608	Soybean meal.	75.0	12.6	94	-	64	-	-	-	19.5	11.2
671	Soybean meal.	86.7	14.3	-	53	64	-	-	-	19.7	12.0
672	Soybean meal.	85.9	18.6	-	61	68	3.2	9.0	-	19.7	11.4
673	Soybean meal.	84.7	17.1	-	82	66	2.6	9.0	-	19.8	12.8
723	Soybean meal.	87.0	11.1	-	73	64	2.1	8.6	-	19.8	12.4
724	Soybean meal.	85.1	11.7	-	64	65	2.4	9.3	-	19.5	13.3
748	Soybean meal.	87.0	11.7	-	56	64	2.1	8.8	-	19.7	12.3
749	Soybean meal.	85.8	7.6	-	68	69	3.0	8.6	-	19.5	12.3
750	Soybean meal.	88.1	8.8	-	63	65	2.5	9.3	-	19.8	13.1
751	Soybean meal.	85.7	17.8	-	46	73	2.6	8.6	-	19.8	12.7
799	Soybean meal.	86.0	10.5	-	108	66	3.0	8.0	-	19.8	13.0

TABLE 1 cont.

Code	Description of Feedingstuff	Nitrogen		Ether Extract		Crude Fibre		Neutral Detergent Fibre		Gross Phosphorus		Energy	TME	TME _n
		Total	Nitrogen	Ether Extract	Crude Fibre	Neutral Detergent Fibre	Ash	Calcium	Phosphorus	Total	Gross Energy			
800	Soybean meal.	89.7	59.2	-	132	67	2.5	9.1	19.8	12.6	11.4			
801	Soybean meal.	86.9	10.0	-	110	66	2.5	8.8	19.7	12.4	11.2			
802	Soybean meal.	87.7	10.0	-	161	69	3.9	7.6	19.7	12.6	11.7			
803	Soybean meal.	88.5	6.8	-	119	66	2.7	8.8	19.6	12.5	11.4			
804	Soybean meal.	88.2	9.4	-	154	68	2.5	9.1	19.8	12.8	11.8			
823	Soybean meal.	86.9	18.1	-	112	68	3.2	8.2	19.7	12.4	11.5			
824	Soybean meal.	91.0	3.8	-	101	67	2.6	9.5	19.7	13.5	12.3			
825	Soybean meal.	90.0	8.4	-	112	65	2.6	9.6	19.9	13.2	12.0			
826	Soybean meal.	88.5	7.3	-	114	63	2.5	9.4	19.9	12.5	11.4			
837	Soybean meal.	87.3	18.0	-	69	65	2.8	9.8	19.6	12.1	11.2			
505	Sunflower seed meal.	66.4	24.0	189	-	79	-	-	-	19.4	9.4	8.9		
516	Sunflower seed meal.	72.7	23.7	153	-	81	-	-	-	19.4	10.5	9.7		
566	Sunflower seed meal.	78.2	14.9	128	-	88	-	-	-	19.3	10.4	9.2		
MS	Sunflower seed meal.	74.8	16.0	108	-	79	-	-	-	19.7	10.8	-		
513	Triticale. whole ground.	27.7	19.9	37	-	21	-	-	-	18.6	15.2	14.4		
514	Triticale. whole ground.	27.2	17.0	41	-	20	-	-	-	18.6	15.2	14.6		
515	Triticale. whole ground.	24.9	9.6	31	-	16	-	-	-	18.4	15.6	15.0		
583	Triticale. whole ground.	26.8	15.5	38	-	18	-	-	-	18.2	14.6	14.1		
427	Triticale. whole ground. (cv. Palouse).	-	-	-	-	-	-	-	-	18.2	14.7	-		
588	Triticale. whole ground. (cv. Welsh).	26.4	11.1	39	-	20	-	-	-	18.4	14.9	14.3		
361	Wheat. whole ground.	23.5	27.7	39	-	23	-	-	-	18.5	15.6	-		
381	Wheat. whole ground.	25.7	17.5	30	-	20	-	-	-	18.0	15.8	-		
400	Wheat. whole ground.	25.8	-	-	-	-	-	-	-	18.6	15.0	14.6		
471	Wheat. whole ground.	27.1	19.0	26	-	18	-	-	-	18.7	16.5	15.6		
472	Wheat. whole ground.	24.9	15.8	17	-	25	-	-	-	18.7	16.2	15.2		
679	Wheat. whole ground.	30.0	16.3	-	117	21	.4	-	7.2	18.7	15.6	15.0		
680	Wheat. whole ground.	29.8	21.8	-	72	21	.3	7.2	18.7	15.3	14.8			
681	Wheat. whole ground.	30.5	16.4	-	114	18	.3	6.1	18.7	15.7	15.0			
682	Wheat. whole ground.	27.3	15.1	-	112	20	.4	6.6	18.6	15.1	14.5			
683	Wheat. whole ground.	24.0	16.6	-	125	19	.3	6.1	18.4	15.3	14.7			
684	Wheat. whole ground.	28.9	20.4	-	94	22	.4	6.9	18.6	14.9	14.5			
685	Wheat. whole ground.	30.4	15.8	-	132	21	.4	6.4	18.7	15.3	14.7			
686	Wheat. whole ground.	27.5	22.4	-	94	21	.5	6.8	18.6	15.3	14.8			
687	Wheat. whole ground.	29.0	15.4	-	127	22	.4	7.0	18.5	15.6	15.0			
688	Wheat. whole ground.	30.9	14.2	-	139	20	.3	7.0	18.6	15.1	14.7			

TABLE 1 cont.

Code	Description of Feedingstuff	Nitrogen	Ether Extract	Crude Fibre	Neutral Detergent Fibre	Ash	Calcium	Total Phosphorus	Gross Energy	TME	TME _n
366	Wheat, whole ground, (cv. Glenlea).	28.4	16.0	26	-	20	-	-	18.4	15.8	-
391	Wheat, whole ground, (cv. Glenlea).	26.2	15.0	25	-	18	-	-	18.6	16.4	-
418	Wheat, whole ground, Australian, (cv. Cootamundra).	-	-	-	-	-	-	-	18.5	15.8	-
417	Wheat, whole ground, Australian, (cv. Dubba).	-	-	-	-	-	-	-	18.6	15.7	-
419	Wheat, whole ground, Australian, (cv. Gunnedah).	-	-	-	-	-	-	-	18.4	15.3	-
416	Wheat, whole ground, Australian, (cv. Wagga).	-	-	-	-	-	-	-	18.4	15.5	-
428	Wheat, whole ground, purple.	-	-	-	-	-	-	-	18.5	15.3	-
773	Wheat, whole ground, No. 1 CW, Red Spring.	31.7	15.6	-	134	22	.5	6.1	18.6	15.0	14.4
778	Wheat, whole ground, No. 1 CW, Red Spring.	32.0	16.3	-	144	20	.3	6.5	18.8	15.7	14.9
784	Wheat, whole ground, No. 1 CW, Red Spring.	27.6	16.2	-	103	19	.6	5.3	18.6	15.7	15.0
779	Wheat, whl. grd., No. 1 CW, Red Spring, 78 kg/hL.	28.9	15.8	-	128	20	.4	5.8	18.7	15.6	14.8
782	Wheat, whl. grd., No. 1 CW, Red Spring, 80 kg/hL.	28.6	15.5	-	132	20	.6	6.3	18.6	15.5	14.9
783	Wheat, whl. grd., No. 2 CW, Red Spring, 77 kg/hL.	28.8	15.5	-	133	21	.5	6.5	18.7	16.1	15.4
772	Wheat, whl. grd., No. 2 CW, R. S., 37% vitr. kern.	26.2	15.4	-	138	21	.6	5.8	18.6	15.3	14.8
781	Wheat, whl. grd., No. 2 CW, R. S., 4% midge damage.	27.9	15.0	-	119	21	.6	5.8	18.6	15.5	14.7
776	Wheat, whl. grd., No. 2 CW, R. S., immat. shrunken.	37.3	14.3	-	142	23	.6	6.4	18.9	15.3	14.7
780	Wheat, whl. grd., No. 3 CW, R. S., 20% vitr. kern.	31.2	15.1	-	82	22	.5	6.0	18.7	15.9	15.1
438	Wheat bran.	27.2	37.7	122	-	63	-	-	19.0	9.0	8.7
542	Wheat bran. Ontario wheat.	25.2	27.7	131	-	66	-	-	18.9	8.1	7.5
520	Wheat middlings.	26.7	39.3	95	-	27	-	-	19.1	14.0	14.2
539	Wheat middlings. Ontario wheat.	25.7	42.4	37	-	26	-	-	18.9	15.0	13.5
439	Wheat shorts.	24.9	32.1	86	-	52	-	-	19.0	11.7	11.2
509	Wheat shorts.	30.6	41.4	56	-	35	-	-	18.9	10.8	9.4
511	Wheat shorts.	30.6	44.1	112	-	51	-	-	19.3	10.6	9.0
676	Wheat shorts.	30.0	12.4	-	347	52	1.0	13.1	19.2	10.5	9.7
677	Wheat shorts.	31.5	34.2	-	310	54	1.1	14.0	19.6	10.8	9.8
728	Wheat shorts.	30.8	41.8	-	345	53	.8	11.4	19.2	10.6	9.9
729	Wheat shorts.	32.1	48.5	-	324	57	.9	12.7	19.3	10.5	9.6
730	Wheat shorts.	31.5	44.5	-	292	55	1.0	12.2	19.2	10.6	9.7
756	Wheat shorts.	30.2	39.3	-	292	49	.8	11.2	19.2	10.8	10.0
757	Wheat shorts.	29.9	41.6	-	281	45	.9	10.2	19.2	11.2	10.6
758	Wheat shorts.	29.9	45.9	-	340	56	.8	12.2	19.1	9.9	8.9
809	Wheat shorts.	31.5	34.8	-	332	56	.8	11.9	19.3	9.8	9.4
810	Wheat shorts.	30.5	31.7	-	336	54	.8	12.6	19.0	10.2	9.4
811	Wheat shorts.	31.6	42.1	-	321	54	.9	12.5	19.4	10.7	10.3

TABLE 1 cont.

Code	Description of Feedingstuff	Nitrogen	Ether Extract	Crude Fibre	Neutral Detergent Fibre			Ash	Calcium	Phos- phorus	Gross Energy	TME	TME _n
					Total	Detergent	Fibre						
832	Wheat shorts.	31.4	35.9	-	270	51	.8	11.4	19.3	10.8	9.9		
833	Wheat shorts.	31.4	33.1	-	315	52	.8	11.2	19.2	11.3	10.2		
834	Wheat shorts.	31.1	40.5	-	340	55	.8	11.5	19.4	10.7	9.8		
835	Wheat shorts.	31.6	39.8	-	326	56	.9	12.6	19.5	10.5	9.6		
508	Whey powder.	19.6	2.2	4	-	89	-	-	15.4	5.8	2.7		
535	Whey powder.	20.3	2.5	3	-	94	-	-	15.5	5.9	3.1		
602	Yeast. dried.	82.0	1.5	80	-	78	-	-	21.2	13.5	13.4		
503	Yeast. dried, brewer's.	72.0	4.0	55	-	104	-	-	19.5	11.2	10.4		
541	Yeast. dried, brewer's.	64.2	6.9	57	-	82	-	-	18.9	12.8	12.0		
574	Yeast. dried, brewer's.	63.2	30.3	65	-	84	-	-	19.4	12.8	11.6		
572	Yucca. dried.	6.0	1.2	175	-	48	-	-	16.9	10.9	10.5		
638	Yucca. dried.	3.2	4.4	-	260	19	5.6	.5	17.3	9.6	10.1		
640	Yucca. dried.	2.3	4.1	-	57	32	1.1	.8	16.5	15.6	15.2		

1 Blanks (-) = variable not measured; ND = variable not detected in a reportable amount.

2 1 MJ = .239 Mcal.

3 Space limitations make the following abbreviations necessary:

c.v. = cultivar

immat. = immature

inf. var. = inferior variety or varieties

R. S. = Red Spring

vitr. kern. = vitreous kernels

whl. grd. = whole ground

4 By microscopic analysis found to contain: lambsquarter and red root pig weeds 53%; brassica sp. 37%; stink weed 8%; miscellaneous 2%.

5 By microscopic analysis found to contain: empty seed pods, stems, chaff 32%; wild oats 27%; rapeseed 25%; wheat and barley 9%; lambsquarter, red root pig weeds with trace of stickseed and seeds of composite family 4%; wild buckwheat 1%; injurious weeds 1%; rodent excreta 1%.

TABLE 2

Total amino acids in the dry matter of several feedingstuffs (g/kg)¹

Code	Description	ALA	ARG	ASP	CYS	GLU	GLY	HIS	ILE	LEU	LYS	MET	PHE	PRO	SER	THR	TYR	VAL
655	Alfalfa.	7.4	6.4	21.8	2.4	13.9	6.6	3.2	5.3	10.1	7.6	2.5	6.8	13.1	7.4	6.7	4.2	7.3
656	Alfalfa.	9.4	9.4	25.6	-	17.9	8.3	3.8	6.9	13.5	9.3	-	8.8	13.3	8.9	8.2	5.5	9.0
653	Alfalfa.	7.6	6.6	24.5	2.3	14.6	7.0	3.3	5.6	10.5	7.9	2.0	7.2	12.2	7.8	6.9	4.3	7.6
839	Alfalfa.	10.5	7.4	21.3	4.3	14.2	7.6	3.7	5.8	11.2	8.8	2.3	7.2	8.2	7.6	6.9	4.7	7.2
657	Alfalfa.	8.7	7.5	23.2	2.2	15.2	7.6	4.0	6.2	11.7	8.5	2.4	7.8	11.7	8.1	7.4	4.7	8.2
652	Alfalfa.	8.7	7.7	23.3	2.3	15.6	7.6	4.0	6.2	11.7	8.6	2.6	7.7	11.4	8.1	7.5	4.6	8.0
654	Alfalfa.	9.0	7.8	23.4	4.4	15.5	7.9	4.0	6.1	12.0	8.9	2.6	7.8	9.7	8.2	7.6	4.9	8.1
649	Algae.	34.7	24.0	40.4	12.2	39.1	22.9	6.0	14.5	33.3	21.5	8.8	19.6	20.3	17.7	21.7	13.9	22.4
650	Algae.	31.4	22.7	39.7	11.3	39.1	21.8	5.8	15.0	33.0	19.1	8.4	19.5	19.7	16.9	20.2	13.3	22.3
689	Barley.	5.4	7.2	8.7	3.3	34.9	5.5	3.4	4.3	9.4	5.7	2.3	6.8	14.6	6.3	4.7	4.3	5.7
690	Barley.	5.1	6.3	9.0	4.1	31.3	5.2	3.0	3.9	8.9	5.0	2.0	6.3	13.1	5.4	4.3	3.6	4.8
691	Barley.	3.8	5.1	7.6	2.5	20.4	4.0	2.3	2.8	6.4	3.9	1.9	4.1	8.2	4.0	3.3	2.4	3.2
692	Barley.	5.0	6.7	9.1	3.1	30.8	5.1	3.0	4.0	8.7	4.8	2.2	6.2	13.1	5.3	4.3	3.5	4.9
693	Barley.	4.5	5.6	8.5	2.9	25.6	4.6	2.6	3.3	7.6	4.4	1.7	4.9	10.6	4.6	3.8	3.0	4.2
694	Barley.	4.6	6.0	8.8	2.9	26.7	4.8	2.8	3.4	7.8	4.7	1.5	5.3	11.1	4.9	3.8	3.1	4.2
695	Barley.	5.1	7.3	9.5	3.8	33.7	5.4	3.2	4.3	9.2	5.1	2.5	6.7	14.6	5.8	4.6	3.9	4.8
696	Barley.	4.8	6.6	9.2	3.4	30.1	5.0	2.9	3.7	8.4	4.9	2.4	5.9	11.9	6.7	4.2	3.4	4.7
697	Barley.	4.7	6.6	9.3	3.0	29.5	5.0	3.0	3.8	8.6	4.8	3.6	6.0	12.4	5.2	4.3	3.3	4.7
698	Barley.	5.9	8.7	10.5	3.0	37.3	6.0	3.4	5.0	10.6	5.8	2.3	8.1	17.2	6.3	5.0	4.4	7.0
769	Barley.	5.6	7.4	8.9	3.4	35.1	5.7	3.4	4.5	10.0	5.4	2.2	7.6	15.8	6.2	4.8	4.4	6.3
765	Barley.	5.2	6.7	7.9	3.2	28.7	5.2	3.0	4.1	8.7	4.9	2.0	6.2	12.7	5.1	4.2	3.8	5.6
762	Barley.	5.8	7.3	8.9	3.3	34.3	5.7	3.3	4.6	9.9	5.3	2.2	7.2	15.5	6.3	4.9	4.4	6.4
775	Barley.	4.5	5.7	7.2	3.0	23.5	4.6	2.6	3.2	7.4	4.3	1.8	5.0	10.3	4.6	3.7	3.1	4.6
777	Barley.	5.7	7.4	8.8	4.0	41.4	5.6	3.5	4.9	10.9	5.2	2.7	8.4	19.5	6.6	4.8	4.9	6.8
767	Barley.	4.4	5.7	7.0	3.0	22.4	4.4	2.5	3.4	7.2	4.3	1.7	5.0	9.7	4.5	3.6	3.1	4.7
768	Barley.	4.7	5.7	7.3	3.1	26.2	4.5	2.6	3.6	7.8	4.3	1.8	5.5	11.5	4.9	3.8	3.2	5.1
761	Barley.	5.0	6.4	8.5	3.4	28.8	5.0	2.9	3.6	8.4	4.7	2.0	6.0	12.7	5.6	4.3	3.7	5.0
763	Barley.	4.9	6.1	7.8	2.9	27.1	4.9	2.8	3.8	8.2	4.6	1.9	5.8	11.9	5.2	4.1	3.5	5.3
766	Barley.	6.3	8.5	9.8	3.6	42.0	6.3	3.8	5.4	11.6	6.0	2.4	9.1	19.7	7.1	5.3	5.1	7.4
666	Blood meal.	73.3	52.8	111.8	18.9	100.2	49.4	55.8	17.6	120.7	79.9	8.5	63.9	50.3	57.9	42.2	31.1	73.8
667	Blood meal.	94.0	49.9	128.5	11.9	100.8	49.9	73.4	8.6	143.9	104.8	13.2	81.7	43.5	58.7	51.4	33.6	94.3
717	Blood meal.	79.6	40.5	105.3	10.6	87.1	40.8	58.1	6.8	120.9	89.9	14.0	70.4	35.9	53.7	48.4	30.0	70.8
718	Blood meal.	68.1	47.0	104.5	17.6	93.5	45.2	53.6	15.3	112.8	76.5	9.4	59.4	40.6	54.2	39.3	28.5	69.0
719	Blood meal.	72.4	42.9	101.1	13.6	87.2	40.5	56.3	11.8	112.3	82.8	12.6	62.2	35.7	49.0	43.2	29.2	65.3
739	Blood meal.	76.9	40.2	100.6	12.1	85.6	39.2	56.8	6.8	117.8	87.8	14.3	67.9	36.0	51.9	47.3	29.6	69.4

TABLE 2 cont.

Code	Description	ALA	ARG	ASP	CYS	GLU	GLY	HIS	ILE	LEU	LYS	MET	PHE	PRO	SER	THR	TYR	VAL
740	Blood meal.	69.7	45.7	101.2	17.0	90.6	43.5	55.5	14.7	111.9	77.4	9.3	58.9	39.6	50.4	38.0	28.2	68.7
741	Blood meal.	72.7	42.0	99.9	12.4	87.8	40.7	56.6	10.5	114.1	82.4	11.6	62.8	35.4	48.7	42.5	29.2	68.8
742	Blood meal.	76.4	41.8	103.9	11.8	85.4	41.9	61.7	8.1	119.1	86.7	11.4	65.9	35.4	47.8	41.4	27.7	71.1
792	Blood meal.	80.4	49.3	116.4	11.8	93.9	42.7	68.3	8.5	132.6	95.4	11.7	73.4	39.4	53.6	45.7	30.9	78.5
793	Blood meal.	71.4	53.7	109.0	17.3	97.6	43.8	59.5	15.8	120.8	82.1	9.5	64.3	43.2	56.1	41.3	30.4	72.5
794	Blood meal.	74.4	49.3	114.9	13.7	95.9	44.5	60.7	10.6	122.8	84.2	8.5	64.1	37.9	47.6	37.0	27.7	71.9
795	Blood meal.	78.1	46.2	108.8	12.4	91.8	38.8	61.1	7.4	126.1	94.4	14.4	73.8	37.6	56.0	50.7	31.5	73.6
817	Blood meal.	76.5	53.5	111.7	16.3	98.4	47.5	62.5	14.3	125.2	85.5	8.8	65.2	42.3	55.2	40.7	31.1	77.3
643	Coconut meal.	9.1	28.7	18.6	4.6	39.3	9.5	3.9	6.7	12.4	7.1	4.9	9.7	7.8	9.5	6.3	5.0	10.3
838	Corn.	7.6	5.2	7.7	2.5	18.1	3.9	2.9	3.0	12.2	3.2	2.3	4.7	9.3	4.8	3.6	4.0	4.4
853	Corn.	7.4	5.0	7.6	2.6	18.8	3.8	3.1	3.0	12.6	3.5	2.2	4.5	9.2	4.6	3.6	4.0	4.1
855	Corn.	7.6	5.6	7.6	3.0	19.5	3.8	3.2	3.3	13.4	3.3	2.4	4.9	9.9	4.8	3.7	4.0	4.6
856	Corn.	7.8	5.4	7.8	3.0	19.8	3.9	3.2	3.3	13.6	3.4	1.6	5.0	10.1	4.9	3.9	4.3	4.6
857	Corn.	8.9	6.2	8.3	2.9	22.3	4.3	3.5	3.8	15.2	3.7	2.6	5.6	11.3	5.5	4.2	4.7	5.2
858	Corn.	8.0	5.5	7.7	2.7	20.1	4.0	3.3	3.6	13.8	3.4	2.3	5.1	10.3	4.8	3.7	4.4	4.8
859	Corn.	8.7	5.7	7.9	2.9	21.2	4.0	3.3	3.7	14.5	3.3	2.6	5.4	10.7	5.0	4.0	4.5	5.1
860	Corn.	8.4	5.5	8.0	2.9	21.5	4.0	3.3	3.6	14.9	3.4	2.5	5.4	10.8	5.2	3.9	4.6	5.0
861	Corn.	8.2	5.7	7.7	2.8	20.9	4.0	3.3	3.5	14.1	3.4	2.5	5.3	10.0	5.3	4.0	4.5	4.9
862	Corn.	7.5	5.5	7.4	2.6	19.0	3.7	3.1	3.3	13.2	3.3	2.5	4.8	9.5	4.5	3.6	4.1	4.6
646	Corn germ meal.	13.9	16.3	17.3	4.4	32.2	14.0	7.4	7.4	18.1	10.1	4.1	10.1	13.0	11.2	8.7	7.0	12.5
651	Corn gluten feed.	22.2	11.2	17.6	5.9	47.5	9.7	6.9	8.3	33.2	7.0	5.3	13.3	24.9	13.5	9.6	10.6	11.8
674	Corn gluten feed.	15.3	11.3	15.8	4.9	35.2	9.9	7.1	6.3	21.7	7.9	3.5	8.6	18.4	10.2	8.3	6.7	10.1
726	Corn gluten feed.	14.1	11.6	14.8	5.4	32.2	9.8	6.7	6.4	19.7	7.6	3.8	8.4	17.1	9.9	7.7	6.4	9.9
727	Corn gluten feed.	14.9	12.3	15.4	5.5	34.2	10.4	6.9	6.7	20.5	7.7	4.2	8.9	17.5	10.5	8.4	6.7	10.5
754	Corn gluten feed.	17.0	13.5	17.5	6.4	39.4	11.3	7.6	6.8	23.6	8.1	4.7	9.7	21.1	12.0	9.2	8.1	11.1
755	Corn gluten feed.	16.8	12.9	17.4	6.4	38.7	11.0	7.6	6.8	23.3	8.1	4.7	9.5	21.3	11.6	9.1	7.7	11.1
805	Corn gluten feed.	17.6	14.2	18.5	6.3	42.0	11.4	8.2	7.5	25.5	8.8	4.8	9.9	22.2	11.7	9.3	8.2	11.8
806	Corn gluten feed.	16.5	13.6	17.9	6.2	39.7	11.1	7.9	7.2	23.7	8.7	4.6	9.5	20.6	11.4	9.2	7.6	11.3
827	Corn gluten feed.	15.0	12.8	16.4	5.6	34.8	10.1	7.3	6.8	21.4	8.0	4.4	8.7	18.8	10.1	8.1	7.2	10.5
828	Corn gluten feed.	15.5	10.7	15.2	5.9	36.8	9.0	7.2	5.9	22.5	7.2	4.1	8.1	21.3	10.3	8.2	7.3	9.1
675	Corn gluten meal.	54.7	19.0	38.2	10.1	135.5	15.8	11.6	21.7	102.5	9.7	13.1	38.1	58.4	33.4	20.3	31.2	24.5
725	Corn gluten meal.	60.5	21.3	43.1	14.1	151.7	18.0	13.7	23.5	114.6	11.7	18.5	41.9	63.4	37.6	22.7	35.6	26.4
752	Corn gluten meal.	53.3	17.4	38.1	12.4	135.3	15.1	11.6	18.7	101.1	9.8	16.8	36.1	54.6	33.4	19.6	30.7	22.6
753	Corn gluten meal.	52.1	19.5	40.1	13.1	121.7	16.3	10.5	19.7	73.2	10.1	18.2	39.7	60.1	34.2	21.9	33.4	24.2
807	Corn gluten meal.	66.2	23.2	45.6	12.9	152.2	17.8	13.9	25.1	115.1	11.5	16.9	46.1	69.2	40.0	24.2	38.2	30.0
808	Corn gluten meal.	69.3	24.5	47.0	12.9	162.4	18.5	14.7	27.3	123.7	12.3	17.7	48.6	70.0	40.2	24.8	32.3	32.3

TABLE 2 cont.

Code	Description	ALA	ARG	ASP	CYS	GLU	GLY	HIS	ILE	LEU	LYS	MET	PHE	PRO	SER	THR	TYR	VAL
829	Corn gluten meal.	67.4	24.7	45.8	13.4	158.1	19.1	14.9	25.3	122.0	11.9	15.5	46.4	70.7	39.0	24.5	39.4	31.4
830	Corn gluten meal.	67.3	26.8	47.4	14.4	150.8	21.7	15.2	25.9	115.9	13.3	21.0	46.8	72.0	40.6	24.5	40.1	31.0
831	Corn gluten meal.	65.0	26.1	46.0	13.4	154.7	20.2	14.9	25.8	114.8	12.9	20.5	45.9	68.6	39.1	23.7	39.2	30.7
663	Feather meal.	40.4	65.5	64.1	46.0	102.0	71.0	8.6	38.5	71.5	27.5	6.8	40.7	113.1	96.3	41.8	25.8	54.8
664	Feather meal.	44.0	69.0	65.8	44.1	104.0	74.5	7.4	42.3	77.0	23.6	5.7	46.1	137.6	111.1	43.6	28.1	65.1
665	Feather meal.	42.7	62.6	61.0	44.2	93.6	66.2	7.2	40.0	72.5	22.2	5.7	43.7	121.6	103.5	41.5	24.8	59.2
714	Feather meal.	39.9	62.3	64.1	45.6	110.3	63.3	11.0	34.8	72.1	29.1	7.3	39.9	80.1	97.5	41.3	26.1	53.2
715	Feather meal.	40.2	62.6	60.5	52.9	104.2	69.6	6.5	38.7	70.9	20.2	5.4	42.0	94.6	112.0	42.7	24.9	57.1
716	Feather meal.	40.1	61.6	60.1	45.2	100.4	66.8	7.8	38.9	72.0	23.4	5.9	41.7	91.8	103.8	41.0	24.1	58.4
737	Feather meal.	40.8	59.4	62.6	44.4	105.7	60.0	13.0	32.3	71.5	30.3	7.3	39.4	74.4	89.3	39.5	24.8	53.9
738	Feather meal.	39.8	62.3	59.5	61.8	102.1	67.3	7.1	37.6	70.2	20.6	6.0	41.6	90.2	104.6	40.9	25.0	57.7
790	Feather meal.	40.9	73.3	66.1	47.4	116.7	65.5	11.2	38.6	77.6	30.1	7.3	43.5	89.2	104.5	44.3	27.7	60.0
791	Feather meal.	40.3	72.9	64.1	52.9	110.4	66.7	7.6	40.2	76.2	24.1	6.6	44.8	96.9	110.8	44.8	27.0	61.0
815	Feather meal.	41.9	73.5	65.2	44.2	118.2	66.8	10.6	38.4	75.9	30.5	7.7	41.8	84.7	100.0	43.4	27.7	58.1
816	Feather meal.	41.1	72.4	63.3	52.9	110.8	70.6	7.7	41.2	74.4	24.8	5.9	43.5	93.5	109.0	43.0	26.9	60.2
662	Fish meal.	43.5	42.5	60.8	6.0	81.9	65.9	12.3	20.0	41.9	41.5	17.8	22.4	35.7	32.5	24.2	17.3	26.0
712	Fish meal.	41.2	43.3	65.3	6.2	88.7	54.0	14.1	22.4	45.1	47.8	20.0	23.7	30.3	32.4	27.7	21.3	26.0
713	Fish meal.	43.9	45.7	67.2	6.6	91.3	61.7	14.0	22.6	46.6	49.3	20.8	23.4	34.8	35.4	28.6	21.0	27.1
734	Fish meal.	41.7	48.5	68.3	8.1	92.4	40.1	16.5	26.2	53.5	55.2	23.0	26.7	27.7	30.4	30.4	23.0	33.2
735	Fish meal.	43.4	43.4	68.5	7.2	90.0	49.9	16.3	24.9	50.1	51.4	21.1	26.0	31.4	31.2	29.2	22.0	29.7
736	Fish meal.	40.7	42.4	62.2	9.4	87.4	53.8	13.2	23.6	47.1	45.5	20.2	23.8	33.1	35.0	26.5	20.7	28.0
788	Fish meal.	43.4	50.5	68.1	7.4	94.6	57.9	14.5	23.2	48.8	52.2	20.7	24.6	33.2	35.3	29.8	22.4	28.1
789	Fish meal.	42.9	46.7	65.8	6.9	89.1	51.2	16.4	24.9	49.7	52.7	21.3	25.9	31.4	31.5	29.1	22.6	30.2
814	Fish meal.	42.5	43.6	65.2	7.1	84.4	48.2	17.0	25.0	48.7	51.8	20.3	26.0	30.0	27.8	29.2	21.8	30.8
620	Fish meal.	40.4	41.8	65.0	5.7	90.0	45.1	16.1	26.9	49.0	52.5	18.3	25.8	30.0	26.3	28.1	21.4	31.8
621	Fish meal.	42.0	42.4	62.4	5.9	89.4	48.1	15.8	25.4	47.6	51.5	19.0	25.1	31.2	26.3	27.7	20.9	29.8
622	Fish meal.	40.8	42.2	62.7	5.6	88.3	47.1	15.7	25.2	47.3	51.8	18.6	24.8	30.3	26.4	27.0	20.4	30.1
623	Fish meal.	41.2	42.5	62.8	5.9	89.2	47.1	15.7	25.8	47.7	52.6	19.3	24.9	30.6	26.4	27.4	20.6	30.7
624	Fish meal.	40.5	42.8	63.8	5.9	88.8	45.0	15.8	27.0	49.2	52.6	19.2	26.5	30.7	27.4	28.7	22.2	31.2
625	Fish meal.	40.6	42.9	63.8	6.0	89.1	46.2	15.3	25.6	48.1	51.3	18.1	25.4	31.2	27.3	28.8	21.4	30.7
626	Fish meal.	44.1	42.1	62.7	5.1	88.2	51.9	16.0	25.0	46.7	50.5	16.6	24.7	32.4	27.1	27.8	19.3	30.2
627	Fish meal.	43.4	41.0	62.6	5.4	86.9	50.7	15.5	23.9	46.0	49.3	17.0	24.3	32.1	26.8	27.1	19.0	29.2
628	Fish meal.	43.0	41.1	62.3	5.3	86.4	50.3	15.7	23.9	45.9	49.2	17.0	24.2	31.8	26.6	27.3	18.9	29.0
629	Fish meal.	43.5	44.8	67.7	5.9	92.9	48.1	16.0	27.7	51.2	54.1	18.8	27.5	32.5	28.4	30.6	22.3	33.0
630	Fish meal.	42.4	43.5	65.2	5.6	92.0	47.1	15.9	27.2	50.1	52.9	18.4	26.6	31.7	28.0	29.2	22.1	32.5
631	Fish meal.	44.1	44.7	68.9	6.1	93.6	49.1	16.6	27.8	51.9	54.9	19.2	27.7	33.0	28.8	30.4	23.0	33.3
632	Fish meal.	40.4	39.7	57.9	5.1	79.4	47.4	19.1	22.5	44.5	46.7	15.7	23.0	30.4	26.5	24.9	18.6	28.2
633	Fish meal.	43.0	42.1	62.1	5.5	84.2	50.5	20.1	24.2	46.7	49.6	16.2	24.3	32.6	27.3	27.4	19.5	29.6
634	Fish meal.	43.7	42.8	62.8	5.6	84.9	51.5	20.4	24.8	47.5	50.3	16.5	24.9	33.0	27.6	27.4	19.8	30.7

TABLE 2 cont.

Code	Description	ALA	ARG	ASP	CYS	GLU	GLY	HIS	ILE	LEU	LYS	MET	PHE	PRO	SER	THR	TYR	VAL
658	Meat meal.	46.0	42.7	55.3	5.4	71.2	80.3	11.5	15.8	37.4	32.1	8.1	20.6	51.6	24.7	19.7	13.7	26.3
659	Meat meal.	40.5	35.6	44.3	4.0	56.9	75.5	9.0	11.0	29.3	25.2	5.9	16.6	46.2	20.5	15.1	10.0	19.9
660	Meat meal.	37.6	34.3	43.7	4.6	56.4	69.5	9.1	11.2	28.1	25.5	6.6	15.8	42.4	19.3	14.8	10.0	18.0
661	Meat meal.	42.3	39.0	48.4	4.5	65.3	81.5	10.0	14.0	31.5	27.8	6.8	17.5	48.8	21.9	17.1	11.3	22.2
709	Meat meal.	35.9	33.2	43.2	4.9	57.9	64.4	9.0	12.0	28.5	24.3	7.6	14.9	38.5	21.3	15.6	10.8	18.1
710	Meat meal.	42.3	36.7	49.7	4.8	63.2	72.8	12.3	12.0	34.7	29.0	7.6	18.0	43.3	22.6	17.6	11.6	21.1
711	Meat meal.	37.5	34.3	44.8	4.6	57.9	68.6	10.0	11.7	28.9	26.9	7.2	15.5	40.5	19.8	15.8	10.1	17.5
731	Meat meal.	38.6	37.4	47.7	6.1	63.5	72.3	9.8	13.3	31.3	27.8	7.8	16.7	44.3	23.1	17.1	11.5	20.3
732	Meat meal.	37.1	33.8	44.3	5.7	57.7	67.5	9.1	11.6	28.7	25.6	7.3	15.1	40.1	20.4	15.6	10.3	18.1
733	Meat meal.	38.5	35.9	46.3	6.1	59.7	69.6	10.2	12.7	30.1	27.7	7.6	16.2	42.4	20.9	16.4	10.7	18.6
785	Meat meal.	38.2	36.6	45.7	5.7	61.6	71.2	9.6	13.3	31.8	26.5	7.5	17.0	42.9	22.6	17.5	11.7	20.7
786	Meat meal.	43.6	39.5	51.7	5.9	67.0	77.2	12.2	13.8	37.7	30.5	7.8	19.8	46.2	24.4	19.2	13.1	24.4
787	Meat meal.	38.1	35.2	43.7	5.1	57.6	72.7	9.3	11.9	28.7	25.6	7.2	15.6	42.8	19.7	15.5	10.2	18.5
812	Meat meal.	8.5	13.4	14.6	4.5	35.0	9.8	5.5	4.9	10.8	9.0	3.4	6.5	10.5	7.5	6.1	5.0	6.7
813	Meat meal.	40.7	41.1	48.7	5.6	63.8	74.5	11.0	13.6	32.5	29.6	7.6	17.4	45.3	21.5	17.4	11.5	21.5
699	Oats.	5.2	7.3	10.2	3.3	21.3	5.5	2.5	3.9	8.3	5.0	2.0	5.4	5.8	4.9	3.6	3.6	5.2
700	Oats.	5.3	6.8	9.8	6.4	20.1	5.4	2.3	4.0	8.1	4.8	1.8	5.4	5.6	4.7	3.7	3.4	5.3
701	Oats.	4.9	7.9	9.8	5.4	20.0	5.2	2.4	3.8	7.9	4.8	1.4	5.1	5.7	4.6	3.5	3.4	5.0
702	Oats.	5.4	7.4	10.3	9.5	22.7	5.6	2.6	4.1	8.8	5.2	2.1	5.8	6.1	5.2	3.7	3.8	5.7
703	Oats.	6.6	10.5	12.9	11.0	26.8	6.8	3.1	5.0	10.4	6.2	2.2	6.8	7.3	6.3	4.6	4.6	6.5
764	Oats.	5.8	8.4	10.8	3.7	24.7	6.1	2.9	4.2	9.0	5.3	1.9	5.9	6.3	6.1	4.4	4.0	5.5
774	Oats.	5.5	7.6	9.9	4.0	23.6	5.8	2.7	3.8	8.8	5.1	2.1	5.7	6.1	5.8	4.0	3.7	5.2
770	Oats.	4.7	6.0	8.5	2.7	18.7	5.0	2.3	3.1	7.1	4.5	1.2	4.5	4.9	5.0	3.3	3.1	4.2
760	Oats.	4.9	6.5	9.0	3.7	19.1	5.3	2.3	3.0	7.2	4.6	2.0	4.6	5.2	5.2	3.6	3.4	4.0
759	Oats.	5.5	7.4	10.0	3.7	22.4	5.6	2.6	3.5	8.3	5.0	1.8	5.3	6.0	5.7	3.9	3.6	4.7
771	Oats.	5.6	7.7	9.7	4.0	28.2	5.9	2.9	4.1	9.2	5.0	2.1	6.1	8.3	6.0	4.1	3.9	5.5
796	Rapeseed meal.	18.7	26.5	32.4	9.8	68.6	20.8	11.0	14.4	29.2	22.1	8.5	16.1	24.4	18.7	18.3	11.7	18.6
797	Rapeseed meal.	17.9	25.8	31.0	9.5	66.0	20.0	10.6	13.6	27.8	21.5	8.2	15.5	23.5	17.9	17.6	11.3	18.0
798	Rapeseed meal.	17.3	23.9	29.3	9.3	64.0	19.5	10.3	13.2	27.4	20.9	8.0	15.0	22.4	17.1	17.4	11.0	17.5
818	Rapeseed meal.	18.7	27.2	33.1	10.7	70.6	21.0	11.3	14.5	29.1	23.8	8.6	16.2	24.5	18.7	18.6	11.9	18.6
819	Rapeseed meal.	17.5	25.0	30.3	9.5	65.7	19.7	10.5	13.5	27.3	21.4	8.3	15.2	22.6	17.4	17.4	11.1	17.0
820	Rapeseed meal.	19.3	28.9	33.0	10.9	76.5	21.5	11.9	15.2	30.7	24.8	9.1	16.9	26.1	19.4	18.5	12.3	19.4
821	Rapeseed meal.	16.9	25.2	30.3	18.0	65.5	19.8	11.2	14.1	27.5	23.1	14.9	22.2	17.4	17.3	10.8	17.7	10.8
822	Rapeseed meal.	17.0	24.4	29.7	9.4	63.1	19.2	10.4	13.5	26.5	21.9	8.1	14.7	22.4	17.2	17.1	10.9	17.9
668	Rapeseed meal.	19.7	27.2	33.1	9.9	74.7	21.8	11.6	15.6	31.0	25.1	7.9	17.0	26.7	19.8	19.2	12.2	20.2
669	Rapeseed meal.	16.4	22.5	28.4	9.5	60.3	19.1	9.7	12.6	25.4	21.0	8.2	14.2	21.2	16.2	15.9	10.2	16.6
670	Rapeseed meal.	18.9	26.2	32.4	9.9	70.4	20.8	11.1	15.0	29.5	23.9	8.0	16.4	24.4	18.5	18.8	11.7	19.0
720	Rapeseed meal.	17.0	24.2	31.7	10.4	68.5	19.7	11.1	13.3	27.6	21.9	8.5	14.9	22.2	18.1	16.9	20.7	16.6

TABLE 2. cont.

Code	Description	ALA	ARG	ASP	CYS	GLU	GLY	HIS	ILE	LEU	LYS	MET	PHE	PRO	SER	THR	TYR	VAL
721	Rapeseed meal.	16.1	22.2	29.4	9.6	64.0	18.7	10.3	12.3	25.5	21.6	7.2	13.6	21.3	17.1	16.2	10.4	15.2
722	Rapeseed meal.	16.5	23.8	30.6	9.7	65.4	19.0	10.4	12.6	26.6	21.1	7.2	14.4	21.2	17.4	16.4	10.2	15.4
743	Rapeseed meal.	16.2	23.6	30.9	10.0	65.2	19.4	10.6	13.2	26.2	22.1	7.8	14.1	21.8	17.5	16.8	10.7	16.9
744	Rapeseed meal.	17.1	25.2	31.2	11.5	72.3	19.9	11.3	13.9	27.7	22.6	10.4	15.0	23.0	17.8	16.9	10.7	17.1
745	Rapeseed meal.	15.3	20.8	28.0	9.4	59.4	18.1	10.1	12.5	24.5	20.5	8.3	13.2	20.3	16.0	15.6	10.0	15.5
746	Rapeseed meal.	16.9	25.2	31.5	10.8	71.9	19.9	11.1	13.7	27.7	22.4	8.8	15.0	23.1	17.9	16.7	10.8	17.2
747	Rapeseed meal.	17.6	25.8	32.4	11.4	75.1	20.4	11.6	14.4	28.6	23.3	9.0	15.8	23.4	18.3	17.3	11.1	17.8
641	Rice bran.	8.0	10.4	11.7	3.2	17.8	7.0	3.5	4.5	9.3	6.1	2.5	5.8	5.7	6.3	4.8	4.7	6.9
642	Rice bran.	9.5	13.0	13.2	3.3	18.7	8.2	4.1	4.9	9.9	7.1	3.5	6.8	7.2	5.6	4.8	8.0	
647	Rice bran.	9.1	13.6	14.8	3.8	21.3	8.1	4.5	4.6	10.6	6.2	3.1	6.6	7.0	5.4	5.3	7.3	
648	Rice bran.	9.7	9.5	14.7	3.7	20.5	8.4	4.1	4.9	11.2	7.1	3.5	6.9	7.0	7.8	6.1	5.3	7.6
644	Rice bran.	10.9	15.3	14.8	3.5	20.6	9.5	4.8	5.5	11.1	8.2	3.5	7.5	7.6	8.1	6.4	5.0	9.2
704	Screenings.	8.6	14.6	15.9	8.1	29.0	9.7	4.6	6.9	13.2	10.4	3.7	7.7	10.1	8.4	7.5	5.5	8.7
705	Screenings.	8.1	11.9	14.2	2.6	27.0	8.2	3.4	5.8	11.5	6.3	2.9	6.9	7.8	7.4	5.9	4.2	7.3
706	Screenings.	5.3	8.1	8.6	3.0	39.1	6.0	3.5	4.9	9.8	4.5	2.3	6.8	13.2	6.7	4.3	4.0	5.9
707	Screenings.	6.6	5.7	8.4	1.8	18.6	5.0	2.4	3.9	9.1	4.3	2.0	4.7	6.9	5.0	4.1	3.0	5.1
708	Screenings.	6.2	9.2	11.2	3.8	27.0	6.5	3.1	4.4	10.0	5.5	2.3	6.8	7.2	6.7	4.6	4.7	5.9
639	Sesame seed meal.	19.5	52.1	38.3	9.5	81.4	21.7	11.1	14.1	28.5	11.3	12.7	18.3	13.9	22.6	15.3	15.0	17.7
645	Sesame seed meal.	21.1	52.7	36.6	10.5	79.2	21.2	10.5	14.7	28.4	10.6	13.7	19.7	15.5	20.7	16.0	15.5	19.2
678	Shrimp meal.	23.5	25.0	39.6	4.8	52.4	20.1	10.0	15.5	24.2	23.2	8.9	133.6	18.1	22.8	18.2	25.0	19.9
840	Single cell protein.	50.8	33.1	71.4	5.3	72.4	39.6	13.6	30.9	52.3	44.4	16.9	25.4	22.2	24.9	33.9	22.4	38.0
671	Soybean meal.	22.2	39.4	61.8	7.8	93.8	21.6	13.6	20.4	39.2	32.0	7.6	25.7	26.0	27.2	20.2	17.9	21.4
672	Soybean meal.	22.4	39.0	61.8	7.8	94.2	21.8	13.6	20.0	39.3	32.1	7.8	25.6	26.1	27.5	20.2	17.9	21.0
673	Soybean meal.	26.8	46.8	73.8	8.7	111.7	26.1	16.5	25.8	48.2	38.6	8.7	31.1	31.2	33.2	24.7		28.7
723	Soybean meal.	22.8	39.9	63.2	8.0	98.0	22.7	14.7	20.2	40.7	34.1	7.7	25.8	26.3	28.8	21.1	17.7	20.6
724	Soybean meal.	22.2	39.0	61.6	8.5	96.0	21.7	14.3	21.1	39.9	33.4	8.2	25.3	25.3	28.2	20.5	17.3	20.6
748	Soybean meal.	22.0	38.7	63.3	8.9	98.1	22.0	14.2	18.2	39.1	33.0	8.4	24.6	26.4	28.6	20.5	17.2	17.7
749	Soybean meal.	20.4	35.0	59.3	9.0	91.0	20.4	13.1	16.6	36.4	31.0	8.1	22.8	24.0	26.6	19.3	15.8	17.1
750	Soybean meal.	22.3	38.7	63.8	9.5	99.1	22.2	14.2	18.3	39.1	33.1	8.4	24.8	26.5	28.9	20.5	17.6	18.8
751	Soybean meal.	21.4	36.6	61.8	9.0	96.8	21.5	13.7	18.1	38.2	32.3	7.7	24.3	25.1	28.1	20.1	17.0	18.7
799	Soybean meal.	24.9	42.7	66.1	8.9	103.9	23.8	14.9	22.6	44.2	35.7	8.4	28.1	29.1	30.0	22.4	19.5	23.8
800	Soybean meal.	25.4	44.7	68.2	9.4	110.2	24.7	15.5	23.1	45.6	37.2	8.4	29.4	30.5	31.9	23.1	20.7	23.8
801	Soybean meal.	24.5	42.1	64.2	9.4	102.6	23.7	14.7	22.1	43.2	35.4	8.6	27.7	28.5	29.8	22.1	19.9	23.1
802	Soybean meal.	24.4	44.8	67.9	9.2	104.0	23.9	15.1	23.0	43.8	35.8	8.7	28.0	28.8	30.6	22.1	20.0	23.5

TABLE 2 cont.

Code	Description	ALA	ARG	ASP	CYS	GLU	GLY	HIS	ILE	LEU	LYS	MET	PHE	PRO	SER	THR	TYR	VAL
803	Soybean meal.	23.8	44.6	66.7	9.5	102.7	23.1	14.7	22.3	42.5	35.4	8.5	27.5	28.5	30.0	22.3	19.4	23.2
804	Soybean meal.	24.5	46.1	68.6	9.6	105.3	23.6	15.3	22.5	43.4	36.1	8.7	27.9	29.0	31.0	22.7	20.1	23.2
823	Soybean meal.	24.4	46.0	65.9	8.8	102.7	23.5	15.1	23.3	43.7	35.7	8.0	28.1	28.7	29.6	21.9	19.7	24.0
824	Soybean meal.	22.7	42.9	63.0	9.6	97.6	21.9	14.3	22.4	40.8	33.9	8.9	26.5	26.9	28.0	20.6	17.8	23.1
825	Soybean meal.	25.9	49.2	70.6	9.6	110.4	24.8	16.1	24.9	46.5	38.3	8.5	30.0	30.7	32.0	23.1	20.3	26.1
826	Soybean meal.	24.7	46.8	68.5	9.3	104.9	23.7	15.2	23.6	44.2	36.4	8.2	28.4	29.3	30.5	22.3	19.9	24.2
679	Wheat.	6.5	8.6	11.3	4.5	62.0	7.9	4.8	6.0	13.1	5.6	3.0	9.3	20.1	8.9	5.5	4.9	6.5
680	Wheat.	5.9	8.6	9.6	3.8	53.0	7.0	4.3	5.3	11.1	5.7	2.3	7.9	17.0	7.9	4.8	4.7	6.4
681	Wheat.	6.2	8.7	10.4	4.6	58.9	7.5	4.6	5.7	12.3	5.3	3.4	9.0	17.8	8.6	5.3	4.8	6.6
682	Wheat.	5.7	7.7	9.4	4.3	52.0	6.8	4.1	5.1	11.1	4.9	2.9	7.5	6.2	7.6	4.7	4.3	5.7
683	Wheat.	5.2	6.6	8.5	3.9	43.5	6.0	3.7	4.3	9.8	4.4	2.7	6.4	13.8	6.7	4.3	3.7	4.9
684	Wheat.	5.7	8.1	9.2	3.7	52.5	6.8	4.2	4.7	10.7	5.5	2.5	7.8	17.2	7.7	4.6	4.7	5.5
685	Wheat.	6.8	9.3	11.4	4.6	62.2	8.1	4.9	6.4	13.3	5.7	2.5	9.4	19.9	8.9	5.4	5.2	7.2
686	Wheat.	5.4	7.6	8.6	3.8	49.2	6.5	4.0	4.7	10.5	5.3	2.2	7.2	16.1	7.6	4.5	4.1	5.7
687	Wheat.	6.1	8.3	10.8	4.3	58.6	7.3	4.6	5.8	12.3	5.2	1.2	8.8	18.8	8.2	5.0	4.9	6.3
688	Wheat.	6.0	8.9	10.5	4.7	55.9	7.1	4.4	5.6	11.8	5.0	3.0	8.1	18.4	8.2	5.0	4.6	6.0
773	Wheat.	6.3	8.7	9.7	5.0	58.8	7.3	4.5	5.7	12.5	5.1	2.3	9.1	20.3	8.7	5.2	4.9	7.0
778	Wheat.	6.2	8.5	9.6	4.5	59.1	7.3	4.5	5.7	12.6	5.1	2.7	9.2	20.3	8.7	5.1	5.1	6.8
784	Wheat.	6.4	8.6	9.2	4.2	113.8	7.2	4.2	5.5	12.0	5.0	2.8	8.7	19.1	8.0	5.0	5.2	7.0
779	Wheat.	5.7	8.0	8.9	4.3	52.8	6.8	4.1	5.1	11.4	4.8	2.5	8.2	18.0	8.1	4.7	4.7	6.2
782	Wheat.	6.3	8.3	9.2	4.1	58.5	7.1	4.1	5.5	12.0	4.8	2.5	8.7	19.0	8.0	5.0	5.0	6.9
783	Wheat.	6.5	8.6	9.5	4.3	117.4	7.3	4.3	5.5	12.1	5.0	2.9	8.8	19.4	8.1	5.1	5.0	6.9
772	Wheat.	5.3	7.4	8.2	4.2	46.0	6.3	3.8	4.7	10.4	4.5	2.6	7.2	15.9	7.0	4.5	4.4	5.9
781	Wheat.	5.6	7.7	8.4	4.2	49.5	6.6	3.9	4.8	10.8	4.6	2.6	7.7	17.0	7.8	4.8	4.6	5.9
776	Wheat.	7.2	10.3	11.1	5.1	66.2	8.6	5.2	6.5	14.6	5.9	3.8	10.9	24.1	10.3	6.0	6.4	8.1
780	Wheat.	6.1	8.7	9.8	4.5	56.8	7.2	4.4	5.4	12.2	5.0	2.7	8.8	19.4	8.6	5.1	5.2	6.5
676	Wheat shorts.	9.1	13.8	14.2	4.3	37.9	10.2	5.3	5.5	11.9	8.6	2.7	7.4	12.3	7.7	6.2	5.4	8.1
677	Wheat shorts.	10.9	16.4	17.0	4.4	41.0	12.0	6.2	6.3	13.6	10.3	3.1	8.4	13.4	9.0	7.2	6.1	9.8
728	Wheat shorts.	8.3	12.6	12.7	4.2	34.3	9.1	4.8	5.2	11.0	7.5	2.9	7.0	11.3	8.1	5.8	5.1	7.6
729	Wheat shorts.	9.3	14.0	42.8	4.4	35.9	10.2	5.3	5.4	11.8	8.3	2.8	7.4	11.8	8.7	6.6	5.5	8.1
730	Wheat shorts.	8.3	12.7	13.0	4.3	34.5	9.3	4.9	5.1	11.0	7.4	2.8	6.9	11.4	8.1	6.0	5.1	7.4
756	Wheat shorts.	8.4	12.4	13.2	4.5	36.6	9.1	4.9	4.8	11.2	7.5	2.7	7.0	12.3	8.4	6.0	5.2	7.2
757	Wheat shorts.	8.8	12.8	13.6	4.7	38.4	9.5	5.0	5.1	11.6	7.9	2.8	7.2	12.9	8.6	6.2	5.5	7.4
758	Wheat shorts.	8.0	11.8	37.4	4.7	33.2	9.0	4.7	4.3	10.3	7.0	3.0	6.5	11.0	7.9	5.7	4.9	6.5
809	Wheat shorts.	9.5	14.7	13.8	4.9	36.5	10.1	5.4	5.7	12.5	8.3	3.4	7.8	12.4	7.7	6.0	5.7	8.4
810	Wheat shorts.	9.7	14.7	15.5	4.5	40.0	10.1	5.5	6.3	13.5	8.4	3.0	8.2	13.4	8.9	6.5	5.9	8.8
811	Wheat shorts.	9.3	15.0	14.2	4.5	36.4	10.1	5.3	5.9	12.1	8.4	3.4	7.6	12.5	8.2	6.3	5.4	8.8
832	Wheat shorts.	9.7	15.4	14.5	4.5	39.3	10.4	5.5	6.0	12.5	8.4	3.3	8.0	13.2	8.9	6.5	5.8	8.5

TABLE 2 cont.

Code	Description	ALA	ARG	ASP	CYS	GLU	GLY	HIS	ILE	LEU	LYS	MET	PHE	PRO	SER	THR	TYR	VAL
833	Wheat shorts.	10.0	15.8	14.6	4.6	41.6	10.8	5.6	6.0	12.9	8.7	3.2	8.2	13.9	9.6	6.7	6.0	8.8
834	Whcat shorts.	9.9	16.1	14.9	4.6	39.8	10.8	5.6	5.9	12.8	8.7	3.4	8.2	13.4	9.2	6.6	5.9	8.8
835	Wheat shorts.	10.0	16.2	14.7	4.5	40.7	10.8	5.6	6.0	13.0	8.8	3.2	8.2	13.2	9.2	6.8	5.9	8.8
638	Yucca.	5.5	12.7	1.6	0.6	14.1	6.1	4.5	ND	7.7	9.5	ND	ND	ND	7.2	5.3	ND	ND
640	Yucca.	0.8	1.3	0.9	ND	2.4	0.5	0.4	ND	ND	ND	ND	ND	0.5	0.4	ND	ND	ND

1 Blanks (-) = variable not measured; ND = variable not detected in a reportable amount.

TABLE 3

True available amino acids in the dry matter of several feedingstuffs (g/kg)¹

Code	Description	ALA	ARG	ASP	CYS	GLU	HIS	ILE	LEU	LYS	MET	PHE	PRO	SER	THR	TYR	VAL
655	Alfalfa.	4.9	4.9	16.8	0.7	9.3	2.0	3.8	7.5	4.7	1.9	5.1	11.0	4.8	4.3	2.8	5.2
656	Alfalfa.	7.7	8.3	21.5	-	14.7	2.8	5.9	11.7	5.1	-	7.5	11.5	6.8	6.5	4.6	7.4
653	Alfalfa.	6.1	5.8	20.6	1.0	12.1	2.4	4.7	8.9	6.1	1.7	6.0	10.5	5.3	5.3	3.4	6.2
839	Alfalfa.	7.8	5.2	15.3	2.4	8.9	2.6	3.9	7.9	5.1	1.4	5.1	5.4	4.6	4.3	3.1	4.7
657	Alfalfa.	6.3	6.3	19.1	0.6	11.3	3.2	4.8	9.3	4.4	1.8	6.1	9.7	5.9	5.6	3.4	6.2
652	Alfalfa.	7.0	7.0	19.7	0.9	13.1	3.3	5.1	10.1	5.2	2.1	6.3	10.2	6.4	6.1	4.0	6.7
654	Alfalfa.	6.4	6.2	18.9	2.3	11.1	3.0	4.4	9.2	4.1	1.8	5.9	7.2	5.5	5.3	3.5	5.8
649	Algae.	27.1	19.7	30.4	10.1	29.8	4.7	11.0	25.8	16.8	6.9	15.4	15.4	12.5	15.9	10.6	17.1
650	Algae.	27.2	20.9	34.0	9.5	34.3	5.1	12.9	29.1	16.6	7.4	17.2	16.7	13.7	16.6	11.6	19.2
689	Barley.	3.9	6.1	6.7	2.5	31.8	3.0	3.5	7.9	4.4	1.9	5.8	13.0	4.9	3.7	3.5	4.7
690	Barley.	3.9	5.4	7.0	3.9	28.8	2.7	3.3	7.9	3.7	1.6	5.6	12.3	4.7	3.3	3.2	3.9
691	Barley.	2.5	4.1	5.5	2.1	17.9	1.8	2.1	5.2	2.6	1.5	3.3	7.1	3.2	2.1	1.9	2.2
692	Barley.	3.7	5.6	6.9	2.8	28.1	2.6	3.2	7.5	3.5	1.8	5.4	12.1	4.4	3.2	3.0	3.8
693	Barley.	3.4	4.8	6.7	2.3	23.1	2.3	2.7	6.5	3.2	1.2	4.2	9.5	3.9	2.9	2.6	3.3
694	Barley.	3.7	5.4	7.6	2.8	25.0	2.5	3.0	7.1	3.8	1.3	4.9	10.6	4.5	3.1	3.0	3.6
695	Barley.	3.9	6.4	7.8	3.3	31.1	2.8	3.7	8.1	3.9	2.2	6.0	13.4	5.0	3.6	3.5	3.9
696	Barley.	3.7	5.7	7.4	3.1	27.8	2.6	3.1	7.4	3.8	0.7	5.3	11.1	6.1	3.4	3.1	3.9
697	Barley.	3.8	6.0	7.7	2.8	27.6	2.7	3.3	7.8	4.0	3.2	5.4	11.7	4.7	3.5	3.1	4.1
698	Barley.	4.6	7.9	8.7	2.5	35.0	3.2	4.3	9.3	4.7	2.0	7.5	16.2	5.2	4.1	3.9	5.9
769	Barley.	3.8	6.0	6.3	2.6	31.3	2.9	3.4	8.3	4.0	1.6	6.5	14.6	5.0	3.6	3.6	4.8
765	Barley.	3.9	5.6	6.1	2.2	25.9	2.6	3.4	7.5	4.1	1.6	5.4	10.8	4.1	3.2	3.3	4.7
762	Barley.	4.4	6.0	6.9	2.5	31.4	2.8	3.8	8.5	4.4	1.9	6.3	14.2	5.2	3.9	3.8	5.2
775	Barley.	3.2	4.5	5.2	2.2	20.6	2.1	2.5	6.1	3.4	1.4	4.3	9.2	3.5	2.7	2.5	3.6
777	Barley.	4.3	6.3	6.5	3.2	38.2	2.8	4.1	9.5	3.8	2.2	7.6	18.3	5.7	3.6	4.2	5.7
767	Barley.	2.8	4.2	4.8	1.9	19.0	2.1	2.4	5.5	3.2	1.3	3.9	8.0	2.9	2.4	2.4	3.3
768	Barley.	3.3	4.8	5.3	2.4	23.5	2.3	2.8	6.6	3.4	1.4	4.7	10.6	4.0	2.8	2.6	4.0
761	Barley.	3.6	5.1	6.3	2.9	25.8	2.5	2.6	6.9	3.6	1.5	5.1	11.6	4.7	3.3	3.0	3.8
763	Barley.	3.7	5.1	5.9	2.1	24.6	2.4	3.1	6.9	3.7	1.5	5.0	10.9	4.3	3.1	3.0	4.3
766	Barley.	5.1	7.2	7.8	2.7	39.0	3.4	4.6	10.2	5.2	1.9	8.3	18.2	5.8	4.2	4.4	6.2
666	Blood meal.	67.7	46.5	101.0	15.5	87.0	48.4	15.5	110.6	69.0	7.9	59.4	44.2	52.0	37.9	28.5	67.2
667	Blood meal.	89.7	45.5	121.6	9.7	91.0	67.2	7.3	136.1	93.1	12.6	77.9	40.0	55.4	48.0	31.6	89.2
717	Blood meal.	64.9	31.5	81.5	6.9	66.3	44.8	4.4	96.3	68.4	11.6	57.2	28.3	43.3	38.0	24.4	53.9
718	Blood meal.	62.8	41.1	92.9	13.2	81.7	47.5	12.4	102.8	66.8	8.6	54.9	34.5	47.8	34.4	25.7	61.7
719	Blood meal.	68.0	39.0	93.7	11.5	80.1	51.6	10.1	103.8	75.3	11.9	58.1	32.9	45.8	40.0	27.4	59.7
739	Blood meal.	66.5	32.3	84.8	9.3	68.4	46.3	4.6	100.5	67.0	12.6	58.9	30.9	44.6	39.5	25.4	57.1

TABLE 3 cont.

Code	Description	ALA	ARG	ASP	CYS	GLU	HIS	ILE	LEU	LYS	MET	PHE	PRO	SER	THR	TYR	VAL
740	Blood meal.	64.6	40.2	90.4	12.9	79.8	48.9	12.1	101.9	69.6	8.6	54.5	34.4	45.2	33.5	25.8	61.4
741	Blood meal.	64.9	35.6	86.7	9.9	74.5	47.7	8.5	100.7	66.6	10.7	56.0	31.6	43.5	37.3	26.7	59.1
742	Blood meal.	71.8	35.2	95.9	9.2	74.8	54.4	6.4	110.7	70.8	10.8	62.0	33.0	44.1	37.4	25.5	65.2
759	Blood meal.	75.9	41.5	107.5	8.6	83.4	60.5	6.5	123.8	85.3	11.1	69.3	36.0	49.2	41.4	28.6	72.0
792	Blood meal.	66.8	47.9	99.0	13.5	87.0	52.7	13.3	111.4	73.5	8.8	60.1	38.1	51.0	36.8	28.1	65.7
793	Blood meal.	66.5	42.0	99.2	9.3	82.0	52.4	7.6	107.9	72.4	7.7	57.3	32.3	41.4	31.5	24.0	61.3
794	Blood meal.	68.2	38.4	91.6	8.8	76.7	51.7	5.3	109.1	77.7	12.6	64.9	31.9	48.4	42.6	27.7	61.2
795	Blood meal.	71.8	46.9	101.5	12.2	87.7	56.2	12.2	115.8	76.0	8.4	61.1	37.3	50.2	36.5	28.9	70.9
817	Blood meal.																
643	Coconut meal.	7.1	24.8	14.1	2.4	31.6	2.9	5.6	10.0	5.6	4.3	8.4	5.7	7.0	4.1	3.8	8.6
838	Corn.	7.3	5.0	7.5	2.4	17.7	2.8	3.0	12.0	3.3	2.3	4.6	9.0	4.6	3.4	4.0	4.3
853	Corn.	7.0	5.4	7.0	2.3	18.3	3.1	2.7	12.2	3.7	2.1	4.4	9.0	4.4	3.3	3.9	3.8
855	Corn.	7.0	5.6	6.7	2.4	18.5	3.1	2.8	12.6	3.6	2.2	4.5	9.3	4.2	3.2	3.7	4.0
856	Corn.	7.4	5.8	7.4	2.8	19.4	3.3	3.0	13.2	3.8	1.4	4.8	9.9	4.8	3.6	4.3	4.3
857	Corn.	8.6	6.6	7.9	2.8	21.9	3.6	3.5	14.8	4.2	2.4	5.4	11.1	5.3	4.0	4.6	4.8
858	Corn.	7.5	5.7	6.9	2.3	19.3	3.3	3.2	13.2	3.6	2.1	4.8	9.9	4.4	3.2	4.1	4.3
859	Corn.	7.9	5.7	6.5	2.4	19.7	3.2	3.2	13.6	3.1	2.2	4.9	10.0	4.4	3.3	4.1	4.5
860	Corn.	8.0	5.4	7.5	2.8	21.2	3.4	3.5	14.6	3.6	2.4	5.2	10.7	5.2	3.7	4.6	4.8
861	Corn.	7.8	5.6	7.0	2.6	20.1	3.2	3.2	13.7	3.2	2.3	5.1	9.7	5.1	3.5	4.4	4.3
862	Corn.	6.9	5.3	6.4	2.3	18.1	3.0	3.0	12.7	2.9	2.3	4.6	9.1	4.1	3.0	3.9	4.0
646	Corn germ meal.	11.9	14.7	13.4	2.5	28.0	6.2	6.4	15.8	8.4	3.6	8.9	10.6	9.0	6.8	5.8	10.7
651	Corn gluten feed.	20.3	10.0	14.4	4.6	43.6	6.0	7.3	30.9	5.2	4.7	12.2	22.5	11.8	7.8	9.6	10.2
674	Corn gluten feed.	14.1	10.5	13.6	3.6	32.4	6.3	5.6	20.3	6.4	3.0	7.9	16.3	9.0	7.0	6.0	9.1
726	Corn gluten feed.	11.5	10.0	10.8	3.3	27.0	5.4	4.9	17.0	5.0	3.0	6.9	14.3	8.1	5.5	5.1	7.8
727	Corn gluten feed.	12.1	10.6	10.9	3.5	28.6	5.6	5.1	17.7	5.1	3.3	7.4	14.5	8.4	6.1	5.5	8.3
754	Corn gluten feed.	14.6	11.7	13.5	4.3	34.2	6.4	5.5	21.1	5.5	4.0	8.3	18.2	9.8	6.9	7.0	9.2
755	Corn gluten feed.	14.5	11.0	13.3	4.2	33.1	6.2	5.5	20.8	5.8	4.0	8.2	17.9	9.4	6.7	6.5	9.0
805	Corn gluten feed.	15.8	13.0	15.1	3.8	38.1	7.2	6.7	23.7	6.5	4.2	9.0	19.8	10.3	7.6	7.3	10.6
806	Corn gluten feed.	14.0	11.9	13.5	3.4	34.2	6.6	6.0	21.0	6.2	3.9	8.1	17.2	9.2	6.9	6.4	9.3
827	Corn gluten feed.	13.2	11.6	13.2	3.4	31.0	6.1	5.8	19.4	5.6	3.7	7.6	16.4	8.6	6.4	6.3	9.3
828	Corn gluten feed.	13.3	9.8	11.5	3.7	31.6	6.1	4.6	20.0	5.5	3.5	6.9	17.8	8.3	6.2	6.2	7.2
675	Corn gluten meal.	54.1	18.7	37.1	9.1	133.9	11.2	21.3	101.6	9.2	12.9	37.7	57.4	32.5	19.4	30.7	24.0
725	Corn gluten meal.	58.7	20.1	40.4	12.9	148.0	13.0	22.2	112.3	10.0	18.0	40.6	61.5	36.0	20.9	34.4	24.9
752	Corn gluten meal.	51.9	16.1	35.9	11.0	131.9	10.9	17.7	99.1	8.7	16.4	35.2	52.7	31.9	18.2	29.8	21.4
753	Corn gluten meal.	50.9	18.6	38.1	11.7	117.2	9.8	18.9	71.5	9.4	17.8	38.8	58.6	33.0	20.5	32.6	23.1
807	Corn gluten meal.	64.5	21.9	42.4	10.6	148.1	13.0	24.0	112.9	9.8	16.4	44.9	67.1	38.1	22.2	37.1	28.7
808	Corn gluten meal.	67.3	23.0	43.5	10.4	157.9	13.8	26.2	121.3	10.6	17.1	47.3	67.9	38.3	22.8	39.5	30.9

TABLE 3 cont.

Code	Description	ALA	ARG	ASP	CYS	GLU	HIS	ILE	LEU	LYS	MET	PHE	PRO	SER	THR	TYR	VAL
829	Corn gluten meal.	65.5	24.2	42.7	11.4	153.5	14.2	24.0	119.4	10.6	14.9	45.0	68.7	37.1	22.5	38.1	29.8
830	Corn gluten meal.	64.8	26.1	43.7	12.2	145.1	14.3	24.2	112.4	11.5	20.3	45.0	69.4	38.6	22.4	38.5	28.9
831	Corn gluten meal.	63.2	25.6	43.2	11.4	150.7	14.1	24.4	112.1	11.5	19.8	44.5	66.8	37.5	22.0	37.9	29.1
663	Feather meal.	34.5	57.5	43.6	27.0	75.8	6.9	34.5	62.7	19.9	5.7	36.8	94.0	78.9	32.3	21.8	46.7
664	Feather meal.	35.8	57.6	36.0	20.8	66.4	5.0	36.3	64.3	13.9	4.3	40.2	109.3	92.1	30.7	22.1	53.7
665	Feather meal.	34.0	52.8	33.2	22.8	60.6	5.0	34.0	62.2	12.9	4.5	36.8	95.4	79.8	29.1	19.4	47.8
714	Feather meal.	30.1	48.3	34.6	23.0	69.5	7.9	27.7	56.9	17.9	5.4	32.9	50.9	71.9	27.6	19.4	39.6
715	Feather meal.	32.8	51.6	34.3	31.1	72.1	4.6	33.0	58.6	12.6	3.9	35.9	67.4	89.1	30.6	19.6	46.3
716	Feather meal.	31.2	51.0	33.5	23.1	68.6	5.5	32.3	57.7	15.3	4.3	34.4	63.8	81.1	28.8	18.7	45.5
737	Feather meal.	31.7	47.2	36.6	24.8	69.9	9.8	25.8	57.3	20.2	5.4	32.9	48.8	67.4	27.1	19.2	40.8
738	Feather meal.	31.9	51.8	34.7	41.0	71.4	5.0	32.0	57.8	13.1	4.6	35.6	63.7	82.9	29.9	19.8	46.4
790	Feather meal.	30.9	58.6	36.7	23.8	76.1	8.2	31.3	61.8	19.1	5.3	36.3	59.1	79.4	29.9	21.2	45.8
791	Feather meal.	34.2	64.2	44.0	34.4	85.0	5.7	35.5	66.0	17.5	5.2	40.1	75.9	93.1	35.5	22.9	51.8
815	Feather meal.	32.7	59.8	36.1	18.1	77.8	7.8	31.9	61.4	19.1	6.2	35.3	55.3	74.8	29.4	21.5	45.5
816	Feather meal.	34.9	62.7	40.6	32.2	82.2	5.9	36.9	64.5	16.8	4.3	38.8	71.1	90.1	32.8	22.3	51.3
662	Fish meal.	39.1	38.3	48.9	3.6	71.3	10.2	17.8	37.7	35.1	16.5	20.1	31.4	27.8	20.7	15.2	22.9
712	Fish meal.	36.9	39.5	56.5	4.5	80.3	12.5	20.1	41.1	42.2	18.6	21.3	27.0	29.0	24.9	19.5	23.0
713	Fish meal.	40.8	41.8	60.6	5.2	85.0	12.6	20.9	43.8	44.7	19.8	21.7	32.4	32.8	26.4	19.5	25.0
734	Fish meal.	38.6	43.5	62.3	6.2	86.2	14.7	24.3	50.6	48.8	21.9	24.9	25.1	27.8	21.6	30.8	30.8
735	Fish meal.	37.0	37.3	57.2	5.3	76.7	13.5	20.7	43.1	42.8	18.2	22.0	26.4	26.8	25.1	19.0	24.8
736	Fish meal.	36.8	38.9	54.8	7.4	78.9	11.6	21.3	43.2	40.9	18.7	21.5	29.0	31.8	24.1	18.8	25.0
788	Fish meal.	40.8	47.7	61.8	5.9	89.4	13.2	22.0	46.7	49.2	19.9	23.2	31.1	33.0	27.9	21.4	26.5
789	Fish meal.	40.5	43.0	60.0	5.6	84.1	14.8	23.5	47.4	47.1	20.5	24.5	29.5	29.4	27.1	21.5	28.5
814	Fish meal.	38.8	39.3	57.2	4.7	77.1	14.7	23.0	45.3	43.6	19.0	23.9	26.6	24.8	25.8	20.1	28.3
620	Fish meal.	36.9	39.4	57.2	4.3	83.6	14.5	25.4	46.2	47.1	17.2	24.0	27.0	23.8	25.7	20.1	29.9
621	Fish meal.	38.3	39.8	55.3	4.7	82.6	14.1	24.1	45.0	45.0	17.9	23.4	27.9	23.8	25.2	19.7	27.9
622	Fish meal.	37.9	40.2	56.7	4.7	83.4	14.2	24.0	45.1	47.0	17.2	23.4	28.0	24.5	25.0	19.5	28.3
623	Fish meal.	37.7	39.9	56.5	5.0	83.6	14.4	24.6	45.5	48.6	18.0	23.5	27.1	24.2	25.3	19.6	28.9
624	Fish meal.	37.1	40.4	57.3	4.9	82.7	14.3	25.6	46.8	48.0	17.7	25.0	27.4	25.1	26.6	21.1	29.3
625	Fish meal.	37.2	40.2	57.4	4.9	83.2	13.8	24.4	45.8	46.7	16.5	23.9	27.8	24.9	26.7	20.3	29.0
626	Fish meal.	39.9	39.5	53.9	3.8	80.3	14.2	23.5	43.8	43.8	14.9	22.9	29.4	24.2	25.1	18.1	28.2
627	Fish meal.	40.1	38.9	53.4	4.2	80.8	14.0	22.6	43.4	45.0	15.3	22.6	28.9	24.0	24.5	18.0	27.5
628	Fish meal.	40.2	39.4	54.9	4.0	81.2	14.3	22.8	43.6	44.8	15.4	22.8	29.6	24.2	25.2	18.0	27.6
629	Fish meal.	40.2	42.4	61.0	4.7	86.7	14.5	26.1	48.4	49.8	17.2	25.8	29.6	26.0	28.0	21.0	30.8
630	Fish meal.	39.7	41.7	58.7	4.6	86.3	14.2	25.7	47.6	46.9	16.9	25.0	29.4	25.7	27.0	21.0	30.6
631	Fish meal.	40.6	42.0	62.4	4.6	87.4	15.0	26.2	49.0	48.1	17.3	25.8	30.0	26.3	28.1	21.7	31.0
632	Fish meal.	37.6	37.5	51.6	3.8	73.9	17.4	21.0	41.3	41.0	21.3	28.2	23.9	22.8	17.3	26.3	27.5
633	Fish meal.	40.5	40.4	55.9	4.3	79.3	18.6	22.7	44.0	45.4	14.6	22.7	30.4	24.8	25.2	18.4	27.8
634	Fish meal.	41.1	41.0	56.8	4.3	79.9	18.6	23.4	44.9	43.7	14.8	23.3	31.2	25.4	25.4	18.7	28.9

TABLE 3 cont.

Code	Description	ALA	ARG	ASP	CYS	GLU	HIS	ILE	LEU	LYS	MET	PHE	PRO	SER	THR	TYR	VAL
658	Meat meal.	41.0	39.4	45.3	3.3	62.7	10.0	14.3	34.1	27.2	7.4	18.8	45.4	21.3	17.1	12.4	23.9
659	Meat meal.	31.2	28.2	30.8	1.2	43.5	7.4	8.6	23.9	18.4	4.8	13.6	34.7	15.1	11.5	8.0	15.9
660	Meat meal.	31.8	29.8	36.2	2.6	47.6	7.8	9.5	24.4	21.1	5.9	13.6	35.0	15.5	12.4	8.4	15.2
661	Meat meal.	36.7	35.0	39.3	2.4	56.5	8.8	12.5	28.2	23.6	6.2	15.6	41.7	18.3	14.8	10.1	19.6
709	Meat meal.	29.5	28.5	31.4	1.9	46.9	7.3	9.8	24.0	18.7	6.6	12.5	30.5	16.7	12.5	9.1	14.5
710	Meat meal.	36.2	31.8	39.4	2.4	53.4	10.6	10.0	30.4	23.6	6.7	15.7	36.2	19.0	14.8	10.0	18.0
711	Meat meal.	31.2	29.0	37.3	3.0	49.2	8.6	9.9	25.3	22.7	6.4	13.4	32.8	16.5	13.3	8.8	14.7
731	Meat meal.	33.0	32.7	37.8	3.7	54.0	8.4	11.3	27.5	23.2	6.9	14.5	36.6	19.2	14.3	10.0	17.2
732	Meat meal.	30.9	27.8	36.0	3.0	48.2	7.7	9.5	24.6	21.0	6.4	12.8	32.5	16.4	12.9	8.7	15.0
733	Meat meal.	32.9	30.7	38.4	3.6	48.9	8.8	10.6	26.2	23.6	6.8	14.0	35.3	17.0	13.7	9.1	15.4
785	Meat meal.	33.7	32.8	38.1	3.2	53.8	8.5	11.7	28.6	22.5	6.7	15.2	37.4	19.5	15.1	10.3	18.1
786	Meat meal.	38.8	35.4	43.1	3.5	59.0	10.8	12.2	34.2	26.3	7.0	17.9	40.5	21.2	16.9	11.9	21.9
787	Meat meal.	32.7	30.3	36.8	3.1	49.6	8.0	10.3	25.5	21.7	6.5	13.7	36.0	16.5	13.1	9.0	16.1
812	Meat meal.	2.6	8.5	4.4	1.3	24.9	4.1	3.0	6.6	4.0	2.5	4.2	3.2	3.3	2.9	3.5	3.5
813	Meat meal.	36.4	36.1	41.9	3.0	55.7	9.9	11.8	28.6	25.6	6.9	15.4	40.4	18.3	14.7	10.1	18.9
699	Oats.	4.3	6.9	8.7	2.9	20.0	2.4	3.5	7.6	4.3	1.6	5.1	5.6	4.4	3.1	3.5	4.5
700	Oats.	4.7	6.7	8.7	6.1	19.3	2.2	3.7	7.7	4.3	1.6	5.2	5.4	4.4	3.3	3.4	4.8
701	Oats.	4.0	7.5	8.5	4.9	18.4	2.2	3.4	7.1	4.1	1.1	4.7	5.1	3.9	2.9	3.2	4.3
702	Oats.	4.9	7.4	9.8	9.1	22.3	2.6	3.9	8.4	4.9	1.9	5.6	5.8	4.8	3.5	3.7	5.3
703	Oats.	6.0	10.4	11.9	10.8	26.1	3.1	4.8	10.0	5.7	2.0	6.6	7.2	6.0	4.3	4.6	6.1
764	Oats.	4.5	7.4	8.8	2.4	22.3	2.6	3.5	7.8	4.4	1.6	5.2	5.2	5.0	3.3	3.3	4.4
774	Oats.	4.7	7.0	8.4	3.0	21.8	2.4	3.2	7.9	4.4	1.9	5.3	5.1	4.9	3.2	3.3	4.5
770	Oats.	3.7	5.6	6.9	2.1	16.9	2.1	2.6	6.3	3.7	0.9	4.2	4.3	4.4	2.6	2.7	3.5
760	Oats.	4.2	6.1	7.9	3.0	17.8	2.2	2.6	6.5	4.2	1.8	4.2	4.6	4.6	3.0	3.1	3.5
759	Oats.	4.6	6.8	8.6	3.1	20.9	2.3	3.1	7.6	4.2	1.6	4.9	5.4	5.1	3.2	3.4	4.1
771	Oats.	4.9	7.2	8.3	3.1	26.8	2.7	3.7	8.4	4.3	1.7	5.7	7.7	5.4	3.5	3.6	5.0
796	Rapeseed meal.	16.2	24.2	26.9	7.1	63.1	9.8	12.2	25.9	17.8	7.7	14.3	20.3	15.8	14.8	10.1	15.5
797	Rapeseed meal.	15.4	23.3	25.2	6.1	59.9	9.3	11.6	24.4	17.1	7.4	13.6	19.2	14.6	14.2	9.6	15.0
798	Rapeseed meal.	14.8	21.6	23.7	6.1	58.3	9.0	11.2	24.2	16.7	7.2	13.2	18.4	13.9	14.1	9.3	14.8
818	Rapeseed meal.	16.4	24.8	28.0	8.9	65.3	10.0	12.6	26.2	19.4	7.9	14.6	20.5	15.9	15.3	10.4	15.6
819	Rapeseed meal.	14.8	22.3	24.5	5.4	59.4	9.1	11.4	23.9	17.0	7.4	13.4	18.2	13.9	9.4	14.2	
820	Rapeseed meal.	17.0	26.2	28.1	9.0	70.9	10.6	13.2	27.7	20.0	8.4	15.2	22.3	16.6	15.4	10.8	
821	Rapeseed meal.	14.2	22.5	24.2	15.6	58.9	9.9	11.9	24.2	18.7	14.0	12.8	17.8	14.1	13.6	9.0	14.5
822	Rapeseed meal.	14.9	22.2	24.7	6.7	57.9	9.2	11.7	23.8	18.0	7.4	13.2	18.7	14.5	14.1	9.3	15.3
668	Rapeseed meal.	17.5	25.0	28.2	6.0	69.6	10.3	13.8	28.0	21.0	7.2	15.4	23.0	17.0	16.1	10.7	17.7
669	Rapeseed meal.	13.1	19.4	22.3	6.8	52.7	8.2	10.1	20.9	15.6	7.3	11.7	16.3	12.1	8.0	13.2	
670	Rapeseed meal.	16.1	23.4	26.4	7.1	63.7	9.5	12.8	25.6	19.5	7.1	14.2	19.9	14.9	15.0	9.9	15.9
720	Rapeseed meal.	14.2	21.7	26.0	8.0	62.2	9.7	10.8	23.8	17.4	7.6	12.6	18.2	15.1	13.4	8.9	

TABLE 3 cont.

Code	Description	ALA	ARG	ASP	CYS	GLU	HIS	ILE	LEU	LYS	MET	PHE	PRO	SER	THR	TYR	VAL
721	Rapeseed meal.	13.6	19.6	24.3	7.2	58.4	9.1	10.1	22.5	17.2	6.4	11.8	17.5	14.3	12.9	8.7	12.3
722	Rapeseed meal.	13.8	21.4	25.1	7.1	59.5	9.2	10.4	23.1	16.9	6.3	12.4	17.4	14.5	13.2	8.5	12.3
743	Rapeseed meal.	14.0	21.3	26.1	7.4	60.2	9.4	11.2	23.3	18.1	7.1	12.4	18.1	14.9	13.7	9.2	14.0
744	Rapeseed meal.	14.5	22.5	25.9	8.7	66.3	10.0	11.6	24.4	18.3	9.4	13.0	19.2	15.1	13.4	9.0	14.0
745	Rapeseed meal.	12.6	18.6	22.5	7.5	53.7	8.8	10.2	21.2	16.0	7.7	11.4	16.2	13.0	12.1	8.3	12.4
746	Rapeseed meal.	14.5	22.8	26.7	8.6	66.3	9.8	11.5	24.5	18.3	8.1	13.2	19.2	15.1	13.7	9.2	14.3
747	Rapeseed meal.	15.0	22.9	26.8	8.8	68.2	10.3	11.9	24.9	18.2	8.1	13.8	19.0	15.3	13.7	9.4	14.6
641	Rice bran.	5.9	8.8	8.2	1.9	14.4	2.9	3.5	7.0	4.5	1.9	4.4	3.9	4.3	3.2	3.6	5.3
642	Rice bran.	7.0	11.0	8.6	1.8	14.4	3.0	3.6	6.7	4.7	2.7	4.8	4.3	4.9	3.6	3.5	5.8
647	Rice bran.	7.3	12.5	11.8	2.9	18.3	3.9	3.7	8.6	5.0	2.5	5.4	5.2	5.7	4.2	4.4	5.9
648	Rice bran.	7.4	7.9	10.6	2.3	16.3	3.4	3.6	8.4	5.7	2.6	5.2	5.1	5.9	4.4	4.1	5.7
644	Rice bran.	8.5	13.3	10.2	2.0	17.0	4.0	4.2	8.0	5.9	2.8	5.6	5.3	6.0	4.4	3.9	7.2
704	Screenings.	6.8	12.3	12.5	6.6	24.2	3.6	5.3	10.5	7.9	2.9	6.2	8.1	6.7	5.7	4.4	6.6
705	Screenings.	6.8	10.9	12.0	1.9	24.0	2.6	4.7	10.0	4.8	2.4	6.1	6.7	6.3	4.6	3.7	6.1
706	Screenings.	4.5	7.5	8.0	2.8	36.9	3.3	4.5	9.0	3.8	2.0	6.1	12.1	6.3	3.7	3.7	5.2
707	Screenings.	5.2	4.8	6.0	0.9	16.0	1.9	3.2	7.5	3.1	1.6	3.9	5.3	3.6	2.9	2.5	3.9
708	Screenings.	5.4	8.8	9.9	2.8	25.8	3.0	4.0	9.4	4.9	2.0	6.5	6.6	6.2	3.9	4.4	5.5
639	Sesame seed meal.	16.8	47.2	32.8	7.4	75.5	9.6	12.7	25.5	9.5	11.7	16.7	12.0	20.3	13.0	13.6	15.7
645	Sesame seed meal.	19.5	49.6	33.2	8.9	75.3	9.6	13.7	26.3	9.7	13.1	18.7	14.3	19.0	14.2	14.5	17.9
678	Shrimp meal.	21.1	23.3	34.6	3.8	46.3	9.1	14.8	23.1	20.9	8.6	132.3	16.2	19.9	16.5	23.5	18.6
840	Single cell protein.	46.1	31.3	63.9	3.6	66.0	12.3	28.6	48.0	39.5	15.0	23.6	19.6	21.8	30.0	20.5	34.2
671	Soybean meal.	19.5	37.0	57.3	6.5	88.4	12.6	18.7	35.9	28.2	7.0	23.9	24.0	25.2	18.1	16.7	19.4
672	Soybean meal.	20.1	36.2	57.6	6.6	89.9	12.7	18.6	36.4	28.2	7.3	23.9	24.3	25.4	18.2	16.6	19.2
673	Soybean meal.	24.6	44.0	69.4	7.6	106.9	15.4	24.4	45.3	35.6	8.2	29.6	29.5	31.2	22.6	20.7	27.0
723	Soybean meal.	19.9	36.2	58.0	6.5	91.8	13.6	18.1	37.0	30.7	7.0	23.7	24.5	26.5	18.6	16.3	18.1
724	Soybean meal.	20.1	36.5	57.7	7.4	91.8	13.4	19.6	37.4	30.7	7.7	23.8	24.1	26.7	18.6	16.4	18.9
748	Soybean meal.	19.4	34.8	57.6	7.3	91.3	12.9	16.2	35.4	29.5	7.7	22.5	24.0	25.8	17.8	15.8	15.3
749	Soybean meal.	17.8	30.1	54.1	7.3	84.9	11.7	14.6	32.8	26.2	7.5	20.7	21.7	23.7	16.5	14.4	14.7
750	Soybean meal.	20.4	35.7	59.7	8.4	94.6	13.1	16.9	36.5	30.5	8.0	23.3	24.9	27.1	18.7	16.6	17.4
751	Soybean meal.	19.3	32.0	57.8	7.9	92.0	12.4	16.5	35.3	28.4	7.1	22.6	23.5	26.1	18.1	15.9	16.9
799	Soybean meal.	22.6	38.6	61.2	7.0	98.4	13.7	21.0	41.2	32.0	7.7	26.4	27.4	28.0	20.3	18.3	22.0
800	Soybean meal.	22.8	40.2	62.4	7.3	103.9	14.1	21.3	42.0	33.0	7.8	27.4	28.4	29.4	20.4	19.2	21.6
801	Soybean meal.	21.9	38.0	58.7	7.3	96.7	13.4	20.3	39.8	31.5	7.9	26.7	27.5	19.6	18.6	21.0	21.0
802	Soybean meal.	21.7	41.5	62.6	7.1	98.3	14.2	21.4	40.5	32.2	7.9	26.3	27.2	28.4	19.6	18.8	21.7

TABLE 3. cont.

Code	Description	ALA	ARG	ASP	CYS	GLU	HIS	ILE	LEU	LYS	MET	PHE	PRO	SER	THR	TYR	VAL
803	Soybean meal.	21.3	40.7	60.9	7.4	96.7	13.6	20.6	39.1	32.2	7.9	25.6	26.3	27.7	19.8	18.1	21.1
804	Soybean meal.	22.3	42.4	63.9	7.6	100.7	14.3	21.1	40.7	32.8	8.2	26.3	27.5	29.1	20.6	19.0	21.5
823	Soybean meal.	21.8	42.8	60.4	6.6	96.2	14.0	21.3	40.0	31.9	7.5	26.1	26.6	27.1	19.2	18.3	21.7
824	Soybean meal.	20.9	38.9	59.1	8.4	93.1	13.2	21.1	38.5	29.3	8.5	25.1	25.7	26.3	19.0	16.9	21.7
825	Soybean meal.	23.4	45.3	65.4	7.6	104.4	14.8	23.1	43.1	34.2	7.9	28.1	28.8	29.8	20.8	19.1	24.0
826	Soybean meal.	22.4	43.0	63.5	7.8	99.5	14.1	22.0	41.1	32.6	7.6	26.7	27.6	28.4	20.0	18.7	22.4
679	Wheat.	5.7	8.2	9.9	4.3	60.7	4.5	5.5	12.4	4.6	2.7	8.9	19.8	8.5	4.9	4.6	5.9
680	Wheat.	4.7	7.6	8.1	3.2	50.8	4.1	4.6	9.8	4.6	2.1	7.2	15.9	6.6	3.9	4.1	5.4
681	Wheat.	5.3	8.0	9.0	4.2	57.4	4.3	5.1	11.6	4.4	3.1	8.5	17.4	8.1	4.6	4.6	5.8
682	Wheat.	4.1	6.2	6.8	3.8	49.1	3.5	4.1	9.4	3.1	2.5	6.5	4.9	6.2	3.4	3.5	4.4
683	Wheat.	4.5	6.1	7.4	3.6	42.4	3.4	3.9	9.2	3.6	2.4	6.1	13.6	6.3	3.7	3.6	4.3
684	Wheat.	4.1	6.8	7.1	3.0	49.8	3.8	3.9	9.1	4.0	2.0	6.9	16.1	6.4	3.5	3.8	4.4
685	Wheat.	5.4	8.0	9.3	4.2	59.7	4.4	5.6	12.0	4.4	2.1	8.6	19.0	8.0	4.4	4.6	6.1
686	Wheat.	4.2	6.8	7.2	3.4	47.1	3.8	4.2	9.4	4.3	1.8	6.6	15.3	6.8	3.8	3.6	5.0
687	Wheat.	5.4	7.8	9.7	4.2	57.5	4.3	5.4	11.8	4.4	1.0	8.5	18.7	8.0	4.5	4.8	5.7
688	Wheat.	4.7	7.8	8.4	4.1	53.6	3.9	4.9	10.6	3.9	2.6	7.4	17.6	7.3	4.0	4.1	5.0
773	Wheat.	5.1	7.6	8.0	4.3	56.5	4.1	5.0	11.2	4.3	1.9	8.4	19.5	7.8	4.2	4.3	6.1
778	Wheat.	5.1	7.6	8.0	4.1	57.2	4.2	5.1	11.5	4.4	2.4	8.7	19.8	8.1	4.4	4.6	6.0
784	Wheat.	4.4	6.7	5.8	2.9	109.8	3.6	4.3	9.9	3.1	2.1	7.6	17.5	6.4	3.6	4.2	5.4
779	Wheat.	4.5	6.9	7.1	3.6	50.5	4.4	10.2	4.4	4.1	2.1	7.5	17.2	7.2	3.8	4.1	5.2
782	Wheat.	5.3	7.3	7.7	3.5	56.6	3.8	4.9	11.0	4.1	2.2	8.2	18.4	7.3	4.2	4.6	6.0
783	Wheat.	5.8	7.9	8.6	4.0	116.3	4.1	5.1	11.5	4.6	2.6	8.4	19.0	7.7	4.8	4.8	6.3
772	Wheat.	4.3	6.6	6.8	3.7	44.2	3.5	4.1	9.5	3.8	2.3	6.7	15.4	6.3	3.8	3.9	5.2
781	Wheat.	4.7	6.7	7.0	3.6	47.7	3.6	4.3	9.9	4.0	2.3	7.1	16.3	7.1	4.1	4.2	5.1
776	Wheat.	6.1	9.3	9.3	4.5	64.0	4.7	5.9	13.5	5.0	3.4	10.3	23.5	9.5	5.1	5.8	7.2
780	Wheat.	5.3	7.6	8.4	4.1	55.2	4.1	4.9	11.3	4.4	2.5	8.3	18.9	7.8	4.4	4.8	5.8
676	Wheat shorts.	7.6	12.1	11.8	3.4	35.2	4.7	4.9	10.4	7.2	2.3	6.6	11.1	6.4	5.2	4.7	6.9
677	Wheat shorts.	9.1	14.6	14.3	3.3	37.7	5.4	5.6	11.8	8.7	2.5	7.4	12.1	7.6	5.9	5.3	8.6
728	Wheat shorts.	6.3	10.6	9.7	2.6	30.7	4.0	4.1	9.0	5.8	2.2	5.8	9.9	6.6	4.4	4.3	5.9
729	Wheat shorts.	7.1	11.9	39.9	3.2	32.4	4.4	4.3	9.7	6.5	2.2	6.1	10.5	7.4	5.2	4.7	6.5
730	Wheat shorts.	6.1	10.6	9.8	3.2	30.6	4.0	3.9	8.9	5.6	2.1	5.7	10.0	6.6	4.5	4.1	5.7
756	Wheat shorts.	7.0	10.7	11.0	3.3	34.1	4.3	4.0	9.8	6.4	2.1	6.1	11.4	7.4	5.0	4.6	6.0
757	Wheat shorts.	6.7	10.5	10.4	3.6	34.7	4.0	4.0	9.5	6.2	2.2	6.0	11.5	7.1	4.6	4.6	5.6
758	Wheat shorts.	6.0	9.3	34.0	3.3	29.2	3.7	3.2	8.0	5.3	2.5	5.2	9.5	6.1	4.1	3.9	4.7
809	Wheat shorts.	7.2	12.1	10.2	2.9	32.1	4.1	4.6	10.3	5.6	2.7	6.5	10.9	5.9	4.3	4.6	6.8
810	Wheat shorts.	7.8	12.4	12.5	2.6	36.2	4.6	5.3	11.5	6.5	2.4	7.0	12.0	7.3	4.9	5.1	7.2
811	Wheat shorts.	7.5	13.0	11.4	2.9	33.2	4.4	5.0	10.3	6.6	2.8	6.6	11.3	6.9	5.1	4.6	7.5
832	Wheat shorts.	7.8	14.2	11.7	2.4	36.0	4.8	5.0	10.6	7.4	2.5	6.9	12.1	7.5	5.2	5.0	7.0

TABLE 3 cont.

Code	Description	ALA	ARG	ASP	CYS	GLU	HIS	ILE	LEU	LYS	MET	PHE	PRO	SER	THR	TYR	VAL
833	Wheat shorts.	8.2	14.4	11.9	3.5	38.4	4.8	5.0	11.1	7.6	2.5	7.2	13.1	8.6	5.6	5.3	7.4
834	Wheat shorts.	8.2	14.8	12.3	3.3	36.8	4.9	5.0	11.0	7.7	2.7	7.2	12.6	8.0	5.4	5.2	7.5
835	Wheat shorts.	8.2	14.8	12.0	3.1	37.6	4.8	5.1	11.2	7.8	2.6	7.2	12.3	8.0	5.6	5.2	7.4
638	Yucca.	4.2	11.6	ND	ND	11.6	3.9	ND	6.2	7.9	ND	ND	ND	5.4	3.8	ND	ND
640	Yucca.	0.5	1.3	0.5	ND	2.1	0.4	ND	ND	ND	ND	ND	ND	0.3	0.2	ND	ND

1 Blanks (-) = variable not measured; ND = variable not detected in a reportable amount.

TABLE 4

Bioavailability of amino acids in the dry matter of several feedingstuffs (%).^{1,2}

Code	Description	ALA	ARG	ASP	CYS	GLU	HIS	ILE	LEU	LYS	MET	PHE	PRO	SER	THR	TYR	VAL
655	Alfalfa.	66.1	76.6	77.0	30.1	66.6	63.4	72.1	73.9	61.2	77.4	75.3	83.7	64.8	64.3	66.5	72.0
656	Alfalfa.	81.3	87.7	84.0	-	82.0	74.6	85.0	86.4	54.8	-	85.7	86.4	76.9	78.9	82.8	81.3
653	Alfalfa.	81.1	86.6	84.1	44.5	82.7	73.4	83.8	84.5	77.5	83.1	84.3	85.9	75.6	76.8	79.4	81.8
839	Alfalfa.	74.1	70.5	71.7	54.1	62.3	68.4	68.4	70.5	57.7	62.1	71.1	66.6	60.3	62.0	64.7	65.5
657	Alfalfa.	72.8	83.2	82.5	27.0	74.4	80.2	76.4	78.9	51.5	73.3	78.8	82.4	73.5	75.2	73.5	76.3
652	Alfalfa.	80.6	91.0	84.5	40.2	83.7	83.8	83.3	86.2	61.2	79.2	81.8	89.9	78.7	80.9	85.3	83.4
654	Alfalfa.	71.3	79.1	80.7	51.4	71.4	74.2	72.6	76.5	46.1	68.2	76.0	74.4	67.5	69.3	71.3	72.0
649	Algae.	78.1	82.1	75.1	82.6	76.2	77.8	76.2	77.4	78.2	78.0	78.3	75.8	70.3	73.6	76.5	76.3
650	Algae.	86.7	92.1	85.7	83.8	87.7	86.9	86.2	88.3	86.8	88.7	88.4	84.4	81.2	82.3	87.1	85.8
689	Barley.	72.4	84.4	76.9	75.7	91.0	89.3	82.1	83.8	76.7	84.1	85.2	89.2	78.3	78.5	81.8	81.8
690	Barley.	75.7	85.5	78.2	95.3	92.1	87.7	83.9	88.2	74.9	81.1	89.1	94.0	88.0	78.6	88.5	81.6
691	Barley.	65.7	81.8	72.5	84.6	87.4	80.2	75.5	81.9	67.6	80.2	81.3	87.4	78.7	64.8	79.5	70.5
692	Barley.	73.2	83.8	75.9	88.9	91.0	85.7	80.6	85.9	72.6	79.6	87.3	92.4	82.7	75.0	85.1	78.8
693	Barley.	75.0	85.4	78.8	82.2	90.2	87.2	81.1	86.2	73.7	74.8	85.2	89.2	83.7	75.9	85.4	79.8
694	Barley.	79.8	90.7	85.7	97.8	93.5	90.4	87.4	91.5	82.2	82.8	91.4	95.1	94.0	81.9	94.0	85.4
695	Barley.	77.2	87.7	81.6	88.3	92.3	86.6	86.3	88.5	77.7	85.7	89.3	91.8	85.5	77.2	89.2	82.7
696	Barley.	76.4	87.1	80.7	91.3	92.4	89.4	83.6	88.1	78.1	29.5	89.2	93.5	90.4	79.6	90.0	82.7
697	Barley.	80.0	90.3	82.7	95.4	93.4	90.4	87.6	91.1	83.3	90.8	91.3	94.1	89.9	80.7	93.5	86.2
698	Barley.	78.2	90.9	82.8	80.6	93.9	92.7	86.1	88.0	81.5	83.8	91.8	94.0	83.3	81.9	88.3	83.8
769	Barley.	68.1	81.1	70.4	75.8	89.3	85.4	76.4	82.5	73.2	72.7	85.9	91.8	81.2	73.4	81.7	77.0
765	Barley.	76.4	83.3	77.3	70.0	90.4	87.1	82.3	85.5	84.2	80.1	86.9	84.8	79.5	76.0	86.0	83.5
762	Barley.	76.1	82.6	78.0	77.4	91.4	86.1	82.2	86.2	83.4	83.6	87.7	91.9	83.0	78.4	86.2	81.3
775	Barley.	71.1	79.5	72.5	71.5	87.7	79.3	76.1	81.6	78.0	77.3	84.5	89.2	76.7	71.2	81.0	78.8
777	Barley.	75.3	84.8	74.1	81.9	92.4	79.0	83.2	87.3	72.6	83.1	90.2	94.1	85.4	75.3	85.9	83.3
767	Barley.	63.5	73.4	67.9	62.3	84.8	82.0	71.2	76.5	73.7	74.9	78.6	83.0	64.9	67.2	76.3	69.3
768	Barley.	71.7	83.6	73.2	78.4	89.7	86.2	78.2	84.9	78.5	78.3	86.0	91.5	81.4	73.4	79.6	79.4
761	Barley.	70.5	79.4	74.4	86.3	89.4	85.6	74.3	82.3	75.8	76.1	84.4	91.8	83.5	76.5	82.5	75.4
763	Barley.	74.9	82.9	75.6	72.2	90.6	85.8	80.6	84.4	79.5	76.2	86.3	91.4	82.2	76.0	83.7	80.7
766	Barley.	80.4	85.2	79.6	73.5	92.9	88.9	85.3	88.1	86.1	80.4	90.4	92.5	81.5	78.6	87.3	83.6
666	Blood meal.	92.3	88.2	90.3	82.2	86.8	86.8	87.9	91.7	86.3	92.9	92.9	87.7	89.8	89.7	91.6	91.0
667	Blood meal.	95.4	91.2	94.6	81.4	90.3	91.5	84.3	94.6	88.9	95.6	95.4	92.1	94.3	93.3	93.9	94.6
717	Blood meal.	81.5	77.7	77.3	65.4	76.1	77.1	63.8	79.7	76.2	83.1	81.3	78.8	80.7	78.5	81.3	76.1
718	Blood meal.	92.3	87.3	88.9	74.9	87.3	88.6	81.3	91.1	87.3	91.5	92.3	85.0	88.2	87.5	90.2	89.4
719	Blood meal.	93.8	91.0	92.7	84.8	92.0	91.5	85.3	92.5	91.0	94.9	93.3	92.1	93.5	92.5	93.9	91.4
739	Blood meal.	86.4	80.4	84.4	76.3	80.0	81.6	67.7	85.3	76.3	87.9	86.7	85.8	85.9	83.4	86.0	82.3

TABLE 4 cont.

Code	Description	ALA	ARG	ASP	CYS	GLU	HIS	ILE	LEU	LYS	MET	PHE	PRO	SER	THR	TYR	VAL
740	Blood meal.	92.8	88.1	89.3	75.8	88.1	82.2	91.0	89.9	92.6	92.5	86.8	89.8	88.1	91.5	89.4	
741	Blood meal.	89.3	84.7	86.8	79.4	84.9	84.3	80.6	88.3	91.8	89.2	89.1	89.4	87.6	91.3	85.9	
742	Blood meal.	93.9	84.1	92.3	78.3	87.6	88.3	78.1	93.0	81.7	94.1	94.1	93.0	92.4	90.5	91.9	
792	Blood meal.	94.4	84.1	92.3	72.7	88.8	88.5	76.3	93.4	89.3	94.9	94.5	91.4	91.8	90.5	92.8	
793	Blood meal.	93.5	89.2	90.8	78.0	89.1	88.5	84.2	92.2	89.5	92.2	93.5	88.2	90.9	89.3	92.3	
794	Blood meal.	89.4	85.3	86.3	67.4	85.5	86.3	71.1	87.9	86.0	90.2	89.4	85.1	87.0	85.2	86.7	
795	Blood meal.	87.3	83.0	84.2	70.4	84.6	71.1	86.5	82.3	87.5	88.1	85.0	86.4	84.1	87.8	83.2	
817	Blood meal.	93.8	87.8	90.8	74.7	89.1	90.0	85.0	92.5	88.8	94.7	93.7	88.3	91.0	89.9	92.7	
643	Coconut meal.	78.0	86.5	75.8	52.9	80.2	74.8	83.6	80.0	79.7	88.1	86.2	74.1	74.1	65.5	76.7	
838	Corn.	97.0	95.9	97.1	97.0	98.1	96.8	99.5	103.1	100.5	97.8	97.1	96.8	93.5	98.3	97.6	
853	Corn.	94.2	109.7	91.8	89.5	97.3	99.7	90.5	96.6	105.9	92.9	96.8	97.6	96.5	92.4	97.2	
855	Corn.	92.2	100.5	88.5	80.5	94.9	98.6	86.8	94.2	107.8	92.6	92.4	94.1	87.4	85.0	93.0	
856	Corn.	95.4	106.8	94.5	93.8	97.6	102.3	92.3	97.2	110.1	91.3	96.2	97.4	96.7	92.7	98.3	
857	Corn.	96.2	107.0	94.7	94.7	98.2	103.6	93.6	97.3	113.1	94.5	96.8	98.5	96.6	95.1	97.3	
858	Corn.	92.9	103.8	89.4	86.8	95.7	99.9	95.7	95.4	105.3	91.1	93.1	95.7	91.6	87.1	94.3	
859	Corn.	90.5	100.0	82.2	83.8	92.9	96.4	84.9	93.9	93.0	86.8	91.1	94.0	87.2	83.6	87.4	
860	Corn.	95.2	99.7	93.6	98.9	98.3	102.3	95.2	98.0	106.2	95.0	96.8	99.5	99.1	95.1	98.9	
861	Corn.	94.6	97.6	90.7	90.7	96.4	97.1	89.7	97.0	93.7	93.3	96.2	96.7	95.5	89.5	89.6	
862	Corn.	92.3	95.8	86.3	87.9	95.2	96.8	90.9	96.1	87.5	91.6	94.4	96.2	91.8	83.7	94.7	
646	Corn germ meal.	85.8	90.3	77.5	57.8	86.8	83.7	86.2	87.2	83.0	85.6	88.0	81.5	80.3	78.6	83.5	
651	Corn gluten feed.	91.3	89.1	82.1	77.3	91.7	86.7	87.3	93.1	75.1	89.4	91.4	90.4	87.2	81.1	90.2	
674	Corn gluten feed.	91.8	92.8	85.9	73.1	92.0	89.1	90.3	93.2	80.9	87.7	91.5	88.7	88.7	84.6	90.2	
726	Corn gluten feed.	81.6	85.9	72.8	61.9	83.7	81.2	77.1	86.3	65.8	78.0	83.0	83.6	81.0	70.9	79.1	
727	Corn gluten feed.	81.5	86.0	70.9	62.3	83.6	81.5	76.7	86.1	65.6	79.6	82.8	82.8	80.2	72.1	81.4	
754	Corn gluten feed.	85.9	86.3	77.2	67.4	86.7	83.3	80.0	89.2	68.7	85.0	85.6	86.0	81.9	75.3	86.9	
755	Corn gluten feed.	86.1	85.3	76.5	65.1	85.7	81.9	81.0	88.9	72.6	85.4	86.7	84.1	80.6	73.6	84.7	
805	Corn gluten feed.	89.7	91.8	81.8	61.3	90.8	87.5	89.4	92.9	74.3	88.2	90.8	89.0	87.7	81.5	89.8	
806	Corn gluten feed.	84.9	87.8	75.4	54.0	86.2	82.8	83.0	88.6	71.2	84.6	85.6	83.5	80.7	75.1	83.5	
827	Corn gluten feed.	88.4	90.5	80.7	59.9	89.1	87.6	86.0	90.8	69.4	85.1	87.8	87.2	85.5	78.8	87.2	
828	Corn gluten feed.	86.0	92.1	75.5	62.0	85.9	84.8	78.3	88.9	75.8	84.9	85.8	83.4	80.6	75.5	84.9	
675	Corn gluten meal.	98.9	98.3	96.9	90.6	98.8	96.4	98.1	99.1	94.8	98.5	98.9	98.2	97.5	95.5	98.4	
725	Corn gluten meal.	97.1	94.3	93.7	91.5	97.5	95.0	94.7	98.0	84.9	97.4	97.0	95.7	92.2	96.6	94.1	
752	Corn gluten meal.	97.4	92.4	94.2	89.0	97.5	94.2	94.9	98.0	88.5	97.5	97.3	96.4	95.5	92.7	97.0	
753	Corn gluten meal.	97.7	95.6	95.0	88.9	96.3	94.2	95.7	97.8	92.7	97.9	97.8	97.4	96.3	93.9	97.6	
807	Corn gluten meal.	97.5	94.4	93.0	82.2	97.3	95.8	98.1	85.6	97.1	97.4	96.9	95.4	91.9	97.1	95.5	
808	Corn gluten meal.	97.2	93.8	92.6	80.8	97.2	93.9	95.8	98.0	85.7	96.2	97.3	97.0	95.3	91.6	96.8	

TABLE 4 cont.

Code	Description	ALA	ARG	ASP	CYS	GLU	HIS	ILE	LEU	LYS	MET	PHE	PRO	SER	THR	TYR	VAL
829	Corn gluten meal.	97.2	97.8	93.3	85.0	97.1	95.0	94.9	97.8	88.6	96.2	97.2	97.1	95.1	91.8	96.8	94.8
830	Corn gluten meal.	96.3	97.4	92.3	84.5	96.2	94.3	93.4	96.9	86.5	96.5	96.2	96.3	95.0	91.4	96.0	93.3
831	Corn gluten meal.	97.1	98.2	93.9	85.5	97.4	94.9	94.8	97.7	89.0	96.4	97.0	97.3	95.8	93.1	96.8	94.7
663	Feather meal.	85.5	87.8	68.0	58.6	74.3	79.5	89.7	87.7	72.3	83.9	90.4	83.2	81.9	77.3	84.5	85.3
664	Feather meal.	81.3	83.5	54.7	47.1	63.9	68.0	85.9	83.5	58.7	74.5	87.0	79.5	82.9	70.4	78.8	82.6
665	Feather meal.	79.7	84.4	54.4	51.6	64.7	69.3	84.9	85.8	57.9	79.6	84.1	78.5	77.1	70.1	78.5	80.8
666	Feather meal.	75.4	77.5	54.1	50.5	63.0	71.7	79.7	78.9	61.4	73.7	82.7	63.6	73.7	66.7	74.3	74.5
714	Feather meal.	81.4	82.5	56.6	58.9	69.1	69.9	85.3	82.7	62.5	71.7	85.7	71.2	79.5	71.7	78.6	81.1
715	Feather meal.	77.7	82.8	55.8	51.2	68.3	70.4	82.9	80.2	65.4	72.1	82.5	69.6	78.2	70.1	77.3	78.0
716	Feather meal.	77.5	79.4	58.4	55.8	66.1	75.6	80.1	80.1	66.6	73.3	83.5	65.5	75.4	68.7	77.2	75.7
737	Feather meal.	80.2	83.1	58.4	66.4	70.0	70.4	85.2	82.3	63.9	76.2	85.4	70.6	79.3	73.2	79.2	80.4
738	Feather meal.	75.6	80.0	55.5	50.3	65.2	73.5	81.3	79.7	63.4	71.9	83.6	66.3	76.0	67.4	76.5	76.4
790	Feather meal.	84.8	88.1	68.6	65.0	77.0	75.3	88.2	86.6	72.7	78.0	89.4	78.4	84.0	79.4	84.7	85.0
791	Feather meal.	78.0	81.4	55.4	41.0	65.8	74.1	82.9	80.9	62.7	80.3	84.5	65.2	74.8	67.7	77.4	78.3
815	Feather meal.	84.8	86.6	64.2	60.9	74.2	77.2	89.7	86.6	67.7	73.9	89.3	76.0	82.7	82.9	85.2	85.2
816	Feather meal.	90.0	90.1	59.8	87.1	83.5	88.9	90.1	84.5	92.6	89.6	92.6	87.8	85.6	85.2	87.7	88.2
662	Fish meal.	89.6	91.3	86.5	71.5	90.5	88.4	89.6	91.3	88.3	93.0	89.7	89.1	89.5	89.7	91.5	88.6
712	Fish meal.	92.9	91.6	90.2	77.9	93.1	89.4	92.6	93.9	90.8	95.2	92.8	93.1	92.7	92.3	93.0	92.3
713	Fish meal.	92.6	89.6	91.2	77.3	93.3	88.9	92.8	94.5	88.4	95.2	93.2	90.6	91.6	91.6	93.8	92.9
734	Fish meal.	85.2	85.9	83.5	72.8	85.2	82.6	83.2	86.0	83.3	86.2	84.6	84.3	85.9	85.8	86.6	83.6
735	Fish meal.	90.5	91.6	88.2	78.7	90.3	87.8	90.3	91.7	89.8	92.5	90.3	87.7	90.9	91.0	91.1	89.3
736	Fish meal.	93.9	94.5	90.8	80.0	94.5	90.7	94.5	95.6	94.2	95.9	94.5	93.7	93.5	93.6	95.4	94.3
788	Fish meal.	94.4	92.1	91.2	81.1	94.4	90.3	94.3	95.3	89.4	96.1	94.8	94.0	93.5	93.1	95.1	94.2
789	Fish meal.	91.3	90.2	87.7	66.0	91.2	86.5	92.1	93.0	84.2	93.8	91.7	88.8	89.2	88.5	92.3	91.8
814	Fish meal.	91.4	94.2	88.1	76.1	92.9	90.5	94.3	94.2	89.8	94.1	93.1	90.0	90.3	91.4	93.9	93.9
620	Fish meal.	91.2	93.9	88.5	81.0	92.4	89.5	94.6	94.4	87.5	94.5	93.2	89.4	90.2	91.0	94.2	93.5
621	Fish meal.	93.0	95.2	90.5	84.1	94.4	90.4	95.2	95.4	90.7	92.7	94.4	92.3	92.8	92.7	95.6	94.0
622	Fish meal.	91.6	93.8	90.0	85.0	93.7	93.7	95.1	95.4	92.4	93.3	94.2	88.8	91.9	92.4	95.1	94.2
623	Fish meal.	91.6	94.2	89.7	82.4	93.1	93.1	95.0	95.1	91.4	92.3	94.2	89.1	91.4	92.7	95.1	94.0
624	Fish meal.	91.8	93.8	89.9	81.2	93.4	93.4	95.2	95.3	91.0	91.0	94.2	89.2	91.0	92.7	94.9	94.4
626	Fish meal.	90.5	94.0	86.0	74.5	91.0	88.6	94.0	93.8	86.6	89.8	92.6	90.5	89.4	90.1	94.0	93.4
627	Fish meal.	92.3	95.0	85.4	77.5	93.0	90.1	94.6	94.3	91.2	90.1	93.0	89.9	89.8	90.5	94.9	94.0
628	Fish meal.	93.7	95.8	88.1	76.6	94.1	91.1	95.2	95.0	91.2	90.4	94.2	93.0	91.2	92.4	95.2	94.9
629	Fish meal.	92.5	94.6	90.1	79.9	93.3	90.7	94.2	94.6	92.0	91.3	93.9	91.2	91.4	91.8	94.4	93.6
630	Fish meal.	93.5	95.9	90.0	80.5	93.8	89.6	94.6	94.9	88.7	91.9	94.0	92.7	92.0	92.7	95.1	94.3
631	Fish meal.	92.1	94.1	90.6	76.2	93.3	89.8	94.2	94.4	87.6	90.2	93.4	90.8	91.3	92.6	94.5	93.2
632	Fish meal.	93.0	94.5	89.0	75.2	93.1	91.1	93.1	93.8	88.3	89.2	92.8	92.7	90.3	91.6	93.3	93.3
633	Fish meal.	94.2	96.1	90.0	77.8	94.2	92.8	93.9	94.3	91.6	90.1	93.3	93.2	90.9	92.2	94.5	93.7
634	Fish meal.	94.1	95.8	90.5	77.4	94.1	91.3	94.5	94.5	86.8	89.8	93.5	94.4	91.9	92.6	94.4	94.0

TABLE 4 cont.

Code	Description	ALA	ARG	ASP	CYS	GLU	HIS	ILE	LEU	LYS	MET	PHE	PRO	SER	THR	TYR	VAL
658	Meat meal.	89.2	92.3	82.0	61.5	88.1	87.6	90.8	91.1	84.6	91.7	91.2	88.1	86.4	86.9	90.8	90.6
659	Meat meal.	77.1	79.3	69.6	30.5	76.4	82.6	78.1	81.7	73.2	81.4	82.0	75.2	73.5	76.2	80.6	80.0
660	Meat meal.	84.8	86.8	82.8	56.9	84.4	86.3	84.3	86.7	83.0	89.4	86.3	82.7	80.6	83.8	84.4	84.6
661	Meat meal.	86.7	89.8	81.3	52.8	86.5	87.7	88.8	89.5	85.0	90.2	89.3	85.3	83.6	86.8	89.7	88.5
709	Meat meal.	82.2	86.0	72.7	37.7	80.9	80.5	81.3	84.1	77.0	87.2	83.5	79.2	78.6	80.0	83.8	80.0
710	Meat meal.	85.6	86.8	79.3	50.6	84.6	86.5	83.5	87.7	81.6	89.0	87.3	83.7	83.9	84.1	86.8	85.4
711	Meat meal.	83.2	84.4	83.2	64.0	84.9	86.4	84.4	87.4	84.6	89.7	86.3	81.2	83.3	84.3	86.6	83.9
731	Meat meal.	85.4	87.3	79.4	59.9	85.1	85.8	85.4	87.7	83.6	89.1	87.2	82.7	83.1	83.7	86.6	84.7
732	Meat meal.	83.3	82.2	81.1	53.3	83.6	84.3	82.2	85.9	82.0	87.8	85.0	81.0	80.7	82.7	84.4	82.8
733	Meat meal.	85.4	85.4	82.9	60.1	81.9	86.9	83.7	86.9	85.1	89.6	86.7	83.1	81.2	83.1	85.0	83.1
785	Meat meal.	88.2	89.7	83.4	56.2	87.4	88.5	87.5	90.0	85.1	89.1	89.3	87.4	86.2	86.3	88.6	87.6
786	Meat meal.	89.0	89.8	83.4	59.0	88.0	88.7	88.6	90.8	86.3	90.2	90.1	87.7	87.0	87.9	90.4	89.8
787	Meat meal.	86.0	86.2	84.2	59.3	86.1	86.2	86.3	88.7	84.8	90.5	87.3	84.1	83.7	84.8	88.2	86.9
812	Meat meal.	30.2	62.9	30.3	30.0	71.2	73.3	59.8	61.3	44.8	73.7	64.7	30.3	44.3	48.0	66.1	52.2
813	Meat meal.	89.3	87.7	86.1	54.0	87.3	89.9	86.2	88.2	86.3	90.8	88.1	89.2	85.1	84.5	87.3	88.1
699	Oats.	83.9	94.5	84.9	86.2	93.8	96.0	90.1	91.8	85.0	77.1	94.8	95.6	90.5	86.0	95.5	86.3
700	Oats.	88.8	98.7	89.1	95.1	96.2	96.8	93.0	95.0	88.6	89.7	97.4	95.8	94.1	90.2	100.8	91.1
701	Oats.	81.7	94.2	86.6	91.1	92.4	92.4	88.4	90.0	85.5	77.7	92.6	90.1	86.1	83.3	93.4	85.9
702	Oats.	89.9	99.3	94.7	95.1	98.2	101.7	94.3	95.6	94.5	89.5	97.4	95.1	94.1	93.2	99.1	92.4
703	Oats.	91.1	98.5	92.6	97.8	97.7	99.6	94.9	96.1	92.9	90.3	97.3	98.0	96.3	92.6	100.3	94.1
764	Oats.	77.2	88.3	81.7	65.6	90.1	88.6	83.8	86.2	82.1	83.3	88.3	83.2	81.0	76.7	85.9	80.4
774	Oats.	84.3	92.3	84.7	74.9	92.2	87.6	85.4	90.1	85.8	89.2	92.7	83.2	85.1	79.4	89.5	87.1
770	Oats.	78.7	92.9	80.9	77.0	90.5	89.0	83.3	88.5	82.4	80.3	91.5	89.0	87.1	79.0	88.3	83.9
760	Oats.	85.5	93.1	88.2	81.4	93.4	92.6	87.2	90.5	90.5	92.3	91.3	89.4	88.0	84.9	92.0	85.8
759	Oats.	84.3	91.4	85.6	83.0	93.2	85.7	87.2	91.2	83.6	88.1	92.1	90.7	89.0	81.9	93.7	88.0
771	Oats.	86.9	93.3	85.7	78.7	94.9	92.3	89.2	91.7	85.9	93.5	92.9	93.4	92.9	89.7	92.9	90.8
796	Rapeseed meal.	86.7	91.3	83.0	73.0	92.0	88.7	84.6	88.7	80.8	90.0	88.7	83.4	84.5	81.1	86.4	83.2
797	Rapeseed meal.	85.6	90.3	81.3	64.7	90.8	87.7	85.6	87.9	79.4	90.2	87.9	82.0	81.6	80.6	85.2	83.5
798	Rapeseed meal.	85.4	90.1	80.8	65.5	91.2	87.9	85.0	88.2	79.9	89.5	87.9	82.2	81.4	80.8	84.6	84.4
818	Rapeseed meal.	87.7	91.4	84.6	83.9	92.5	88.7	86.6	89.9	81.3	91.6	89.9	84.0	84.8	82.4	87.2	84.3
819	Rapeseed meal.	84.7	89.1	80.9	57.1	90.4	86.8	84.7	87.5	79.1	89.0	88.1	80.5	80.0	79.9	85.1	83.1
820	Rapeseed meal.	88.1	90.7	85.2	83.1	92.7	89.3	86.8	90.0	80.4	92.2	89.7	85.4	85.5	83.3	87.6	85.8
821	Rapeseed meal.	83.7	89.5	80.1	86.7	89.9	88.0	84.2	87.8	80.9	93.9	86.2	79.9	81.1	78.6	83.7	82.3
822	Rapeseed meal.	87.3	90.8	83.0	71.8	88.7	86.4	89.6	81.9	92.3	89.9	83.4	84.1	82.4	85.5	85.5	83.2
668	Rapeseed meal.	88.9	91.8	85.4	60.7	93.1	88.6	88.8	90.3	83.5	91.1	90.3	86.3	86.0	83.4	87.7	87.7
669	Rapeseed meal.	79.8	86.0	78.7	71.1	87.5	84.9	80.4	82.4	74.1	88.2	76.9	74.6	75.1	78.2	79.3	83.7
670	Rapeseed meal.	85.0	89.5	81.4	72.2	90.5	86.0	85.1	86.6	81.3	88.0	86.6	81.8	80.5	79.9	84.1	83.7
720	Rapeseed meal.	83.6	89.5	82.0	76.6	90.8	86.8	81.2	86.2	79.5	89.0	85.1	81.8	83.2	79.3	80.7	83.3

TABLE 4 cont.

Code	Description	ALA	ARG	ASP	CYS	GLU	HIS	TLE	LEU	LYS	MET	PHE	PRO	SER	THR	TYR	VAL
721	Rapeseed meal.	84.5	88.2	82.6	74.9	91.3	88.4	82.6	88.1	79.9	88.2	86.4	82.3	83.7	79.9	83.7	80.8
722	Rapeseed meal.	83.9	89.9	81.9	73.8	90.9	88.0	82.0	86.9	80.3	87.7	86.2	82.0	83.2	80.6	82.9	79.4
743	Rapeseed meal.	86.5	90.3	84.5	74.2	92.3	88.6	84.7	89.0	81.8	90.6	88.1	82.8	85.4	81.7	85.6	83.0
744	Rapeseed meal.	84.9	89.3	83.2	76.2	91.6	88.0	83.5	87.9	81.1	90.9	87.0	83.3	84.7	79.6	83.9	81.8
745	Rapeseed meal.	82.4	89.4	80.2	79.8	90.3	86.9	81.8	86.7	78.0	93.3	86.3	79.6	81.2	77.7	82.3	79.6
746	Rapeseed meal.	85.5	90.8	84.7	79.0	92.3	88.2	83.9	88.6	81.6	91.8	87.9	83.0	84.5	81.9	85.1	83.1
747	Rapeseed meal.	84.8	89.1	82.7	77.6	90.8	88.7	83.1	87.2	78.4	90.2	87.1	81.3	83.4	79.4	84.7	82.2
641	Rice bran.	74.7	84.8	70.0	61.3	80.8	82.5	77.2	75.1	73.4	75.5	76.7	68.7	68.2	65.6	77.5	76.5
642	Rice bran.	73.5	84.8	64.9	55.3	77.0	73.9	72.5	67.4	66.8	77.7	70.6	63.5	68.2	63.1	74.6	72.8
647	Rice bran.	80.3	91.4	80.2	76.6	85.8	86.6	80.7	81.3	79.7	79.9	82.4	80.4	80.9	77.7	84.1	80.8
648	Rice bran.	76.9	82.7	72.1	63.6	79.3	81.8	74.8	75.0	80.7	72.6	75.3	73.6	74.9	71.5	77.3	74.4
644	Rice bran.	78.0	87.3	68.7	57.1	82.5	84.2	75.9	72.1	71.2	79.1	74.3	70.3	73.8	69.1	78.4	78.8
704	Screenings.	78.7	84.1	78.5	81.8	83.3	78.7	77.5	79.9	76.0	77.9	80.5	80.6	79.1	76.8	81.1	76.1
705	Screenings.	83.8	91.6	84.6	74.1	88.8	76.6	81.5	86.8	75.0	81.4	88.1	85.8	84.5	77.8	88.4	83.9
706	Screenings.	85.6	92.3	93.2	94.3	94.6	91.1	92.3	85.6	87.0	90.2	91.7	94.0	87.9	91.7	88.5	
707	Screenings.	78.7	83.0	71.5	49.0	85.9	81.9	80.0	82.6	71.2	78.2	82.3	77.1	71.9	70.3	82.4	77.5
708	Screenings.	88.2	96.0	88.7	73.2	95.7	95.8	91.7	93.9	88.7	87.7	95.4	92.4	92.6	86.1	94.6	92.2
639	Sesame seed meal.	86.2	90.6	85.7	78.2	92.9	87.2	90.1	89.6	84.0	92.1	91.3	86.9	89.5	84.8	90.5	88.7
645	Sesame seed meal.	92.3	94.2	90.8	85.2	95.0	91.4	93.8	92.6	91.2	96.1	95.1	92.0	91.7	89.1	94.0	93.2
678	Shrimp meal.	89.9	93.5	87.4	78.6	88.3	91.0	95.1	95.2	90.0	96.2	99.1	89.3	87.2	91.1	93.9	93.5
840	Single cell protein.	90.9	94.6	89.5	66.9	91.1	90.1	92.4	91.8	89.0	88.7	92.9	88.2	87.6	88.6	91.5	90.0
671	Soybean meal.	88.1	93.8	92.7	83.9	94.3	92.6	92.0	91.7	88.0	92.0	92.6	92.3	89.5	92.8	90.5	
672	Soybean meal.	89.6	92.8	93.3	85.1	94.7	93.0	92.9	92.6	87.7	93.8	93.6	93.0	92.3	89.9	93.1	91.5
673	Soybean meal.	91.6	94.0	94.1	87.4	95.7	93.6	94.8	94.1	92.2	94.0	95.0	94.8	94.0	91.2	94.9	94.2
723	Soybean meal.	87.3	90.7	91.7	81.9	93.6	92.7	89.6	90.9	89.9	91.7	93.3	92.0	87.9	91.9	88.0	
724	Soybean meal.	90.6	93.5	93.6	86.9	95.6	93.9	93.0	93.8	91.8	93.2	94.3	95.1	94.8	91.1	94.5	91.9
748	Soybean meal.	88.3	89.9	90.9	81.8	93.1	91.0	89.1	90.5	89.3	91.6	91.3	91.2	90.2	86.5	91.7	86.4
749	Soybean meal.	87.1	85.9	91.2	81.3	93.3	88.9	87.7	90.1	84.7	92.3	90.9	90.5	89.1	85.8	91.1	85.6
750	Soybean meal.	91.6	92.3	93.6	88.7	95.4	92.8	92.5	93.4	92.2	94.7	93.8	93.6	93.1	91.2	94.2	92.3
751	Soybean meal.	90.2	87.4	93.5	87.8	95.0	90.7	91.6	92.5	87.8	92.6	93.0	93.4	92.7	90.3	93.5	89.9
799	Soybean meal.	90.7	90.4	92.6	78.4	94.7	91.9	93.0	93.2	89.8	92.2	94.0	94.4	93.3	90.3	94.0	92.3
800	Soybean meal.	89.9	90.0	91.4	77.8	94.3	90.9	92.1	92.1	88.6	92.3	93.1	92.0	88.1	92.9	90.9	
801	Soybean meal.	89.7	90.4	91.5	78.2	94.2	91.2	92.2	92.1	89.1	92.0	93.5	92.4	88.7	93.7	90.9	
802	Soybean meal.	89.0	92.7	92.2	77.9	94.5	93.1	92.6	90.0	91.3	92.6	93.7	94.6	92.8	88.6	93.8	92.2

TABLE 4 cont.

Code	Description	ALA	ARG	ASP	CYS	GLU	HIS	ILE	LEU	LYS	MET	PHE	PRO	SER	THR	TYR	VAL
803	Soybean meal.	89.3	91.3	91.4	77.8	94.1	91.9	92.1	91.9	90.9	92.8	93.0	92.6	92.1	88.6	93.4	91.0
804	Soybean meal.	91.2	92.0	93.1	79.0	95.6	93.4	93.8	93.7	90.8	94.4	94.9	93.7	90.8	94.6	92.6	92.6
823	Soybean meal.	89.3	93.1	91.6	75.0	93.6	92.8	91.6	91.5	89.3	92.9	92.8	91.3	88.0	93.2	90.3	90.3
824	Soybean meal.	92.1	90.6	93.8	87.6	95.4	92.7	94.3	94.1	86.3	95.0	94.7	95.7	94.1	92.3	94.7	93.8
825	Soybean meal.	90.4	92.2	92.7	79.2	94.6	92.0	92.8	92.7	89.4	93.6	93.6	93.8	93.1	90.1	93.9	92.2
826	Soybean meal.	90.8	91.9	92.7	84.2	94.8	92.5	93.1	92.9	89.7	93.5	93.9	94.3	92.9	89.8	93.9	92.7
679	Wheat.	87.4	95.8	88.1	95.3	97.9	94.6	92.0	95.1	82.9	89.3	95.7	98.7	96.2	88.9	95.3	90.3
680	Wheat.	79.5	87.8	84.5	84.3	95.9	94.3	87.9	88.5	80.2	88.9	90.4	93.6	84.5	81.0	85.9	85.5
681	Wheat.	86.0	91.8	86.7	92.2	97.5	93.2	90.0	93.7	91.6	94.7	97.6	93.6	86.9	94.6	88.6	88.6
682	Wheat.	72.2	80.7	73.1	86.7	94.4	85.9	80.8	84.9	64.4	85.7	87.0	79.6	82.1	71.6	80.9	76.8
683	Wheat.	86.2	92.4	86.4	93.7	97.4	92.1	89.8	94.0	81.2	90.7	94.7	98.3	94.4	86.6	95.7	87.8
684	Wheat.	72.4	84.6	76.6	81.1	94.7	91.3	82.4	84.9	72.4	81.6	88.0	93.4	82.8	76.6	81.4	80.0
685	Wheat.	80.3	86.0	81.9	90.2	96.0	89.7	87.3	89.9	76.4	85.0	91.5	95.7	89.6	80.9	88.9	84.7
686	Wheat.	78.0	89.9	84.0	87.7	95.8	95.3	88.7	90.1	80.6	84.5	91.4	95.3	90.1	85.0	86.8	87.3
687	Wheat.	88.0	94.3	89.5	98.1	98.1	93.1	93.3	96.0	84.0	80.9	96.6	99.2	97.9	89.5	97.6	91.5
688	Wheat.	79.3	87.7	80.6	87.3	95.9	88.3	87.5	90.2	78.6	88.7	91.8	95.9	89.1	79.2	89.0	83.8
773	Wheat.	80.9	86.7	81.9	86.1	96.0	90.8	87.7	90.2	84.4	83.4	92.7	95.8	89.6	81.7	87.5	86.3
778	Wheat.	82.9	89.1	83.6	90.7	96.8	94.0	89.1	91.6	86.9	87.7	93.8	97.4	92.7	86.2	90.0	87.9
784	Wheat.	68.9	78.4	62.8	69.1	96.5	84.1	77.5	82.5	61.8	76.2	86.7	92.1	80.1	71.6	81.9	76.6
779	Wheat.	79.6	86.6	79.8	84.9	95.6	91.2	85.7	89.5	84.2	83.9	93.9	95.6	89.6	80.8	87.6	84.4
782	Wheat.	84.9	88.4	83.7	84.6	96.8	92.1	89.5	92.2	85.2	86.4	93.8	96.4	91.2	84.3	91.5	87.4
783	Wheat.	89.9	92.2	90.1	92.7	99.1	95.3	93.2	95.0	92.6	90.8	95.5	98.3	95.7	92.6	95.5	91.3
772	Wheat.	82.5	88.9	82.3	87.8	96.2	92.1	87.4	91.1	85.1	86.3	92.4	96.8	90.4	85.4	90.1	88.2
781	Wheat.	83.6	87.7	84.2	86.5	96.4	91.6	90.0	91.7	86.1	88.7	92.8	95.7	91.1	85.9	91.7	87.7
776	Wheat.	84.2	90.3	84.0	88.1	96.7	90.0	89.7	92.2	84.3	90.0	94.6	97.4	92.1	84.5	91.8	88.5
780	Wheat.	86.6	88.0	86.3	89.7	97.2	92.9	91.7	92.9	87.2	92.0	94.5	97.0	91.5	87.0	93.6	89.3
676	Wheat shorts.	83.5	87.9	83.0	80.5	93.0	87.5	88.0	87.4	83.2	83.8	88.1	90.5	83.2	82.8	87.5	85.9
677	Wheat shorts.	83.2	89.4	84.0	75.2	92.0	87.2	87.7	87.0	84.8	81.6	87.5	90.5	84.3	81.5	86.4	87.3
728	Wheat shorts.	75.8	84.1	76.5	63.3	89.6	82.7	78.4	81.6	78.4	75.4	83.2	87.6	82.0	75.4	83.2	78.1
729	Wheat shorts.	77.1	84.7	93.3	73.0	90.1	82.4	79.9	82.7	77.9	82.7	83.0	83.0	84.6	78.9	85.2	80.3
730	Wheat shorts.	73.4	83.5	75.5	73.8	88.7	80.7	76.8	80.7	75.0	76.1	81.4	87.7	81.7	74.7	80.5	77.4
756	Wheat shorts.	83.7	85.9	82.8	73.0	93.2	87.3	84.0	87.0	84.9	76.3	87.4	92.8	88.0	83.9	87.9	84.3
757	Wheat shorts.	76.9	82.0	76.6	76.0	90.2	80.2	78.1	81.9	78.2	80.5	82.8	89.1	81.7	75.1	83.3	75.9
758	Wheat shorts.	74.0	78.7	90.9	70.6	88.0	78.8	73.7	78.0	75.4	82.9	80.1	86.0	77.6	72.1	79.7	72.5
809	Wheat shorts.	76.3	82.5	73.6	58.9	87.9	76.4	81.6	82.3	68.0	79.4	82.8	87.6	76.8	70.4	81.4	80.6
810	Wheat shorts.	79.8	84.3	80.3	59.2	90.5	83.0	84.4	85.1	78.0	80.7	85.8	89.4	81.8	76.2	85.6	82.2
811	Wheat shorts.	80.5	86.8	80.3	64.1	91.0	84.0	86.2	85.6	78.3	81.8	86.8	90.6	84.0	80.2	85.5	85.5
832	Wheat shorts.	81.2	92.0	80.7	52.9	91.6	87.1	83.3	84.6	87.8	77.1	86.4	91.6	84.4	80.0	86.3	82.1

TABLE 4 cont.

Code	Description	ALA	ARG	ASP	CYS	GLU	HIS	ILE	LEU	LYS	MET	PHE	PRO	SER	THR	TYR	VAL	
833	Wheat shorts.	81.9	90.9	81.2	75.3	92.4	86.0	84.1	85.8	87.6	78.9	87.6	94.3	89.3	83.0	88.3	84.1	
834	Wheat shorts.	83.0	92.3	82.5	71.0	92.5	88.0	84.5	86.2	88.7	81.1	88.6	94.3	87.6	87.6	82.1	87.9	85.0
835	Wheat shorts.	81.9	91.3	81.6	69.3	92.4	85.8	84.3	86.5	88.5	79.8	88.0	93.0	87.4	82.0	82.0	88.2	84.6
638	Yucca.	77.0	91.5	-	-	82.2	86.6	-	80.7	83.2	-	-	-	-	75.1	72.8	-	
640	Yucca.	62.1	97.8	51.6	-	86.0	89.9	-	66.0	69.0	-	-	-	-	62.6	36.4	-	

1 Data of Table 3 expressed as percentages of data of Table 2.

2 Blanks (-) = variable cannot be calculated.

BIBLIOGRAPHY

The references are arranged by year and, within years, alphabetically by author's surnames. Abstracts are identified by asterisks. Some references may have been overlooked and readers are asked to bring them to the attention of the author for inclusion in subsequent updates or revisions.

1917

- 1 Armsby, H.P., 1917. The Nutrition of Farm Animals. The MacMillan Co., New York, U.S.A.

1924

- 2 Mitchell, H.H., 1924. A method for determining the biological value of protein. J. Biol. Chem. 58:873-903.

1926

- 3 Kellner, O., 1926. The Scientific Feeding of Animals (Translated by W. Goodwin). Duckworth, London, U.K.

1966

- 4 Harris, L.E., 1966. Biological Energy Interrelationships and Glossary of Energy Terms. N.A.S.-N.R.C. Publ. 1411, Washington, U.S.A.

1970

- 5 Guillaume, J., and J.D. Summers, 1970. Maintenance energy requirement of the rooster and influence of plane of nutrition on metabolizable energy. Can. J. Anim. Sci. 50:363-369.

1975

- 6 Sibbald, I.R., 1975. The effect of level of feed intake on metabolizable energy values measured with adult roosters. Poultry Sci. 54:1990-1997.
- 7 *Sibbald, I.R., 1975. The relationship between feed intake and metabolizable energy values. Poultry Sci. 54:1818.
- 8 Sibbald, I.R., 1975. The measurement of metabolizable energy for poultry feedingstuffs. Proc. Arkansas Nutr. Conf., Fayetteville, U.S.A. pp. 53-68.
- 9 Sibbald, I.R., 1975. The measurement of apparent and true metabolizable energy in poultry feedingstuffs. Proc. North Carolina Nutr. Conf., Charlotte, U.S.A. pp. 43-47.
- 10 Sibbald, I.R., 1975. Energy values in feed formulation. Proc. North Carolina Nutr. Conf., Charlotte, U.S.A. pp. 50-53.

- 11 Sibbald, I.R., and K. Price, 1975. Variation in the metabolizable energy values of diets and dietary components fed to adult roosters. *Poultry Sci.* 54:448-456.

1976

- 12 Sibbald, I.R., 1976. A bioassay for true metabolizable energy in feedingstuffs. *Poultry Sci.* 55:303-308.
- 13 Sibbald, I.R., 1976. The effect of cold pelleting on the true metabolizable energy values of cereal grains fed to adult roosters and a comparison of observed with predicted metabolizable energy values. *Poultry Sci.* 55:970-974.
- 14 Sibbald, I.R., 1976. The true metabolizable energy values of several feedingstuffs measured with roosters, laying hens, turkeys and broiler hens. *Poultry Sci.* 55:1459-1463.
- 15 Sibbald, I.R., 1976. The effect of duration of starvation of the assay bird on true metabolizable energy values. *Poultry Sci.* 55:1578-1579.
- 16 Sibbald, I.R., 1976. Energy in poultry nutrition, a bioassay for true metabolizable energy/L'énergie dans l'alimentation de la volaille. Essai biologique sur l'énergie métabolisable vraie. Canadex 450.55, Agriculture Canada, Ottawa, Canada.
- 17 Sibbald, I.R., 1976. Energy in poultry nutrition, terminology and definitions/L'énergie dans l'alimentation de la volaille. Terminologie et définitions. Canadex 450.55, Agriculture Canada, Ottawa, Canada.
- 18 Sibbald, I.R., 1976. The true metabolizable energy values of some feedingstuffs/Valeurs de l'énergie métabolisable vraie (EMV) de certains aliments du bétail. Canadex 450.60, Agriculture Canada, Ottawa, Canada.
- 19 Sibbald, I.R., 1976. Shorter method to determine ME values. *Feed Management* 27(1):17-19.
- 20 Sibbald, I.R., 1976. The measurement of true metabolizable energy in poultry feedingstuffs. Proc. Guelph Nutr. Conf., Toronto, Canada. pp. 102-114.
- 21 Sibbald, I.R., and K. Price, 1976. True metabolizable energy values for poultry of Canadian barleys measured by bioassay and predicted from physical and chemical data. *Can. J. Anim. Sci.* 56:775-782.
- 22 Sibbald, I.R., and K. Price, 1976. Measurement of the true metabolizable energy values of poultry feeds and ingredients. Proc. 7th Symp. Energy Metab. in Farm Animals, Vichy, France. pp. 269-272.

1977

- 23 Featherston, W.R., 1977. Poultry nutrition research 1976. 25th Pfizer Annu. Res. Conf., Minneapolis, U.S.A. pp. 31-59.
- 24 Halloran, H.R., 1977. Metabolizable energy values of feed fats for broilers and turkeys. Proc. Maryland Nutr. Conf., Washington, U.S.A. pp. 26-31.
- 25 Johns, D.C., and I.C. Edmundson, 1977. Metabolisable energy determinations. Proc. 23rd Annu. Poult. Conv., Massey Univ., Palmerston North, N.Z. pp. 84-91.
- 26 Muztar, A.J., S.J. Slinger and J.H. Burton, 1977. Metabolisable energy content of freshwater plants in chickens and ducks. Poultry Sci. 56:1893-1899.
- 27 Patchell, M.R., and I.C. Edmundson, 1977. Metabolisable energy. 3. Metabolisable energy of New Zealand ingredients. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 48-50.
- 28 Patchell, M.R., and I.C. Edmundson, 1977. Metabolisable energy. 4/5. Metabolisable energy of New Zealand ingredients. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 51-52.
- 29 Patchell, M.R., and I.C. Edmundson, 1977. Metabolisable energy. 6. Metabolisable energy of New Zealand ingredients. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. p. 71.
- 30 Salmon, R.E., and J.B. O'Neil, 1977. Effects of frost damage on physical and chemical characteristics and true metabolizable energy of wheat. Can. J. Anim. Sci. 57:755-760.
- 31 Sibbald, I.R., 1977. A test of the additivity of true metabolizable energy values of feedingstuffs. Poultry Sci. 56:363-366.
- 32 Sibbald, I.R., 1977. The true metabolizable energy values of some feedingstuffs. Poultry Sci. 56:380-382.
- 33 Sibbald, I.R., 1977. The true metabolizable energy values for poultry of rapeseed and of the meal and oil derived therefrom. Poultry Sci. 56:1652-1656.
- 34 Sibbald, I.R., 1977. The effect of level of feed input on true metabolizable energy values. Poultry Sci. 56:1662-1663.
- 35 Sibbald, I.R., 1977. The effect of steam pelleting on the true metabolizable energy values of poultry diets. Poultry Sci. 56:1686-1688.

- 36 *Sibbald, I.R., 1977. Effect of levels of input and of dietary inclusion on TME values. *Poultry Sci.* 56:1757.
- 37 Sibbald, I.R., 1977. The true metabolizable energy system. Part 1. *Feedstuffs* 49(42):21-22.
- 38 Sibbald, I.R., 1977. The true metabolizable energy system. Part 2. *Feedstuffs* 49(43):23-24.
- 39 Sibbald, I.R., 1977. Definition, measurement and usefulness of true metabolizable energy. Proc. 36th Semi-Annu. Meet. Am. Feed Manuf. Assoc. Nutr. Council, Memphis, U.S.A. pp. 32-35.
- 40 Sibbald, I.R., 1977. The true metabolizable energy system of feed evaluation. Proc. Maryland Nutr. Conf., Washington, U.S.A. pp. 32-35.
- 41 Sibbald, I.R., and J.K.G. Kramer, 1977. The true metabolizable energy values of fats and fat mixtures. *Poultry Sci.* 56:2079-2086.
- 42 Sibbald, I.R., and K. Price, 1977. True and apparent metabolizable energy values for poultry of Canadian wheats and oats measured by bioassay and predicted from physical and chemical data. *Can. J. Anim. Sci.* 57:365-374.
- 43 Sibbald, I.R., and K. Price, 1977. The true metabolizable energy values of the seeds of Brassica campestris, B. hirta and B. napus. *Poultry Sci.* 56:1329-1331.
- 44 Sibbald, I.R., and K. Price, 1977. The effect of level of dietary inclusion and of calcium on the true metabolizable energy values of fats. *Poultry Sci.* 56:2070-2078.
- 45 Terpstra, K., 1977. Discussion on the determination of the digestibility of amino acids in monogastric animals. Proc. 2nd Int. Symp. Protein Metabolism and Nutrition. Cent. Agric. Publ. Doc., Wageningen, The Netherlands. pp. 90-91.

1978

- 46 Anonymous, 1978. Poultry Research Centre Report for Year Ending 31st March 1978. Agric. Res. Council, London, U.K. pp. 9-15.
- 47 *Boldaji, F., W.B. Roush, H.S. Nakaue and G.H. Arscott, 1978. True metabolizable energy values of different varieties of wheat grown in Oregon, measured with normal size and dwarf White Leghorn roosters. *Poultry Sci.* 57:1119-1120.
- 48 Cuesta, M.P., 1978. Efficiencia energetica de los alimentos en la produccion avia. Proc. 3rd World Congr. Anim. Feed 8:139-160.
- 49 Edmundson, I.C., M.R. Patchell and R.D. King, 1978. Metabolisable energy 7. Assays of ingredients and diets. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 58-60.

- 50 Farrell, D.J., 1978. Rapid determination of metabolizable energy of foods using cockerels. *Br. Poult. Sci.* 19:303-308.
- 51 Farrell, D.J., 1978. Metabolizable energy of some local ingredients using a rapid bioassay. *Proc. 2nd Australasian Poult. and Stockfeed Conv.*, Sydney, Australia. pp. 204-208.
- 52 Farrell, D.J., 1978. Energy systems in poultry, how well do they predict biological performance. *Proc. 24th Annu. Poult. Conv.*, Massey Univ., Palmerston North, N.Z. pp. 96-105.
- 53 Johns, D.C., 1978. Chick nutrition 165 - Full-fat soybean meal. *Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ.*, Palmerston North, N.Z. p. 16.
- 54 Kessler, J.W., and O.P. Thomas, 1978. An evaluation of the true metabolizable energy system. *Proc. Maryland Nutr. Conf.*, Washington, U.S.A. pp. 18-21.
- 55 King, R.D., 1978. Layer nutrition 94 - Whole barley choice feeding. *Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ.*, Palmerston North, N.Z. pp. 19-21.
- 56 King, R.D., 1978. Comparisons of dietary costs of AME and TME based formulations. *Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ.*, Palmerston North, N.Z. pp. 80-90.
- 57 Likuski, H.J.A., and H.G. Dorrell, 1978. A bioassay for rapid determination of amino acid availability values. *Poultry Sci.* 57:1658-1660.
- 58 *Miles, R.D., A.S. Arafa, R.J. Bloomer and R.H. Harms, 1978. The true metabolizable energy of turkey poult diets containing dietary fillers. *Poultry Sci.* 57:1107.
- 59 Mutzar, A.J., H.J. Likuski and S.J. Slinger, 1978. Metabolizable energy content of Tower and Candle rapeseeds and rapeseed meals determined in two laboratories. *Can. J. Anim. Sci.* 58:485-492.
- 60 Sell, J.L., 1978. Poultry nutrition research 1977. *26th Pfizer Annu. Res. Conf.*, Las Vegas, U.S.A. pp. 27-70.
- 61 Sibbald, I.R., 1978. The effect of the duration of the time interval between assays on true metabolizable energy values measured with adult roosters. *Poultry Sci.* 57:455-460.
- 62 Sibbald, I.R., 1978. The true metabolizable energy values of mixtures of tallow with either soybean oil or lard. *Poultry Sci.* 57:473-477.
- 63 Sibbald, I.R., 1978. The effect of the age of the assay bird on the true metabolizable energy values of feedingstuffs. *Poultry Sci.* 57:1008-1012.

- 64 Sibbald, I.R., 1978. Scientists study metabolizable energy variations in swine and poultry diets. Feedstuffs 50(48):20-22.
- 65 Sibbald, I.R., 1978. True metabolizable energy determination and application. Proc. Georgia Nutr. Conf., Atlanta, U.S.A. pp. 105-109.
- 66 Sibbald, I.R., and J.K.G. Kramer, 1978. The effect of the basal diet on the true metabolizable energy value of fat. Poultry Sci. 57:685-691.
- 67 Sibbald, I.R., and K. Price, 1978. The metabolic and endogenous energy losses of adult roosters. Poultry Sci. 57:556-557.
- 68 *Tennesaca, L.G., and J.L. Sell, 1978. Influence of indigestible material on energy excretion and true metabolizable energy of corn. Poultry Sci. 57:1167.

1979

- 69 Cid Diaz, J.M., 1979. La Energia Metabolisable Aparente Y Verdadera, En Avicultura: Situacion Actual. Academia de Ciencias Veterinarias de Madrid, Spain.
- 70 *Coon, C., M. Scheibel and K. Kelley, 1979. The heat increment and net available and true metabolisable energy of poultry feeds. Poultry Sci. 58:1046.
- 71 Edmundson, I.C., 1979. True metabolisable energy trial ME/9. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 82-84.
- 72 Edmundson, I.C., 1979. True metabolisable energy trial ME/10. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 104-105.
- 73 Edmundson, I.C., 1979. True metabolisable energy trial ME/11. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. p. 106.
- 74 Edmundson, I.C., 1979. True metabolisable energy trial ME/12(a). Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. p. 107.
- 75 Farrell, D.J., 1979. Energy systems for pigs and poultry: a review. J. Aust. Inst. Agric. Sci. 45:21-34.
- 76 Halloran, H.R., and I.R. Sibbald, 1979. Metabolizable energy values of fats measured by several procedures. Poultry Sci. 58:1299-1307.
- 77 Hartel, H., 1979. Methods to calculate the energy of mixed feeds. Proc. 2nd Eur. Symp. Poult. Nutr., Beekbergen, The Netherlands. pp. 6-13.
- 78 Hulan, H.W., F.G. Proudfoot and C.G. Zarkadas, 1979. The nutritional value and quality of squid (Illex illecebrosus) meal as source of dietary protein for broiler chicken. Br. J. Nutr. 41:163-173.

- 79 Jones, J.D., and I.R. Sibbald, 1979. The true metabolizable energy values for poultry of fractions of rapeseed (Brassica napus cv. Tower). *Poultry Sci.* 58:385-391.
- 80 *Kessler, J.W., and O.P. Thomas, 1979. An evaluation of the "true metabolizable energy" system. *Poultry Sci.* 58:1072-1073.
- 81 Kessler, J.W., and O.P. Thomas, 1979. A further evaluation of the true metabolizable energy system. *Proc. Maryland Nutr. Conf.*, Washington, U.S.A. pp. 83-87.
- 82 Kratzer, F.H., 1979. Poultry nutrition research 1978. 27th Pfizer Annu. Res. Conf., St. Louis, U.S.A. pp. 7-52.
- 83 Kussabati, R., 1979. Influence of dietary intake level on the metabolizable energy and digestibility of lipids in the growing chicks and the adult cockerel. *Proc. 2nd Eur. Symp. Poult. Nutr.*, Beekbergen, The Netherlands. pp. 14-18.
- 84 Lopez, C.V., 1979. Energia metabolizable aparante y verdadera de materias primas utilizadas en raciones para aves en Columbia. *Rev. Inst. Columb. Agropécu.* 14(Suppl.):312.
- 85 Martinez, R.I.A., 1979. Determinacion de energia metabolizable verdadera de materias primas Colombianas en aves. (Thesis summary). *Rev. Inst. Columb. Agropécu.* 14(Suppl.):307.
- 86 Mutzar, A.J., and S.J. Slinger, 1979. Some factors affecting the true metabolizable energy values of soybean meal and rapeseed meals. *Can. J. Anim. Sci.* 59:317-321.
- 87 Mutzar, A.J., and S.J. Slinger, 1979. Effect of length of excreta collection period and feed input level on the true metabolizable energy value of rapeseed meal. *Nutr. Rep. Int.* 19:689-694.
- 88 *Roitz, D.O., and H.L. Fuller, 1979. A comparison of methods for determining biological value of oilseed meals. *Poultry Sci.* 58:1099.
- 89 Salmon, R.E., K.K. Klein and E. Larmond, 1979. Low glucosinolate rapeseed meal in turkey broiler diets of varying nutrient density. *Poultry Sci.* 58:1514-1523.
- 90 Scheibel, M.S., C.N. Coon and K.W. Kelley, 1979. The heat increment of feeds used in poultry diets. *Nutr. Rep. Int.* 20:871-881.
- 91 Shires, A., A.R. Robblee, R.T. Hardin and D.R. Clandinin, 1979. Effect of previous diet, body weight, and duration of starvation of the assay bird on the true metabolizable energy value of corn. *Poultry Sci.* 58:602-608.
- 92 Sibbald, I.R., 1979. Passage of feed through the adult rooster. *Poultry Sci.* 58:446-459.

- 93 Sibbald, I.R., 1979. A bioassay for available amino acids and true metabolizable energy in feedingstuffs. *Poultry Sci.* 58:668-675.
- 94 Sibbald, I.R., 1979. The effect of the duration of the excreta collection period on the true metabolizable energy values of feedingstuffs with slow rates of passage. *Poultry Sci.* 58:896-899.
- 95 Sibbald, I.R., 1979. Bioavailable amino acids and true metabolizable energy of cereal grains. *Poultry Sci.* 58:934-939.
- 96 Sibbald, I.R., 1979. Effect of level of feed input, dilution of test material and duration of excreta collection on true metabolizable energy values. *Poultry Sci.* 58:1325-1329.
- 97 Sibbald, I.R., 1979. The effect of the drying procedure on excreta energy values for poultry and other species. *Poultry Sci.* 58:1392-1394.
- 98 Sibbald, I.R., 1979. Bioavailable amino acids in feedingstuffs. p. 6 in A.P. Williams, ed. ARC Protein Evaluation Group News Sheet #5. Agric. Res. Council, Poultry Res. Cent., Roslin, U.K.
- 99 Sibbald, I.R., 1979. Metabolizable energy evaluation of poultry diets. pp. 35-49 in W. Haresign and D. Lewis, eds. Recent Advances in Animal Nutrition - 1979. Butterworths, London, U.K.
- 100 Sibbald, I.R., 1979. True metabolizable energy. Proc. 2nd Eur. Symp. Poult. Nutr., Beekbergen, The Netherlands. pp. 25-30.
- 101 *Tenesaca, G.L., and J.L. Sell, 1979. True metabolizable energy of corn and oats. *Poultry Sci.* 58:1115.

1980

- 102 *Arvat, V., J. Lyons and J.M. Vanderpopuliere, 1980. A comparison of metabolizable energy and true metabolizable energy. *Poultry Sci.* 59:1579.
- 103 Chami, D.B., P. Vohra and F.H. Kratzer, 1980. Evaluation of a method for determination of true metabolizable energy of feed ingredients. *Poultry Sci.* 59:569-571.
- 104 Chand, S., V.R. Sadagopan and V.R. Reddy, 1980. Utilization of solvent extracted mustard-cake in chicks and its true metabolizable energy content. *Ind. J. Anim. Sci.* 50:521-524.
- 105 Christison, G.I., and J.M. Bell, 1980. Evaluation of Terra, a new cultivar of naked oats (Avena nuda), when fed to young pigs and chicks. *Can. J. Anim. Sci.* 60:465-471.
- 106 Dale, N.M., and H.L. Fuller, 1980. Additivity of true metabolizable energy values as measured with roosters, broiler chicks and poult. *Poultry Sci.* 59:1941-1942.

- 107 *Dale, N.M., and H.L. Fuller, 1980. Additivity of true metabolizable energy values as measured with roosters, broiler chicks and poultts. *Poultry Sci.* 59:1561.
- 108 *Dale, N.M., and H.L. Fuller, 1980. A comparison of apparent and true metabolizable energy systems in evaluating the calorie content of broiler diets. *Poultry Sci.* 59:1598-1599.
- 109 *Dudley, W.A., S.M. Kim and M. Winegardener, 1980. The use of gelatin capsules for measuring true metabolizable energy. *Poultry Sci.* 59:1601.
- 110 Edmundson, I.C., 1980. The true metabolisable energy of meat and bone meal determined at different levels. *Proc. South Pacific Poult. Sci. Conv.*, Auckland, N.Z. pp. 20-25.
- 111 Edmundson, I.C., 1980. True metabolisable energy trial ME/12(a). Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 95-98.
- 112 Edmundson, I.C., 1980. True metabolisable energy trial ME/13(c). Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 99-100.
- 113 *Emebo, G.O., and R.H. Roberson, 1980. The true amino acid availability of feedstuffs for poultry. *Poultry Sci.* 59:1604.
- 114 Farrell, D.J., 1980. Feeding standards for poultry. *Proc. Aust. Soc. Anim. Prod.* 13:26-31.
- 115 Farrell, D.J., 1980. True metabolizable energy (TME) and the alternative. pp. 146-153 in D.J. Farrell, ed. *Recent Advances in Animal Nutrition in Australia* 1980. Univ. New England, Armidale, Australia.
- 116 Farrell, D.J., 1980. The case for retaining the apparent metabolizable energy system because of variation in endogenous excreta. *Proc. Poult. Husb. Res. Found. Symp.*, Sydney, Australia. 11 pp.
- 117 Fekete, S., 1980. Energetikai takarmanyértékelés (Irodalmi attékintés). *Magy. Allatorv. Lapja* 35:472-479.
- 118 Fenwick, G.R., and R.F. Curtis, 1980. Rapeseed meal and its use in poultry diets. A review. *Anim. Feed Sci. Technol.* 5:255-298.
- 119 Fuller, H.L., 1980. Poultry nutrition research 1979. 28th Pfizer Annu. Res. Conf., Atlanta, U.S.A. pp. 91-127.
- 120 Halloran, H.R., 1980. Comparison of metabolizable energy methods on identical ingredient samples. *Poultry Sci.* 59:1552-1553.
- 121 Johnsson, G., 1980. Nutritive value of grassmeal for poultry. M. Phil. Thesis, Univ. Edinburgh, Edinburgh, U.K.

- 122 Kessler, J.W., 1980. An investigation of the "true" metabolizable energy system. Ph.D. Thesis, Univ. Maryland, College Park, U.S.A.
- 123 Kessler, J.W., and O.P. Thomas, 1980. The influence of time and cecectomy on the true metabolizable energy. Proc. Maryland Nutr. Conf., Washington, U.S.A. pp. 27-31.
- 124 King, R.D., 1980. Layer nutrition 140 - Within flock performance. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 64-66.
- 125 King, R.D., 1980. Layer nutrition 142 - Dietary energy levels. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 69-70.
- 126 King, R.D., 1980. Layer nutrition 151 - Dietary energy levels. Summary of Final Reports and Current Research, Poult. Res. Center, Massey Univ., Palmerston North, N.Z. p. 117.
- 127 *Luis, E.S., T.W. Sullivan and L.A. Nelson, 1980. The nutritional value of three varieties of proso millet and bird resistant sorghum for poultry. Poultry Sci. 59:1632.
- 128 Mahlotra, N.K., V.R. Sadagopan, V.R. Reddy and B. Panda, 1980. Metabolizable energy and utilization of treated deoiled salseed-meal in chicks. Ind. J. Anim. Sci. 50:437-443.
- 129 Mateos, G.G., and J.L. Sell, 1980. Influence of fat and carbohydrate source on energy utilization by laying hens. Nutr. Rep. Int. 22:79-89.
- 130 Mateos, G.G., and J.L. Sell, 1980. True and apparent metabolizable energy value of fat for laying hens: influence of level of use. Poultry Sci. 59:369-373.
- 131 Muztar, A.J., and S.J. Slinger, 1980. The effect of level of feed input in a short term assay for determining apparent metabolizable energy of some feedingstuffs. Nutr. Rep. Int. 22:117-122.
- 132 Muztar, A.J., and S.J. Slinger, 1980. Effect of body weight and duration of fast on metabolic and endogenous excretion by the mature rooster. Nutr. Rep. Int. 22:147-156.
- 133 Muztar, A.J., and S.J. Slinger, 1980. The effect of fiber on metabolic and endogenous energy losses in mature cockerels. Nutr. Rep. Int. 22:183-186.
- 134 Muztar, A.J., and S.J. Slinger, 1980. Rate of passage of feedstuffs through mature roosters and effect on true metabolizable energy. Nutr. Rep. Int. 22:361-367.
- 135 Muztar, A.J., and S.J. Slinger, 1980. Effect of length of collection period on the metabolizable energy values in short term assays. Nutr. Rep. Int. 22:589-595.

- 136 Muztar, A.J., and S.J. Slinger, 1980. An evaluation of the rapid apparent metabolizable energy assay in relation to feed intake using mature cockerels. Nutr. Rep. Int. 22:745-750.
- 137 Muztar, A.J., and S.J. Slinger, 1980. Effect of level of dietary fiber on nitrogen and amino acid excretion in the fasted mature rooster. Nutr. Rep. Int. 22:863-868.
- 138 Muztar, A.J., and S.J. Slinger, 1980. The effect of dry matter on metabolic and endogenous amino acid excretion in mature cockerels. Nutr. Rep. Int. 22:901-906.
- 139 Muztar, A.J., and S.J. Slinger, 1980. Bioavailable amino acids in corn and alfalfa as measured by applying the true metabolizable energy assay. Poultry Sci. 59:1873-1877.
- 140 Muztar, A.J., S.J. Slinger, H.J.A. Likuski and H.G. Dorrell, 1980. True amino acid availability values for soybean meal and Tower and Candle rapeseed and rapeseed meals determined in two laboratories. Poultry Sci. 59:605-610.
- 141 *Parsons, C.M., and L.M. Potter, 1980. True metabolizable energy and amino acid digestibility of dehulled soybean meals. Poultry Sci. 59:1648-1649.
- 142 Patchell, M.R., 1980. The minimum energy intake of layers. Proc. South Pacific Poult. Sci. Conv., Auckland, N.Z. pp. 163-170.
- 143 Patchell, M.R., 1980. Chick nutrition 186 - Feeding a starter diet for 4 or 8 weeks. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 1-2.
- 144 Patchell, M.R., 1980. Chick nutrition 187a - Feed restriction during rearing. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 5-6.
- 145 Patchell, M.R., 1980. Layer nutrition 123 - Effect of split diets during rearing. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 36-37.
- 146 Patchell, M.R., 1980. Layer nutrition 128 - Energetic efficiencies of maintenance and production of laying diets. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 41-48.
- 147 Patchell, M.R., 1980. Layer nutrition 129 - True metabolisable energy intakes. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 49-51.
- 148 Patchell, M.R., 1980. Layer nutrition 132 - Energy intake. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 52-54.

- 149 Patchell, M.R., 1980. Layer nutrition 133 -- Energy intake. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 55-57.
- 150 Patchell, M.R., 1980. Layer nutrition 134 - Energy intake (sand). Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 58-59.
- 151 Patchell, M.R., 1980. Layer nutrition 141 - Energy intake (sand). Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 67-68.
- 152 Patchell, M.R., 1980. Layer nutrition 143 - Energy intake. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 71-74.
- 153 Patchell, M.R., 1980. Chick nutrition 190 - Energy allowance for growers. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 101-102.
- 154 Patchell, M.R., 1980. Layer nutrition 135 -- The energy requirements of four strains. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 105-108.
- 155 Patchell, M.R., 1980. Layer nutrition 148 - Body weight and energy intake. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 112-114.
- 156 Patchell, M.R., and M. Fraser, 1980. Chick nutrition 191 - Apex meal. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 9-10.
- 157 Shires, A., A.R. Robblee, R.T. Hardin and D.R. Clandinin, 1980. Effect of age of chicken on the true metabolizable energy values of feed ingredients. Poultry Sci. 59:396-403.
- 158 *Shires, A., A.R. Robblee, R.T. Hardin and D.R. Clandinin, 1980. Effect of age on the true metabolizable energy values of feedingstuffs. Can. J. Anim. Sci. 58:810.
- 159 Sibbald, I.R., 1980. Metabolizable energy in poultry nutrition. BioScience 30:736-741.
- 160 Sibbald, I.R., 1980. The clearance time and rate of passage of feed residues. Poultry Sci. 59:374-377.
- 161 Sibbald, I.R., 1980. The effects of dietary cellulose and sand on the combined metabolic plus endogenous energy and amino acid output of adult cockerels. Poultry Sci. 59:836-844.
- 162 Sibbald, I.R., 1980. Gut clearance and true metabolizable energy value of dehydrated alfalfa supplemented with sodium chloride. Poultry Sci. 59:939-940.

- 163 Sibbald, I.R., 1980. The passage of oat and other feed residues through the adult cockerel. *Poultry Sci.* 59:2136-2144.
- 164 Sibbald, I.R., 1980. The effect of heat on the clearance time, true metabolizable energy and true available amino acids of raw soybean flakes. *Poultry Sci.* 59:2358-2360.
- 165 Sibbald, I.R., 1980. A new technique for estimating the ME content of feeds for poultry. pp. 38-43 in W.J. Pidgen, C.C. Balch and M. Graham, eds. *Standardization of Analytical Methodology for Feeds*. Int. Develop. Res. Cent. Publ. 134e, Ottawa, Canada.
- 166 Sibbald, I.R., 1980. Amino acid availability. *Feed Management* 31(10):26.
- 167 Sibbald, I.R., 1980. Amino acid availability. *Poultry Digest* 39:604.
- 168 Sibbald, I.R., 1980. Selection of a bioassay for available energy. Proc. Poult. Husb. Res. Found. Symp., Sydney, Australia. 13 pp.
- 169 Sibbald, I.R., 1980. Selection of a bioassay for available energy. Proc. South Pacific Poult. Sci. Conv., Auckland, N.Z. pp. 10-19.
- 170 Sibbald, I.R., and J.K.G. Kramer, 1980. The effect of the basal diet on the utilization of fat as a source of true metabolizable energy, lipid and fatty acids. *Poultry Sci.* 59:316-324.
- 171 Sibbald, I.R., and J.K.G. Kramer, 1980. The effects of fractions of yellow corn on the true metabolizable energy of beef tallow. *Poultry Sci.* 59:1505-1509.
- 172 Sibbald, I.R., and K. Price, 1980. Variability in metabolic plus endogenous energy losses in adult cockerels and in true metabolizable energy values and rates of passage of alfalfa. *Poultry Sci.* 59:1275-1279.
- 173 Sibbald, I.R., J.P. Barrette and K. Price, 1980. Predicting true metabolizable energy, gross energy, carbohydrate and proximate analysis values by assuming additivity. *Poultry Sci.* 59:805-807.
- 174 Sibbald, I.R., K. Price and J.P. Barrette, 1980. True metabolizable energy for poultry of commercial diets measured by bioassay and predicted from chemical data. *Poultry Sci.* 59:808-811.
- 175 Slinger, S.J., and A.J. Muztar, 1980. Energy measures for poultry and swine, practical application. Proc. Guelph Nutr. Conf., Toronto, Canada. pp. 103-109.
- 176 Van Es, A.J.H., 1980. Feed evaluation, a survey. pp. 85-92 in L.E. Mount, ed. *Energy Metabolism*. Butterworths, London, U.K.
- 177 Wehner, G.R., R.L. Harrold and M. Wanapat, 1980. True metabolizable energy of sprouted wheat. *North Dakota Farm Res. Bull.* 47.

178 Willis, G.M., and D.H. Baker, 1980. Evaluation of turfgrass clippings as a dietary ingredient for the growing chick. *Poultry Sci.* 59:404-411.

1981

179 Allen, N.K., 1981. Nutrition of growing geese. *Proc. Minnesota Nutr. Conf.*, Minneapolis, U.S.A. pp. 119-123.

180 Boldaji, F., W.B. Roush, H.S. Nakaue and G.H. Arscott, 1981. True metabolizable energy values of corn and different varieties of wheat and barley using normal and dwarf single comb White Leghorn roosters. *Poultry Sci.* 60:225-227.

181 Campbell, L.D., B.O. Eggum and I. Jacobsen, 1981. Biological value, amino acid availability and true metabolizable energy of low-glucosinolate rapeseed meal (Canola) determined with rats and/or roosters. *Nutr. Rep. Int.* 24:791-797.

182 Dale, N.M., 1981. Energia metabolizable "Verdadera". *Industria Avicola* 28(9):78-86.

183 Dale, N.M., and H.L. Fuller, 1981. Effect of carrier on the true metabolizable energy of corn oil. *Poultry Sci.* 60:1504-1508.

184 *Dale, N.M., and H.L. Fuller, 1981. Applicability of the true metabolizable energy system in practical feed formulation. *Poultry Sci.* 60:1598.

185 *Dale, N.M., and H.L. Fuller, 1981. Studies on the true metabolizable energy of fats at low-level dietary inclusion. *Poultry Sci.* 60:1643.

186 Dale N.M., and H.L. Fuller, 1981. The use of true metabolizable energy (TME) in formulating poultry rations. *Proc. Georgia Nutr. Conf.*, Atlanta, U.S.A. pp. 50-57.

187 Davidson, J., and S. Graham, 1981. The estimation of metabolizable energy in maize and in wheat by-products using chickens, with observations on problems of prediction from chemical composition. *J. Agric. Sci. Camb.* 96:221-225.

188 Devegowda, G., 1981. Evaluation of feedstuffs for true metabolizable energy and bioavailable amino acids in turkeys. Ph.D. Thesis, Univ. Minnesota, St. Paul, U.S.A.

189 Edmundson, I.C., 1981. True metabolisable energy trial ME/13. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 97-98.

190 Engster, H.M., D.C. Snetsinger and J.W. Kessler, 1981. Commercial application of the TME system. *Proc. 3rd Eur. Symp. Poult. Nutr.*, Peebles, U.K. pp. 56-63.

- 191 Farrell, D.J., 1981. An assessment of quick bioassays for determining the true metabolizable energy and apparent metabolizable energy of poultry feedstuffs. *World's Poult. Sci. J.* 37:72-83.
- 192 *Fuller, H.L., and N.M. Dale, 1981. Endogenous energy losses of fed vs. fasted roosters in true metabolizable energy assays. *Poultry Sci.* 60:1601.
- 193 *Fuller, H.L., and N.M. Dale, 1981. Influence of the ratio of basal fat to test fat on the true metabolizable energy of several fats. *Poultry Sci.* 60:1659.
- 194 Johns, D.C., 1981. Chick nutrition 192 - Maize quality and heat treatment. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 4-6.
- 195 Kessler, J.W., and O.P. Thomas, 1981. The effect of cecectomy and extension of the collection period on the true metabolizable energy values of soybean meal, feather meal, fish meal and blood meal. *Poultry Sci.* 60:2639-2647.
- 196 Kessler, J.W., T.H. Nguyen and O.P. Thomas, 1981. The amino acid excretion values in intact and cecectomized negative control roosters used for determining metabolic plus endogenous urinary energy losses. *Poultry Sci.* 60:1576-1577.
- 197 Kienholz, E.W., and R.E. Moreng, 1981. A new brewery by-product for turkey diets. *Poultry Sci.* 60:1879-1883.
- 198 King, R.D., 1981. Layer nutrition 151 - Dietary energy levels. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 48-49.
- 199 Latshaw., J.D., 1981. Counting calories in poultry feed. *Ohio Poult. Pointers* 20(5). Ohio State Univ., U.S.A.
- 200 Low, A.G., and I.G. Partridge, 1981. The value of in vivo and in vitro digestibility measurements of amino acids as predictors of performance and carcass composition in pigs. *Proc. 6th Int. Symp. Amino Acids*, Serock, Poland. pp. 1-9.
- 201 McNab, J.M., and C. Fisher, 1981. The choice between apparent and true metabolizable energy systems - recent evidence. *Proc. 3rd Eur. Symp. Poult. Nutr.*, Peebles, U.K. pp. 45-55.
- 202 McNab, J.M., J.C. Blair, W.A. Dewar and J.N. Downie, 1981. The effects of different dietary (Na + K + Cl) contents on the performance of young broilers and on the metabolizable energy values of their diets. *Proc. 3rd Eur. Symp. Poult. Nutr.*, Peebles, U.K. pp. 132-136.
- 203 Miles, R.D., A.S. Arafa, R.J. Bloomer and R.H. Harms, 1981. The influence of certain dietary fillers on the true metabolizable energy of diets for turkeys. *Br. Poult. Sci.* 22:411-414.

- 204 Minaar, A.duP., and J. Erasmus, 1981. True metabolizable energy with nitrogen corrections to improve additivity in poultry diets. S. Afr. J. Anim. Sci. 11:279-283.
- 205 Miski, A.M.A., and S. Quazi, 1981. Influence of age and sex of growing broiler chicks and body weight of roosters on their endogenous and metabolic energy losses. Poultry Sci. 60:781-785.
- 206 Mollah, Y., W.L. Bryden, D. Balnave and E.F. Annison, 1981. Studies of low - ME wheats. Proc. 4th Australasian Poult. Stockfeed Conv., Perth, Australia. pp. 264-265.
- 207 Moran, E.T., 1981. Poultry nutrition research 1980. 29th Pfizer Annu. Res. Conf., St. Louis, U.S.A. pp. 91-130.
- 208 Muztar, A.J., and S.J. Slinger, 1981. A study of the metabolic fecal and endogenous amino acid excretion in fasted mature cockerels with time. Nutr. Rep. Int. 23:465-469.
- 209 Muztar, A.J., and S.J. Slinger, 1981. An evaluation of the true metabolizable energy assay for monitoring the apparent metabolizable energy values of poultry diets. Poultry Sci. 60:598-602.
- 210 Muztar, A.J., and S.J. Slinger, 1981. A comparison of the true and apparent metabolizable energy measures using corn and soybean meal samples. Poultry Sci. 60:611-616.
- 211 Muztar, A.J., and S.J. Slinger, 1981. Relationship between body weight and amino acid excretion in fasted mature cockerels. Poultry Sci. 60:790-794.
- 212 Muztar, A.J., and S.J. Slinger, 1981. An evaluation of the nitrogen correction in the true metabolizable energy assay. Poultry Sci. 60:835-839.
- 213 Muztar, A.J., S. Leeson and S.J. Slinger, 1981. Effect of blending and level of inclusion on the metabolizable energy of tallow and Tower rapeseed soapstocks. Poultry Sci. 60:365-372.
- 214 Muztar, A.J., H.J.A. Likuski and S.J. Slinger, 1981. True metabolizable energy values of a number of feedingstuffs and complete diets as determined in two laboratories. Poultry Sci. 60:373-377.
- 215 *Ngou, J.D., P.M. Mairorino and B.L. Reid, 1981. Evaluation of jojoba meal for poultry diets. Poultry Sci. 60:1703.
- 216 Okumura, J., Y. Isshiki and N. Nakahiro, 1981. Some factors affecting urinary and faecal nitrogen loss by chickens fed on a protein-free diet. Br. Poult. Sci. 22:1-7.
- 217 Parsons, C.M., 1981. Evaluation of a procedure for determination of amino acid digestibility and metabolizable energy of feedstuffs for poultry. Ph.D. Thesis, Virginia Polytech. Inst. and State Univ., Blacksburg, U.S.A.

- 218 Parsons, C.M., L.M. Potter and R.D. Brown, Jr., 1981. True metabolizable energy and amino acid digestibility of dehulled soybean meal. *Poultry Sci.* 60:2687-2696.
- 219 *Parsons, C.M., L.M. Potter and R.D. Brown Jr., 1981. Effects of dietary protein and carbohydrate and of intestinal microflora on excretion of amino acids by poultry. *Poultry Sci.* 60:1709-1710.
- 220 *Parsons, C.M., L.M. Potter, R.D. Brown Jr., T.D. Wilkins and B.A. Bliss, 1981. Microbial contribution to dry matter and amino acid content of poultry excreta. *Poultry Sci.* 60:1709.
- 221 Patchell, M.R., 1981. Chick nutrition 190 - Energy allowance for growers. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 1-3.
- 222 Patchell, M.R., 1981. Chick nutrition 202 - Energy allowance for growers. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 21-23.
- 223 Patchell, M.R., 1981. Chick nutrition 203 - Apex paunch meal. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 24-25.
- 224 Patchell, M.R., 1981. Layer nutrition 135 - The energy requirement of four strains. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 28-31.
- 225 Patchell, M.R., 1981. Layer nutrition 148 - Body weight and energy intake. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 37-40.
- 226 Patchell, M.R., 1981. Layer nutrition 159 - Energy level of diets. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 69-72.
- 227 Patchell, M.R., 1981. Layer nutrition 160 - Amino acid levels - energy and state of lay. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 73-76.
- 228 Pearson, R.A., and K.M. Herron, 1981. Effects of energy and protein allowances during lay on the reproductive performance of broiler breeder hens. *Br. Poult. Sci.* 22:227-239.
- 229 du Preez, J.J., J.P. Hayes and J.S. Duckitt, 1981. Endogenous energy, a cause of biased true metabolizable energy values. *S. Afr. J. Anim. Sci.* 11:269-272.
- 230 du Preez, J.J., J.P. Hayes and J.B. Marais, 1981. Groeitoets met kuikens vir die lisiengehalte van vis-, grondboonoliekoek-en sonneblomoliekoekmeel. *S. Afr. J. Anim. Sci.* 11:77-84.

- 231 Raharjo, Y., and D.J. Farrell, 1981. The determination of amino acid digestibility in poultry feedstuffs. Recent Advances in Animal Nutrition in Australia 1981. Univ. New England, Armidale, Australia. pp. 197-218.
- 232 Ranaweera, K.N.P., and W.E. Nano, 1981. True and apparent metabolizable energy of some indigenous feedstuffs and finished feeds determined by modified rooster bioassay techniques. J. Agric. Sci. Camb. 97:403-407.
- 233 *Ravindran, V., E.T. Kornegay, L.M. Potter, K.E. Webb, Jr. and C.M. Parsons, 1981. Nutrient composition and true metabolizable energy values of three tropical feedstuffs. J. Anim. Sci. 53(Suppl. 1):258.
- 234 Reid, B.L., 1981. Fat levels in layer feeds. J. Am. Oil Chem. Soc. 58:306-309.
- 235 Salmon, R.E., and T.I. Szabo, 1981. Dried bee meal as a feedstuff for growing turkeys. Can. J. Anim. Sci. 61:965-968.
- 236 Salmon, R.E., E.E. Gardiner, K.K. Klein and E. Larmond, 1981. Effect of Canola (low glucosinolate rapeseed) meal, protein and nutrient density on performance, carcass grade, and meat yield, and of Canola meal on sensory quality of broilers. Poultry Sci. 60:2519-2528.
- 237 *Salmon, R.E., E.E. Gardiner, K.K. Klein and E. Larmond, 1981. Canola meal in diets for broiler chickens. Poultry Sci. 60:1723.
- 238 *Schang, M.J., and R.M.G. Hamilton, 1981. Comparison of two direct bioassays using adult cockerels for estimating the available energy content of 13 feedingstuffs. Poultry Sci. 60:1726-1727.
- 239 *Schang, M.J., R.M.G. Hamilton, J.D. Summers and M.F. Bonino, 1981. Estimation of available energy in feedstuffs from metabolizable dry matter. Poultry Sci. 60:1726.
- 240 Shires, A., J.M. Bell and A.R. Robblee, 1981. Effect of the age of chickens on the true metabolizable energy values of feedstuffs. 60th Annu. Feeder's Day Rept., Univ. Alberta, Edmonton, Canada. pp. 78-79.
- 241 Sibbald, I.R., 1981. Metabolic plus endogenous energy and nitrogen losses of adult cockerels: the correction used in the bioassay for true metabolizable energy. Poultry Sci. 60:805-811.
- 242 Sibbald, I.R., 1981. Metabolic plus endogenous energy excretion by fowl. Poultry Sci. 60:2672-2677.
- 243 *Sibbald, I.R., 1981. Basic assumptions of the true metabolizable energy (TME) bioassay. Poultry Sci. 60:1733.
- 244 Sibbald, I.R., 1981. Valores de energia de grasas para alimentas aves. Industria Avicola 28(11):8-20.

- 245 Sibbald, I.R., 1981. Bioassays Based on Precision Feeding of Poultry/
Dosage biologique en cage métabolique chez la volaille. Tech. Bull.
No.3, Animal Research Centre, Agriculture Canada, Ottawa, Canada.
34 pp.
- 246 Sibbald, I.R., 1981. Energy values of feeding fats for poultry.
Feedstuffs 53(23):19-20.
- 247 Sibbald, I.R., 1981. Energy values of feeding fats for poultry. Poultry
Digest 40:262-266.
- 248 Sibbald, I.R., 1981. The true metabolizable energy and related bioassays.
Proc. Minnesota Nutr. Conf., Minneapolis, U.S.A. pp. 105-112.
- 249 Sibbald, I.R., 1981. Advantages of the TME system. Proc. Minnesota Nutr.
Conf., Minneapolis, U.S.A. pp. 113-118.
- 250 Sibbald, I.R., 1981. Energy values of feeding fats for poultry. Proc.
Natl. Renderers Assoc. Nutr. Conf., Atlanta, U.S.A.
- 251 Sibbald, I.R., 1981. Sistema para la evaluacion de las materias primas en
las aves. Proc. Symp. Cientifico, Seccion Espanola, Asoc. Mundial
Avicult. Cientif., Barcelona, Spain. pp. 337-361.
- 252 Storey, M.L., 1981. The true metabolizable energy and the factors
affecting the true metabolizable energy of feedstuffs for mature
non-laying female Embden geese. Ph.D. Thesis, Univ. Minnesota,
St. Paul, U.S.A.
- 253 *Storey, M.L., and N.K. Allen, 1981. Effect of duration of fasting on the
true metabolizable energy of corn for Embden geese. Poultry Sci.
60:1739.
- 254 *Storey, M.L., and N.K. Allen, 1981. True metabolizable energy of
feedstuffs for Embden geese. Poultry Sci. 60:1739-1740.
- 255 Tenesaca, L.G., and J.L. Sell, 1981. Influence of an indigestible
material on energy excretion by roosters and on true metabolizable
energy of corn. Poultry Sci. 60:623-630.
- 256 Tyrrell, H.F., L.S. Bull, W.N. Garrett, I.R. Sibbald and R.R. Smith, 1981.
Nutritional Energetics of Domestic Animals and Glossary of Energy
Terms. National Academy Press, Washington, U.S.A. 54 pp.
- 257 Van Es, A.J.H., 1981. Poultry production in relation to energy
utilization and environment. pp. 41-53 in C.W. Scheele and C.H.
Veerkamp, eds. World Poultry Production Where and How? Spelderholt
Inst. for Poult. Res., Beekbergen, The Netherlands.
- 258 Yeong, S.W., 1981. Biological utilization of palm oil by-products by
chickens. Ph.D. Thesis, Univ. Malaysia, Selangor, Malaysia.

1982

- 259 Anonymous, 1982. Energy evaluation. Poultry Research Centre Report for Year Ending 31st March 1982. Agric. Res. Council, London, U.K. pp. 4-5.
- 260 Anonymous, 1982. TME tests urged on non-U.S. feedstuffs. Feedstuffs 54(5):13.
- 261 Anonymous, 1982. Purina introduces new Turkey Chow. Feedstuffs 54(10):57.
- 262 Atwal, A.S., and S.C. Stothers, 1982. Assay of amino acid availability in swine and poultry. Proc. Nutr. Conf. Feed Manuf., Toronto, Canada. pp. 59-65.
- 263 Baghel, R.P.S., and S.P. Netke, 1982. Studies on incorporation of kangni (Setaria italica) in starter chick diets. Indian J. Anim. Sci. 52:411-417.
- 264 Bilgili, S.F., and G.H. Arscott, 1982. The effect of initial fasting period on gut clearances and on endogenous energy outputs of adult roosters. Nutr. Rep. Int. 25:613-619.
- 265 Bilgili, S.F., G.H. Arscott and R. Kellems, 1982. True metabolizable energy value of hydrolyzed Pacific Whiting (Merluccius productus) as determined with gelatin capsules in roosters. Nutr. Rep. Int. 25:639-642.
- 266 *Bilgili, S.F., M.P. Goeger and G.H. Arscott, 1982. Effect of inorganic matter (granite grit) on endogenous energy and true metabolizable energy values. Poultry Sci. 61:1417-1418.
- 267 Campbell, L.D., B.O. Eggum and J. Wolstrup, 1982. Influence of dietary antibiotics on nitrogen, amino acid and energy digestion by rats and adult roosters. Can. J. Anim. Sci. 62:537-545.
- 268 Chubb, L., 1982. Determining the M.E. of poultry feed. AgriTrade May:27-28.
- 269 Dale, N.M., and H.L. Fuller, 1982. Applicability of the true metabolizable energy system in practical feed formulation. Poultry Sci. 61:351-356.
- 270 Dale, N.M., and H.L. Fuller, 1982. Endogenous energy losses of fed versus fasted roosters. Poultry Sci. 61:898-901.
- 271 Dale, N.M., and H.L. Fuller, 1982. True metabolizable energy of fats at low levels of dietary inclusion. Poultry Sci. 61:2415-2420.
- 272 *Dale, N.M., and H.L. Fuller, 1982. True metabolizable energy and fatty acid absorption of fats at low level dietary inclusion. Poultry Sci. 61:1445.

- 273 *Delfino, F.J., A.R. Urias, L.G. Schmid, F.L. Prouty, R.S. Swingle and B.L. Reid, 1982. Variation in chemical composition and TME of grain from different wheat genotypes. *Poultry Sci.* 61:1448.
- 274 Donaldson, W.E., 1982. Poultry nutrition research 1981. 30th Pfizer Annu. Res. Conf., Dallas, U.S.A. pp. 73-100.
- 275 Edmundson, I.C., 1982. True metabolizable energy trial ME/14. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 96-97.
- 276 Edmundson, I.C., 1982. True metabolizable energy trial ME/15. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. p. 98.
- 277 Edmundson, I.C., 1982. True metabolizable energy trial ME/16. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. p. 110.
- 278 *El Husseiny, O., 1982. True metabolizable energy of some Egyptian fats. *Poultry Sci.* 61:1459.
- 279 Farrell, D.J., 1982. True metabolizable energy. Is it a significant advance in poultry energy systems? Proc. 1982 Maryland Nutr. Conf., Washington, U.S.A. pp. 79-87.
- 280 Farrell, D.J., 1982. Amino acid digestibility of Australian feedstuffs determined with cockerels. Proc. Poult. Husb. Res. Found. Symp., Sydney, Australia. 11 pp.
- 281 Fisher, C., 1982. Energy evaluation of poultry rations. pp. 113-139 in W. Haresign and D. Lewis, eds. Recent Advances in Animal Nutrition - 1982. Butterworths, London, U.K.
- 282 Fisher, C., 1982. Energy values of compound poultry feeds. Agric. Res. Council, Poultry Res. Cent., Roslin, U.K. Occasional Publ. #2, 109 pp.
- 283 Fuller, H.L., and N.M. Dale, 1982. Effect of ratio of basal diet fat to test fat on the true metabolizable energy of the test fat. *Poultry Sci.* 61:914-918.
- 284 *Fuller, H.L., and N.M. Dale, 1982. The value of brewers condensed solubles in broiler and layer rations. *Poultry Sci.* 61:1468.
- 285 Gous, R.M., M. Kuyper and C. Dennison, 1982. The relationship between tannic acid content and metabolisable energy concentration of some sorghum cultivars. *S. Afr. J. Anim. Sci.* 12:39-44.
- 286 *Halley, J.T., T.S. Nelson, L.K. Kirby and Z.B. Johnson, 1982. Relation between dry matter digestion and metabolizable energy. *Poultry Sci.* 61:1474.

- 287 Han, Y.W., and C.K. Wu, 1982. Metabolizable energy studies of poultry feeds. 3. Direct and rapid determination of metabolizable energy value of poultry feedstuffs. *J. Northeast Agric. Coll.* (Peoples Republic of China) 4:27-32.
- 288 *Hayes, J.P., and J.J. du Preez, 1982. Digestibility of lysine in heat-damaged fish meal by normal and colostomized roosters. *S. Afr. J. Anim. Sci.* 12:298.
- 289 Hulan, H.W., F.G. Proudfoot and C.G. Zarkadas, 1982. Potato waste meal. I. Compositional analyses. *Can. J. Anim. Sci.* 62:1161-1169.
- 290 King, R.D., 1982. Chick nutrition 193 - Amino acid availability by chick assay. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 1-6.
- 291 King, R.D., 1982. Chick nutrition 207 -- Amino acid/energy ratios for growers. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 13-18.
- 292 King, R.D., 1982. Layer nutrition 171 - Tryptophan requirements. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 49-50.
- 293 King, R.D., 1982. Layer nutrition 169 - Amino acid/energy ratios for growers. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 57-58.
- 294 Kussabati, R., J. Guillaume and B. Leclercq, 1982. The effects of age, dietary fat and bile salts, and feeding rate on apparent and true metabolisable energy values in chickens. *Br. Poult. Sci.* 23:393-403.
- 295 Kussabati, R., J. Guillaume and B. Leclercq, 1982. The effects of endogenous energy, type of diet, and addition of bile salts on true metabolizable energy values in young chicks. *Poultry Sci.* 61:2218-2223.
- 296 Kussabati, R., J. Guillaume, B. Leclercq and J.P. Lafont, 1982. Effects of the intestinal microflora and added bile salts on the metabolizable energy and digestibility of saturated fats in the chicken. *Arch. Geflugelkd.* 46:42-46.
- 297 Lessire, M., and B. Leclercq, 1982. Metabolizable energy value of fats in chicks and adult cockerels. *Anim. Feed Sci. Technol.* 7:365-374.
- 298 Likuski, H.J.A., 1982. Bioavailabilities of amino acids in poultry feedstuffs as determined by precision-fed rooster assays. *Proc. Nutr. Conf. Feed Manuf.*, Toronto, Canada. pp. 50-58.
- 299 Likuski, H.J.A., 1982. Implication of amino acid availability for poultry. *Proc. 3rd Western Nutr. Conf.*, Winnipeg, Canada. pp. 29-38.

- 300 Luis, E.S., T.W. Sullivan and L.A. Nelson, 1982. Nutrient composition and feeding value of proso millets, sorghum grains and corn in broiler diets. *Poultry Sci.* 61:311-320.
- 301 McNab, J.M., 1982. The choice between apparent and true metabolizable energy systems for poultry. Recente toename van het gebruik van olien en velten in de veevoeding. Am. Soybean Assoc. Conf., Arnhem, The Netherlands. pp. 1-8.
- 302 Mollah, Y., W.L. Bryden, D. Balnave and E.F. Annison, 1982. Comparison of methods for the bioassay of the metabolizable energy of wheat-based poultry diets. Proc. Poult. Husb. Res. Found. Symp., Sydney, Australia. 6 pp.
- 303 Muztar, A.J., and S.J. Slinger, 1982. The true metabolizable energy and amino acid content of Candle, Altex and Regent Canola meals. *Can. J. Anim. Sci.* 62:521-525.
- 304 Ngoupayou, J.D.N., P.M. Maiorino and B.L. Reid, 1982. Jojoba meal in poultry diets. *Poultry Sci.* 61:1692-1696.
- 305 Okumura, J., Y. Isshiki and Y. Nakahira, 1982. Influence of dietary cellulose and indigestible dry matter on metabolic and endogenous nitrogen excretion in chickens. *Jpn. Poult. Sci.* 19:300-303.
- 306 Parsons, C.M., L.M. Potter and B.A. Bliss, 1982. True metabolizable energy corrected to nitrogen equilibrium. *Poultry Sci.* 61:2241-2246.
- 307 *Parsons, C.M., L.M. Potter and B.A. Bliss, 1982. True metabolizable energy corrected to nitrogen equilibrium. *Poultry Sci.* 61:1523.
- 308 Parsons, C.M., L.M. Potter and R.D. Brown, Jr., 1982. Effects of dietary protein and intestinal microflora on excretion of amino acids in poultry. *Poultry Sci.* 61:939-946.
- 309 Parsons, C.M., L.M. Potter, R.D. Brown, Jr., T.D. Wilkins and B.A. Bliss, 1982. Microbial contribution to dry matter and amino acid content of poultry excreta. *Poultry Sci.* 61:925-932.
- 310 Patchell, M.R., 1982. Chick nutrition 208 - Systems of rearing. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 19-25.
- 311 Patchell, M.R., 1982. Chick nutrition 209 - Dry and low temperature rendered meat and bone meals. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 26-29.
- 312 Patchell, M.R., 1982. Layer nutrition 158 - Layer phase of CN 190. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. p. 32.

- 313 Patchell, M.R., 1982. Layer nutrition 164 - Strain comparison. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 33-35.
- 314 Patchell, M.R., 1982. Layer nutrition 165 - Energy levels of diets. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 36-37.
- 315 Patchell, M.R., 1982. Layer nutrition 168 - Amino acids + models 3/5. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 44-48.
- 316 Patchell, M.R., 1982. Layer nutrition 170 - Available amino acids. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 51-56.
- 317 Patchell, M.R., 1982. Layer nutrition 172 - Low protein at end of lay. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 59-62.
- 318 Patchell, M.R., 1982. Layer nutrition 173 - Methionine and isoleucine interaction. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 63-65.
- 319 Patchell, M.R., 1982. Layer nutrition 175 - Energy utilization and tallow. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 72-75.
- 320 Patchell, M.R., 1982. Layer nutrition 176 - Amino acid availability. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 76-80.
- 321 Patchell, M.R., 1982. Layer nutrition 178 - Energetic efficiency. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 81-83.
- 322 Patchell, M.R., 1982. Broiler nutrition trial BG 5 - Apex meal. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 90-94.
- 323 Patchell, M.R., 1982. Layer nutrition 184 - Balance trial with available amino acids. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. p. 105.
- 324 Patchell, M.R., 1982. Feed ingredients. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. p. 112.
- 325 Patchell, M.R., 1982. TME and amino acid availability values. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 127-139.

- 326 Pearson, A.R., and K.M. Herron, 1982. Relationship between energy and protein intakes and laying characteristics in individually-caged broiler breeder hens. *Br. Poult. Sci.* 23:145-159.
- 327 Pesti, G.M., and H.M. Edwards, Jr., 1982. Metabolizable energy: Questions of nomenclature and application. *Poultry Digest* 41:228-231.
- 328 Pesti, G.M., and H.M. Edwards, Jr., 1982. Metabolizable energy: questions of nomenclature and application. *Proc. Georgia Nutr. Conf.*, Atlanta, U.S.A. pp. 69-73.
- 329 Salmon, R.E., K.E. Dunkelgod and B.J. Wilson, 1982. Influence of dietary protein concentration and frequency of diet changes on rate of growth, efficiency of food utilization and carcass quality of Large White turkeys. *Br. Poult. Sci.* 23:501-517.
- 330 Schang, M.J., and R.M.G. Hamilton, 1982. Comparison of two direct bioassays using adult cocks and four indirect methods for estimating the metabolizable energy content of different feedingstuffs. *Poultry Sci.* 61:1344-1353.
- 331 Schang, M.J., S. Leeson and J.D. Summers, 1982. Effect of starvation and force feeding on body weight recovery time of adult roosters. *Can. J. Anim. Sci.* 62:307-309.
- 332 Schang, M.J., S. Leeson and J.D. Summers, 1982. Effect of double force feeding on true metabolizable energy. *Can. J. Anim. Sci.* 62:963-965.
- 333 Shannon, D.W.F., 1982. Developments in ME methodology and their implications for practical poultry feeding. *Proc. Poult. Husb. Res. Found. Symp.*, Sydney, Australia. 9 pp.
- 334 *Shen, T., and W.F. Dean, 1982. True metabolizable energy value of corn and soybean meal for ducks. *Poultry Sci.* 61:1543.
- 335 *Shires, A., and J.M. Bell, 1982. Effect of the age of chickens, duration of excreta collection and glucosinolate content on the true metabolizable energy values of rapeseed meal. *Poultry Sci.* 61:1543-1544.
- 336 Sibbald, I.R., 1982. Measurement of bioavailable energy in poultry feedingstuffs. *Can. J. Anim. Sci.* 62:983-1048.
- 337 Sibbald, I.R., 1982. Effects of tea and coffee on excretion of energy and nitrogen by cockerels. *Nutr. Rep. Int.* 25:133-137.
- 338 Sibbald, I.R., 1982. Measurement of mineral bioavailability: extension of true metabolizable energy methodology. *Poultry Sci.* 61:485-487.
- 339 Sibbald, I.R., 1982. The effect of grinding on the true metabolizable energy value of hull-less barley. *Poultry Sci.* 61:2509-2511.

- 340 Sibbald, I.R., 1982. Evolution of the TME system. *Feed International* 3(2):51-52.
- 341 Sibbald, I.R., 1982. Evolution of the TME system. *Feed Management* 33(1):12-18.
- 342 Sibbald, I.R., and P.M. Morse, 1982. Variation among gross energy values measured by two modes of adiabatic oxygen bomb calorimetry. *Poultry Sci.* 61:994-997.
- 343 Sibbald, I.R., and P.M. Morse, 1982. Pooling excreta prior to calorimetry in the bioassay for true metabolizable energy: the effects on estimates of variance. *Poultry Sci.* 61:1853-1858.
- 344 *Sibbald, I.R., and P.M. Morse, 1982. Effects of feed input and excreta collection time on estimates of metabolic plus endogenous energy losses. *Poultry Sci.* 61:1544.
- 345 *Sibbald, I.R., and P.M. Morse, 1982. Effects of a nitrogen correction and of feed input on true metabolizable energy (TME) values. *Poultry Sci.* 61:1545.
- 346 Storey, M.L., and N.K. Allen, 1982. The effect of fasting time of mature, non-laying female Embden geese on true metabolizable energy of corn. *Poultry Sci.* 61:101-106.
- 347 Storey, M.L., and N.K. Allen, 1982. Apparent and true metabolizable energy of feedstuffs for mature, non-laying female Embden geese. *Poultry Sci.* 61:739-745.
- 348 Tenesaca, G., 1982. Contenido de energia metabolizable verdadera en ingredientes utilizados en formulacion de dietas avicolas. *Vi Cielo Internacional de Conferencias Sobre Avicultura.* Sept. 1982, Unidad de Congresos del Centro Medico Nacional, Mexico, D.F. pp. 92-98.
- 349 *Thomas, V.M., R.J. Katz, D.A. Auld, C.F. Petersen and E.A. Sauter, 1982. Feeding value of expeller extracted rape and safflower oilseed meals for broilers. *Poultry Sci.* 61:1555.
- 350 Vohra, P., D.B. Chami and E.O. Oyawoye, 1982. Determination of metabolizable energy by a fast method. *Poultry Sci.* 61:766-769.
- 351 Wehner, G.R., and R.L. Harrold, 1982. The effect of feeding techniques on the true metabolizable energy value of yellow corn. *Poultry Sci.* 61:595-597.
- 352 Wehner, G.R., and R.L. Harrold, 1982. Crop volume of chickens as affected by body size, sex, and breed. *Poultry Sci.* 61:598-600.
- 353 Yamazaki, M., and Z. Zi-Yi, 1982. A note on the effect of temperature on true and apparent metabolisable energy values of a layer diet. *Br. Poultry Sci.* 23:447-450.

- 354 Yanaka, M., and J. Okumura, 1982. Effects of dietary excess tyrosine and level of feed intake on nitrogen and energy utilization in chicks. *Poultry Sci.* 61:2440-2446.

1983

- 355 Anonymous, 1983. Poultry Research Centre Report for the Year Ending 31st March 1984. Agric. Food Res. Council, London, U.K. pp. 4-5.
- 356 Anonymous, 1983. Modificaciones en la prueba para energia metabolizable verdadera. *Avicultura Profesional* 1(1):22-23.
- 357 Anonymous, 1983. An ME challenge goes with this full-fat soya. *Poultry World* 137(34):8.
- 358 Austic, R.E., 1983. Assessment of amino acid availability for poultry. *Proc. Cornell Nutrition Conf.*, Syracuse, U.S.A. pp. 75-78.
- 359 Austic, R.E., 1983. The availability of amino acids as an attribute of feeds. pp. 175-189 in G.E. Robards and R.G. Packham, eds. *Proc. 2nd Symp. Int. Network Feed Inf. Cent.*, Commonw. Agric. Bur., London, U.K.
- 360 Bielorai, R., Z. Harduf, B. Iosif and E. Alumot, 1983. Apparent amino acid absorption from feather meal by chicks. *Br. J. Nutr.* 49:395-399.
- 361 *Caldwell, J.M., J.M. Vandepopuliere, J.J. Lyons and G. Krause, 1983. True metabolizable energy value of two types of methane digester effluent. *Poultry Sci.* 62:1395.
- 362 Campbell, G.L., L.D. Campbell and R. Blair, 1983. Calculation of metabolizable energy for ingredients incorporated at low levels into a reference diet. *Poultry Sci.* 62:705-707.
See also erratum *Poultry Sci.* 62:2289.
- 363 Campbell, L.D., I. Jacobsen, B.O. Eggum and A. Just, 1983. A study of the variability of the endogenous energy output by adult roosters and a determination of the available energy of nine different feedingstuffs. *J. Sci. Food Agric.* 34:221-226.
- 364 Clandinin, D.R., and A.R. Robblee, 1983. Apparent and true metabolizable energy values for low glucosinolate-type rapeseed meal. *Feedstuffs* 55(5):20.
- 365 Coelho, M.dasG.R., H.S. Rostagno, J.B. Fonseca and D.J. da Silva, 1983. Composição química e valores energéticos de alguns alimentos determinados com pintos e galos. VIII Congreso Latinoamericano de Avicultura, Camboriú, Brasil.
- 366 Corazza, L., and W.W. Saylor, 1983. Nutritional value of Artemia salina in broiler starter diets. *Poultry Sci.* 62:846-852.
- 367 *Crissey, S.D., and O.P. Thomas, 1983. The amount of fecal amino acids from roosters fasted, fed non-protein diets, soybean meal or autoclaved soybean meal. *Poultry Sci.* 62:1406.

- 368 *Dale, N.M., and H.L. Fuller, 1983. Corn oil, poultry fat energy levels weighed. *Broiler Ind.* 46(3):62.
- 369 *Dale, N.M., and H.L. Fuller, 1983. Energetic contribution of fats to practical diets. *Poultry Sci.* 62:1342.
- 370 *Dale, N., and H.L. Fuller, 1983. Oven drying vs freeze drying of excreta in true amino acid availability and true metabolizable energy assays. *Poultry Sci.* 62:1407-1408.
- 371 Dale, N.M., and H.L. Fuller, 1983. Nitrogen corrected true metabolizable energy. *Fed. Proc.* 43:668.
- 372 Damron, B.L., 1983. Poultry nutrition research 1982. 30th Pfizer Annu. Res. Conf., Kansas City, U.S.A. pp. 72-105.
- 373 Edmundson, I.C., and G.D. Watts, 1983. True metabolisable energy trial ME/16. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 56-58.
- 374 Fisher, C., 1983. Energy evaluation of poultry rations. pp. 113-139 in W. Haresign and D. Lewis, eds. *Recent Advances in Animal Nutrition*. Butterworths, London, U.K.
- 375 Fisher, C., 1983. The derivation of chemical prediction equations for monitoring energy declarations. pp. 294-306 in D.J. Farrell and P. Vohra, eds. *Recent Advances in Nutrition in Australia*, Univ. New England, Armidale, Australia.
- 376 Fisher, C., 1983. Report on a ring-test to evaluate the proposed EEC equation for declaration of ME values, poultry. Agric. Res. Council, Poultry Res. Cent., Roslin, U.K. Occasional Publ. #3, 34 pp.
- 377 Fisher, C., 1983. Putting an energy value on feed. *Poultry Int.* 22(4):64-67.
- 378 Fraser, D., and I.R. Sibbald, 1983. The effects of precision feeding on behaviour of adult cockerels. *Poultry Sci.* 62:2224-2226.
- 379 Fuller, H.L., and N.M. Dale, 1983. Feeding of brewers' condensed solubles for broilers and laying hens. *Poultry Sci.* 62:914-916.
- 380 Goeger, M.P., and H.S. Nakae, 1983. True metabolizable energy of two fat blends and yellow grease. *Nutr. Rep. Int.* 27:977-981.
- 381 Gomez, M., 1983. Nutritional characteristics of some selected nonconventional feedstuffs: their acceptability, improvement, and potential use in poultry feeds. pp. 71-81 in B. Kiflewahid, G.R. Potts and R.M. Drysdale, eds. *By-product Utilization for Animal Production*. Int. Develop. Res. Cent. Publ. 206e, Ottawa, Canada.

- 382 Gous, R.M., and C. Dennison, 1983. The metabolizable energy content of some South African feedingstuffs evaluated with poultry. S. Afr. J. Anim. Sci. 13:147-153.
- 383 Halloran, H.R., 1983. Calorie vs. joule, conformity or communication. The bottom line. Feedstuffs 55(27):10.
- 384 Halvorson, J.C., M.A. Shehata and P.E. Waibel, 1983. White lupins and triticale as feedstuffs in diets for turkeys. Poultry Sci. 62:1038-1044.
- 385 Hamilton, R.M.G., H.L. Trenholm and B.K. Thompson, 1983. Effects of feeding chicks and laying hens diets that contain vomitoxin (deoxynivalenol) contaminated wheat. Proc. Guelph Nutr. Conf., Toronto, Canada. pp. 48-55.
- 386 Han, I.K., M.K. Song, D.S. Nam, and K.H. Lee, 1983. Studies on the TME bioassay of poultry feedstuffs. I. The effect of age of bird on true metabolizable energy value of corn. Korean J. Anim. Sci. 25:634-638.
- 387 Han, I.K., M.K. Song, D.S. Nam, and K.H. Lee, 1983. Studies on the TME bioassay of poultry feedstuffs. II. The effect of breed and sex of assay bird on true metabolizable energy value of corn. Korean J. Anim. Sci. 25:639-643.
- 388 Han, I.K., M.K. Song, D.S. Nam, and K.H. Lee, 1983. Studies on the TME bioassay of poultry feedstuffs. III. The effect of and level of feed input feeding method on true metabolizable energy value of corn. Korean J. Anim. Sci. 25:644-650.
- 389 Johns, D.C., and K.A.C. James, 1983. Amino acid availability. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 65-67.
- 390 Jonsson, G., and J.M. McNab, 1983. A comparison of methods for estimating the metabolizable energy of a sample of grass meal. Br. Poult. Sci. 49:349-359.
- 391 King, R.D., 1983. Layer nutrition 182 - Malt culms. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 22-23.
- 392 King, R.D., 1983. Layer nutrition 190 - Amino acid dietary levels for growers. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 34-35.
- 393 King, R.D., 1983. Chick nutrition 212 - Amino acid dietary levels for growers. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 44-46.
- 394 King, R.D., 1983. A model analysing the relationship between true metabolisable energy (TME) and nitrogen correction. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 81-87.

- 395 Kussabati, R., B. Prevotel and B. Leclercq, 1983. Factors affecting endogenous energy losses in chickens - assessment of the methods of estimation. Nutr. Rep. Int. 27:221-230.
- 396 Latshaw, J.D., 1983. Methods of measuring the digestibility of energy materials and other nutrients in feed. Ohio Poult. Pointers 22(1). Ohio State Univ., U.S.A.
- 397 Latshaw, J.D., 1983. Measuring digestibility of energy materials. Poultry Digest 42:170.
- 398 Lessire, M., and B. Leclercq, 1983. Metabolizable energy of meat meal for chicken. Arch. Gefluegelkd. 47:1-3.
- 399 Likuski, H.J.A., 1983. Chemical and biological assays used commercially in determining the energy and nutrient contents of feedstuffs. Monsanto Symp. Present and Future Trends Anim. Nutr. Feed Manuf. Technol. Hot Springs, Arkansas, U.S.A. pp. 53-59.
- 400 *McFarland, D.C., and C.N. Coon, 1983. The evaluation of by-product protein digestibility utilizing coomassie blue dye binding. Poultry Sci. 62:1468.
- 401 *McNab, J.M., 1983. The metabolizable energy value of manioc for poultry. World's Poult. Sci. J. 39:241.
- 402 Minaar, A.duP., D.K. Thomson, J.J. du Preez and J.P. Hayes, 1983. Evaluering van verskillende kommersiele prasesseringsmetodes op die voedingswaarde van pluimveeafvalmeel. S. Afr. J. Anim. Sci. 13:154-156.
- 403 Mollah, Y., W.L. Bryden, I.R. Wallis, D. Balnave and E.F. Annison, 1983. Studies on low metabolizable energy wheats for poultry using conventional and rapid assay procedures and the effects of processing. Br. Poult. Sci. 24:81-89.
- 404 Myung, K.H., S.S. Sun and Y.S. Kim, 1983. Studies on calcium bioavailability by true available mineral methodology. 1. Measurement of calcium requirement in breeding Japanese quails. Korean J. Anim. Sci. 25:680-685.
- 405 Ostrowski-Meissner, H.T., 1983. Effect of dietary aflatoxins on protein and energy utilization by two Indonesian breeds of ducklings (Alabio and Tegal). Poultry Sci. 62:599-607.
- 406 Ostrowski-Meissner, H.T., 1983. Some problems of evaluating feed quality for poultry in the tropics. pp. 491-496 in G.E. Robards and R.G. Packham, eds. Proc. 2nd Symp. Int. Network Feed Inf. Cent., Commonw. Agric. Bur., London, U.K.
- 407 Parsons, C.M., D.H. Baker and J.M. Harter, 1983. Distillers dried grains with solubles as a protein source for the chick. Poultry Sci. 62:2445-2451.

- 408 Parsons, C.M., L.M. Potter and R.D. Brown, Jr., 1983. Effects of dietary carbohydrate and of intestinal microflora on excretion of endogenous amino acids by poultry. *Poultry Sci.* 62:483-489.
- 409 *Parsons, C.M., L.M. Potter and B.A. Bliss, 1983. A modified voluntary feed intake bioassay for determination of metabolizable energy with Leghorn roosters. *Poultry Sci.* 62:1480.
- 410 Patchell, M.R., 1983. Layer nutrition 176 - Amino acid availability. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 1-7.
- 411 Patchell, M.R., 1983. Layer nutrition 183 - Available amino acids. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 24-30.
- 412 Patchell, M.R., 1983. Layer nutrition 184-186 - Energy utilization of diets of L/183. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 31-33.
- 413 Patchell, M.R., 1983. Layer nutrition 191 - Isoleucine and lysine. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 36-40.
- 414 Patchell, M.R., 1983. Layer nutrition 193a, b, c -- Energy utilization of diets. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 41-43.
- 415 Patchell, M.R., 1983. Layer nutrition 194 -- Available amino acids. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 76-78.
- 416 Pesti, G.M., and H.M. Edwards, 1983. Metabolizable energy nomenclature for poultry feedingstuffs. *Poultry Sci.* 62:1275-1280.
- 417 *Picard, M., S. Bertrand, M. Duron, and R. Maillard, 1983. Comparative digestibility of amino acids using 5 animal models -- intact cockerels, caecectomised cockerels, rats deprived of large intestine, piglets with an ileo-caecal canulation, piglets with an ileo-rectal shunt. Proc. 4th Eur. Symp. Poult. Nutr. Tours, France. p. 165.
- 418 Ravindran, V., E.T. Kornegay, L.M. Potter, K.E. Webb, Jr. and C.M. Parsons, 1983. True metabolizable energy values of cassava tuber and leaf meals for poultry. *Trop. Agric.(Trinidad)* 60:82-84.
- 419 Reid, B.L., 1983. Tallow for laying hens. *Poultry Tribune* 89(8):36-40.
- 420 Salmon, R.E., 1983. Effect of duration of excreta collection on, and additivity of, true metabolizable energy values of feedstuffs. *Nutr. Rep. Int.* 27:655-660.
- 421 Salmon, R.E., 1983. The effect of the energy protein ratio of the diet, strain and age at slaughter on performance and carcass quality of turkey broilers. *Poultry Sci.* 62:824-831.

- 422 Salmon, R.E., H.L. Classen and R.K. McMillan, 1983. Effect of starter and finisher protein on performance, carcass grade, and meat yield of broilers. *Poultry Sci.* 62:837-845.
- 423 Schang, M., 1983. Nuevo rumbo en la nutricion. *Balanceados Argentinos* 1(10):6-8.
- 424 *Schang, M.J., and F. Borras, 1983. Energia metabolizable verdadera de algunos ingredientes Argentinos. IX Reunion Latinoamericana de Produccion Animal (NR-32), Santiago, Chile.
- 425 Schang, M.J., I.R. Sibbald and R.M.G. Hamilton, 1983. Comparison of two direct bioassays using young chicks and two internal indicators for estimating the metabolizable energy content of feedingstuffs. *Poultry Sci.* 62:117-124.
- 426 Sibbald, I.R., 1983. The T.M.E. System of Feed Evaluation/La méthode d'évaluation des aliments par le calcul de l'énergie métabolisable vraie (E.M.V.). Research Branch Contribution 83-20E/83-20F, Animal Research Centre, Agriculture Canada, Ottawa, Canada. 89 pp.
- 427 Sibbald, I.R., 1983. The current status of TME. pp. 286-293 in D.J. Farrell and P. Vohra, eds. *Recent Advances in Nutrition in Australia*. Univ. New England, Armidale, Australia.
- 428 Sibbald, I.R., and P.M. Morse, 1983. The effects of feed input and excreta collection time on estimates of metabolic plus endogenous energy losses in the bioassay for true metabolizable energy. *Poultry Sci.* 62:68-76.
- 429 Sibbald, I.R., and P.M. Morse, 1983. Effects of the nitrogen correction and of feed intake on true metabolizable energy values. *Poultry Sci.* 62:138-142.
- 430 Sibbald, I.R., and P.M. Morse, 1983. Provision of supplemental feed and the application of a nitrogen correction in bioassays for true metabolizable energy. *Poultry Sci.* 62:1587-1605.
- 431 Sibbald, I.R., and P.M. Morse, 1983. Recent developments in the bioassay for true metabolizable energy in poultry feedingstuffs. pp. 239-242 in G.E. Robards and R.G. Packham, eds. *Proc. 2nd Symp. Int. Network Feed Inf. Cent., Commonw. Agric. Bur., London, U.K.*
- 432 Sibbald, I.R., J.I. Elliot and M. Coté, 1983. Comparison of bioavailable energy values of diets measured with poultry and swine. *Anim. Feed Sci. Technol.* 8:191-196.
- 433 Stevenson, M.H., and W.D. Graham, 1983. The chemical composition and true metabolizable energy content of cassava root meal imported into Northern Ireland. *J. Sci. Food Agric.* 34:1005-1007.
- 434 Thomas, V.M., R.J. Katz, D.A. Auld, C.F. Petersen, E.A. Sauter and E.E. Steele, 1983. Nutritional value of expeller extracted rape and safflower oilseed meals for poultry. *Poultry Sci.* 62:882-886.

- 435 Van Kampen, M., 1983. Effect of water:food intake ratios in laying hens on food metabolizable energy. Br. Poult. Sci. 24:169-172.
- 436 Wallis, I., and D. Balnave, 1983. A comparison of different drying techniques for energy and amino acid analyses of poultry excreta. Br. Poult. Sci. 24:255-260.
- 437 Watts, G.D., 1983. True metabolisable energy trial ME/17. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 59-60.
- 438 Wiseman, J., 1983. Assessment of the digestible and metabolizable energy of fats for non-ruminants. pp. 277-297 in J. Wiseman ed. Fats in Animal Nutrition. Butterworths, London, U.K.
- 439 Yamazaki, M., 1983. A comparison of two methods in determining amino acid availability of feed ingredients. Jpn. J. Zootechn. Sci. 54:729-733.

1984

- 440 Achmanu, 1984. Perbandingan cara pemberian makan secara paksa dan normal terhadap nilai "apparent dan true metabolizable energy." Nuffic-Unibraw/Animal Husbandry Misc. Paper 21. Universitas Brawijaya, Malang, Indonesia. (Comparison of forced and normal feeding on metabolizable energy values in ducks.)
- 441 Achmanu, 1984. Kandungan enersi metabolismis dedak padi pada ayam dan itik. Nuffic-Unibraw/Animal Husbandry Misc. Paper 22. Universitas Brawijaya, Malang, Indonesia. (Metabolizable energy values of ricebran in chickens and ducks.)
- 442 Achmanu, 1984. Pertambahan nilai enersi metabolismis campuran jagung dan dedak padi pada itik. Nuffic-Unibraw/Animal Husbandry Misc. Paper 23. Universitas Brawijaya, Malang, Indonesia. (Additivity of metabolizable energy values of maize and ricebran in ducks.)
- 443 *Al-Athari, A-M.K., and W. Guenther, 1984. A new variety of triticale (Carman) in diets for broiler chickens. Poultry Sci. 63(Suppl. 1):52.
- 444 Almeida, J.A., and E.S. Baptista, 1984. A new approach to the quantitative collection of excreta from birds in a true metabolizable energy bioassay. Poultry Sci. 63:2501-2503.
- 445 Baldwin, R.L., and A.C. Bywater, 1984. Nutritional energetics of animals. Ann. Rev. Nutr. 4:101-114.
- 446 Bonino, M.F., F. Borras, S.M. de Barreiro, M.J. Schang and O.A. Scaglio, 1984. Evaluando raciones. Balanceados Argentinos 3(27):17-25.
- 447 el Boushy, A.R., and A.E. Roodbeen, 1984. Amino acid availability in dry poultry waste in comparison with relevant feedstuffs. Poultry Sci. 63:583-585.

- 448 *Caldwell, J.M., D.H. Thorne and J.M. Vandepopuliere, 1984. Nutritional composition of methane digester effluent. Proc. Abstr. XVII World's Poult. Congr. p. 741.
- 449 *Cantor, A.H., D.E. Reese, D.O. Liptrap, A.H. Hussein, D.M. Nash and T.H. Johnson, 1984. Evaluation of metabolizable energy of corn grown during periods of extreme heat and dryness. Poultry Sci. 63(Suppl. 1):74.
- 450 Carré, B., B. Prevotel, and B. Leclercq, 1984. Cell wall content as a predictor of metabolizable energy value of poultry feedingstuffs. Br. Poult. Sci. 25:561-572.
- 451 Chae, B.J. and I.K. Han, 1984. Studies on the determination of amino acid bioavailability for broiler-type chicks. I. Bioavailable amino acid values of corn, sorghum, soybean meal and fish meal as measured by two different bioassays. Korean J. Anim. Sci. 26:372-379.
- 452 Chae, B.J., I.K. Han, and S.C. Lee, 1984. Studies on the determination of amino acid bioavailability for broiler-type chicks. II. The effect of input levels on the amino acid bioavailability of soybean meal. Korean J. Anim. Sci. 26:380-382.
- 453 Chee, K.M., J.H. Jo, M.H. Kang, and I.K. Han, 1984. Evaluation of various feeding methods and factors affecting true metabolizable energy values of diets in chicken. Korean J. Anim. Sci. 26:54-60.
- 454 *Crissey, S.D., and O.P. Thomas, 1984. The amount of lysine excreted from intact or surgically modified roosters when fed soybean meal, autoclaved soybean meal or starch. Poultry Sci. 63(Suppl. 1):83.
- 455 Dale, N., and H.L. Fuller, 1984. Correlation of protein content of feedstuffs with the magnitude of nitrogen correction in true metabolizable energy determination. Poultry Sci. 63:1008-1012.
- 456 *Dale, N.M., and H.L. Fuller, 1984. Effect of nitrogen correction on repeatability of true metabolizable energy values. Poultry Sci. 63(Suppl. 1):85.
- 457 Dale, N.M., and H.L. Fuller, 1984. The metabolizable energy of corn. Are table values too high. Feedstuffs 56(20):23.
- 458 Dale, N.M., H.L. Fuller and H.R. Halloran, 1984. True metabolizable energy and true amino acid availability of an alfalfa protein concentrate (Pro-Xan). Poultry Sci. 63:388-390.
- 459 *Engster, H.M., 1984. Effect of feeding an energy source on endogenous and metabolic amino acid excretion in determining true amino acid availability of hard and soft wheat. Poultry Sci. 63(Suppl. 1):97.
- 460 Fuller, H.L., and N.M. Dale, 1984. The effective energy value of fats in poultry rations. 12th Annu. Symp. Feed Ind., Natl. Renderers Assoc. pp. 1-14.

- 461 Hamilton, R.M.G., and H.L. Trenholm, 1984. Observations on the chemical and nutritive content of white winter and spring wheats contaminated with deoxynivalenol (vomitoxin). *Anim. Feed Sci. Technol.* 11:293-300.
- 462 Han, I.K., M.K. Song, D.S. Nam, K.H. Lee, and C.H. Kwack, 1984. Studies on the TME bioassay of poultry feedstuffs. IV. The effect of the duration of starvation and excreta collection period on true metabolizable energy value of corn. *Korean J. Anim. Sci.* 26:29-33.
- 463 Han, I.K., M.K. Song, K.H. Lee, and C.H. Kwack, 1984. Studies on the TME bioassay of poultry feedstuffs. V. The effect of mixing ratio of basal diet and test material as corn on true metabolizable energy value. *Korean J. Anim. Sci.* 26:34-39.
- 464 Han, I.K., M.K. Song, K.H. Lee, and C.H. Kwack, 1984. Studies on the TME bioassay of poultry feedstuffs. VI. A comparison of apparent and true metabolizable energy measures using some feedstuffs. *Korean J. Anim. Sci.* 26:40-45.
- 465 *Han, Y., and C.M. Parsons, 1984. High-oil corn for poultry. *Poultry Sci.* 63(Suppl. 1):109-110.
- 466 Hayes, J.P., R.E. Austic and M.L. Scott, 1984. True lysine digestibility of normal and heat-damaged fish meal by colostomized and caecectomized roosters. *Proc. Abstr. XVII World's Poult. Congr.* pp. 303-305.
- 467 King, R.D., 1984. A study of the relationship between nitrogen excretion and true metabolizable energy. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 137-149.
- 468 King, R.D., and K. Sim Poi, 1984. Layer nutrition 202 - Nitrogen excretion evaluation. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 56-69.
- 469 King, R.D., and K. Sim Poi, 1984. Layer nutrition 204 - TME and nitrogen correction. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 115-116.
- 470 *Laurin, D.E., S.P. Touchburn, E.R. Chavez and C.W. Chan, 1984. Methods of measuring energy utilization in broilers: effect of genetic line and presence of supplemental fat. *Poultry Sci.* 63(Suppl. 1):135.
- 471 Leclercq, B., B. Prevotel and B. Carre, 1984. Equations de prédiction de la valeur énergétique des aliments composés destinés aux volailles: étude expérimental. L'alimentation des volailles. *Inst. Nat. Rech. Agron., Nouzilly, France.* pp. 1-69.
- 472 Lilburn, M.S., and L.S. Jensen, 1984. Evaluation of corn fermentation solubles as a feed ingredient for laying hens. *Poultry Sci.* 63:542-547.

- 473 McFarland, D.C., and C.N. Coon, 1984. The evaluation of by-product protein digestibility utilizing coomassie blue dye-binding. *Poultry Sci.* 63:2225-2228.
- 474 McNab, J.M., and C. Fisher, 1984. An assay for true and apparent metabolizable energy. *Proc. Abstr. XVII World's Poult. Congr.* pp. 374-375.
- 475 Miller, M.R., and K.J. Reinecke, 1984. Proper expression of metabolizable energy in avian energetics. *Condor* 86:396-400.
- 476 Mohamed, K., B. Leclercq, A. Anwar, H. El-Alaily and H. Soliman, 1984. A comparative study of metabolizable energy in ducklings and domestic chicks. *Anim. Feed Sci. Technol.* 11:199-209.
- 477 Myung, K.H, S.S. Sun and Y.S. Kim, 1984. Studies on the calcium bioavailability by true available mineral methodology. II. The effects of duration of fasting period and excreta collection and level of calcium input on true available calcium value in breeding Japanese quails. *Korean J. Anim. Sci.* 26:464-468.
- 478 Narahari, D., and P. Kothandaraman, 1984. Chemical composition and nutritional value of para-rubber seed and its products for chickens. *Anim. Feed Sci. Technol.* 10:257-267.
- 479 Nordheim, J.P., and C.N. Coon, 1984. A comparison of four methods for determining available lysine in animal protein meals. *Poultry Sci.* 63:1040-1051.
- 480 Ostrowski-Meissner, H.T., 1984. Effect of contamination of foods by Aspergillus flavus on the nutritive value of protein. *J. Sci. Food Agric.* 35:47-58.
- 481 Ostrowski-Meissner, H.T., 1984. A method for simultaneous measurement of apparent (AME) and true (TME) metabolizable energy with ducks and comparisons of data obtained with drakes and cockerels. *Nutr. Rep. Int.* 29:1239-1248.
- 482 *Ostrowski-Meissner, H.T., 1984. Simultaneous determination of apparent (AME) and true (TME) metabolizable energy on drakes. *Proc. Abstr. XVII World's Poult. Congr.* p. 382.
- 483 Papadopoulos, M.C., 1984. Feather meal: evaluation of the effect of processing conditions by chemical and chick assays. Doctoral Thesis, Agric. Univ., Wageningen, The Netherlands. 139 pp.
- 484 Park, J.H., and I.K. Han, 1984. Studies on the measurement of apparent and true metabolizable energy values for poultry feed ingredients. *Korean J. Anim. Sci.* 26:363-374.
- 485 Park, J.H., and I.K. Han, 1984. Effect of cellulose input on the metabolic fecal plus endogenous urinary energy excretion of broiler chicks. *Korean J. Anim. Sci.* 26:555-557.

- 486 Parsons, C.M., 1984. Influence of caeectomy and source of dietary fibre or starch on excretion of endogenous amino acids by laying hens. Brit. J. Nutr. 51:541-548.
- 487 *Parsons, C.M., 1984. Determination of digestible and bioavailable amino acids in meat meal using intact and cecectomized roosters or chick growth assays. Poultry Sci. 63(Suppl. 1):161-162.
- 488 Parsons, C.M., L.M. Potter, and B.A. Bliss, 1984. A modified voluntary feed intake bioassay for determination of metabolizable energy with Leghorn roosters. Poultry Sci. 63:1610-1616.
- 489 Patchell, M.R., 1984. Layer nutrition 187 - Maintenance requirements of Sykes Strain layers. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ. Palmerston North, N.Z. pp. 11-14.
- 490 Patchell, M.R., 1984. Layer nutrition 192 - Isoleucine and lysine levels in layer diets. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ. Palmerston North, N.Z. pp. 30-36.
- 491 Patchell, M.R., 1984. Layer nutrition 194 - Formulation of diets based on available amino acids. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ. Palmerston North, N.Z. pp. 37-41.
- 492 Patchell, M.R., 1984. Layer nutrition 201 - Isoleucine and lysine. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ. Palmerston North, N.Z. pp. 52-55.
- 493 Patchell, M.R., 1984. Chick nutrition 211 - Microbial recovery product. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ. Palmerston North, N.Z. pp. 74-79.
- 494 Patchell, M.R., 1984. Layer nutrition 205 - Minimum levels of isoleucine and lysine. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ. Palmerston North, N.Z. pp. 117-120.
- 495 Patchell, M.R., 1984. TME and amino acid availability values. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ. Palmerston North, N.Z. pp. 150-182.
- 496 Patchell, M.R., and W.C. Smith, 1984. Energy evaluation comparisons - cockerel, pig and rat. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ. Palmerston North, N.Z. pp. 127-128.
- 497 Pesti, G.M., 1984. Influence of substitution method and of feed intake on bioassays to determine metabolizable energy with chickens. Br. Poult. Sci. 25:495-504.
- 498 *Pesti, G.M., 1984. The effects of substitution method and feed intake on bioassays for metabolizable energy with chickens. Poultry Sci. 63(Suppl. 1):29-30.

- 499 Picard, M., D. Bourdon, and J. Le Dividich, 1984. Methods of estimating digestibility and metabolism of crop residues and agro-industrial by-products in monogastric species in developing countries. pp. 134-153 in T.R. Preston, V.V. Kossila, J. Goodwin and S. Reid, eds. Guidelines for Research on the Better Utilization of Crop Residues and Agro-Industrial By Products in Animal Feeding in Developing Countries. Proc. FAO/ILCA Expert Consulting Committee, Addis Ababa, Ethiopia.
- 500 du Preez, J.J., A.duP. Minnaar and J.S. Duckitt, 1984. An alternative approach to a compulsive change from conventional to rapid methods of evaluating metabolizable energy. World's Poult. Sci. J. 40:121-130.
- 501 Raharjo, Y.C., and D.J. Farrell, 1984. Effects of caecectomy and dietary antibiotics on the digestibility of dry matter and amino acids in poultry feeds determined by excreta analysis. Aust. J. Exp. Agric. Anim. Husb. 24:516-521.
- 502 Reid, B.L., S. Galaviz-Moreno and P.M. Maiorino, 1984. A comparison of glandless and regular cottonseed meals for laying hens. Poultry Sci. 63:1803-1809.
- 503 Salmon, R.E., 1984. True metabolizable energy and total and available amino acids of Candle, Altex and Regent Canola meals. Poultry Sci. 63:135-138.
- 504 Salmon, R.E., 1984. True metabolizable energy and dry matter contents of some feedstuffs. Poultry Sci. 63:381-383.
- 505 Salmon, R.E., 1984. True metabolizable energy and amino acid composition of wheat and triticale and their comparative performance in turkey starter diets. Poultry Sci. 63:1664-1666.
- 506 Salmon, R.E., 1984. Effect of grower and finisher protein on performance, carcass grade and meat yield of turkey broilers. Poultry Sci. 63:1980-1986.
- 507 *Salmon, R.E., 1984. Triticale as a feedstuff for poultry. Can. J. Anim. Sci. 64:197.
- 508 *Salmon, R.E., 1984. True metabolizable energy for poultry. Can. J. Anim. Sci. 64:199.
- 509 Salmon, R.E., 1984. Metabolizable energy of poultry feeds. Weekly Letter, Agriculture Canada, Research Station, Swift Current, Sask. 24(Aug. 10).
- 510 Seemann, M., 1984. Vorschlageiner verbesserten Auswertungsmethodik fur die energetische Futterwertschatzung einzelner Komponenten aus einer Ration. Arch. Gefluegelkd. 48:69-76.
- 511 Shebata, M.A., N.K. Allen and R.D. Goodrich, 1984. Evaluation of dried rumen liquor as a feed ingredient for poultry. Arch. Gefluegelkd. 48:89-92.

- 512 *Sibbald, I.R., 1984. Current techniques for evaluation of monogastric feeds. Eighth Annu. Conf., Feeds and Feeding Systems for Livestock, Malays. Soc. Anim. Prod., Genting Highlands, Malaysia. p. K5-56.
- 513 Sibbald, I.R., 1984. A bioassay for true metabolizable energy corrected to zero nitrogen retention (TME_n)/Dosage biologique de l'énergie métabolisable vraie corrigé pour une rétention nulle d'azote (EMV_n). Canadex 450.55, Agriculture Canada, Ottawa, Canada.
- 514 Sibbald, I.R., 1984. Proximate composition and true metabolizable energy values of poultry feedstuffs/Composition quantitative et énergie métabolisable vraie de certains aliments de la volaille. Canadex 450.60, Agriculture Canada, Ottawa, Canada.
- 515 Sibbald, I.R., and P.M. Morse, 1984. A preliminary investigation of the utilization of true metabolizable energy by chicks. Poultry Sci. 63:954-971.
- 516 Sibbald, I.R., and M.S. Wolynetz, 1984. A longitudinal study of energy and nitrogen excretion by fasted cockerels. Poultry Sci. 63:691-702.
- 517 Sibbald, I.R., and M.S. Wolynetz, 1984. The nutrient content of menhaden fish meal. Poultry Sci. 63:1987-1993.
- 518 *Sibbald, I.R., and M.S. Wolynetz, 1984. A preliminary appraisal of nitrogen balance as an estimate of nitrogen retention in assays for bioavailable energy. Poultry Sci. 63(Suppl. 1):182-183.
- 519 Sibbald, I.R., and M.S. Wolynetz, 1984. Energy in poultry nutrition: terminology and definitions. Canadex 450.55, Agriculture Canada, Ottawa, Canada.
- 520 Teeter, R.G., M.O. Smith, E. Murray and H. Hall, 1984. Force feeding methodology and equipment for poultry. Poultry Sci. 63:573-575.
- 521 Terapuntuwat, S., and I. Tasaki, 1984. Protein and amino acid digestibility, biological value of protein and energy metabolizability of some leaf-protein concentrates in chickens. Jpn. Poult. Sci. 21:64-74.
- 522 Terapuntuwat, S., and I. Tasaki, 1984. Protein utilization and amino acid digestibility of some leaf protein concentrates supplemented with amino acids in chickens. Jpn. Poult. Sci. 21:89-93.
- 523 Touchburn, S.P., D.E. Laurin, C.W. Chan and E.R. Chavez, 1984. Energy metabolism and the excess fat problem in broiler chickens. Proc. Abstr. XVII World's Poult. Congr. pp. 258-259.
- 524 Wallis, I.R., and D. Balnave, 1984. A comparison of amino acid digestibility bioassays for broilers. Br. Poult. Sci. 25:389-399.
- 525 Watts, G.D., 1984. True metabolizable energy trial ME/17. Summary of Final Reports and Current Research, Poult. Res. Centre, Massey Univ., Palmerston North, N.Z. pp. 97-100.

- 526 Wolynetz, M.S., and I.R. Sibbald, 1984. Relationships between apparent and true metabolizable energy and the effects of a nitrogen correction. *Poultry Sci.* 63:1386-1399.
- 527 Yeong, S.W., F. Lim and M. Faizah, 1984. A comparison between local and imported fish meals in poultry diets. Eighth Annu. Conf., Feeds and Feeding Systems for Livestock, Malays. Soc. Anim. Prod., Genting Highlands, Malaysia.
- 528 *Yeong, S.W., F. Lim and M. Faizah, 1984. A comparison of different sources of fish meals in poultry. Eighth Annu. Conf., Feeds and Feeding Systems for Livestock, Malays. Soc. Anim. Prod., Genting Highlands, Malaysia, p. S11-10.

1985

- 529 Anonymous, 1985. ¿ Parqué son toxicos los taninos? *Avicultura Profesional* 3(1):14-15.
- 530 Bielorai, R., B. Iosif, and H. Neumark, 1985. Nitrogen absorption and endogenous nitrogen along the intestinal tract of chicks. *J. Nutr.* 115:568-572.
- 531 Cave, N.A., and V.D. Burrows, 1985. Naked oats in feeding the broiler chicken. *Poultry Sci.* 64:771-773.
- 532 *Dale, N.M., and H.L. Fuller, 1985. Energetic content of yellow corn and several alternative energy sources. *Poultry Sci.* 64(Suppl. 1):13.
- 533 *Dale, N., and H.L. Fuller, 1985. Relationship between proximate composition and true metabolizable energy of yellow corn and dehulled soybean meal. *Poultry Sci.* 64(Suppl. 1):84.
- 534 Dale, N.M., H.L. Fuller, G.M. Pesti, and R.D. Phillips, 1985. Freeze drying versus oven drying of excreta in true metabolizable nitrogen-corrected true metabolizable energy and true amino acid availability bioassays. *Poultry Sci.* 64:362-365.
- 535 Engster, H.M., N.A. Cave, H. Likuski, J.M. McNab, C.A. Parsons and F.E. Pfaff, 1985. A collaborative study to evaluate a precision-fed rooster assay for true amino acid availability of feed ingredients. *Poultry Sci.* 64:487-498.
- 536 Halley, J.T., T.S. Nelson, L.K. Kirby, and Z.B. Johnson, 1985. Relationship between dry matter digestion and metabolizable energy. *Poultry Sci.* 64:1934-1937.
- 537 Halloran, H.R., 1985. Disponibilidad de aminoacidos en materias primas. *Avicultura Profesional* 3:62.
- 538 Halloran, H.R., 1985. Amino acid availability of feed ingredients. *Feedstuffs* 57(22):10-11.

- 539 Hamilton, R.M.G., H.L. Trenholm, B.K. Thompson, and R. Greenhalgh, 1985. The tolerance of White Leghorn and broiler chicks, and turkey poult to diets that contained deoxynivalenol (vomitoxin) - contaminated wheat. *Poultry Sci.* 64:273-286.
- 540 Karunajeewa, H., and B.E. Bartlett, 1985. The effects of replacing soyabean meal in broiler starter diets with white lupin seed meal of high manganese content. *Nutr. Rep. Int.* 31:53-58.
- 541 Kim, Y.S., S.S. Sun, and K.H. Myung, 1985. A comparison of true available calcium with apparent available calcium using 6 calcium supplements for breeding Japanese quails. *Korean J. Anim. Sci.* 27:297-300.
- 542 Kim, Y.S., S.S. Sun, and K.H. Myung, 1985. The effects of level and method of calcium input, age and particle size on true available calcium value in breeding Japanese quails. *Korean J. Anim. Sci.* 27:386-390.
- 543 Kussabati, R., and B. Leclercq, 1985. A simplified rapid method for the determination of apparent and true metabolizable energy values of poultry feed. *Arch. Gefluegelkd.* 49:54-62.
- 544 Laurin, D.E., S.P. Touchburn, E.R. Chavez, and C.W. Chan, 1985. Methods of measuring energy utilization in broilers: effect of genetic line and presence of supplemental dietary fat. *Poultry Sci.* 64:969-978.
- 545 Lessire, M., B. Leclercq, L. Conan, and J.M. Hallouis, 1985. A methodological study of the relationship between metabolizable energy values of two meat meals and their level of inclusion in the diet. *Poultry Sci.* 64:1721-1728.
- 546 Papadopoulos, M.C., 1985. Estimations of amino acid digestibility and availability in feedstuffs for poultry. *World's Poult. Sci. J.* 41:64-71.
- 547 Papadopoulos, M.C., A.R. El Boushy, and E.H. Ketelaars, 1985. Effect of different processing conditions on amino acid digestibility of feather meal determined by chicken assay. *Poultry Sci.* 64:1729-1741.
- 548 Parsons, C.M., 1985. Influence of caeectomy on digestibility of amino acids by roosters fed distillers' dried grains with solubles. *J. Agric. Sci. Camb.* 104:469-472.
- 549 *Parsons, C.M., Y. Han, and C.C. Welch, 1985. Evaluation of amino acid availability assays for poultry: methodology as applied to excreta collection and chick growth bioassays. *Poultry Sci.* 64(Suppl. 1):160.
- 550 *Pesti, G.M., L.O. Faust, H.L. Fuller, and N.M. Dale, 1985. The metabolizable energy of poultry by-product meal as measured by three assays. *Poultry Sci.* 64(Suppl. 1):162.

- 551 Picard, M., and B. Darcy-Vrillon, 1985. Composition en acides amines du tourteau de colza et digestibilité chez les monogastriques. Proc. Colloque de l'ONIDOL-Tourteau de Colza - l'Enjeu. ONIDOL, Paris, France. pp. 54-73.
- 552 Raharjo, Y., and D.J. Farrell, 1985. A new biological method for determining amino acid digestibility in poultry feedstuffs using a simple cannula, and the influence of dietary fibre on endogenous amino acid output. Anim. Feed Sci. Technol. 12:29-45.
- 553 Reid, B.L., 1985. Energetic value of fat for layers evaluated. Feedstuffs 57(9):32-37.
- 554 Sibbald, I.R., 1985. Theory and practical use of TME values in poultry rations. Proc. Feeding Systems of Animals in Temperate Areas. Korea Feed Inf. Cent., Seoul, Korea. pp. 52-57.
- 555 *Sibbald, I.R., 1985. The TME system of feed evaluation. Proc. 33rd Annu. Meet. Am. Assoc. Feed Microsc., Ottawa, Canada. pp. 58-59.
- 556 Sibbald, I.R., and M.S. Wolynetz, 1985. Relationships between estimates of bioavailable energy made with adult cockerels and chicks: effects of feed intake and nitrogen retention. Poultry Sci. 64:127-138.
- 557 Sibbald, I.R., and M.S. Wolynetz, 1985. Estimates of retained nitrogen used to correct estimates of bioavailable energy. Poultry Sci. 64:1506-1513.
- 558 Sibbald, I.R., and M.S. Wolynetz, 1985. The bioavailability of supplementary lysine and its effect on the energy and nitrogen excretion of adult cockerels fed diets diluted with cellulose. Poultry Sci. 64:1972-1975.
- 559 Sibbald, I.R., and M.S. Wolynetz, 1985. The excreta energy and nitrogen losses of adult cockerels during a fast. Poultry Sci. 64:1976-1980
- 560 Soares, J.H., Jr., 1985. Poultry nutrition research 1984. 33rd Pfizer Annu. Res. Conf., Minneapolis, U.S.A. pp. 87-115.
- 561 *Zumbado, M.E., and M.G. Murillo, 1985. Effect of processing on nutritive value of pejibaye meal (*Bactris gasipaes*) for poultry. Poultry Sci. 64:(Suppl. 1):204.

LIBRARY / BIBLIOTHEQUE



AGRICULTURE CANADA OTTAWA K1A 0C5

з 9073 00040139 0

DUE DATE

