

# Apparent Opioid and Stimulant Toxicity Deaths

---

Surveillance of Opioid- and  
Stimulant-Related Harms in Canada



January 2016 to June 2021



Public Health  
Agency of Canada

Agence de la santé  
publique du Canada

Canada

**TO PROMOTE AND PROTECT THE HEALTH OF CANADIANS THROUGH LEADERSHIP,  
PARTNERSHIP, INNOVATION AND ACTION IN PUBLIC HEALTH.**

—Public Health Agency of Canada

Prepared by the Substance-related Overdose and Mortality Surveillance Task Group (SOMS-TG)  
of the Special Advisory Committee on the Epidemic of Opioid Overdoses (SAC)

Également disponible en français sous le titre :

Décès apparemment liés à une intoxication aux opioïdes et aux stimulants

*Surveillance des méfaits associés aux opioïdes et aux stimulants au Canada*

*Janvier 2016 à juin 2021*

To obtain additional information, please contact:

Public Health Agency of Canada

Address Locator 0900C2

Ottawa, ON K1A 0K9

Tel.: 613-957-2991

Toll free: 1-866-225-0709

Fax: 613-941-5366

TTY: 1-800-465-7735

E-mail: [publications-publications@hc-sc.gc.ca](mailto:publications-publications@hc-sc.gc.ca)

© Her Majesty the Queen in Right of Canada, as represented by the Minister of Health, 2021

Publication date: December 2021

This publication may be reproduced for personal or internal use only without permission provided  
the source is fully acknowledged.

Cat.: HP33-3/2021E-2-PDF

ISBN: 978-0-660-40668-8

Pub.: 210384



# Table of Contents

---

<b>Acknowledgements</b>	<b>4</b>
<b>Suggested Citation</b>	<b>4</b>
<b>Key Findings</b>	<b>5</b>
<b>Technical Notes</b>	<b>8</b>
Definitions	8
How apparent opioid and stimulant toxicity deaths are counted	10
Limitations of the data on apparent opioid and/or stimulant toxicity deaths	11
Notes on provincial and territorial data	12
General notes	12
Manner of death	13
Sex and age group	13
Fentanyl, fentanyl analogues, and non-fentanyl opioids	13
Origin of opioid(s)	14
Cocaine, methamphetamine and other stimulants	15
Other psychoactive substances	15
Data suppression	16
<b>Appendix</b>	<b>19</b>
Table 1a. Number and rate (per 100,000 population) of total apparent opioid toxicity deaths by province or territory, 2016 to 2021 (Jan to Jun)	20
Table 1b. Number and rate (per 100,000 population) of total apparent stimulant toxicity deaths by province or territory, 2018 to 2021 (Jan to Jun)	22
Table 2a. Number and rate (per 100,000 population) of accidental apparent opioid toxicity deaths by province or territory, 2016 to 2021 (Jan to Jun)	23
Table 2b. Number and rate (per 100,000 population) of accidental apparent stimulant toxicity deaths by province or territory, 2018 to 2021 (Jan to Jun)	25
Table 3a. Summary of apparent opioid toxicity deaths by manner of death, 2016 to 2021 (Jan to Jun)	26
Table 3b. Summary of apparent stimulant toxicity deaths by manner of death, 2018 to 2021 (Jan to Jun)	26



Table 4a. Summary of total apparent opioid toxicity deaths in Canada, 2016 to 2021 (Jan to Jun)	27
Table 4b. Summary of total apparent stimulant toxicity deaths in Canada (based on available information from five to seven provinces or territories), 2018 to 2021 (Jan to Jun)	28
Table 5a. Number of total apparent opioid toxicity deaths by quarter and province or territory, January 2016 to June 2021	29
Table 5b. Number of total apparent stimulant toxicity deaths by quarter and province or territory, January 2018 to June 2021	31
Table 6a. Number of accidental apparent opioid toxicity deaths by quarter and province or territory, January 2016 to June 2021	32
Table 6b. Number of accidental apparent stimulant toxicity deaths by quarter and province or territory, January 2018 to June 2021	34
Table 7a. Percentage of accidental apparent opioid toxicity deaths by sex and province or territory, 2016 to 2021 (Jan to Jun)	35
Table 7b. Percentage of accidental apparent stimulant toxicity deaths by sex and province or territory, 2018 to 2021 (Jan to Jun)	36
Table 8a. Crude rate (per 100,000) of accidental apparent opioid toxicity deaths by sex and province or territory, 2016 to 2021 (Jan to Jun)	37
Table 8b. Crude rate (per 100,000 population) of accidental apparent stimulant toxicity deaths by sex and province or territory, 2018 to 2021 (Jan to Jun)	38
Table 9a. Percentage of accidental apparent opioid toxicity deaths by age group and province or territory, 2016 to 2021 (Jan to Jun)	39
Table 9b. Percentage of accidental apparent stimulant toxicity deaths by age group and province or territory, 2016 to 2021 (Jan to Jun)	42
Table 10a. Percentage of accidental apparent opioid toxicity deaths by type of opioids and province or territory, 2016 to 2021 (Jan to Jun)	44
Table 10b. Percentage of accidental apparent stimulant toxicity deaths by type of stimulants and province or territory, 2018 to 2021 (Jan to Jun)	46
Table 11a. Percentage of accidental apparent opioid toxicity deaths involving stimulants, by province or territory, 2018 to 2021 (Jan to Jun)	47
Table 11b. Percentage of accidental apparent stimulant toxicity deaths involving opioids, by province or territory, 2018 to 2021 (Jan to Jun)	48
Table 12a. Percentage of accidental apparent opioid toxicity death involving other psychoactive substances, by province or territory, 2016 to 2021 (Jan to Jun)	49



Table 12b. Percentage of accidental apparent stimulant toxicity death involving other psychoactive substances, by province or territory, 2018 to 2021 (Jan to Jun)	50
Table 13. Percentage of accidental opioid toxicity deaths by origin of opioid(s), by province or territory, 2018 to 2021 (Jan to Jun)	51
Table 14a. Number of accidental apparent opioid toxicity deaths by sex and age group, by province or territory, 2018 to 2021 (Jan to Jun)	54
Table 14b. Number of accidental apparent stimulant toxicity deaths by sex and age group, by province or territory, 2018 to 2021 (Jan to Jun)	58
Table 15a. Percentage of accidental apparent opioid toxicity deaths by type of opioids and sex, by province or territory, 2018 to 2021 (Jan to Jun)	61
Table 15b. Percentage of accidental apparent stimulant toxicity deaths by type of stimulants and sex, by province or territory, 2018 to 2021 (Jan to Jun)	63
Table 16a. Percentage of accidental apparent opioid toxicity deaths by type of opioids and age group, by province or territory, 2018 to 2021 (Jan to Jun)	65
Table 16b. Percentage of accidental apparent stimulant toxicity deaths by type of stimulants and age group, by province or territory, 2018 to 2021 (Jan to Jun)	73
Table 17. Percentage of accidental opioid toxicity deaths by type of opioids and origin of opioid(s), by province or territory, 2018 to 2021 (Jan to Jun)	77
Table 18. Percentage of accidental opioid toxicity deaths by sex and origin of opioid(s), by province or territory, 2018 to 2021 (Jan to Jun)	82



## Acknowledgements

---

This update would not be possible without the collaboration and dedication of provincial and territorial (PT) offices of Chief Coroners and Chief Medical Examiners as well as PT public health and health partners and Emergency Medical Services data providers. We would also like to acknowledge the Canadian Institute for Health Information (CIHI) for collecting and providing the data used for reporting opioid and stimulant related poisoning hospitalizations.

## Suggested Citation

---

Special Advisory Committee on the Epidemic of Opioid Overdoses. Opioid and Stimulant-related Harms in Canada. Ottawa: Public Health Agency of Canada; December 2021. <https://health-infobase.canada.ca/substance-related-harms/opioids-stimulants/>



# Key Findings

---

## Context

The COVID-19 outbreak is worsening the already deadly and ongoing public health crisis of opioid overdoses and deaths. It is having a tragic impact on people who use substances, their families, and communities across Canada. People who use substances, such as opioids, cocaine, and methamphetamine, are experiencing a number of increased risks, with several jurisdictions reporting higher rates of fatal overdoses and other harms.

These updates include available data on overdoses and deaths involving opioids and/or stimulants from January 2016 to June 2021, where available. Recognizing that harms related to opioids, stimulants, and other substances extend beyond overdoses (poisonings) and deaths, we continue to work with federal, provincial and territorial partners to build a broad understanding of harms and substances involved to better respond to the crisis. [Additional studies](#) can also help us plan and tailor actions to achieve better possible outcomes.

## **24,626 apparent opioid toxicity deaths between January 2016 and June 2021<sup>1,2</sup>**

- 1,720 apparent opioid toxicity deaths occurred between April and June 2021 (approximately 19 deaths per day), similar to the period from January to March 2021 (1,792 deaths), but representing a 2% increase compared to April to June 2020 (1,680 deaths) and a 66% increase compared to April to June 2019 (1,038 deaths).
- A number of factors have likely contributed to a worsening of the overdose crisis over the course of the pandemic, including the increasingly toxic drug supply, increased feelings of isolation, stress and anxiety and limited availability or accessibility of services for people who use drugs.

## **Majority of deaths in British Columbia, Alberta and Ontario; increases observed in other regions**

- While rates continue to remain high in Western Canada, increases have been observed in other regions such as Ontario and Yukon. Several jurisdictions have observed record-breaking numbers and rates in relation to the wider impacts of the COVID-19 pandemic.
- Between January and June 2021, 90% of all apparent opioid toxicity deaths occurred in British Columbia, Alberta or Ontario.

---

<sup>1</sup> Manitoba data from April 2020 to June 2021 and Newfoundland data from April to June 2021 were not available at the time of this update.

<sup>2</sup> National overall count from January 2016 to June 2021 includes deaths from British Columbia (2018 to 2021) and Quebec (2020 to 2021) related to all illicit drugs including, but not limited to opioids.



### **Most apparent opioid toxicity deaths among males and individuals aged 20 to 49 years**

- Males accounted for the majority of accidental apparent opioid toxicity deaths (75%) from January to June 2021; for both males and females, the majority of deaths were among individuals aged 20 to 49 years.
- Individuals between 30 and 39 years accounted for a higher proportion of accidental opioid toxicity deaths where fentanyl (31%) was involved from January to June 2021.

### **Fentanyl and fentanyl analogues continue to be major drivers of the crisis**

- 87% of accidental apparent opioid toxicity deaths involved fentanyl in 2021 (Jan to Jun).
- 90% of accidental apparent opioid toxicity deaths from January to June 2021 involved a non-pharmaceutical opioid.<sup>3,4</sup>

---

<sup>3</sup> Based on 2021 (Jan to Jun) data on origin of opioids from deaths with completed investigations from seven provinces.

<sup>4</sup> Opioids with a pharmaceutical origin refer to opioids that were manufactured by a pharmaceutical company and approved for medical purposes in humans. Pharmaceutical origin does not indicate how the opioids were obtained (e.g. through personal prescription or by other means).





## Data on apparent opioid and stimulant toxicity deaths based on six reporting provinces and territories

- Available information from six provinces and territories indicates the number of deaths involving stimulants from April to June 2021 remained high and is similar to the period from January to March 2021. Almost all (96%) of those deaths were accidental.
- More than half (59%) of accidental opioid toxicity deaths in 2021 (Jan to Jun) also involved a stimulant, reflecting the polysubstance nature of this crisis.
- 65% of identified accidental apparent stimulant toxicity deaths from January to June 2021 involved cocaine while 50% involved methamphetamines.
- 86% of accidental apparent stimulant toxicity deaths also involved an opioid in 2021 (Jan to Jun).
- Males accounted for the majority of accidental apparent stimulant toxicity deaths (74%) from January to June 2021; for males and females, the majority of deaths were among individuals aged 20 to 49 years.

Data on apparent opioid toxicity deaths and stimulant toxicity deaths are not mutually exclusive. A high proportion of deaths involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.



This update is based on data submitted to the Public Health Agency of Canada on or before November 12, 2021. Due to differences in identifying and reporting cases, comparisons over time and between provinces and territories should be interpreted with caution. Refer to the [Technical notes](#) for more information.



# Technical notes

---

## Definitions

**Apparent opioid toxicity death (AOTD):** A death caused by intoxication/toxicity (poisoning) resulting from substance use, where one or more of the substances is an opioid, regardless of how it was obtained (e.g. illegally or through personal prescription). Other substances may also be involved.

---

**Apparent stimulant toxicity death (ASTD):** A death caused by intoxication/toxicity (poisoning) resulting from substance use, where one or more of the substances is a stimulant, regardless of how it was obtained (e.g. illegally or through personal prescription). Other substances may also be involved.

---

**Ongoing investigation:** Coroners and medical examiners continue to collect information on how and why the death occurred. Data for ongoing investigations are considered preliminary and subject to change.

---

**Completed investigation:** Coroners and medical examiners have collected all available information on how and why a death occurred. The time required to complete an investigation and related administrative processes is case-dependent and can range from approximately three to twenty-four months.

---

**Manner of death – Accident:** Deaths with completed investigations where the coroner or medical examiner determined that the death was unintentional. This category also includes deaths with ongoing investigations where the manner of death was believed to be unintentional or had not been assigned at the time of reporting.

---

**Manner of death – Suicide:** Deaths with completed investigations where the coroner or medical examiner determined that the opioids were consumed with the intent to die. This category also includes deaths with ongoing investigations where suicide was believed to be the manner of death at the time of reporting.

---

**Manner of death – Undetermined:** Deaths with completed investigations where a specific manner of death (e.g. accident, suicide) could not be assigned based on available or competing information. For this manner of death category, provinces and territories report only completed investigations with the exception of British Columbia which also includes data from ongoing investigations.

---

**Opioid origin – Pharmaceutical:** Deaths with completed investigations where all opioids that directly contributed to death were manufactured by a pharmaceutical company and approved for medical purposes in humans. Pharmaceutical origin does not indicate how the opioids were obtained (e.g. through personal prescription or by other means).

---



**Opioid origin – Non-pharmaceutical:** Deaths with completed investigations where all opioids that directly contributed to the death were not manufactured by a pharmaceutical company or not approved for medical purposes in humans.

---

**Opioid origin – Both pharmaceutical and non-pharmaceutical:** Deaths with completed investigations where the opioids that directly contributed to the death were a combination of pharmaceutical and non-pharmaceutical opioids, without any opioids of undetermined origin.

---

**Opioid origin – Undetermined:** Deaths with completed investigations where, for one or more opioids that directly contributed to the death, it was not possible to determine whether the opioid was pharmaceutical or non-pharmaceutical.

---



## How apparent opioid and stimulant toxicity deaths are counted

Counts or record-level information are provided by the provinces and territories that collect data from their respective offices of Chief Coroners or Chief Medical Examiners. Crude and age-adjusted rates are calculated using the most current population data from Statistics Canada. Age-adjusted rates used the 2016 Canadian population as a reference and direct standardization was applied.

**Crude rates:** summarize the situation within a region at a certain time period and have not been adjusted for existing differences by provincial and territorial age distributions (e.g. (number of deaths / population) x 100,000).

**Age-adjusted rates:** have been adjusted for existing differences by provincial and territorial age distributions using the 2016 Canadian population as a reference. These rates assume that all regions have the same age distributions and are useful when comparing between regions and over time.

The data provided by the provinces and territories can include deaths:

- with completed or ongoing investigations
- where manner of death is classified as accident, suicide, or undetermined

These data **do not** include deaths due to:

- the medical consequences of long-term substance use or overuse (for example, alcoholic cirrhosis)
- medical assistance in dying
- trauma where use of the substance(s) contributed to the circumstances of the injury that lead to the death, but was not directly involved in the death
- homicide

However, some provincial and territorial differences remain in the type of data reported and in the time periods for which data are available (refer to [Table A](#)).



## Limitations of the data on apparent opioid and/or stimulant toxicity deaths

Data presented in this update should be interpreted with caution.

- This update is based on data submitted to the Public Health Agency of Canada on or before November 12, 2021. New or revised data reported after this date will be reflected in future updates.
- Data released by provinces and territories may differ due to the availability of updated data, differences in the type of data reported (e.g. manners of death), the use of alternate age groupings, differences in time periods presented and/or population estimates used for calculations, etc.
- As some data are based on ongoing investigations by coroners and medical examiners, they are considered preliminary and subject to change. The time required to complete an investigation and related administrative processes is case-dependent and can range from approximately three to twenty-four months.
- This update is based on data that do not specify how the opioids or stimulant were obtained (e.g. illegally or through personal prescription); the level of toxicity may differ depending on the opioid or stimulant (substance(s) involved, concentration, and dosage).
- Provincial and territorial differences in the death investigation process, death classification method, toxicology testing, and the manners of death reported may impact the interpretation and comparability of the data presented in this update over time and between provinces and territories.
- Data on apparent opioid toxicity deaths and stimulant toxicity deaths are not mutually exclusive. A high proportion of deaths involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.
- National rates of apparent opioid toxicity deaths for 2021 exclude Manitoba.
- Quarterly totals for Canada may not equal the annual totals due to suppressed data for some provinces and territories with low number of deaths.
- Data on apparent stimulant toxicity deaths were only available from five to seven provinces and territories depending on the year. Therefore, national numbers and rates are not provided in this update. Refer to [Table A](#) for more details.



## Notes on provincial and territorial data

Due to differences in identifying and reporting cases, comparisons over time and between provinces and territories should be interpreted with caution.

### General notes

1. Data reported by some provinces and territories do not include all manners of death (accident, suicide, undetermined) or stages of investigation (ongoing, completed); refer to [Table A](#) for more details.
2. Data presented will be updated quarterly, based on results of completed investigations.
3. Rates for provinces and territories with relatively smaller populations may change substantially with even slight changes in the number of deaths.
4. British Columbia data for 2016 and 2017 include deaths with completed investigations only. Overall numbers for British Columbia from 2018 to 2021 include deaths with ongoing investigations related to all illicit drugs, including but not limited to opioids and stimulants, used alone or in combination with prescribed/diverted medication. However, stratified data (e.g. by sex, age group, etc.) are based only on opioid toxicity deaths for which investigations are completed.
5. Quebec data from 2016 to 2019 include deaths with completed investigations only; death investigations were underway for 5% in 2019. Available 2020 and 2021 data from Quebec include unintentional deaths with ongoing investigations related to all illicit drugs including, but not limited to, opioids and stimulants. Preliminary data for drug-related poisonings, for which toxicology information was available, indicate that 50% of deaths between January 2020 and June 2021 involved an opioid.
6. Data from Yukon include deaths with completed investigations only. In 2018, one apparent opioid toxicity death occurred in a different province following an overdose in Yukon. This death is included in the data from the jurisdiction where the death occurred and is not reported in the data from Yukon.
7. Data from Prince Edward Island include accidental deaths with completed investigations only. Only annual totals were available for 2016 data from Prince Edward Island; quarterly data for 2016 were not available at the time of this update.
8. In Ontario, apparent opioid toxicity death data were captured using an enhanced data collection tool by the Office of the Chief Coroner as of May 1, 2017. Prior to this time period, retrospective case information was collected using a different tool.
9. For Newfoundland and Labrador, data on apparent opioid toxicity deaths between January 2016 and December 2019 were based on the detection of opioids as indicated on the toxicological report. As of 2020, data include deaths where opioids directly contributed to the death.
10. Data from Nunavut were not included in national counts or percentages.
11. Manitoba data from April 2020 to June 2021 were not available at the time of this update.
12. Newfoundland and Labrador data from April to June 2021 were not available at the time of this update.



## Manner of death

13. Manner of death is assigned by the coroner or medical examiner during, or following an investigation. The data in this update include accidental, suicide and undetermined deaths.
14. Suicide data were unavailable from Alberta (2018 to 2021), Prince Edward Island and Nunavut.

## Sex and age group

15. For most provinces and territories, data on the sex of the individual was based on biological characteristics or legal documentation.
16. Data on deaths where sex was categorized as “Other” were excluded from analyses by sex, but were included in overall analyses.
17. Due to rounding, percentages may not add to 100%.
18. For Ontario, from January 2016 to April 2017, data on the sex of the individual reflected the sex assigned at birth or was based on biological characteristics at the time of death; as of May 2017, the perceived or projected identity of the individual was reported.
19. Alberta uses data on the sex of the individual based on the medical examiner’s assessment, which is largely based on biological characteristics. In a small subset of cases where the individual was known to identify with a gender different than their biological sex, the medical examiner may indicate their identified gender.
20. Data on deaths where age group was categorized as “Unknown” were excluded from analyses by age group, but were included in overall analyses.

## Fentanyl, fentanyl analogues, and non-fentanyl opioids

21. Refer to [Table B](#) below for details on opioids.
22. Prior to 2018, the percentage of deaths involving fentanyl and/or fentanyl analogues represented a single category. For data reported for 2018 to 2021, some provinces and territories did not report fentanyl analogue information or required additional information to differentiate fentanyl from fentanyl analogues until investigations are completed. Therefore, deaths involving fentanyl analogues may be included in the fentanyl percentages for some jurisdictions.
23. The sum of percentages by type of opioids may not add up to 100% because a death may involve more than one type of opioid.
24. Observed trends of accidental apparent opioid toxicity deaths involving fentanyl or fentanyl analogues should be interpreted with caution until additional data become available. In addition, changes to testing practices during the reporting period may affect observed trends.
25. Given provincial and territorial differences in death classification methods, the term “involving” includes deaths where the substance was either detected and/or directly contributed to the death. Substances can be detected through toxicology testing and may or may not have directly contributed to the death. Direct contribution to the death is based on investigation by coroner or medical examiner.



26. Available 2020 and 2021 data from Quebec on deaths related to drugs where toxicology information was available and fentanyl (or fentanyl analogues) was detected were used to approximate apparent opioid toxicity deaths involving fentanyl (or fentanyl analogues), among deaths where opioids were detected.
27. For Alberta and Ontario, only data on deaths with completed investigations, where relevant toxicology information was available, were included in percentages for fentanyl or fentanyl analogues.

### Origin of opioid(s)

28. The origin of opioid(s) refers to whether the opioids that directly contributed to the death were pharmaceutical, non-pharmaceutical, both or undetermined.
29. Data on origin were only available for deaths with completed investigations from 2018 onward from between seven and eight provinces and territories, depending on the year. Completed investigations represented 77% of apparent opioid toxicity death investigations from these provinces and territories over that period; refer to [Table A](#) for more details.
30. Summary data and trends based on origin of opioid(s) should be interpreted with caution until additional data become available.
31. Origin categorization is based on toxicology results and scene evidence and does not indicate how the substances were prepared, their appearance, or how they were “advertised”; nor should it be used to infer the timing or mode of consumption.
32. Pharmaceutical opioids also include those approved for use in humans in other countries, but not necessarily in Canada.
33. For the purposes of origin categorization, deaths involving fentanyl are categorized as “suspected non-pharmaceutical” when there is: 1) no evidence of a patch, vial, or other pharmaceutical formulation at the scene, or 2) no/unknown evidence of a prescription. These deaths are grouped with deaths involving non-pharmaceutical opioids.
34. Origin categorization represents the best estimate based on the information available and should be interpreted with caution.
35. Origin refers only to the opioid(s) involved in death and should not be used as an indication of prior use of opioids of the same or other origin.
36. British Columbia only reports opioid toxicity deaths involving any illicit opioids, resulting in a high proportion of non-pharmaceutical opioids. For that reason, data on origin of opioid from British Columbia were not included in the national proportions.





## Cocaine, methamphetamine and other stimulants

37. Refer to [Table B](#) below for details on stimulants.
38. Amphetamine is a known metabolite of methamphetamine but can also be consumed separately and directly contribute to a toxicity death. Deaths where amphetamine (without methamphetamine) directly contributed to the death are reported under “other stimulants”. In situations where both methamphetamine and amphetamine were consumed separately, and both directly contributed to death, the death is reported under both methamphetamine and “other stimulants”.
39. Data on apparent stimulant toxicity deaths were available from between five and seven provinces and territories, depending on the year from 2018 to 2021 (Jan to Jun).
40. The sum of percentages by type of stimulant may not add up to 100% because a death may involve more than one type of stimulant.
41. For Ontario, only data on deaths with completed investigations, where relevant toxicology information was available, were included in percentages for cocaine, methamphetamine and other stimulants.
42. Data from Quebec on “other stimulants” include deaths involving methamphetamine.
43. For Alberta, only apparent opioid toxicity deaths with completed investigations are used in the numerator for percentage of deaths involving stimulants. As a result, these values may change when more investigations are completed.

## Other psychoactive substances

44. Refer to [Table B](#) below for details on other psychoactive substances.
45. National-level percentages of accidental apparent opioid toxicity deaths involving other non-opioid substances do not include data from Nunavut as these data were not available.
46. For Alberta, only data on deaths with completed investigations and specific substances causing death listed on the death certificate were included in percentages of accidental apparent opioid toxicity deaths involving other non-opioid substances.
47. For Ontario, only data on deaths with completed investigations, where relevant toxicology information was available, were included in percentages of accidental apparent opioid toxicity deaths involving other non-opioid substances. Data for non-opioid substances from Ontario between January 2016 and April 2017 were based on their detection and do not include alcohol; as of May 1, 2017, data on non-opioid substances are based on their direct effects and include alcohol.



## Data suppression

The suppression of data in this update is based on the preferences of individual provinces or territories to address concerns around releasing small numbers for their jurisdiction.

- Quebec suppressed counts less than five for deaths with ongoing investigations (2020 to 2021).
- Nova Scotia suppressed all counts for age group 0 to 19 years when stratified by sex.
- Prince Edward Island suppressed counts between one and four for quarterly data, and for any data related to sex or age distribution.
- Newfoundland and Labrador suppressed counts between one and four for quarterly data, and data related to substances involved and sex or age distribution.
- Yukon suppressed counts between one and four for data related to sex or age distribution.
- Nunavut suppressed all counts less than five.

In addition, suppression was applied in some instances where all data for a province or territory fell in a single category of a given table or figure.



**Table A. Reporting periods, manners of death, and availability of opioid and stimulant data included in this update by province or territory**

		BC	AB	SK	MB	ON	QC	NB	NS	PE	NL	YT	NT	NU
<b>Available data on apparent toxicity deaths involving opioids as of November 12, 2021</b>														
2016-18	January to December	✓ (C)	✓	✓ (C)	✓	✓	✓ (C)	✓	✓	✓ (C)	✓ (C)	✓ (C)	✓	✓
2019	January to December	✓	✓	✓ (C)	✓	✓	✓ (C)	✓	✓	✓ (C)	✓ (C)	✓ (C)	✓	✓
2020	January to December	✓	✓	✓ (C)	✓ (INC)	✓	✓	✓	✓	✓ (C)	✓ (C)	✓ (C)	✓	✓
2021	January to June	✓	✓	✓ (C)	n/a	✓	✓	✓	✓	✓ (C)	✓ (INC) (C)	✓ (C)	✓	✓
<b>Available data on apparent toxicity deaths involving stimulants as of November 12, 2021</b>														
2018	January to December	✓ (C)	n/a	✓ (C)	n/a	✓	✓ (C)	n/a	✓	n/a	n/a	n/a	n/a	n/a
2019	January to December	✓ (C)	n/a	✓ (C)	✓	✓	✓ (C)	n/a	✓	n/a	n/a	n/a	n/a	n/a
2020	January to December	✓ (C)	n/a	✓ (C)	✓ (INC)	✓	n/a	n/a	✓	n/a	✓ (C)	n/a	✓	n/a
2021	January to June	✓ (C)	n/a	✓ (C)	n/a	✓	n/a	n/a	✓	n/a	✓ (INC) (C)	n/a	✓	n/a
<b>Classification of deaths included in the reported data</b>														
Accident	Completed investigations	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	Ongoing investigations where manner of death was believed to be unintentional	✓	-	n/a	✓	-	✓	✓	✓	n/a	n/a	n/a	✓	n/a
	Ongoing investigations where manner of death had not been assigned at the time of reporting	-	✓	n/a	✓	✓	n/a	✓	✓	n/a	-	-	✓	n/a
Suicide	Completed investigations	✓	✓ (INC)	✓	✓	✓	✓	✓	✓	n/a	✓	✓	✓	✓
	Ongoing investigations where the manner of death was believed to be suicide	✓	n/a	n/a	✓	-	n/a	✓	✓	n/a	n/a	n/a	✓	n/a
Deaths with completed investigations and an undetermined manner of death		✓	✓	✓	✓	✓	✓	✓	✓	n/a	✓	✓	✓	✓
<b>Available data on origin of the opioid(s)</b>														
2018-20	January to December	✓	✓	✓	✓ (INC)	✓	n/a	✓	✓	n/a	✓	n/a	n/a	n/a
2021	January to June	✓	✓	✓	n/a	✓	n/a	✓	✓	✓	✓ (INC)	n/a	n/a	n/a

- ✓ These data have been reported by the province or territory and are reflected in this update, unless otherwise specified
- (C) Data includes deaths with completed investigations only
- (INC) Data was not available for the entire period
- The classification is not used in the province or territory
- n/a Data were not available at the time of this publication



**Table B. Types of opioids and stimulants**

Category	Includes (but are not limited to):	
Fentanyl and fentanyl analogues	<ul style="list-style-type: none"><li>• 3-methylfentanyl</li><li>• acetylfentanyl</li><li>• acrylfentanyl</li><li>• butyrylfentanyl</li><li>• carfentanil</li><li>• crotonyl fentanyl</li><li>• cyclopropyl fentanyl</li></ul>	<ul style="list-style-type: none"><li>• despropionyl-fentanyl</li><li>• fentanyl</li><li>• fluoroisobutyrylfentanyl (FIBF)</li><li>• furanylfentanyl</li><li>• methoxyacetylfentanyl</li><li>• norfentanyl</li></ul>
Non-fentanyl opioids	<ul style="list-style-type: none"><li>• 2-methyl AP-237</li><li>• AH-7921</li><li>• AP-237</li><li>• broporphine</li><li>• buprenorphine metabolites</li><li>• codeine</li><li>• desomorphine</li><li>• dihydrocodeine</li><li>• etodesnitazene</li><li>• heroin</li><li>• hydrocodone</li><li>• hydromorphone</li><li>• isopropyl-U-47700</li><li>• isotonitazene</li><li>• loperamide</li></ul>	<ul style="list-style-type: none"><li>• meperidine</li><li>• methadone</li><li>• metonitazene</li><li>• mitragynine</li><li>• monoacetylmorphine</li><li>• morphine</li><li>• MT-45</li><li>• normeperidine</li><li>• oxycodone</li><li>• tapentadol</li><li>• tramadol</li><li>• U-47700</li><li>• U-49900</li><li>• U-50488</li></ul>
Stimulants	<ul style="list-style-type: none"><li>• amphetamine</li><li>• atomoxetine</li><li>• catha</li><li>• cocaine</li><li>• dexamphetamine</li><li>• ethylphenidate</li><li>• lisdexamphetamine</li><li>• MDA</li><li>• MDMA</li></ul>	<ul style="list-style-type: none"><li>• mephedrone</li><li>• methamphetamine</li><li>• methylphenidate</li><li>• modafinil</li><li>• pemoline</li><li>• phentermine</li><li>• pseudoephedrine</li><li>• TFMPP</li></ul>
Other psychoactive substances	<ul style="list-style-type: none"><li>• Alcohol</li><li>• Benzodiazepines</li><li>• Gabapentinoids</li><li>• Ketamine</li><li>• LSD</li></ul>	<ul style="list-style-type: none"><li>• PCP</li><li>• Psilocin</li><li>• W-18</li><li>• Z-drugs</li></ul>



# Appendix

---

**Table 1a. Number and rate (per 100,000 population) of total apparent opioid toxicity deaths by province or territory, 2016 to 2021 (Jan to Jun)**

Province or territory	2016			2017			2018		
	Number	Crude rate per 100,000 population	Age-adjusted rate per 100,000 population	Number	Crude rate per 100,000 population	Age-adjusted rate per 100,000 population	Number	Crude rate per 100,000 population	Age-adjusted rate per 100,000 population
BC	806	16.6	16.5	1,289	26.1	26.0	1,570	31.3	31.2
AB	602	14.3	13.8	744	17.5	17.0	805	18.7	18.3
SK	83	7.3	7.5	85	7.4	7.6	126	10.8	11.4
MB	88	6.7	6.8	106	7.9	8.1	93	6.9	7.0
ON	867	6.2	6.3	1,265	9.0	9.0	1,479	10.3	10.4
QC	258	3.1	3.1	281	3.4	3.4	209	2.5	2.5
NB	34	4.5	4.4	38	5.1	5.1	30	3.9	3.9
NS	53	5.6	5.7	64	6.7	6.9	54	5.6	5.8
PE	5	3.4	3.2	5	3.3	3.6	8	5.2	5.4
NL	18	3.4	3.6	33	6.2	6.5	12	2.3	2.3
YT	7	18.2	17.5	7	17.7	16.6	5	12.3	12.4
NT	5	11.2	12.2	1	2.2	1.9	2	4.4	3.8
NU	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>2,826</b>	<b>7.8</b>	<b>7.8</b>	<b>3,918</b>	<b>10.7</b>	<b>10.8</b>	<b>4,393</b>	<b>11.9</b>	<b>11.9</b>



Province or territory	2019			2020			2021 (Jan to Jun)		
	Number	Crude rate per 100,000 population	Age-adjusted rate per 100,000 population	Number	Crude rate per 100,000 population	Age-adjusted rate per 100,000 population	Number	Crude rate per 100,000 population	Age-adjusted rate per 100,000 population
BC	1,012	19.9	19.8	1,752	34.0	33.9	1,028	39.4	39.4
AB	624	14.3	14.0	1,160	26.2	26.1	708	31.9	31.5
SK	117	10.0	10.6	259	22.0	23.0	96	16.3	17.4
MB	62	4.5	4.6	22	1.6	1.6	n/a	n/a	n/a
ON	1,530	10.5	10.6	2,430	16.5	16.6	1,414	19.1	19.2
QC	203	2.4	2.4	547	6.4	6.5	212	4.9	5.0
NB	35	4.5	4.6	45	5.7	5.9	14	3.5	3.7
NS	57	5.9	5.9	50	5.1	5.3	23	4.6	4.9
PE	5	3.2	3.3	8	5.0	4.9	3	3.7	4.2
NL	18	3.4	3.5	24	4.6	4.9	5	1.9	2.3
YT	4	9.7	10.0	6	14.2	14.2	10	46.5	39.3
NT	1	2.2	2.1	3	6.6	5.7	2	8.8	7.6
NU	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>3,668</b>	<b>9.8</b>	<b>9.9</b>	<b>6,306</b>	<b>16.5</b>	<b>16.8</b>	<b>3,515</b>	<b>19.1</b>	<b>19.3</b>

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases.

Not available (n/a) Data were not available at the time of this publication.

Includes deaths related to all illicit drugs including, but not limited to, opioids and stimulants.

**Note(s):**

Apparent opioid toxicity death (AOTD): A death caused by intoxication/toxicity (poisoning) resulting from substance use, where one or more of the substances is an opioid, regardless of how it was obtained (e.g. illegally or through personal prescription). Other substances may also be involved.

2020 data from Manitoba are based on January to March.

2021 data from Newfoundland and Labrador are based on January to March.

Data on apparent opioid toxicity deaths and stimulant toxicity deaths are not mutually exclusive. A high proportion of deaths involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.

**Table 1b. Number and rate (per 100,000 population) of total apparent stimulant toxicity deaths by province or territory, 2018 to 2021 (Jan to Jun)**

Province or territory	2018			2019			2020			2021 (Jan to Jun)		
	Number	Crude rate per 100,000 population	Age-adjusted rate per 100,000 population	Number	Crude rate per 100,000 population	Age-adjusted rate per 100,000 population	Number	Crude rate per 100,000 population	Age-adjusted rate per 100,000 population	Number	Crude rate per 100,000 population	Age-adjusted rate per 100,000 population
BC	878	17.5	17.5	414	8.1	8.1	570	11.0	11.1	81	3.1	3.0
SK	84	7.2	7.5	95	8.1	8.3	180	15.3	15.9	75	12.7	13.5
MB	n/a	n/a	n/a	81	5.9	6.0	25	1.8	1.9	n/a	n/a	n/a
ON	919	6.4	6.5	1,024	7.0	7.1	1,702	11.5	11.7	948	12.8	12.9
QC	162	1.9	2.0	138	1.6	1.7	n/a	n/a	n/a	n/a	n/a	n/a
NS	25	2.6	2.8	29	3.0	3.1	35	3.6	3.7	18	3.6	3.7
NL	n/a	n/a	n/a	n/a	n/a	n/a	10	1.9	2.2	3	1.2	1.1
NT	n/a	n/a	n/a	n/a	n/a	n/a	1	2.2	1.9	0	0.0	0.0

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases.

Not available (n/a) Data were not available at the time of this publication.

**Note(s):**

Apparent stimulant toxicity death (ASTD): A death caused by intoxication/toxicity (poisoning) resulting from substance use, where one or more of the substances is a stimulant, regardless of how it was obtained (e.g. illegally or through personal prescription). Other substances may also be involved.

2020 data from Manitoba are based on January to March.

2021 data from Newfoundland and Labrador are based on January to March.

Data on apparent opioid toxicity deaths and stimulant toxicity deaths are not mutually exclusive. A high proportion of deaths involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.





**Table 2a. Number and rate (per 100,000 population) of accidental apparent opioid toxicity deaths by province or territory, 2016 to 2021 (Jan to Jun)**

Province or territory	2016			2017			2018		
	Number	Crude rate per 100,000 population	Age-adjusted rate per 100,000 population	Number	Crude rate per 100,000 population	Age-adjusted rate per 100,000 population	Number	Crude rate per 100,000 population	Age-adjusted rate per 100,000 population
BC	790	16.3	16.1	1,268	25.7	25.6	1,540	30.7	30.6
AB	547	13.0	12.4	683	16.1	15.5	799	18.6	18.2
SK	76	6.7	6.9	74	6.4	6.6	114	9.8	10.3
MB	61	4.6	4.7	87	6.5	6.6	78	5.8	5.8
ON	726	5.2	5.2	1,127	8.0	8.0	1,319	9.2	9.3
QC	173	2.1	2.1	187	2.3	2.3	148	1.8	1.8
NB	27	3.5	3.6	32	4.2	4.4	23	3.0	3.0
NS	40	4.2	4.4	58	6.1	6.3	49	5.1	5.3
PE	5	3.4	3.2	5	3.3	3.6	8	5.2	5.4
NL	13	2.5	2.6	23	4.4	4.7	10	1.9	1.9
YT	5	13.0	11.8	6	15.1	13.4	4	9.9	9.5
NT	4	9.0	10.3	1	2.2	1.9	2	4.4	3.8
NU	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>2,467</b>	<b>6.8</b>	<b>6.8</b>	<b>3,551</b>	<b>9.7</b>	<b>9.8</b>	<b>4,094</b>	<b>11.0</b>	<b>11.1</b>



Province or territory	2019			2020			2021 (Jan to Jun)		
	Number	Crude rate per 100,000 population	Age-adjusted rate per 100,000 population	Number	Crude rate per 100,000 population	Age-adjusted rate per 100,000 population	Number	Crude rate per 100,000 population	Age-adjusted rate per 100,000 population
BC	961	18.9	18.8	1,697	32.9	32.8	986	37.8	37.8
AB	617	14.1	13.9	1,144	25.9	25.7	705	31.7	31.4
SK	109	9.3	9.9	246	20.9	21.8	90	15.3	16.3
MB	54	3.9	4.0	20	1.4	1.5	n/a	n/a	n/a
ON	1,408	9.7	9.7	2,303	15.6	15.7	1,325	17.9	18.0
QC	154	1.8	1.8	547	6.4	6.5	212	4.9	5.0
NB	27	3.5	3.5	38	4.9	4.9	13	3.3	3.4
NS	45	4.6	4.8	42	4.3	4.5	22	4.4	4.7
PE	5	3.2	3.3	8	5.0	4.9	3	3.7	4.2
NL	13	2.5	2.5	18	3.5	3.8	5	1.9	2.3
YT	4	9.7	10.0	6	14.2	14.2	10	46.5	39.3
NT	0	0.0	0.0	3	6.6	5.7	2	8.8	7.6
NU	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>3,397</b>	<b>9.0</b>	<b>9.1</b>	<b>6,072</b>	<b>16.0</b>	<b>16.2</b>	<b>3,373</b>	<b>18.3</b>	<b>18.5</b>

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases.

Not available (n/a) Data were not available at the time of this publication.

Includes deaths related to all illicit drugs including, but not limited to, opioids and stimulants.

#### Note(s):

Apparent opioid toxicity death (AOTD): A death caused by intoxication/toxicity (poisoning) resulting from substance use, where one or more of the substances is an opioid, regardless of how it was obtained (e.g. illegally or through personal prescription). Other substances may also be involved.

2020 data from Manitoba are based on January to March.

2021 data from Newfoundland and Labrador are based on January to March.

Data on apparent opioid toxicity deaths and stimulant toxicity deaths are not mutually exclusive. A high proportion of deaths involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.



**Table 2b. Number and rate (per 100,000 population) of accidental apparent stimulant toxicity deaths by province or territory, 2018 to 2021 (Jan to Jun)**

Province or territory	2018			2019			2020			2021 (Jan to Jun)		
	Number	Crude rate per 100,000 population	Age-adjusted rate per 100,000 population	Number	Crude rate per 100,000 population	Age-adjusted rate per 100,000 population	Number	Crude rate per 100,000 population	Age-adjusted rate per 100,000 population	Number	Crude rate per 100,000 population	Age-adjusted rate per 100,000 population
BC	866	17.3	17.2	398	7.8	7.8	565	11.0	11.0	81	3.1	3.0
SK	77	6.6	6.8	93	7.9	8.2	177	15.0	15.6	72	12.2	13.0
MB	n/a	n/a	n/a	73	5.3	5.4	24	1.7	1.8	n/a	n/a	n/a
ON	888	6.2	6.3	989	6.8	6.8	1,663	11.3	11.4	908	12.2	12.4
QC	144	1.7	1.7	130	1.5	1.6	n/a	n/a	n/a	n/a	n/a	n/a
NS	23	2.4	2.5	27	2.8	2.8	32	3.3	3.4	18	3.6	3.7
NL	n/a	n/a	n/a	n/a	n/a	n/a	10	1.9	2.2	3	1.2	1.1
NT	n/a	n/a	n/a	n/a	n/a	n/a	1	2.2	1.9	0	0.0	0.0

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases.

Not available (n/a) Data were not available at the time of this publication.

**Note(s):**

Apparent stimulant toxicity death (ASTD): A death caused by intoxication/toxicity (poisoning) resulting from substance use, where one or more of the substances is a stimulant, regardless of how it was obtained (e.g. illegally or through personal prescription). Other substances may also be involved.

2020 data from Manitoba are based on January to March.

2021 data from Newfoundland and Labrador are based on January to March.

Data on apparent opioid toxicity deaths and stimulant toxicity deaths are not mutually exclusive. A high proportion of deaths involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.



**Table 3a. Summary of apparent opioid toxicity deaths by manner of death, 2016 to 2021 (Jan to Jun)**

Manner of death	2016		2017		2018		2019		2020		2021 (Jan to Jun)	
	Count	%	Count	%	Count	%	Count	%	Count	%	Count	%
Total	2,826	100%	3,918	100%	4,393	100%	3,668	100%	6,306	100%	3,515	100%
Accidental	2,467	87%	3,551	91%	4,094	93%	3,397	93%	6,072	96%	3,373	96%
Suicide	264	9%	276	7%	218	5%	195	5%	114	2%	35	1%
Undetermined	95	3%	91	2%	81	2%	76	2%	120	2%	107	3%

**Note(s):**

Apparent opioid toxicity death (AOTD): A death caused by intoxication/toxicity (poisoning) resulting from substance use, where one or more of the substances is an opioid, regardless of how it was obtained (e.g. illegally or through personal prescription). Other substances may also be involved.

Data on apparent opioid toxicity deaths and stimulant toxicity deaths are not mutually exclusive. A high proportion of deaths involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.

**Table 3b. Summary of apparent stimulant toxicity deaths by manner of death, 2018 to 2021 (Jan to Jun)**

Manner of death	2018	2019	2020	2021 (Jan to Jun)
Total	100%	100%	100%	100%
Accidental	97%	96%	98%	96%
Suicide	2%	3%	1%	1%
Undetermined	1%	1%	1%	3%

**Note(s):**

Apparent stimulant toxicity death (ASTD): A death caused by intoxication/toxicity (poisoning) resulting from substance use, where one or more of the substances is a stimulant, regardless of how it was obtained (e.g. illegally or through personal prescription). Other substances may also be involved.

Data on apparent opioid toxicity deaths and stimulant toxicity deaths are not mutually exclusive. A high proportion of deaths involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.

**Table 4a. Summary of total apparent opioid toxicity deaths in Canada, 2016 to 2021 (Jan to Jun)**

		2016	2017	2018	2019	2020	2021 (Jan to Jun)
<b>Total</b>	Number	2,826	3,918	4,393	3,668	6,306	3,515
	Crude rate	7.8	10.7	11.9	9.8	16.6	19.1
	Age-adjusted rate	7.8	10.8	11.9	9.9	16.8	19.3
<b>Percent by sex</b>	Male	70%	74%	72%	71%	75%	74%
	Female	30%	26%	28%	29%	25%	26%
<b>Percent by age group</b>	0 to 19	2%	2%	2%	2%	2%	2%
	20 to 29	18%	20%	20%	19%	18%	19%
	30 to 39	27%	27%	27%	28%	27%	29%
	40 to 49	21%	21%	21%	20%	23%	22%
	50 to 59	23%	21%	21%	21%	21%	19%
	60 or more	9%	9%	9%	10%	9%	9%
<b>Percent involving</b>  (categories not mutually exclusive)	Fentanyl	52%	69%	69%	59%	81%	86%
	Fentanyl analogues	n/a	n/a	9%	21%	11%	14%
	Non fentanyl opioids	59%	39%	46%	44%	31%	23%
	Stimulants	n/a	n/a	55%	55%	60%	58%
	Other psychoactive substances	68%	60%	44%	44%	40%	40%

**Note(s):**

Apparent opioid toxicity death (AOTD): A death caused by intoxication/toxicity (poisoning) resulting from substance use, where one or more of the substances is an opioid, regardless of how it was obtained (e.g. illegally or through personal prescription). Other substances may also be involved.

Data on apparent opioid toxicity deaths and stimulant toxicity deaths are not mutually exclusive. A high proportion of deaths involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.



**Table 4b. Summary of total apparent stimulant toxicity deaths in Canada (based on available information from five to seven provinces or territories), 2018 to 2021 (Jan to Jun)**

		2018	2019	2020	2021 (Jan to Jun)
<b>Number of reporting provinces or territories</b>		5	6	7	6
<b>Percent by sex</b>	Male	75%	74%	77%	73%
	Female	25%	26%	23%	27%
<b>Percent by age group</b>	0 to 19	1%	2%	2%	1%
	20 to 29	17%	18%	16%	18%
	30 to 39	26%	28%	28%	29%
	40 to 49	23%	22%	24%	23%
	50 to 59	24%	22%	22%	22%
	60 or more	8%	9%	8%	8%
<b>Percent involving</b>  (categories not mutually exclusive)	Cocaine	70%	66%	67%	65%
	Methamphetamine	43%	45%	49%	51%
	Other stimulants	7%	7%	5%	5%
	Opioids	73%	68%	83%	86%
	Other psychoactive substances	30%	28%	22%	18%

**Note(s):**

Apparent stimulant toxicity death (ASTD): A death caused by intoxication/toxicity (poisoning) resulting from substance use, where one or more of the substances is a stimulant, regardless of how it was obtained (e.g. illegally or through personal prescription). Other substances may also be involved.

Data on apparent opioid toxicity deaths and stimulant toxicity deaths are not mutually exclusive. A high proportion of deaths involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.



**Table 5a. Number of total apparent opioid toxicity deaths by quarter and province or territory, January 2016 to June 2021**

Province or territory	2016				2017				2018			
	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4
BC	175	158	153	320	351	363	313	262	401	372	417	380
AB	137	140	159	166	162	182	183	217	194	206	210	195
SK	25	17	24	17	25	20	22	18	34	28	34	30
MB	18	20	17	33	34	34	19	19	21	17	24	31
ON	214	218	195	240	238	311	414	302	309	325	397	448
QC	66	64	68	60	66	79	68	68	54	50	42	63
NB	12	4	10	8	10	8	8	12	8	4	9	9
NS	17	14	17	5	19	17	13	15	15	18	10	11
PE	n/a	n/a	n/a	n/a	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NL	Suppr.	Suppr.	8	5	11	6	9	7	Suppr.	Suppr.	Suppr.	Suppr.
YT	0	2	2	3	4	0	1	2	0	0	3	2
NT	2	0	1	2	0	0	1	0	1	0	1	0
NU	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>666</b>	<b>637</b>	<b>654</b>	<b>859</b>	<b>920</b>	<b>1,020</b>	<b>1,051</b>	<b>922</b>	<b>1,037</b>	<b>1,020</b>	<b>1,147</b>	<b>1,169</b>



Province or territory	2019				2020				2021	
	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2
BC	305	248	228	231	279	491	481	501	519	509
AB	161	188	150	125	160	316	349	335	356	352
SK	25	34	23	35	26	73	80	80	58	38
MB	10	16	11	25	22	n/a	n/a	n/a	n/a	n/a
ON	451	468	241	370	462	621	627	720	727	687
QC	50	53	51	49	116	148	167	116	99	113
NB	6	7	15	7	5	10	12	18	10	4
NS	14	18	13	12	9	12	10	19	12	11
PE	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NL	8	6	Suppr.	Suppr.	Suppr.	9	6	Suppr.	5	n/a
YT	1	0	0	3	2	0	3	1	5	5
NT	1	0	0	0	0	0	0	3	1	1
NU	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>1,032</b>	<b>1,038</b>	<b>732</b>	<b>857</b>	<b>1,081</b>	<b>1,680</b>	<b>1,735</b>	<b>1,793</b>	<b>1,792</b>	<b>1,720</b>

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases.

Not available (n/a) Data were not available at the time of this publication.

Includes deaths related to all illicit drugs including, but not limited to, opioids and stimulants.

Q1 = January to March; Q2 = April to June; Q3 = July to September; Q4 = October to December

#### Note(s):

Apparent opioid toxicity death (AOTD): A death caused by intoxication/toxicity (poisoning) resulting from substance use, where one or more of the substances is an opioid, regardless of how it was obtained (e.g. illegally or through personal prescription). Other substances may also be involved.

Data on apparent opioid toxicity deaths and stimulant toxicity deaths are not mutually exclusive. A high proportion of deaths involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.





**Table 5b. Number of total apparent stimulant toxicity deaths by quarter and province or territory, January 2018 to June 2021**

Province or territory	2018				2019				2020				2021	
	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2
BC	273	235	231	139	43	51	152	168	162	188	106	114	67	14
SK	23	24	18	19	23	31	18	23	21	51	49	59	41	34
MB	n/a	n/a	n/a	n/a	15	23	16	27	25	n/a	n/a	n/a	n/a	n/a
ON	187	207	240	285	277	302	175	270	309	476	440	477	500	448
QC	36	47	49	30	37	33	30	38	n/a	n/a	n/a	n/a	n/a	n/a
NS	4	8	7	6	9	8	7	5	7	10	7	11	8	10
NL	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	Suppr.	6	0	Suppr.	Suppr.	n/a
NT	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	1	0	0	0	0	0

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases.

Not available (n/a) Data were not available at the time of this publication.

Q1 = January to March; Q2 = April to June; Q3 = July to September; Q4 = October to December

**Note(s):**

Apparent stimulant toxicity death (ASTD): A death caused by intoxication/toxicity (poisoning) resulting from substance use, where one or more of the substances is a stimulant, regardless of how it was obtained (e.g. illegally or through personal prescription). Other substances may also be involved.

Data on apparent opioid toxicity deaths and stimulant toxicity deaths are not mutually exclusive. A high proportion of deaths involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.

**Table 6a. Number of accidental apparent opioid toxicity deaths by quarter and province or territory, January 2016 to June 2021**

Province or territory	2016				2017				2018			
	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4
BC	169	158	147	316	347	358	305	258	399	365	404	372
AB	118	130	144	155	152	161	171	199	193	206	207	193
SK	25	14	22	15	22	17	19	16	31	25	30	28
MB	11	14	13	23	26	27	18	16	15	14	22	27
ON	174	180	174	198	191	276	380	280	262	293	362	402
QC	50	44	43	36	45	52	47	43	35	39	27	47
NB	11	3	6	7	8	6	7	11	7	4	5	7
NS	14	8	15	3	17	16	12	13	14	16	8	11
PE	n/a	n/a	n/a	n/a	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NL	Suppr.	Suppr.	Suppr.	Suppr.	7	Suppr.	7	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
YT	0	2	0	3	3	0	1	2	0	0	2	2
NT	1	0	1	2	0	0	1	0	1	0	1	0
NU	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>573</b>	<b>553</b>	<b>565</b>	<b>758</b>	<b>818</b>	<b>913</b>	<b>968</b>	<b>838</b>	<b>957</b>	<b>962</b>	<b>1,068</b>	<b>1,089</b>



Province or territory	2019				2020				2021	
	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2
BC	291	237	212	221	270	476	463	488	499	487
AB	160	184	149	124	152	312	348	332	354	351
SK	23	33	22	31	24	68	78	76	55	35
MB	7	13	10	24	20	n/a	n/a	n/a	n/a	n/a
ON	415	441	217	335	435	593	590	685	692	633
QC	36	45	36	37	116	148	167	116	99	113
NB	5	6	11	5	4	9	9	16	9	4
NS	10	14	10	11	5	12	10	15	11	11
PE	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NL	5	Suppr.	Suppr.	Suppr.	Suppr.	8	Suppr.	Suppr.	5	n/a
YT	1	0	0	3	2	0	3	1	5	5
NT	0	0	0	0	0	0	0	3	1	1
NU	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>953</b>	<b>973</b>	<b>667</b>	<b>791</b>	<b>1,028</b>	<b>1,626</b>	<b>1,668</b>	<b>1,732</b>	<b>1,730</b>	<b>1,640</b>

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases.

Not available (n/a) Data were not available at the time of this publication.

Includes deaths related to all illicit drugs including, but not limited to, opioids and stimulants.

Q1 = January to March; Q2 = April to June; Q3 = July to September; Q4 = October to December

**Note(s):**

Apparent opioid toxicity death (AOTD): A death caused by intoxication/toxicity (poisoning) resulting from substance use, where one or more of the substances is an opioid, regardless of how it was obtained (e.g. illegally or through personal prescription). Other substances may also be involved.

Data on apparent opioid toxicity deaths and stimulant toxicity deaths are not mutually exclusive. A high proportion of deaths involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.

**Table 6b. Number of accidental apparent stimulant toxicity deaths by quarter and province or territory, January 2018 to June 2021**

Province or territory	2018				2019				2020				2021	
	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2
BC	272	233	224	137	38	49	148	163	160	188	105	112	67	14
SK	23	20	15	19	22	30	18	23	19	50	49	59	38	34
MB	n/a	n/a	n/a	n/a	12	21	14	26	24	n/a	n/a	n/a	n/a	n/a
ON	184	202	230	272	264	293	168	264	301	464	428	470	487	421
QC	31	43	43	27	34	33	28	35	n/a	n/a	n/a	n/a	n/a	n/a
NS	4	7	6	6	9	8	6	4	5	10	7	10	8	10
NL	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	Suppr.	6	0	Suppr.	Suppr.	n/a
NT	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	1	0	0	0	0	0

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases.

Not available (n/a) Data were not available at the time of this publication.

Q1 = January to March; Q2 = April to June; Q3 = July to September; Q4 = October to December

**Note(s):**

Apparent stimulant toxicity death (ASTD): A death caused by intoxication/toxicity (poisoning) resulting from substance use, where one or more of the substances is a stimulant, regardless of how it was obtained (e.g. illegally or through personal prescription). Other substances may also be involved.

Data on apparent opioid toxicity deaths and stimulant toxicity deaths are not mutually exclusive. A high proportion of deaths involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.

**Table 7a. Percentage of accidental apparent opioid toxicity deaths by sex and province or territory, 2016 to 2021 (Jan to Jun)**

Province or territory	2016		2017		2018		2019		2020		2021 (Jan to Jun)	
	Male	Female	Male	Female	Male	Female	Male	Female	Male	Female	Male	Female
BC	80%	20%	82%	18%	80%	20%	75%	25%	82%	18%	90%	10%
AB	73%	27%	76%	24%	74%	26%	74%	26%	77%	23%	75%	25%
SK	59%	41%	57%	43%	59%	41%	64%	36%	66%	34%	66%	34%
MB	61%	39%	67%	33%	55%	45%	69%	31%	55%	45%	n/a	n/a
ON	71%	29%	75%	25%	73%	27%	73%	27%	77%	23%	74%	26%
QC	69%	31%	68%	32%	63%	37%	70%	30%	75%	25%	77%	23%
NB	63%	37%	59%	41%	43%	57%	48%	52%	55%	45%	69%	31%
NS	78%	23%	71%	29%	67%	33%	71%	29%	69%	31%	73%	27%
PE	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NL	62%	38%	70%	30%	50%	50%	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
YT	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NT	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NU	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>74%</b>	<b>26%</b>	<b>77%</b>	<b>23%</b>	<b>74%</b>	<b>26%</b>	<b>72%</b>	<b>28%</b>	<b>76%</b>	<b>24%</b>	<b>75%</b>	<b>25%</b>

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases.

Not available (n/a) Data were not available at the time of this publication.

Includes deaths related to all illicit drugs including, but not limited to, opioids and stimulants.

**Note(s):**

Apparent opioid toxicity death (AOTD): A death caused by intoxication/toxicity (poisoning) resulting from substance use, where one or more of the substances is an opioid, regardless of how it was obtained (e.g. illegally or through personal prescription). Other substances may also be involved.

2020 data from Manitoba are based on January to March.

2021 data from Newfoundland and Labrador are based on January to March.

Data on apparent opioid toxicity deaths and stimulant toxicity deaths are not mutually exclusive. A high proportion of deaths involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.





**Table 7b. Percentage of accidental apparent stimulant toxicity deaths by sex and province or territory, 2018 to 2021 (Jan to Jun)**

Province or territory	2018		2019		2020		2021 (Jan to Jun)	
	Male	Female	Male	Female	Male	Female	Male	Female
BC	79%	21%	74%	26%	81%	19%	85%	15%
SK	58%	42%	65%	35%	67%	33%	65%	35%
MB	n/a	n/a	75%	25%	54%	46%	n/a	n/a
ON	74%	26%	75%	25%	77%	23%	73%	27%
QC	78%	22%	75%	25%	n/a	n/a	n/a	n/a
NS	74%	26%	85%	15%	81%	19%	72%	28%
NL	n/a	n/a	n/a	n/a	Suppr.	Suppr.	Suppr.	Suppr.
NT	n/a	n/a	n/a	n/a	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>76%</b>	<b>24%</b>	<b>74%</b>	<b>26%</b>	<b>77%</b>	<b>23%</b>	<b>74%</b>	<b>26%</b>

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases.

Not available (n/a) Data were not available at the time of this publication.

**Note(s):**

Apparent stimulant toxicity death (ASTD): A death caused by intoxication/toxicity (poisoning) resulting from substance use, where one or more of the substances is a stimulant, regardless of how it was obtained (e.g. illegally or through personal prescription). Other substances may also be involved.

2020 data from Manitoba are based on January to March.

2021 data from Newfoundland and Labrador are based on January to March.

Data on apparent opioid toxicity deaths and stimulant toxicity deaths are not mutually exclusive. A high proportion of deaths involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.

**Table 8a. Crude rate (per 100,000) of accidental apparent opioid toxicity deaths by sex and province or territory, 2016 to 2021 (Jan to Jun)**

Province or territory	2016		2017		2018		2019		2020		2021 (Jan to Jun)	
	Male	Female	Male	Female	Male	Female	Male	Female	Male	Female	Male	Female
BC	26.4	6.4	42.9	8.9	35.6	8.7	14.1	4.7	22.1	4.6	7.4	0.8
AB	18.9	7.1	24.3	7.8	27.3	9.8	20.7	7.5	39.5	12.1	47.2	16.1
SK	7.9	5.5	7.2	5.6	11.4	8.2	11.8	6.7	27.3	14.4	19.9	10.6
MB	5.6	3.6	8.7	4.3	6.4	5.2	5.4	2.5	1.6	1.3	n/a	n/a
ON	7.5	3.0	12.1	4.0	13.6	5.0	14.2	5.2	24.2	7.2	26.8	9.1
QC	2.9	1.3	3.1	1.4	2.2	1.3	2.5	1.1	9.5	3.2	7.6	2.3
NB	4.5	2.6	5.0	3.4	2.6	3.3	3.4	3.6	5.4	4.3	4.6	2.0
NS	6.7	1.9	8.8	3.5	7.0	3.3	6.7	2.6	6.0	2.6	6.6	2.4
PE	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NL	3.0	1.9	6.1	2.6	1.9	1.9	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
YT	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NT	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NU	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>10.1</b>	<b>3.6</b>	<b>15.0</b>	<b>4.5</b>	<b>14.6</b>	<b>5.2</b>	<b>11.3</b>	<b>4.3</b>	<b>20.4</b>	<b>6.3</b>	<b>20.3</b>	<b>6.8</b>

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases.

Not available (n/a) Data were not available at the time of this publication.

Includes deaths related to all illicit drugs including, but not limited to, opioids and stimulants.

**Note(s):**

Apparent opioid toxicity death (AOTD): A death caused by intoxication/toxicity (poisoning) resulting from substance use, where one or more of the substances is an opioid, regardless of how it was obtained (e.g. illegally or through personal prescription). Other substances may also be involved.

2020 data from Manitoba are based on January to March.

2021 data from Newfoundland and Labrador are based on January to March.

Data on apparent opioid toxicity deaths and stimulant toxicity deaths are not mutually exclusive. A high proportion of deaths involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.





**Table 8b. Crude rate (per 100,000 population) of accidental apparent stimulant toxicity deaths by sex and province or territory, 2018 to 2021 (Jan to Jun)**

Province or territory	2018		2019		2020		2021 (Jan to Jun)	
	Male	Female	Male	Female	Male	Female	Male	Female
BC	27.6	7.2	11.7	4.0	18.0	4.0	5.4	0.9
SK	7.7	5.6	10.2	5.7	20.0	9.9	15.8	8.5
MB	n/a	n/a	8.0	2.6	1.9	1.6	n/a	n/a
ON	9.3	3.2	10.3	3.4	17.5	5.2	18.2	6.5
QC	2.7	0.7	2.3	0.8	n/a	n/a	n/a	n/a
NS	3.6	1.2	4.8	0.8	5.4	1.2	5.4	2.0
NL	n/a	n/a	n/a	n/a	Suppr.	Suppr.	Suppr.	Suppr.
NT	n/a	n/a	n/a	n/a	Suppr.	Suppr.	Suppr.	Suppr.

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases.

Not available (n/a) Data were not available at the time of this publication.

**Note(s):**

Apparent stimulant toxicity death (ASTD): A death caused by intoxication/toxicity (poisoning) resulting from substance use, where one or more of the substances is a stimulant, regardless of how it was obtained (e.g. illegally or through personal prescription). Other substances may also be involved.

2020 data from Manitoba are based on January to March.

2021 data from Newfoundland and Labrador are based on January to March.

Data on apparent opioid toxicity deaths and stimulant toxicity deaths are not mutually exclusive. A high proportion of deaths involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.





**Table 9a. Percentage of accidental apparent opioid toxicity deaths by age group and province or territory, 2016 to 2021 (Jan to Jun)**

## 2016

Province or territory	Age group					
	0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	3%	21%	28%	23%	21%	5%
AB	2%	21%	37%	18%	17%	5%
SK	0%	12%	45%	13%	21%	9%
MB	5%	30%	26%	16%	15%	8%
ON	2%	19%	24%	22%	25%	8%
QC	2%	13%	19%	25%	28%	12%
NB	0%	11%	22%	30%	26%	11%
NS	3%	20%	30%	13%	23%	13%
PE	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NL	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
YT	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NT	0%	50%	0%	0%	25%	25%
NU	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>2%</b>	<b>20%</b>	<b>28%</b>	<b>21%</b>	<b>22%</b>	<b>7%</b>

## 2017

Province or territory	Age group					
	0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	2%	19%	28%	23%	21%	7%
AB	3%	24%	32%	19%	17%	5%
SK	4%	28%	23%	20%	16%	8%
MB	0%	31%	32%	22%	7%	8%
ON	2%	21%	27%	22%	20%	8%
QC	2%	15%	25%	18%	27%	13%
NB	3%	22%	28%	19%	19%	9%
NS	5%	10%	26%	24%	29%	5%
PE	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NL	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
YT	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NT	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NU	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>2%</b>	<b>21%</b>	<b>28%</b>	<b>21%</b>	<b>20%</b>	<b>7%</b>



## 2018

Province or territory	Age group					
	0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	2%	19%	27%	21%	23%	8%
AB	2%	25%	30%	20%	18%	7%
SK	0%	24%	25%	20%	20%	11%
MB	0%	26%	28%	20%	20%	7%
ON	2%	20%	29%	22%	20%	7%
QC	1%	14%	28%	22%	23%	13%
NB	0%	9%	22%	22%	35%	13%
NS	2%	22%	24%	20%	18%	12%
PE	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NL	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
YT	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NT	50%	50%	0%	0%	0%	0%
NU	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>2%</b>	<b>21%</b>	<b>28%</b>	<b>21%</b>	<b>21%</b>	<b>8%</b>

## 2019

Province or territory	Age group					
	0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	3%	19%	29%	21%	20%	8%
AB	2%	23%	31%	21%	18%	6%
SK	1%	19%	23%	24%	28%	6%
MB	2%	28%	31%	17%	19%	4%
ON	1%	21%	29%	20%	22%	8%
QC	2%	12%	17%	26%	25%	18%
NB	0%	7%	30%	19%	22%	22%
NS	2%	16%	22%	27%	18%	16%
PE	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NL	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
YT	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NT	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NU	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>2%</b>	<b>20%</b>	<b>28%</b>	<b>20%</b>	<b>21%</b>	<b>8%</b>



## 2020


Province or territory	Age group					
	0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	2%	17%	24%	26%	22%	9%
AB	2%	20%	29%	23%	19%	7%
SK	3%	20%	28%	21%	20%	9%
MB	0%	20%	45%	25%	5%	5%
ON	1%	19%	29%	23%	20%	8%
QC	2%	13%	20%	22%	25%	17%
NB	0%	13%	16%	29%	24%	18%
NS	0%	24%	21%	36%	12%	7%
PE	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NL	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
YT	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NT	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NU	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>2%</b>	<b>18%</b>	<b>27%</b>	<b>23%</b>	<b>21%</b>	<b>9%</b>

## 2021 (Jan to Jun)

Province or territory	Age group					
	0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	2%	14%	25%	20%	21%	19%
AB	2%	22%	32%	20%	16%	8%
SK	1%	20%	28%	23%	23%	4%
MB	n/a	n/a	n/a	n/a	n/a	n/a
ON	1%	18%	30%	22%	20%	8%
QC	Suppr.	12%	24%	25%	21%	Suppr.
NB	0%	15%	15%	38%	15%	15%
NS	5%	27%	23%	23%	18%	5%
PE	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NL	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
YT	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NT	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NU	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>1%</b>	<b>19%</b>	<b>30%</b>	<b>22%</b>	<b>19%</b>	<b>9%</b>

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases.

Not available (n/a) Data were not available at the time of this publication.

 Includes deaths related to all illicit drugs including, but not limited to, opioids and stimulants.

### Note(s):

Apparent opioid toxicity death (AOTD): A death caused by intoxication/toxicity (poisoning) resulting from substance use, where one or more of the substances is an opioid, regardless of how it was obtained (e.g. illegally or through personal prescription). Other substances may also be involved.

2020 data from Manitoba are based on January to March.

2021 data from Newfoundland and Labrador are based on January to March.

Data on apparent opioid toxicity deaths and stimulant toxicity deaths are not mutually exclusive. A high proportion of deaths involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.



**Table 9b. Percentage of accidental apparent stimulant toxicity deaths by age group and province or territory, 2016 to 2021 (Jan to Jun)**

**2018**

Province or territory	Age group					
	0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	1%	17%	25%	22%	25%	9%
SK	1%	21%	32%	27%	13%	5%
MB	n/a	n/a	n/a	n/a	n/a	n/a
ON	1%	17%	29%	24%	22%	7%
QC	2%	17%	15%	22%	30%	15%
NS	4%	35%	22%	13%	22%	4%
NL	n/a	n/a	n/a	n/a	n/a	n/a
NT	n/a	n/a	n/a	n/a	n/a	n/a
<b>Canada</b>	<b>1%</b>	<b>17%</b>	<b>26%</b>	<b>23%</b>	<b>24%</b>	<b>8%</b>

**2019**

Province or territory	Age group					
	0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	2%	18%	26%	22%	23%	10%
SK	1%	22%	37%	19%	16%	5%
MB	1%	25%	32%	29%	12%	1%
ON	1%	17%	28%	22%	23%	9%
QC	3%	15%	19%	19%	28%	15%
NS	4%	15%	26%	19%	19%	19%
NL	n/a	n/a	n/a	n/a	n/a	n/a
NT	n/a	n/a	n/a	n/a	n/a	n/a
<b>Canada</b>	<b>2%</b>	<b>18%</b>	<b>27%</b>	<b>22%</b>	<b>22%</b>	<b>9%</b>

## 2020

Province or territory	Age group					
	0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	1%	16%	24%	27%	22%	10%
SK	2%	18%	34%	21%	19%	6%
MB	4%	17%	29%	33%	13%	4%
ON	1%	16%	29%	23%	22%	8%
QC	n/a	n/a	n/a	n/a	n/a	n/a
NS	3%	22%	19%	22%	22%	13%
NL	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NT	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>1%</b>	<b>16%</b>	<b>28%</b>	<b>24%</b>	<b>22%</b>	<b>8%</b>

## 2021 (Jan to Jun)

Province or territory	Age group					
	0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	0%	14%	21%	23%	20%	22%
SK	1%	25%	31%	18%	24%	1%
MB	n/a	n/a	n/a	n/a	n/a	n/a
ON	0%	17%	29%	24%	22%	7%
QC	n/a	n/a	n/a	n/a	n/a	n/a
NS	0%	39%	22%	11%	17%	11%
NL	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NT	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>0%</b>	<b>18%</b>	<b>29%</b>	<b>23%</b>	<b>21%</b>	<b>8%</b>

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases.

Not available (n/a) Data were not available at the time of this publication.

### Note(s):

Apparent stimulant toxicity death (ASTD): A death caused by intoxication/toxicity (poisoning) resulting from substance use, where one or more of the substances is a stimulant, regardless of how it was obtained (e.g. illegally or through personal prescription). Other substances may also be involved.

2020 data from Manitoba are based on January to March.

2021 data from Newfoundland and Labrador are based on January to March.

Data on apparent opioid toxicity deaths and stimulant toxicity deaths are not mutually exclusive. A high proportion of deaths involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.

**Table 10a. Percentage of accidental apparent opioid toxicity deaths by type of opioids and province or territory, 2016 to 2021 (Jan to Jun)**

Province or territory	2016			2017			2018		
	Fentanyl	Analogues	Non-fentanyl	Fentanyl	Analogues	Non-fentanyl	Fentanyl	Analogues	Non-fentanyl
BC	80%	3%	55%	91%	11%	29%	93%	5%	38%
AB	63%	n/a	37%	81%	n/a	19%	72%	20%	28%
SK	11%	n/a	89%	20%	n/a	80%	41%	4%	68%
MB	46%	n/a	54%	51%	n/a	49%	38%	n/a	62%
ON	45%	n/a	55%	69%	n/a	31%	70%	9%	44%
QC	23%	n/a	88%	27%	n/a	93%	18%	4%	94%
NB	15%	n/a	93%	22%	n/a	91%	4%	n/a	100%
NS	18%	n/a	83%	10%	n/a	90%	16%	8%	96%
PE	20%	n/a	80%	0%	n/a	100%	13%	n/a	88%
NL	Suppr.	n/a	92%	26%	n/a	74%	0%	n/a	100%
YT	80%	n/a	20%	83%	n/a	17%	100%	n/a	0%
NT	25%	n/a	75%	0%	n/a	100%	0%	n/a	100%
NU	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>57%</b>	<b>n/a</b>	<b>55%</b>	<b>73%</b>	<b>n/a</b>	<b>35%</b>	<b>73%</b>	<b>10%</b>	<b>43%</b>

Province or territory	2019			2020			2021 (Jan to Jun)		
	Fentanyl	Analogues	Non-fentanyl	Fentanyl	Analogues	Non-fentanyl	Fentanyl	Analogues	Non-fentanyl
BC	88%	11%	33%	91%	33%	24%	94%	36%	21%
AB	77%	10%	31%	87%	9%	20%	90%	20%	12%
SK	37%	11%	71%	63%	57%	50%	87%	74%	26%
MB	59%	11%	54%	65%	0%	40%	n/a	n/a	n/a
ON	57%	35%	36%	89%	2%	24%	92%	6%	20%
QC	16%	1%	93%	30%	3%	84%	42%	6%	72%
NB	19%	n/a	89%	11%	n/a	89%	15%	n/a	85%
NS	9%	2%	96%	24%	7%	86%	9%	9%	95%
PE	0%	n/a	100%	50%	0%	50%	0%	0%	100%
NL	Suppr.	n/a	85%	Suppr.	Suppr.	78%	Suppr.	0%	Suppr.
YT	75%	n/a	25%	67%	0%	17%	100%	89%	11%
NT	Suppr.	Suppr.	Suppr.	100%	Suppr.	Suppr.	50%	0%	100%
NU	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>62%</b>	<b>22%</b>	<b>41%</b>	<b>83%</b>	<b>11%</b>	<b>29%</b>	<b>87%</b>	<b>14%</b>	<b>22%</b>

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases.

Not available (n/a) Data were not available at the time of this publication.

Includes deaths related to all illicit drugs including, but not limited to, opioids and stimulants.

Analogue = Fentanyl analogues; Non-fentanyl = Non-fentanyl opioids

**Note(s):**

Apparent opioid toxicity death (AOTD): A death caused by intoxication/toxicity (poisoning) resulting from substance use, where one or more of the substances is an opioid, regardless of how it was obtained (e.g. illegally or through personal prescription). Other substances may also be involved.

2020 data from Manitoba are based on January to March.

2021 data from Newfoundland and Labrador are based on January to March.

Until investigations are completed, some provinces and territories report fentanyl information only or report data on fentanyl and fentanyl analogues together. Therefore, some deaths involving fentanyl analogues may be included in the fentanyl percentages.

Data on apparent opioid toxicity deaths and stimulant toxicity deaths are not mutually exclusive. A high proportion of deaths involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.



**Table 10b. Percentage of accidental stimulant toxicity deaths by type of stimulants and province or territory, 2018 to 2021 (Jan to Jun)**

Province or territory	2018			2019			2020			2021 (Jan to Jun)		
	Cocaine	Meth	Other stimulants	Cocaine	Meth	Other stimulants	Cocaine	Meth	Other stimulants	Cocaine	Meth	Other stimulants
BC	68%	50%	3%	62%	52%	4%	60%	58%	5%	72%	51%	10%
SK	51%	55%	14%	25%	77%	9%	32%	73%	3%	15%	88%	4%
MB	n/a	n/a	n/a	55%	59%	5%	46%	63%	0%	n/a	n/a	n/a
ON	75%	36%	5%	73%	39%	4%	73%	44%	5%	69%	48%	4%
QC	62%	n/a	38%	62%	n/a	38%	n/a	n/a	n/a	n/a	n/a	n/a
NS	91%	13%	9%	67%	33%	15%	84%	0%	22%	94%	11%	0%
NL	n/a	n/a	n/a	n/a	n/a	n/a	Suppr.	Suppr.	60%	Suppr.	Suppr.	Suppr.
NT	n/a	n/a	n/a	n/a	n/a	n/a	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>70%</b>	<b>43%</b>	<b>7%</b>	<b>66%</b>	<b>45%</b>	<b>7%</b>	<b>67%</b>	<b>49%</b>	<b>5%</b>	<b>65%</b>	<b>50%</b>	<b>4%</b>

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases.

Not available (n/a) Data were not available at the time of this publication.

Meth: methamphetamine

**Note(s):**

Apparent stimulant toxicity death (ASTD): A death caused by intoxication/toxicity (poisoning) resulting from substance use, where one or more of the substances is a stimulant, regardless of how it was obtained (e.g. illegally or through personal prescription). Other substances may also be involved.

2020 data from Manitoba are based on January to March.

2021 data from Newfoundland and Labrador are based on January to March.

Data on apparent opioid toxicity deaths and stimulant toxicity deaths are not mutually exclusive. A high proportion of deaths involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.





**Table 11a. Percentage of accidental apparent opioid toxicity deaths involving stimulants, by province or territory, 2018 to 2021 (Jan to Jun)**

Province or territory	Year			
	2018	2019	2020	2021 (Jan to Jun)
BC	72%	74%	73%	71%
AB	57%	60%	61%	55%
SK	44%	50%	61%	59%
MB	n/a	59%	75%	n/a
ON	48%	52%	59%	61%
QC	n/a	n/a	n/a	n/a
NS	31%	38%	38%	45%
NL	n/a	n/a	44%	Suppr.
NT	n/a	n/a	0%	0%
<b>Canada</b>	<b>57%</b>	<b>57%</b>	<b>62%</b>	<b>59%</b>

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases.

Not available (n/a) Data were not available at the time of this publication.

**Note(s):**

Apparent opioid toxicity death (AOTD): A death caused by intoxication/toxicity (poisoning) resulting from substance use, where one or more of the substances is an opioid, regardless of how it was obtained (e.g. illegally or through personal prescription). Other substances may also be involved.

2020 data from Manitoba are based on January to March.

2021 data from Newfoundland and Labrador are based on January to March.

Data on apparent opioid toxicity deaths and stimulant toxicity deaths are not mutually exclusive. A high proportion of deaths involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.



**Table 11b. Percentage of accidental apparent stimulant toxicity deaths involving opioids, by province or territory, 2018 to 2021 (Jan to Jun)**

Province or territory	Year			
	2018	2019	2020	2021 (Jan to Jun)
BC	91%	88%	88%	93%
SK	65%	58%	85%	74%
MB	n/a	44%	63%	n/a
ON	71%	74%	82%	87%
QC	n/a	n/a	n/a	n/a
NS	65%	63%	50%	56%
NL	n/a	n/a	80%	Suppr.
NT	n/a	n/a	0%	0%
<b>Canada</b>	<b>74%</b>	<b>69%</b>	<b>83%</b>	<b>86%</b>

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases.

Not available (n/a) Data were not available at the time of this publication.

**Note(s):**

Apparent stimulant toxicity death (ASTD): A death caused by intoxication/toxicity (poisoning) resulting from substance use, where one or more of the substances is a stimulant, regardless of how it was obtained (e.g. illegally or through personal prescription). Other substances may also be involved.

2020 data from Manitoba are based on January to March.

2021 data from Newfoundland and Labrador are based on January to March.

Data on apparent opioid toxicity deaths and stimulant toxicity deaths are not mutually exclusive. A high proportion of deaths involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.




**Table 12a. Percentage of accidental apparent opioid toxicity death involving other psychoactive substances, by province or territory, 2016 to 2021 (Jan to Jun)**

Province or territory	Year					
	2016	2017	2018	2019	2020	2021 (Jan to Jun)
BC	31%	33%	35%	41%	41%	55%
AB	65%	72%	73%	77%	81%	83%
SK	95%	91%	43%	52%	49%	78%
MB	95%	99%	51%	34%	20%	n/a
ON	93%	66%	23%	20%	17%	14%
QC	80%	82%	89%	86%	49%	54%
NB	93%	100%	100%	100%	97%	100%
NS	90%	83%	84%	84%	81%	82%
PE	100%	100%	88%	100%	88%	67%
NL	100%	91%	100%	100%	44%	Suppr.
YT	100%	100%	100%	75%	0%	11%
NT	75%	100%	100%	Suppr.	33%	0%
NU	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>66%</b>	<b>59%</b>	<b>43%</b>	<b>43%</b>	<b>40%</b>	<b>41%</b>

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases.

Not available (n/a) Data were not available at the time of this publication.

 Includes deaths related to all illicit drugs including, but not limited to, opioids and stimulants.

**Note(s):**

Apparent opioid toxicity death (AOTD): A death caused by intoxication/toxicity (poisoning) resulting from substance use, where one or more of the substances is an opioid, regardless of how it was obtained (e.g. illegally or through personal prescription). Other substances may also be involved.

2020 data from Manitoba are based on January to March.

2021 data from Newfoundland and Labrador are based on January to March.

Data on apparent opioid toxicity deaths and stimulant toxicity deaths are not mutually exclusive. A high proportion of deaths involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.



**Table 12b. Percentage of accidental apparent stimulant toxicity death involving other psychoactive substances, by province or territory, 2018 to 2021 (Jan to Jun)**

Province or territory	Year			
	2018	2019	2020	2021 (Jan to Jun)
BC	32%	37%	37%	52%
SK	43%	39%	41%	65%
MB	n/a	27%	21%	n/a
ON	16%	15%	14%	10%
QC	86%	81%	n/a	n/a
NS	65%	59%	56%	50%
NL	n/a	n/a	Suppr.	0%
NT	n/a	n/a	0%	0%
<b>Canada</b>	<b>30%</b>	<b>28%</b>	<b>22%</b>	<b>18%</b>

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases.

Not available (n/a) Data were not available at the time of this publication.

**Note(s):**

Apparent stimulant toxicity death (ASTD): A death caused by intoxication/toxicity (poisoning) resulting from substance use, where one or more of the substances is a stimulant, regardless of how it was obtained (e.g. illegally or through personal prescription). Other substances may also be involved.

2020 data from Manitoba are based on January to March.

2021 data from Newfoundland and Labrador are based on January to March.

Data on apparent opioid toxicity deaths and stimulant toxicity deaths are not mutually exclusive. A high proportion of deaths involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.



**Table 13. Percentage of accidental opioid toxicity deaths by origin of opioid(s), by province or territory, 2018 to 2021 (Jan to Jun)**

**2018**

Province or territory	Origin			
	Pharmaceutical only	Non-pharmaceutical only*	Both	Undetermined
BC	0%	62%	0%	38%
AB	16%	78%	1%	5%
SK	58%	35%	4%	3%
MB	n/a	n/a	n/a	n/a
ON	23%	65%	9%	3%
QC	n/a	n/a	n/a	n/a
NB	96%	0%	4%	0%
NS	85%	6%	8%	0%
PE	n/a	n/a	n/a	n/a
NL	100%	0%	0%	0%
YT	n/a	n/a	n/a	n/a
NT	n/a	n/a	n/a	n/a
NU	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>25%</b>	<b>66%</b>	<b>6%</b>	<b>3%</b>

**2019**

Province or territory	Origin			
	Pharmaceutical only	Non-pharmaceutical only*	Both	Undetermined
BC	0%	66%	0%	33%
AB	16%	77%	0%	6%
SK	60%	31%	5%	5%
MB	39%	52%	4%	6%
ON	18%	68%	11%	2%
QC	n/a	n/a	n/a	n/a
NB	81%	11%	4%	4%
NS	93%	5%	2%	0%
PE	n/a	n/a	n/a	n/a
NL	77%	23%	0%	0%
YT	n/a	n/a	n/a	n/a
NT	n/a	n/a	n/a	n/a
NU	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>23%</b>	<b>66%</b>	<b>7%</b>	<b>3%</b>



## 2020

Province or territory	Origin			
	Pharmaceutical only	Non-pharmaceutical only*	Both	Undetermined
BC	0%	76%	0%	24%
AB	9%	84%	1%	6%
SK	32%	52%	9%	7%
MB	32%	63%	0%	5%
ON	10%	78%	10%	1%
QC	n/a	n/a	n/a	n/a
NB	83%	8%	0%	8%
NS	79%	13%	8%	0%
PE	n/a	n/a	n/a	n/a
NL	83%	17%	0%	0%
YT	n/a	n/a	n/a	n/a
NT	n/a	n/a	n/a	n/a
NU	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>13%</b>	<b>76%</b>	<b>7%</b>	<b>3%</b>



## 2021 (Jan to Jun)

Province or territory	Origin			
	Pharmaceutical only	Non-pharmaceutical only*	Both	Undetermined
BC	0%	79%	0%	21%
AB	5%	89%	0%	5%
SK	11%	76%	9%	3%
MB	n/a	n/a	n/a	n/a
ON	6%	82%	11%	1%
QC	n/a	n/a	n/a	n/a
NB	78%	22%	0%	0%
NS	93%	7%	0%	0%
PE	100%	0%	0%	0%
NL	80%	20%	0%	0%
YT	n/a	n/a	n/a	n/a
NT	n/a	n/a	n/a	n/a
NU	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>7%</b>	<b>83%</b>	<b>7%</b>	<b>2%</b>

\* Includes deaths with suspected non-pharmaceutical fentanyl alone or with any other non-pharmaceutical opioid(s)

Only applies to deaths with completed investigations.

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases.

Not available (n/a) Data were not available at the time of this publication.

### Note(s):

Apparent opioid toxicity death (AOTD): A death caused by intoxication/toxicity (poisoning) resulting from substance use, where one or more of the substances is an opioid, regardless of how it was obtained (e.g. illegally or through personal prescription). Other substances may also be involved.

2020 data from Manitoba are based on January to March.

2021 data from Newfoundland and Labrador are based on January to March.

Data on apparent opioid toxicity deaths and stimulant toxicity deaths are not mutually exclusive. A high proportion of deaths involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.



**Table 14a. Number of accidental apparent opioid toxicity deaths by sex and age group, by province or territory, 2018 to 2021 (Jan to Jun)**

**2018**

Province or territory	Sex	Age group					
		0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	Female	4	43	64	44	45	19
	Male	18	167	230	192	205	70
AB	Female	9	51	55	46	36	12
	Male	6	146	181	112	105	40
SK	Female	0	11	12	8	13	3
	Male	0	16	16	15	10	10
MB	Female	n/a	n/a	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a	n/a	n/a
ON	Female	7	77	105	74	66	31
	Male	14	185	277	216	201	66
QC	Female	0	5	9	13	16	12
	Male	1	15	32	20	18	7
NB	Female	0	2	3	1	5	2
	Male	0	0	2	4	3	1
NS	Female	Suppr.	3	5	4	1	2
	Male	Suppr.	8	7	6	8	4
PE	Female	n/a	n/a	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a	n/a	n/a
NL	Female	n/a	n/a	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a	n/a	n/a
YT	Female	n/a	n/a	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a	n/a	n/a
NT	Female	n/a	n/a	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a	n/a	n/a
NU	Female	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Male	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>Female</b>	<b>20</b>	<b>192</b>	<b>253</b>	<b>190</b>	<b>182</b>	<b>81</b>
	<b>Male</b>	<b>39</b>	<b>537</b>	<b>745</b>	<b>565</b>	<b>550</b>	<b>198</b>



## 2019

Province or territory	Sex	Age group					
		0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	Female	9	21	36	22	27	6
	Male	5	69	103	76	70	33
AB	Female	7	40	45	28	33	10
	Male	6	99	145	100	79	25
SK	Female	0	8	7	12	11	1
	Male	1	13	18	14	19	5
MB	Female	1	4	5	2	4	1
	Male	0	11	12	7	6	1
ON	Female	6	76	114	69	100	21
	Male	15	213	296	206	207	85
QC	Female	1	0	6	9	19	11
	Male	2	19	20	31	19	17
NB	Female	0	1	4	1	5	3
	Male	0	1	4	4	1	3
NS	Female	Suppr.	0	4	3	3	3
	Male	Suppr.	7	6	9	5	4
PE	Female	n/a	n/a	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a	n/a	n/a
NL	Female	n/a	n/a	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a	n/a	n/a
YT	Female	n/a	n/a	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a	n/a	n/a
NT	Female	n/a	n/a	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a	n/a	n/a
NU	Female	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Male	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>Female</b>	<b>24</b>	<b>150</b>	<b>221</b>	<b>146</b>	<b>202</b>	<b>56</b>
	<b>Male</b>	<b>29</b>	<b>432</b>	<b>604</b>	<b>447</b>	<b>406</b>	<b>173</b>



## 2020

Province or territory	Sex	Age group					
		0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	Female	5	22	28	32	22	12
	Male	7	97	139	145	126	50
AB	Female	7	59	69	69	50	13
	Male	16	170	259	197	169	66
SK	Female	4	18	20	16	18	8
	Male	3	30	50	35	30	14
MB	Female	0	3	3	3	0	0
	Male	0	1	6	2	1	1
ON	Female	10	103	164	127	95	41
	Male	22	327	510	398	371	135
QC	Female	Suppr.	20	32	29	30	Suppr.
	Male	Suppr.	52	79	93	107	Suppr.
NB	Female	0	3	4	4	3	3
	Male	0	2	2	7	6	4
NS	Female	Suppr.	2	3	5	1	2
	Male	Suppr.	8	6	10	4	1
PE	Female	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Male	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NL	Female	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Male	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
YT	Female	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Male	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NT	Female	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Male	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NU	Female	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Male	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>Female</b>	<b>26</b>	<b>230</b>	<b>323</b>	<b>285</b>	<b>219</b>	<b>79</b>
	<b>Male</b>	<b>48</b>	<b>687</b>	<b>1,051</b>	<b>887</b>	<b>814</b>	<b>271</b>




## 2021 (Jan to Jun)

Province or territory	Sex	Age group					
		0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	Female	0	2	4	1	3	1
	Male	2	13	22	20	19	19
AB	Female	3	46	64	35	17	13
	Male	11	106	165	105	98	42
SK	Female	1	8	11	4	6	1
	Male	0	10	14	17	15	3
MB	Female	n/a	n/a	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a	n/a	n/a
ON	Female	3	72	104	69	66	29
	Male	11	173	288	222	205	83
QC	Female	Suppr.	8	10	15	10	Suppr.
	Male	Suppr.	18	41	37	35	Suppr.
NB	Female	0	0	1	2	1	0
	Male	0	2	1	3	1	2
NS	Female	Suppr.	2	1	2	1	0
	Male	Suppr.	4	4	3	3	1
PE	Female	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Male	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NL	Female	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Male	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
YT	Female	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Male	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NT	Female	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Male	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NU	Female	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Male	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>Female</b>	<b>7</b>	<b>138</b>	<b>195</b>	<b>128</b>	<b>104</b>	<b>44</b>
	<b>Male</b>	<b>24</b>	<b>326</b>	<b>535</b>	<b>407</b>	<b>376</b>	<b>150</b>

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases.

Not available (n/a) Data were not available at the time of this publication.

 Includes deaths related to all illicit drugs including, but not limited to, opioids and stimulants.

### Note(s):

Apparent opioid toxicity death (AOTD): A death caused by intoxication/toxicity (poisoning) resulting from substance use, where one or more of the substances is an opioid, regardless of how it was obtained (e.g. illegally or through personal prescription). Other substances may also be involved.

2020 data from Manitoba are based on January to March.

2021 data from Newfoundland and Labrador are based on January to March.

Data on apparent opioid toxicity deaths and stimulant toxicity deaths are not mutually exclusive. A high proportion of deaths involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.



**Table 14b. Number of accidental apparent stimulant toxicity deaths by sex and age group, by province or territory, 2018 to 2021 (Jan to Jun)**

**2018**

Province or territory	Sex	Age group					
		0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	Female	2	31	57	37	41	13
	Male	10	114	161	156	179	65
SK	Female	1	5	11	9	6	0
	Male	0	11	14	12	4	4
MB	Female	n/a	n/a	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a	n/a	n/a
ON	Female	4	53	71	49	46	11
	Male	8	100	183	164	151	48
QC	Female	0	9	5	10	6	1
	Male	3	16	16	21	37	20
NS	Female	Suppr.	2	2	1	0	0
	Male	Suppr.	6	3	2	5	1
NL	Female	n/a	n/a	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a	n/a	n/a
NT	Female	n/a	n/a	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a	n/a	n/a

**2019**

Province or territory	Sex	Age group					
		0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	Female	5	19	27	21	24	8
	Male	2	52	76	68	66	30
SK	Female	0	10	13	5	4	1
	Male	1	10	21	13	11	4
MB	Female	0	7	7	2	2	0
	Male	1	11	16	19	7	1
ON	Female	3	47	79	55	59	9
	Male	10	125	199	158	164	81
QC	Female	0	7	10	6	7	2
	Male	4	13	15	19	29	18
NS	Female	Suppr.	0	3	1	0	0
	Male	Suppr.	4	4	4	5	5
NL	Female	n/a	n/a	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a	n/a	n/a
NT	Female	n/a	n/a	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a	n/a	n/a



## 2020

Province or territory	Sex	Age group					
		0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	Female	2	20	25	28	18	12
	Male	5	69	113	123	108	42
SK	Female	2	15	16	11	12	2
	Male	2	17	44	26	21	9
MB	Female	1	2	3	5	0	0
	Male	0	2	4	3	3	1
ON	Female	7	64	133	93	70	21
	Male	16	208	351	297	299	104
QC	Female	n/a	n/a	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a	n/a	n/a
NS	Female	Suppr.	2	1	1	0	1
	Male	Suppr.	5	5	6	7	3
NL	Female	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Male	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NT	Female	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Male	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.



## 2021 (Jan to Jun)

Province or territory	Sex	Age group					
		0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	Female	0	2	4	1	4	1
	Male	0	9	13	18	12	17
SK	Female	1	10	8	3	3	0
	Male	0	8	14	10	14	1
MB	Female	n/a	n/a	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a	n/a	n/a
ON	Female	3	47	86	51	44	11
	Male	1	110	181	166	152	56
QC	Female	n/a	n/a	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a	n/a	n/a
NS	Female	Suppr.	3	1	0	1	0
	Male	Suppr.	4	3	2	2	2
NL	Female	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Male	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NT	Female	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Male	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases.

Not available (n/a) Data were not available at the time of this publication.

### Note(s):

Apparent stimulant toxicity death (ASTD): A death caused by intoxication/toxicity (poisoning) resulting from substance use, where one or more of the substances is a stimulant, regardless of how it was obtained (e.g. illegally or through personal prescription). Other substances may also be involved.

2020 data from Manitoba are based on January to March.

2021 data from Newfoundland and Labrador are based on January to March.

Data on apparent opioid toxicity deaths and stimulant toxicity deaths are not mutually exclusive. A high proportion of deaths involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.

**Table 15a. Percentage of accidental apparent opioid toxicity deaths by type of opioids and sex, by province or territory, 2018 to 2021 (Jan to Jun)**

Province or territory	Type of opioids	2018		2019		2020		2021 (Jan to Jun)	
		Male	Female	Male	Female	Male	Female	Male	Female
BC	Fentanyl	80%	20%	75%	25%	83%	17%	89%	11%
	Fentanyl analogues	86%	14%	81%	19%	83%	17%	82%	18%
	Non-fentanyl opioids	76%	24%	67%	33%	76%	24%	82%	18%
AB	Fentanyl	78%	22%	76%	24%	78%	22%	74%	26%
	Fentanyl analogues	78%	22%	75%	25%	81%	19%	78%	22%
	Non-fentanyl opioids	61%	39%	65%	35%	66%	34%	68%	32%
SK	Fentanyl	68%	32%	70%	30%	72%	28%	65%	35%
	Fentanyl analogues	Suppr.	Suppr.	67%	33%	75%	25%	64%	36%
	Non-fentanyl opioids	54%	46%	62%	38%	59%	41%	74%	26%
MB	Fentanyl	n/a	n/a	88%	13%	54%	46%	n/a	n/a
	Fentanyl analogues	n/a	n/a	83%	17%	0%	0%	n/a	n/a
	Non-fentanyl opioids	n/a	n/a	52%	48%	50%	50%	n/a	n/a
ON	Fentanyl	75%	25%	75%	25%	78%	22%	75%	25%
	Fentanyl analogues	76%	24%	77%	23%	80%	20%	69%	31%
	Non-fentanyl opioids	70%	30%	63%	37%	67%	33%	68%	32%
QC	Fentanyl	77%	23%	87%	13%	70%	30%	84%	16%
	Fentanyl analogues	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Non-fentanyl opioids	61%	39%	68%	32%	67%	33%	68%	32%
NB	Fentanyl	Suppr.	Suppr.	80%	20%	75%	25%	50%	50%
	Fentanyl analogues	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	43%	57%	46%	54%	52%	48%	73%	27%
NS	Fentanyl	75%	25%	75%	25%	80%	20%	Suppr.	Suppr.
	Fentanyl analogues	75%	25%	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Non-fentanyl opioids	68%	32%	70%	30%	67%	33%	71%	29%



Province or territory	Type of opioids	2018		2019		2020		2021 (Jan to Jun)	
		Male	Female	Male	Female	Male	Female	Male	Female
PE	Fentanyl	n/a	n/a	n/a	n/a	Suppr.	Suppr.	Suppr.	Suppr.
	Fentanyl analogues	n/a	n/a	n/a	n/a	Suppr.	Suppr.	Suppr.	Suppr.
	Non-fentanyl opioids	n/a	n/a	n/a	n/a	Suppr.	Suppr.	Suppr.	Suppr.
NL	Fentanyl	n/a	n/a	n/a	n/a	Suppr.	Suppr.	Suppr.	Suppr.
	Fentanyl analogues	n/a	n/a	n/a	n/a	Suppr.	Suppr.	Suppr.	Suppr.
	Non-fentanyl opioids	n/a	n/a	n/a	n/a	Suppr.	Suppr.	Suppr.	Suppr.
YT	Fentanyl	n/a	n/a	n/a	n/a	Suppr.	Suppr.	Suppr.	Suppr.
	Fentanyl analogues	n/a	n/a	n/a	n/a	Suppr.	Suppr.	Suppr.	Suppr.
	Non-fentanyl opioids	n/a	n/a	n/a	n/a	Suppr.	Suppr.	Suppr.	Suppr.
NT	Fentanyl	n/a	n/a	n/a	n/a	Suppr.	Suppr.	Suppr.	Suppr.
	Fentanyl analogues	n/a	n/a	n/a	n/a	Suppr.	Suppr.	Suppr.	Suppr.
	Non-fentanyl opioids	n/a	n/a	n/a	n/a	Suppr.	Suppr.	Suppr.	Suppr.
NU	Fentanyl	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Fentanyl analogues	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Non-fentanyl opioids	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
Canada	<b>Fentanyl</b>	<b>77%</b>	<b>23%</b>	<b>76%</b>	<b>24%</b>	<b>79%</b>	<b>21%</b>	<b>76%</b>	<b>24%</b>
	<b>Fentanyl analogues</b>	<b>79%</b>	<b>21%</b>	<b>77%</b>	<b>23%</b>	<b>80%</b>	<b>20%</b>	<b>73%</b>	<b>27%</b>
	<b>Non-fentanyl opioids</b>	<b>68%</b>	<b>32%</b>	<b>64%</b>	<b>36%</b>	<b>67%</b>	<b>33%</b>	<b>70%</b>	<b>30%</b>

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases.

Not available (n/a) Data were not available at the time of this publication.

Includes deaths related to all illicit drugs including, but not limited to, opioids and stimulants.

#### Note(s):

Apparent opioid toxicity death (AOTD): A death caused by intoxication/toxicity (poisoning) resulting from substance use, where one or more of the substances is an opioid, regardless of how it was obtained (e.g. illegally or through personal prescription). Other substances may also be involved.

2020 data from Manitoba are based on January to March.

2021 data from Newfoundland and Labrador are based on January to March.

Data on apparent opioid toxicity deaths and stimulant toxicity deaths are not mutually exclusive. A high proportion of deaths involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.





**Table 15b. Percentage of accidental apparent stimulant toxicity deaths by type of stimulants and sex, by province or territory, 2018 to 2021 (Jan to Jun)**

Province or territory	Type of stimulants	2018		2019		2020		2021 (Jan to Jun)	
		Male	Female	Male	Female	Male	Female	Male	Female
BC	Cocaine	78%	22%	74%	26%	83%	17%	88%	12%
	Methamphetamine	77%	23%	75%	25%	80%	20%	83%	17%
	Other stimulants	88%	12%	71%	29%	79%	21%	75%	25%
SK	Cocaine	67%	33%	83%	17%	68%	32%	73%	27%
	Methamphetamine	52%	48%	58%	42%	67%	33%	67%	33%
	Other stimulants	45%	55%	63%	38%	50%	50%	67%	33%
MB	Cocaine	n/a	n/a	75%	25%	82%	18%	n/a	n/a
	Methamphetamine	n/a	n/a	81%	19%	40%	60%	n/a	n/a
	Other stimulants	n/a	n/a	Suppr.	Suppr.	Suppr.	Suppr.	n/a	n/a
ON	Cocaine	73%	27%	75%	25%	76%	24%	73%	27%
	Methamphetamine	72%	28%	72%	28%	75%	25%	75%	25%
	Other stimulants	77%	23%	60%	40%	66%	34%	68%	32%
QC	Cocaine	81%	19%	75%	25%	n/a	n/a	n/a	n/a
	Methamphetamine	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
	Other stimulants	75%	25%	76%	24%	n/a	n/a	n/a	n/a
NS	Cocaine	71%	29%	89%	11%	81%	19%	71%	29%
	Methamphetamine	67%	33%	89%	11%	0%	0%	50%	50%
	Other stimulants	Suppr.	Suppr.	75%	25%	71%	29%	0%	0%

Province or territory	Type of stimulants	2018		2019		2020		2021 (Jan to Jun)	
		Male	Female	Male	Female	Male	Female	Male	Female
NL	Cocaine	n/a	n/a	n/a	n/a	Suppr.	Suppr.	Suppr.	Suppr.
	Methamphetamine	n/a	n/a	n/a	n/a	Suppr.	Suppr.	Suppr.	Suppr.
	Other stimulants	n/a	n/a	n/a	n/a	Suppr.	Suppr.	Suppr.	Suppr.
NT	Cocaine	n/a	n/a	n/a	n/a	Suppr.	Suppr.	Suppr.	Suppr.
	Methamphetamine	n/a	n/a	n/a	n/a	Suppr.	Suppr.	Suppr.	Suppr.
	Other stimulants	n/a	n/a	n/a	n/a	Suppr.	Suppr.	Suppr.	Suppr.
Canada	<b>Cocaine</b>	<b>76%</b>	<b>24%</b>	<b>75%</b>	<b>25%</b>	<b>78%</b>	<b>22%</b>	<b>74%</b>	<b>26%</b>
	<b>Methamphetamine</b>	<b>74%</b>	<b>26%</b>	<b>72%</b>	<b>28%</b>	<b>75%</b>	<b>25%</b>	<b>75%</b>	<b>25%</b>
	<b>Other stimulants</b>	<b>75%</b>	<b>25%</b>	<b>69%</b>	<b>31%</b>	<b>69%</b>	<b>31%</b>	<b>69%</b>	<b>31%</b>

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases.

Not available (n/a) Data were not available at the time of this publication.

#### Note(s):

Apparent stimulant toxicity death (ASTD): A death caused by intoxication/toxicity (poisoning) resulting from substance use, where one or more of the substances is a stimulant, regardless of how it was obtained (e.g. illegally or through personal prescription). Other substances may also be involved.

2020 data from Manitoba are based on January to March.

2021 data from Newfoundland and Labrador are based on January to March.

Data on apparent opioid toxicity deaths and stimulant toxicity deaths are not mutually exclusive. A high proportion of deaths involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.

**Table 16a. Percentage of accidental apparent opioid toxicity deaths by type of opioids and age group, by province or territory, 2018 to 2021 (Jan to Jun)**

**2018**

Province or territory	Type of opioids	Age group					
		0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	Fentanyl	2%	20%	27%	22%	22%	8%
	Fentanyl analogues	9%	21%	28%	16%	19%	7%
	Non-fentanyl opioids	1%	16%	24%	20%	27%	11%
AB	Fentanyl	2%	27%	33%	20%	14%	4%
	Fentanyl analogues	3%	27%	32%	19%	15%	4%
	Non-fentanyl opioids	1%	15%	22%	21%	29%	12%
SK	Fentanyl	0%	30%	36%	19%	13%	2%
	Fentanyl analogues	0%	40%	60%	0%	0%	0%
	Non-fentanyl opioids	0%	19%	18%	21%	26%	17%
MB	Fentanyl	n/a	n/a	n/a	n/a	n/a	n/a
	Fentanyl analogues	n/a	n/a	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	n/a	n/a	n/a	n/a	n/a	n/a
ON	Fentanyl	1%	22%	33%	22%	19%	4%
	Fentanyl analogues	2%	27%	35%	22%	14%	1%
	Non-fentanyl opioids	2%	16%	25%	23%	23%	12%
QC	Fentanyl	0%	12%	35%	19%	19%	15%
	Fentanyl analogues	0%	33%	33%	0%	33%	0%
	Non-fentanyl opioids	0%	14%	27%	22%	24%	13%
NB	Fentanyl	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Fentanyl analogues	n/a	n/a	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	0%	9%	22%	22%	35%	13%
NS	Fentanyl	13%	25%	25%	25%	13%	0%
	Fentanyl analogues	0%	75%	0%	25%	0%	0%
	Non-fentanyl opioids	0%	21%	26%	21%	19%	13%



Province or territory	Type of opioids	Age group					
		0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
PE	Fentanyl	n/a	n/a	n/a	n/a	n/a	n/a
	Fentanyl analogues	n/a	n/a	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	n/a	n/a	n/a	n/a	n/a	n/a
NL	Fentanyl	n/a	n/a	n/a	n/a	n/a	n/a
	Fentanyl analogues	n/a	n/a	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	n/a	n/a	n/a	n/a	n/a	n/a
YT	Fentanyl	n/a	n/a	n/a	n/a	n/a	n/a
	Fentanyl analogues	n/a	n/a	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	n/a	n/a	n/a	n/a	n/a	n/a
NT	Fentanyl	n/a	n/a	n/a	n/a	n/a	n/a
	Fentanyl analogues	n/a	n/a	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	n/a	n/a	n/a	n/a	n/a	n/a
NU	Fentanyl	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Fentanyl analogues	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Non-fentanyl opioids	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
Canada	<b>Fentanyl</b>	<b>2%</b>	<b>22%</b>	<b>31%</b>	<b>21%</b>	<b>19%</b>	<b>6%</b>
	<b>Fentanyl analogues</b>	<b>3%</b>	<b>27%</b>	<b>32%</b>	<b>19%</b>	<b>15%</b>	<b>3%</b>
	<b>Non-fentanyl opioids</b>	<b>1%</b>	<b>16%</b>	<b>24%</b>	<b>22%</b>	<b>25%</b>	<b>12%</b>



## 2019

Province or territory	Type of opioids	Age group					
		0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	Fentanyl	3%	19%	32%	19%	20%	7%
	Fentanyl analogues	4%	19%	27%	23%	19%	8%
	Non-fentanyl opioids	2%	15%	22%	23%	26%	13%
AB	Fentanyl	2%	25%	34%	20%	15%	3%
	Fentanyl analogues	6%	22%	24%	24%	17%	6%
	Non-fentanyl opioids	2%	15%	24%	23%	26%	10%
SK	Fentanyl	3%	25%	43%	15%	13%	3%
	Fentanyl analogues	0%	33%	42%	8%	17%	0%
	Non-fentanyl opioids	0%	17%	13%	27%	35%	8%
MB	Fentanyl	0%	44%	38%	19%	0%	0%
	Fentanyl analogues	0%	67%	0%	33%	0%	0%
	Non-fentanyl opioids	3%	14%	24%	17%	34%	7%
ON	Fentanyl	1%	25%	32%	20%	17%	5%
	Fentanyl analogues	1%	21%	31%	20%	21%	7%
	Non-fentanyl opioids	2%	15%	23%	19%	30%	12%
QC	Fentanyl	9%	22%	26%	22%	17%	4%
	Fentanyl analogues	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Non-fentanyl opioids	1%	11%	16%	26%	25%	20%
NB	Fentanyl	0%	0%	60%	40%	0%	0%
	Fentanyl analogues	n/a	n/a	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	0%	8%	25%	17%	25%	25%
NS	Fentanyl	0%	50%	0%	25%	25%	0%
	Fentanyl analogues	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Non-fentanyl opioids	2%	14%	23%	26%	19%	16%



Province or territory	Type of opioids	Age group					
		0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
PE	Fentanyl	n/a	n/a	n/a	n/a	n/a	n/a
	Fentanyl analogues	n/a	n/a	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	n/a	n/a	n/a	n/a	n/a	n/a
NL	Fentanyl	n/a	n/a	n/a	n/a	n/a	n/a
	Fentanyl analogues	n/a	n/a	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	n/a	n/a	n/a	n/a	n/a	n/a
YT	Fentanyl	n/a	n/a	n/a	n/a	n/a	n/a
	Fentanyl analogues	n/a	n/a	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	n/a	n/a	n/a	n/a	n/a	n/a
NT	Fentanyl	n/a	n/a	n/a	n/a	n/a	n/a
	Fentanyl analogues	n/a	n/a	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	n/a	n/a	n/a	n/a	n/a	n/a
NU	Fentanyl	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Fentanyl analogues	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Non-fentanyl opioids	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
Canada	<b>Fentanyl</b>	<b>2%</b>	<b>24%</b>	<b>33%</b>	<b>20%</b>	<b>17%</b>	<b>5%</b>
	<b>Fentanyl analogues</b>	<b>2%</b>	<b>22%</b>	<b>30%</b>	<b>20%</b>	<b>20%</b>	<b>7%</b>
	<b>Non-fentanyl opioids</b>	<b>2%</b>	<b>14%</b>	<b>22%</b>	<b>22%</b>	<b>28%</b>	<b>13%</b>



## 2020

Province or territory	Type of opioids	Age group					
		0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	Fentanyl	2%	17%	25%	26%	21%	8%
	Fentanyl analogues	2%	13%	29%	29%	19%	8%
	Non-fentanyl opioids	1%	13%	20%	23%	26%	19%
AB	Fentanyl	2%	19%	32%	24%	18%	6%
	Fentanyl analogues	1%	11%	31%	32%	22%	3%
	Non-fentanyl opioids	3%	15%	24%	22%	25%	12%
SK	Fentanyl	3%	24%	34%	21%	15%	3%
	Fentanyl analogues	3%	21%	37%	20%	15%	4%
	Non-fentanyl opioids	2%	15%	21%	20%	28%	14%
MB	Fentanyl	0%	23%	54%	15%	8%	0%
	Fentanyl analogues	0%	0%	0%	0%	0%	0%
	Non-fentanyl opioids	0%	13%	25%	50%	0%	13%
ON	Fentanyl	1%	19%	30%	23%	19%	6%
	Fentanyl analogues	0%	15%	30%	18%	25%	13%
	Non-fentanyl opioids	1%	14%	25%	23%	25%	14%
QC	Fentanyl	Suppr.	16%	27%	29%	17%	Suppr.
	Fentanyl analogues	Suppr.	Suppr.	Suppr.	63%	Suppr.	Suppr.
	Non-fentanyl opioids	3%	16%	23%	19%	22%	18%
NB	Fentanyl	0%	25%	25%	50%	0%	0%
	Fentanyl analogues	n/a	n/a	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	0%	9%	15%	27%	27%	21%
NS	Fentanyl	0%	30%	40%	0%	10%	20%
	Fentanyl analogues	0%	33%	33%	0%	33%	0%
	Non-fentanyl opioids	0%	22%	17%	42%	14%	6%



Province or territory	Type of opioids	Age group					
		0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
PE	Fentanyl	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Fentanyl analogues	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Non-fentanyl opioids	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NL	Fentanyl	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Fentanyl analogues	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Non-fentanyl opioids	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
YT	Fentanyl	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Fentanyl analogues	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Non-fentanyl opioids	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NT	Fentanyl	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Fentanyl analogues	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Non-fentanyl opioids	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NU	Fentanyl	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Fentanyl analogues	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Non-fentanyl opioids	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
Canada	<b>Fentanyl</b>	<b>2%</b>	<b>19%</b>	<b>30%</b>	<b>24%</b>	<b>19%</b>	<b>6%</b>
	<b>Fentanyl analogues</b>	<b>2%</b>	<b>15%</b>	<b>31%</b>	<b>26%</b>	<b>19%</b>	<b>6%</b>
	<b>Non-fentanyl opioids</b>	<b>1%</b>	<b>14%</b>	<b>23%</b>	<b>22%</b>	<b>24%</b>	<b>15%</b>





## 2021 (Jan to Jun)


Province or territory	Type of opioids	Age group					
		0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	Fentanyl	1%	15%	24%	19%	21%	20%
	Fentanyl analogues	0%	16%	26%	26%	16%	16%
	Non-fentanyl opioids	5%	23%	32%	14%	9%	18%
AB	Fentanyl	1%	22%	33%	18%	17%	8%
	Fentanyl analogues	0%	21%	25%	23%	22%	7%
	Non-fentanyl opioids	5%	20%	30%	17%	9%	20%
SK	Fentanyl	1%	22%	31%	26%	17%	4%
	Fentanyl analogues	1%	22%	34%	27%	13%	1%
	Non-fentanyl opioids	0%	4%	26%	13%	43%	13%
MB	Fentanyl	n/a	n/a	n/a	n/a	n/a	n/a
	Fentanyl analogues	n/a	n/a	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	n/a	n/a	n/a	n/a	n/a	n/a
ON	Fentanyl	1%	19%	30%	22%	20%	7%
	Fentanyl analogues	3%	18%	23%	29%	20%	6%
	Non-fentanyl opioids	1%	10%	19%	25%	31%	14%
QC	Fentanyl	Suppr.	14%	44%	28%	Suppr.	Suppr.
	Fentanyl analogues	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Non-fentanyl opioids	Suppr.	13%	24%	29%	15%	Suppr.
NB	Fentanyl	0%	0%	50%	0%	50%	0%
	Fentanyl analogues	n/a	n/a	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	0%	18%	9%	45%	9%	18%
NS	Fentanyl	50%	50%	0%	0%	0%	0%
	Fentanyl analogues	50%	50%	0%	0%	0%	0%
	Non-fentanyl opioids	5%	24%	24%	24%	19%	5%



Province or territory	Type of opioids	Age group					
		0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
PE	Fentanyl	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Fentanyl analogues	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Non-fentanyl opioids	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NL	Fentanyl	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Fentanyl analogues	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Non-fentanyl opioids	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
YT	Fentanyl	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Fentanyl analogues	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Non-fentanyl opioids	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NT	Fentanyl	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Fentanyl analogues	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Non-fentanyl opioids	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NU	Fentanyl	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Fentanyl analogues	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Non-fentanyl opioids	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>Fentanyl</b>	<b>1%</b>	<b>20%</b>	<b>31%</b>	<b>21%</b>	<b>19%</b>	<b>8%</b>
	<b>Fentanyl analogues</b>	<b>1%</b>	<b>20%</b>	<b>27%</b>	<b>26%</b>	<b>19%</b>	<b>7%</b>
	<b>Non-fentanyl opioids</b>	<b>2%</b>	<b>13%</b>	<b>23%</b>	<b>24%</b>	<b>23%</b>	<b>15%</b>

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases.

Not available (n/a) Data were not available at the time of this publication.

 Includes deaths related to all illicit drugs including, but not limited to, opioids and stimulants.

**Note(s):**

Apparent opioid toxicity death (AOTD): A death caused by intoxication/toxicity (poisoning) resulting from substance use, where one or more of the substances is an opioid, regardless of how it was obtained (e.g. illegally or through personal prescription). Other substances may also be involved.

2020 data from Manitoba are based on January to March.

2021 data from Newfoundland and Labrador are based on January to March.

Data on apparent opioid toxicity deaths and stimulant toxicity deaths are not mutually exclusive. A high proportion of deaths involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.

**Table 16b. Percentage of accidental apparent stimulant toxicity deaths by type of stimulants and age group, by province or territory, 2018 to 2021 (Jan to Jun)**

**2018**

Province or territory	Type of stimulants	Age group					
		0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	Cocaine	0%	15%	25%	22%	27%	10%
	Methamphetamine	1%	18%	28%	23%	22%	8%
	Other stimulants	16%	40%	32%	0%	12%	0%
SK	Cocaine	3%	26%	23%	26%	13%	10%
	Methamphetamine	0%	21%	38%	26%	12%	2%
	Other stimulants	0%	18%	9%	45%	27%	0%
MB	Cocaine	n/a	n/a	n/a	n/a	n/a	n/a
	Methamphetamine	n/a	n/a	n/a	n/a	n/a	n/a
	Other stimulants	n/a	n/a	n/a	n/a	n/a	n/a
ON	Cocaine	2%	16%	28%	24%	24%	7%
	Methamphetamine	1%	20%	32%	26%	17%	4%
	Other stimulants	9%	21%	28%	14%	23%	5%
QC	Cocaine	1%	18%	15%	18%	34%	15%
	Methamphetamine	n/a	n/a	n/a	n/a	n/a	n/a
	Other stimulants	4%	16%	15%	27%	24%	15%
NS	Cocaine	5%	33%	19%	14%	24%	5%
	Methamphetamine	33%	33%	33%	0%	0%	0%
	Other stimulants	0%	50%	50%	0%	0%	0%
NL	Cocaine	n/a	n/a	n/a	n/a	n/a	n/a
	Methamphetamine	n/a	n/a	n/a	n/a	n/a	n/a
	Other stimulants	n/a	n/a	n/a	n/a	n/a	n/a
NT	Cocaine	n/a	n/a	n/a	n/a	n/a	n/a
	Methamphetamine	n/a	n/a	n/a	n/a	n/a	n/a
	Other stimulants	n/a	n/a	n/a	n/a	n/a	n/a
Canada	<b>Cocaine</b>	<b>1%</b>	<b>17%</b>	<b>26%</b>	<b>23%</b>	<b>26%</b>	<b>9%</b>
	<b>Methamphetamine</b>	<b>1%</b>	<b>19%</b>	<b>30%</b>	<b>24%</b>	<b>19%</b>	<b>6%</b>
	<b>Other stimulants</b>	<b>7%</b>	<b>23%</b>	<b>22%</b>	<b>19%</b>	<b>21%</b>	<b>7%</b>



2019

Province or territory	Type of stimulants	Age group					
		0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	Cocaine	1%	19%	25%	20%	24%	12%
	Methamphetamine	2%	17%	27%	24%	23%	7%
	Other stimulants	0%	50%	21%	21%	7%	0%
SK	Cocaine	0%	13%	48%	22%	13%	4%
	Methamphetamine	1%	24%	36%	18%	15%	6%
	Other stimulants	0%	13%	50%	13%	25%	0%
MB	Cocaine	0%	20%	33%	25%	20%	3%
	Methamphetamine	0%	30%	35%	30%	5%	0%
	Other stimulants	25%	0%	50%	25%	0%	0%
ON	Cocaine	1%	16%	28%	21%	24%	10%
	Methamphetamine	2%	20%	31%	22%	20%	6%
	Other stimulants	3%	38%	18%	18%	15%	10%
QC	Cocaine	1%	16%	19%	20%	29%	15%
	Methamphetamine	n/a	n/a	n/a	n/a	n/a	n/a
	Other stimulants	6%	14%	20%	18%	26%	16%
NS	Cocaine	0%	17%	22%	22%	22%	17%
	Methamphetamine	0%	33%	22%	11%	11%	22%
	Other stimulants	25%	0%	25%	50%	0%	0%
NL	Cocaine	n/a	n/a	n/a	n/a	n/a	n/a
	Methamphetamine	n/a	n/a	n/a	n/a	n/a	n/a
	Other stimulants	n/a	n/a	n/a	n/a	n/a	n/a
NT	Cocaine	n/a	n/a	n/a	n/a	n/a	n/a
	Methamphetamine	n/a	n/a	n/a	n/a	n/a	n/a
	Other stimulants	n/a	n/a	n/a	n/a	n/a	n/a
Canada	<b>Cocaine</b>	<b>1%</b>	<b>17%</b>	<b>27%</b>	<b>21%</b>	<b>24%</b>	<b>11%</b>
	<b>Methamphetamine</b>	<b>2%</b>	<b>20%</b>	<b>30%</b>	<b>23%</b>	<b>20%</b>	<b>6%</b>
	<b>Other stimulants</b>	<b>5%</b>	<b>25%</b>	<b>23%</b>	<b>19%</b>	<b>18%</b>	<b>10%</b>



## 2020

Province or territory	Type of stimulants	Age group					
		0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	Cocaine	0%	16%	23%	27%	23%	10%
	Methamphetamine	2%	15%	27%	27%	22%	9%
	Other stimulants	10%	52%	14%	14%	10%	0%
SK	Cocaine	0%	18%	27%	21%	27%	7%
	Methamphetamine	3%	19%	38%	19%	16%	5%
	Other stimulants	0%	33%	17%	17%	33%	0%
MB	Cocaine	9%	9%	27%	27%	18%	9%
	Methamphetamine	0%	20%	40%	33%	7%	0%
	Other stimulants	0%	0%	0%	0%	0%	0%
ON	Cocaine	1%	16%	27%	24%	24%	9%
	Methamphetamine	1%	18%	33%	23%	20%	4%
	Other stimulants	11%	22%	29%	20%	14%	4%
QC	Cocaine	n/a	n/a	n/a	n/a	n/a	n/a
	Methamphetamine	n/a	n/a	n/a	n/a	n/a	n/a
	Other stimulants	n/a	n/a	n/a	n/a	n/a	n/a
NS	Cocaine	0%	22%	19%	22%	22%	15%
	Methamphetamine	0%	0%	0%	0%	0%	0%
	Other stimulants	14%	29%	14%	14%	29%	0%
NL	Cocaine	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Methamphetamine	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Other stimulants	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NT	Cocaine	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Methamphetamine	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Other stimulants	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
Canada	<b>Cocaine</b>	<b>1%</b>	<b>16%</b>	<b>26%</b>	<b>25%</b>	<b>24%</b>	<b>9%</b>
	<b>Methamphetamine</b>	<b>2%</b>	<b>17%</b>	<b>32%</b>	<b>24%</b>	<b>20%</b>	<b>6%</b>
	<b>Other stimulants</b>	<b>10%</b>	<b>31%</b>	<b>24%</b>	<b>18%</b>	<b>15%</b>	<b>3%</b>



## 2021 (Jan to Jun)

Province or territory	Type of stimulants	Age group					
		0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	Cocaine	0%	12%	14%	24%	22%	28%
	Methamphetamine	0%	12%	29%	22%	22%	15%
	Other stimulants	0%	25%	25%	25%	13%	13%
SK	Cocaine	0%	27%	9%	0%	55%	9%
	Methamphetamine	2%	22%	35%	21%	21%	0%
	Other stimulants	0%	33%	0%	67%	0%	0%
MB	Cocaine	n/a	n/a	n/a	n/a	n/a	n/a
	Methamphetamine	n/a	n/a	n/a	n/a	n/a	n/a
	Other stimulants	n/a	n/a	n/a	n/a	n/a	n/a
ON	Cocaine	0%	15%	28%	24%	24%	8%
	Methamphetamine	0%	18%	31%	24%	20%	6%
	Other stimulants	0%	19%	39%	13%	26%	3%
QC	Cocaine	n/a	n/a	n/a	n/a	n/a	n/a
	Methamphetamine	n/a	n/a	n/a	n/a	n/a	n/a
	Other stimulants	n/a	n/a	n/a	n/a	n/a	n/a
NS	Cocaine	0%	41%	24%	12%	12%	12%
	Methamphetamine	0%	0%	0%	0%	0%	0%
	Other stimulants	0%	0%	0%	0%	0%	0%
NL	Cocaine	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Methamphetamine	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Other stimulants	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NT	Cocaine	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Methamphetamine	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
	Other stimulants	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
Canada	<b>Cocaine</b>	<b>0%</b>	<b>16%</b>	<b>26%</b>	<b>23%</b>	<b>24%</b>	<b>10%</b>
	<b>Methamphetamine</b>	<b>0%</b>	<b>18%</b>	<b>31%</b>	<b>24%</b>	<b>20%</b>	<b>6%</b>
	<b>Other stimulants</b>	<b>0%</b>	<b>21%</b>	<b>33%</b>	<b>19%</b>	<b>21%</b>	<b>5%</b>

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases.

Not available (n/a) Data were not available at the time of this publication.

### Note(s):

Apparent stimulant toxicity death (ASTD): A death caused by intoxication/toxicity (poisoning) resulting from substance use, where one or more of the substances is a stimulant, regardless of how it was obtained (e.g. illegally or through personal prescription). Other substances may also be involved.

2020 data from Manitoba are based on January to March.

2021 data from Newfoundland and Labrador are based on January to March.

Data on apparent opioid toxicity deaths and stimulant toxicity deaths are not mutually exclusive. A high proportion of deaths involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.

**Table 17. Percentage of accidental opioid toxicity deaths by type of opioids and origin of opioid(s), by province or territory, 2018 to 2021 (Jan to Jun)**

**2018**

Province or territory	Type of opioids	Origin			
		Pharmaceutical only	Non-pharmaceutical only	Both	Undetermined
BC	Fentanyl	0%	100%	0%	0%
	Fentanyl analogues	0%	100%	0%	0%
	Non-fentanyl opioids	0%	0%	0%	100%
AB	Fentanyl	0%	100%	0%	0%
	Fentanyl analogues	0%	100%	0%	0%
	Non-fentanyl opioids	56%	22%	4%	18%
SK	Fentanyl	13%	87%	0%	0%
	Fentanyl analogues	0%	100%	0%	0%
	Non-fentanyl opioids	87%	9%	0%	4%
MB	Fentanyl	n/a	n/a	n/a	n/a
	Fentanyl analogues	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	n/a	n/a	n/a	n/a
ON	Fentanyl	3%	97%	0%	0%
	Fentanyl analogues	0%	100%	0%	0%
	Non-fentanyl opioids	65%	23%	5%	6%
QC	Fentanyl	n/a	n/a	n/a	n/a
	Fentanyl analogues	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	n/a	n/a	n/a	n/a
NB	Fentanyl	0%	100%	0%	0%
	Fentanyl analogues	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	100%	0%	0%	0%
NS	Fentanyl	14%	86%	0%	0%
	Fentanyl analogues	0%	100%	0%	0%
	Non-fentanyl opioids	94%	2%	4%	0%
PE	Fentanyl	n/a	n/a	n/a	n/a
	Fentanyl analogues	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	n/a	n/a	n/a	n/a
NL	Fentanyl	n/a	n/a	n/a	n/a
	Fentanyl analogues	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	n/a	n/a	n/a	n/a
YT	Fentanyl	n/a	n/a	n/a	n/a
	Fentanyl analogues	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	n/a	n/a	n/a	n/a
NT	Fentanyl	n/a	n/a	n/a	n/a
	Fentanyl analogues	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	n/a	n/a	n/a	n/a
NU	Fentanyl	Suppr.	Suppr.	Suppr.	Suppr.
	Fentanyl analogues	Suppr.	Suppr.	Suppr.	Suppr.
	Non-fentanyl opioids	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>Fentanyl</b>	<b>2%</b>	<b>98%</b>	<b>0%</b>	<b>0%</b>
	<b>Fentanyl analogues</b>	<b>0%</b>	<b>100%</b>	<b>0%</b>	<b>0%</b>
	<b>Non-fentanyl opioids</b>	<b>67%</b>	<b>20%</b>	<b>4%</b>	<b>8%</b>



2019

Province or territory	Type of opioids	Origin			
		Pharmaceutical only	Non-pharmaceutical only	Both	Undetermined
BC	Fentanyl	0%	100%	0%	0%
	Fentanyl analogues	0%	100%	0%	0%
	Non-fentanyl opioids	0%	0%	0%	100%
AB	Fentanyl	0%	100%	0%	0%
	Fentanyl analogues	0%	100%	0%	0%
	Non-fentanyl opioids	53%	24%	2%	21%
SK	Fentanyl	5%	95%	0%	0%
	Fentanyl analogues	0%	100%	0%	0%
	Non-fentanyl opioids	90%	4%	0%	6%
MB	Fentanyl	3%	97%	0%	0%
	Fentanyl analogues	0%	100%	0%	0%
	Non-fentanyl opioids	76%	10%	3%	10%
ON	Fentanyl	3%	97%	0%	0%
	Fentanyl analogues	0%	100%	0%	0%
	Non-fentanyl opioids	74%	16%	4%	6%
QC	Fentanyl	n/a	n/a	n/a	n/a
	Fentanyl analogues	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	n/a	n/a	n/a	n/a
NB	Fentanyl	20%	80%	0%	0%
	Fentanyl analogues	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	92%	4%	0%	4%
NS	Fentanyl	33%	67%	0%	0%
	Fentanyl analogues	0%	100%	0%	0%
	Non-fentanyl opioids	100%	0%	0%	0%
PE	Fentanyl	n/a	n/a	n/a	n/a
	Fentanyl analogues	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	n/a	n/a	n/a	n/a
NL	Fentanyl	n/a	n/a	n/a	n/a
	Fentanyl analogues	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	n/a	n/a	n/a	n/a
YT	Fentanyl	n/a	n/a	n/a	n/a
	Fentanyl analogues	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	n/a	n/a	n/a	n/a
NT	Fentanyl	n/a	n/a	n/a	n/a
	Fentanyl analogues	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	n/a	n/a	n/a	n/a
NU	Fentanyl	Suppr.	Suppr.	Suppr.	Suppr.
	Fentanyl analogues	Suppr.	Suppr.	Suppr.	Suppr.
	Non-fentanyl opioids	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>Fentanyl</b>	<b>2%</b>	<b>98%</b>	<b>0%</b>	<b>0%</b>
	<b>Fentanyl analogues</b>	<b>0%</b>	<b>100%</b>	<b>0%</b>	<b>0%</b>
	<b>Non-fentanyl opioids</b>	<b>73%</b>	<b>15%</b>	<b>3%</b>	<b>9%</b>





## 2020

Province or territory	Type of opioids	Origin			
		Pharmaceutical only	Non-pharmaceutical only	Both	Undetermined
BC	Fentanyl	0%	100%	0%	0%
	Fentanyl analogues	0%	100%	0%	0%
	Non-fentanyl opioids	0%	0%	0%	100%
AB	Fentanyl	0%	100%	0%	0%
	Fentanyl analogues	0%	100%	0%	0%
	Non-fentanyl opioids	47%	19%	4%	31%
SK	Fentanyl	1%	99%	0%	0%
	Fentanyl analogues	0%	100%	0%	0%
	Non-fentanyl opioids	80%	5%	2%	14%
MB	Fentanyl	0%	100%	0%	0%
	Fentanyl analogues	0%	0%	0%	0%
	Non-fentanyl opioids	75%	13%	0%	13%
ON	Fentanyl	1%	99%	0%	0%
	Fentanyl analogues	0%	100%	0%	0%
	Non-fentanyl opioids	78%	12%	4%	6%
QC	Fentanyl	n/a	n/a	n/a	n/a
	Fentanyl analogues	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	n/a	n/a	n/a	n/a
NB	Fentanyl	0%	100%	0%	0%
	Fentanyl analogues	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	91%	0%	0%	9%
NS	Fentanyl	20%	80%	0%	0%
	Fentanyl analogues	0%	100%	0%	0%
	Non-fentanyl opioids	100%	0%	0%	0%
PE	Fentanyl	n/a	n/a	n/a	n/a
	Fentanyl analogues	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	n/a	n/a	n/a	n/a
NL	Fentanyl	25%	75%	0%	0%
	Fentanyl analogues	0%	100%	0%	0%
	Non-fentanyl opioids	100%	0%	0%	0%
YT	Fentanyl	n/a	n/a	n/a	n/a
	Fentanyl analogues	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	n/a	n/a	n/a	n/a
NT	Fentanyl	n/a	n/a	n/a	n/a
	Fentanyl analogues	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	n/a	n/a	n/a	n/a
NU	Fentanyl	Suppr.	Suppr.	Suppr.	Suppr.
	Fentanyl analogues	Suppr.	Suppr.	Suppr.	Suppr.
	Non-fentanyl opioids	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>Fentanyl</b>	<b>1%</b>	<b>99%</b>	<b>0%</b>	<b>0%</b>
	<b>Fentanyl analogues</b>	<b>0%</b>	<b>100%</b>	<b>0%</b>	<b>0%</b>
	<b>Non-fentanyl opioids</b>	<b>73%</b>	<b>11%</b>	<b>3%</b>	<b>12%</b>



## 2021 (Jan to Jun)

Province or territory	Type of opioids	Origin			
		Pharmaceutical only	Non-pharmaceutical only	Both	Undetermined
BC	Fentanyl	0%	100%	0%	0%
	Fentanyl analogues	0%	100%	0%	0%
	Non-fentanyl opioids	0%	0%	0%	100%
AB	Fentanyl	0%	100%	0%	0%
	Fentanyl analogues	0%	100%	0%	0%
	Non-fentanyl opioids	42%	14%	2%	42%
SK	Fentanyl	0%	100%	0%	0%
	Fentanyl analogues	0%	100%	0%	0%
	Non-fentanyl opioids	78%	4%	0%	17%
MB	Fentanyl	n/a	n/a	n/a	n/a
	Fentanyl analogues	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	n/a	n/a	n/a	n/a
ON	Fentanyl	1%	99%	0%	0%
	Fentanyl analogues	0%	100%	0%	0%
	Non-fentanyl opioids	80%	10%	6%	5%
QC	Fentanyl	n/a	n/a	n/a	n/a
	Fentanyl analogues	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	n/a	n/a	n/a	n/a
NB	Fentanyl	0%	100%	0%	0%
	Fentanyl analogues	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	100%	0%	0%	0%
NS	Fentanyl	0%	100%	0%	0%
	Fentanyl analogues	0%	100%	0%	0%
	Non-fentanyl opioids	100%	0%	0%	0%
PE	Fentanyl	0%	0%	0%	0%
	Fentanyl analogues	0%	0%	0%	0%
	Non-fentanyl opioids	100%	0%	0%	0%



Province or territory	Type of opioids	Origin			
		Pharmaceutical only	Non-pharmaceutical only	Both	Undetermined
NL	Fentanyl	0%	100%	0%	0%
	Fentanyl analogues	0%	0%	0%	0%
	Non-fentanyl opioids	100%	0%	0%	0%
YT	Fentanyl	n/a	n/a	n/a	n/a
	Fentanyl analogues	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	n/a	n/a	n/a	n/a
NT	Fentanyl	n/a	n/a	n/a	n/a
	Fentanyl analogues	n/a	n/a	n/a	n/a
	Non-fentanyl opioids	n/a	n/a	n/a	n/a
NU	Fentanyl	Suppr.	Suppr.	Suppr.	Suppr.
	Fentanyl analogues	Suppr.	Suppr.	Suppr.	Suppr.
	Non-fentanyl opioids	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>Fentanyl</b>	<b>0%</b>	<b>100%</b>	<b>0%</b>	<b>0%</b>
	<b>Fentanyl analogues</b>	<b>0%</b>	<b>100%</b>	<b>0%</b>	<b>0%</b>
	<b>Non-fentanyl opioids</b>	<b>74%</b>	<b>9%</b>	<b>4%</b>	<b>13%</b>

Only applies to deaths with completed investigations.

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases.

Not available (n/a) Data were not available at the time of this publication.

**Note(s):**

Apparent opioid toxicity death (AOTD): A death caused by intoxication/toxicity (poisoning) resulting from substance use, where one or more of the substances is an opioid, regardless of how it was obtained (e.g. illegally or through personal prescription). Other substances may also be involved.

2020 data from Manitoba are based on January to March.

2021 data from Newfoundland and Labrador are based on January to March.

Data on apparent opioid toxicity deaths and stimulant toxicity deaths are not mutually exclusive. A high proportion of deaths involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.



**Table 18. Percentage of accidental opioid toxicity deaths by sex and origin of opioid(s), by province or territory, 2018 to 2021 (Jan to Jun)**

**2018**

Province or territory	Sex	Origin			
		Pharmaceutical only	Non-pharmaceutical only	Both	Undetermined
BC	Female	0%	53%	0%	47%
	Male	0%	64%	0%	36%
AB	Female	28%	65%	1%	7%
	Male	11%	83%	1%	4%
SK	Female	77%	17%	4%	2%
	Male	45%	48%	4%	3%
MB	Female	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a
ON	Female	31%	59%	6%	3%
	Male	20%	67%	11%	3%
QC	Female	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a
NB	Female	100%	0%	0%	0%
	Male	90%	0%	10%	0%
NS	Female	81%	6%	13%	0%
	Male	88%	6%	6%	0%
PE	Female	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a
NL	Female	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a
YT	Female	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a
NT	Female	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a
NU	Female	Suppr.	Suppr.	Suppr.	Suppr.
	Male	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>Female</b>	<b>36%</b>	<b>55%</b>	<b>5%</b>	<b>4%</b>
	<b>Male</b>	<b>20%</b>	<b>70%</b>	<b>7%</b>	<b>3%</b>



## 2019

Province or territory	Sex	Origin			
		Pharmaceutical only	Non-pharmaceutical only	Both	Undetermined
BC	Female	1%	56%	0%	43%
	Male	0%	70%	0%	30%
AB	Female	24%	66%	0%	10%
	Male	13%	81%	1%	5%
SK	Female	69%	26%	3%	3%
	Male	55%	33%	6%	6%
MB	Female	71%	24%	0%	6%
	Male	24%	65%	5%	5%
ON	Female	29%	58%	10%	4%
	Male	14%	72%	12%	1%
QC	Female	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a
NB	Female	100%	0%	0%	0%
	Male	62%	23%	8%	8%
NS	Female	100%	0%	0%	0%
	Male	90%	6%	3%	0%
PE	Female	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a
NL	Female	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a
YT	Female	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a
NT	Female	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a
NU	Female	Suppr.	Suppr.	Suppr.	Suppr.
	Male	Suppr.	Suppr.	Suppr.	Suppr.
Canada	Female	<b>34%</b>	<b>54%</b>	<b>6%</b>	<b>5%</b>
	Male	<b>18%</b>	<b>71%</b>	<b>8%</b>	<b>3%</b>



2020

Province or territory	Sex	Origin			
		Pharmaceutical only	Non-pharmaceutical only	Both	Undetermined
BC	Female	0%	67%	0%	33%
	Male	0%	78%	0%	22%
AB	Female	17%	75%	1%	7%
	Male	7%	87%	1%	6%
SK	Female	43%	39%	10%	8%
	Male	26%	59%	9%	6%
MB	Female	33%	56%	0%	11%
	Male	30%	70%	0%	0%
ON	Female	18%	68%	13%	2%
	Male	8%	81%	9%	1%
QC	Female	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a
NB	Female	76%	6%	0%	18%
	Male	89%	11%	0%	0%
NS	Female	100%	0%	0%	0%
	Male	70%	19%	11%	0%
PE	Female	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a
NL	Female	75%	25%	0%	0%
	Male	85%	15%	0%	0%
YT	Female	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a
NT	Female	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a
NU	Female	Suppr.	Suppr.	Suppr.	Suppr.
	Male	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>Female</b>	<b>22%</b>	<b>65%</b>	<b>9%</b>	<b>4%</b>
	<b>Male</b>	<b>10%</b>	<b>80%</b>	<b>7%</b>	<b>3%</b>



## 2021 (Jan to Jun)

Province or territory	Sex	Origin			
		Pharmaceutical only	Non-pharmaceutical only	Both	Undetermined
BC	Female	0%	64%	0%	36%
	Male	0%	81%	0%	19%
AB	Female	9%	87%	0%	3%
	Male	4%	90%	0%	6%
SK	Female	13%	81%	6%	0%
	Male	10%	73%	10%	7%
MB	Female	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a
ON	Female	9%	77%	11%	2%
	Male	4%	84%	11%	0%
QC	Female	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a
NB	Female	75%	25%	0%	0%
	Male	80%	20%	0%	0%
NS	Female	100%	0%	0%	0%
	Male	90%	10%	0%	0%
PE	Female	100%	0%	0%	0%
	Male	100%	0%	0%	0%
NL	Female	100%	0%	0%	0%
	Male	67%	33%	0%	0%
YT	Female	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a
NT	Female	n/a	n/a	n/a	n/a
	Male	n/a	n/a	n/a	n/a
NU	Female	Suppr.	Suppr.	Suppr.	Suppr.
	Male	Suppr.	Suppr.	Suppr.	Suppr.
<b>Canada</b>	<b>Female</b>	<b>12%</b>	<b>79%</b>	<b>7%</b>	<b>2%</b>
	<b>Male</b>	<b>6%</b>	<b>84%</b>	<b>7%</b>	<b>2%</b>

Only applies to deaths with completed investigations.

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases.

Not available (n/a) Data were not available at the time of this publication.

### Note(s):

Apparent opioid toxicity death (AOTD): A death caused by intoxication/toxicity (poisoning) resulting from substance use, where one or more of the substances is an opioid, regardless of how it was obtained (e.g. illegally or through personal prescription). Other substances may also be involved.

2020 data from Manitoba are based on January to March.

2021 data from Newfoundland and Labrador are based on January to March.

Data on apparent opioid toxicity deaths and stimulant toxicity deaths are not mutually exclusive. A high proportion of deaths involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.