

Management Plan for the Eastern Wolf (*Canis lupus lycaon*) in Canada

Eastern Wolf



2021



Government
of Canada

Gouvernement
du Canada

Canada

Recommended citation:

Environment and Climate Change Canada. 2021. Management Plan for the Eastern Wolf (*Canis lupus lycaon*) in Canada, *Species at Risk Act* Management Plan Series, Environment and Climate Change Canada, Ottawa, vii + 57 p.

Official version

The official version of the recovery documents is the one published in PDF. All hyperlinks were valid as of date of publication.

Non-official version

The non-official version of the recovery documents is published in HTML format and all hyperlinks were valid as of date of publication.

For copies of the management plan, or for additional information on species at risk, including the Committee on the Status of Endangered Wildlife in Canada (COSEWIC) Status Reports, residence descriptions, action plans, and other related recovery documents, please visit the [Species at Risk Public Registry](https://www.canada.ca/en/environment-climate-change/services/species-risk-public-registry.html).¹

Cover illustration: © Michael Runtz

Également disponible en français sous le titre
« Plan de gestion pour le loup de l'Est (*Canis lupus lycaon*) au Canada »

© Her Majesty the Queen in Right of Canada, represented by the Minister of Environment and Climate Change, 2021. All rights reserved.

ISBN 978-0-660-36977-8

Catalogue no. En3-5/113-2020E-PDF

Content (excluding the illustrations) may be used without permission, with appropriate credit to the source.

¹ www.canada.ca/en/environment-climate-change/services/species-risk-public-registry.html

MANAGEMENT PLAN FOR THE EASTERN WOLF (*Canis lupus lycaon*) IN CANADA

2021

Under the Accord for the Protection of Species at Risk (1996), the federal, provincial, and territorial governments agreed to work together on legislation, programs, and policies to protect wildlife species at risk throughout Canada.

In the spirit of cooperation of the Accord, the Government of Ontario and the Government of Québec provided comments in the development of the Eastern Wolf Management Plan.

As of June 15, 2016, in Ontario only, the Eastern Wolf was re-assessed as the Algonquin Wolf. The Algonquin Wolf was up-listed as threatened under the Ontario's *Endangered Species Act, 2007* which requires Ontario to develop a recovery Strategy by 2018. As the requirements under Ontario's *Endangered Species Act, 2007* are different than those under the federal Species at Risk Act, for information please refer to the following website (<https://www.ontario.ca/laws/statute/07e06>) or contact the Ontario Ministry of Natural Resources and Forestry at: recovery.planning@ontario.ca

Preface

The federal, provincial, and territorial government signatories under the [Accord for the Protection of Species at Risk \(1996\)](#)² agreed to establish complementary legislation and programs that provide for effective protection of species at risk throughout Canada. Under the *Species at Risk Act* (S.C. 2002, c. 29) (SARA), the federal competent ministers are responsible for the preparation of management plans for listed species of special concern and are required to report on progress within five years after the publication of the final document on the Species at Risk Public Registry.

The Minister of Environment and Climate Change and Minister responsible for the Parks Canada Agency is the competent minister under SARA for the Eastern Wolf and has prepared this management plan, as per section 65 of SARA. To the extent possible, it has been prepared in cooperation with the governments of Ontario and Quebec, as per section 66(1) of SARA.

Success in the conservation of this species depends on the commitment and cooperation of many different constituencies that will be involved in implementing the directions set out in this plan and will not be achieved by Environment and Climate Change Canada and the Parks Canada Agency, or any other jurisdiction alone. All Canadians are invited to join in supporting and implementing this plan for the benefit of the Eastern Wolf and Canadian society as a whole.

Implementation of this management plan is subject to appropriations, priorities, and budgetary constraints of the participating jurisdictions and organizations.

² www.canada.ca/en/environment-climate-change/services/species-risk-act-accord-funding.html#2

Acknowledgements

The development of this management plan was facilitated by Sylvain Giguère and Pierre-André Bernier (Canadian Wildlife Service – Quebec Region) as well as Christina Rohe (Canadian Wildlife Service – Ontario Region). The contributions of Véronique Brondex (Canadian Wildlife Service – National Capital Region), Ken Corcoran (formerly with the Canadian Wildlife Service – Ontario Region), Lesley Dunn, Elizabeth Rezek, Krista Holmes, Andrea Kettle, Elisabeth Shapiro and Lee Voisin (Canadian Wildlife Service – Ontario Region), Luc Bélanger, Sandrine Bureau and Marie-Josée Couture (Canadian Wildlife Service – Quebec Region), Emmanuel Dalpé-Charron and Isabelle Gauthier (Quebec Department of Forests, Wildlife and Parks [MFFP] – Direction générale de la gestion de la faune et des habitats), Édith Cadieux and Jean-François Dumont (MFFP – Direction générale Secteur central), Éric Jaccard (MFFP – Direction générale Secteur métropolitain et sud et gestion des forêts), Yannick Bilodeau and Olivier Cameron-Trudel (MFFP – Direction générale Secteur sud-ouest), Jay Fitzsimmons, Leanne Jennings and Glenn Desy (Ontario Ministry of Natural Resources and Forestry [OMNRF], Species Conservation Policy Branch), Maria de Almeida (OMNRF – Wildlife Policy Section), Jim Saunders and Brian Naylor (OMNRF – Crown Forests and Lands Policy Branch), Brent Patterson (OMNRF – Wildlife Research and Monitoring Section), Peter Bedrossian (Department of National Defence), Denis Masse and Jean-Louis Provencher (Parks Canada Agency), Chief Harry St. Denis (Wolf Lake Anishinabeg Community), Chief Lance Haymond (Kebaowek Anishinabeg community) and Amélie Dastous (Nionwentsio Office) are acknowledged and appreciated.

The first drafts of this management plan were prepared by John Benson (Environmental and Life Sciences Program, Trent University) and Karen Loveless (Montana Department of Fish, Wildlife and Parks) with the assistance of Linda Rutledge (Trent University) and Brent Patterson (Ontario Ministry of Natural Resources and Forestry). The initial contributions of Madeline Austen, Paul Watton and Angela McConnell (Canadian Wildlife Service – Ontario Region), Susan Humphrey, Barbara Slezak and Kari Van Allen (formerly with the Canadian Wildlife Service – Ontario Region), Greg Wilson (Canadian Wildlife Service – Prairie and Northern Region), Paul Johanson, Wendy Dunford and Véronique Brondex (Canadian Wildlife Service – National Capital Region), Denis Masse, Sylvain Paradis and Claude Samson (Parks Canada Agency), Brad Allison, Maria de Almeida, Amelia Argue, Patrick Hubert, Al Hyde, Lauren Krushenske, Donald Lewis, Kathryn Markham, Brian Naylor, Bruce Ranta, Chris Risley and Jim Saunders (Ontario Ministry of Natural Resources and Forestry), Sandy Dobbyn, Paul Gelok, Jennifer Hoare and Ed Morris (Ontario Parks), Michel Hénault, Antoine St-Louis and Nathalie Tessier (Quebec Department of Forests, Wildlife and Parks [MFFP]), Astrid Vik Stronen (University of Montreal) and Hélène Jolicoeur and Mario Villemure (independent) are also acknowledged and appreciated.

Lastly, the contributions of the citizens, non-governmental organizations, Indigenous groups, scientists and various interest groups who provided guidance, comments and

recommendations to improve this management plan during the consultation processes are also acknowledged.

Executive Summary

The Eastern Wolf (*Canis lupus lycaon*) is listed as a species of special concern under Schedule 1 of the *Species at Risk Act* (SARA), which requires the drafting of a management plan. The taxonomic status of the Eastern Wolf has been a subject of debate, and recent progress in genetic research has led to a better understanding of the origins of several species and hybrids of the genus *Canis* in North America. These new genetic analyses indicate that the Eastern Wolf is not a subspecies of the Grey Wolf (*Canis lupus lycaon*). In May 2015, the Eastern Wolf was recognized as the species *Canis* sp. cf. *lycaon* and was designated Threatened by the Committee on the Status of Endangered Wildlife in Canada (COSEWIC 2015). This management plan also considers the Eastern Wolf as a distinct species. However, it is important to note that the scientific name and legal status of the species under SARA remains “*Canis lupus lycaon* - *Special Concern*” until a decision is taken by the Governor in Council to change the species’ scientific name and legal status.

Historically, the Eastern Wolf inhabited the deciduous forests of the eastern United States and southeastern Canada. According to current knowledge, it is believed that the Eastern Wolf is present in certain parts of central Ontario and southern Quebec, but owing to the intensive hybridization of the Eastern Wolf with the Coyote (*Canis latrans*), the Grey Wolf (*Canis lupus*) and hybrids of the genus *Canis*, the limits of its range are difficult to determine. Hence, the distribution of hybrid individuals that have Eastern Wolf genetic material (i.e., “Great Lakes–Boreal Wolf” [*C. lupus* x *C. sp. cf. lycaon*] and “Eastern Coyote” [*C. latrans* x *C. sp. cf. lycaon*]) could extend westward in Canada to certain parts of Manitoba and Saskatchewan and eastward to New Brunswick, as well as to parts of Minnesota, Michigan and Wisconsin in the United States. Recent genetic research suggests that the Eastern Wolf may be closely related to the Red Wolf (*Canis rufus*), an endangered species in the southeastern United States.

Over the course of its life, Eastern Wolf uses various types of habitats. It is generally found in extensive forested areas (e.g., over 100 km²). Eastern Wolf populations are limited by prey availability, and higher densities are observed in areas with high prey abundance, particularly White-tailed Deer (*Odocoileus virginianus*), Moose (*Alces alces*) and Beaver (*Castor canadensis*). Hybridization of the Eastern Wolf with the Grey Wolf is also considered a limiting factor, as is the territoriality of large canid packs, which reduces the potential expansion of the Eastern Wolf population beyond already occupied sites.

The main threats to the Canadian Eastern Wolf population include, but are not limited to hunting, trapping and poaching, roads, and hybridization with the Eastern Coyote. The impact of other threats is considered low to negligible (e.g., residential and commercial development, agriculture and recreational activities) or as yet unknown (e.g., diseases and parasites associated with other animals and changes in habitats and availability of prey induced by climate change).

The management objective is to achieve and maintain a viable Eastern Wolf population within the species' current range in Canada. To achieve this management objective, the density of the Eastern Wolf population in the Algonquin Provincial Park area will, at a minimum, have to be maintained at the current level, which is estimated at approximately 2.1 individuals per 100 km² and, until more detailed basic knowledge is obtained on the other occupied sites within the Canadian area of occupancy, the presence of the species at these sites will have to be maintained. Connectivity between the occupied sites and other regions where suitable habitat is found must also be ensured to facilitate dispersal of individuals and maintain genetic cohesion of the species in Canada. Additional research will be required to improve understanding of the Eastern Wolf's distribution, abundance and effective population size, to identify connectivity requirements, to clarify taxonomic uncertainties and to manage technical challenges associated with the need to identify the species through molecular analyses. When the success of the plan is assessed, the management objective may be re-examined in light of this new information.

Four broad strategies and a number of conservation measures are proposed in order to achieve the objective of this management plan. Although some of them may have a positive or negative effect on other species, it may be possible to reduce or mitigate any negative effects by adopting an ecosystem approach when implementing the species management activities.

Table of Contents

Preface.....	ii
Acknowledgements	iii
Executive Summary	v
1. COSEWIC Species Assessment Information.....	1
2. Species Status Information	2
3. Species information	2
3.1 Species History.....	2
3.2 Species Description	4
3.3 Species Population and Distribution	4
3.4 Needs of the Eastern Wolf.....	9
3.4.1 Limiting Factors	11
4. Threats.....	12
4.1 Threat Assessment.....	12
4.2 Description of Threats	18
4.2.1 Main threats.....	18
4.2.2 Threats of low or negligible impact	21
4.2.3 Potential threats.....	22
5. Management Objective	23
6. Broad Strategies and Conservation Measures	24
6.1 Actions Already Completed or Currently Underway.....	24
6.2 Broad Strategies.....	28
6.3 Conservation Measures.....	28
6.4 Narrative to Support Conservation Measures and Implementation Schedule ..	33
7. Measuring Progress.....	34
8. References	35
Appendix A: Additional Information on the Effects of Trapping on the Eastern Wolf Population in Canada	54
Appendix B: Effects on the Environment and Other Species	55
Appendix C: Season Closures for Hunting and Trapping Wolf and Coyote in Ontario, September 2016.....	57

1. COSEWIC* Species Assessment Information

Date of Assessment: May 2001

Common Name: Eastern Wolf

Scientific Name: *Canis lupus lycaon*

COSEWIC Status: Special concern

Reason for Designation: This wolf may be a separate species. Its exact range is not known, partly because it hybridizes with grey wolves. Although there is no evidence of decline in either number or geographic range over the last 20 years, it may be undergoing hybridization with coyotes, possibly exacerbated by habitat changes and high levels of harvesting. Identification of this taxon requires a molecular analysis.

Canadian Occurrence: Ontario, Quebec

COSEWIC Status History: Designated Data Deficient in April 1999. Status re-examined in May 2001 and designated Special Concern.

Date of Assessment: May 2015 (In a higher risk category)

Common Name: Eastern Wolf

Scientific Name: *Canis sp. cf. lycaon*³

COSEWIC Status: Threatened

Reason for Designation: This species is an intermediate-sized canid with a generally reddish-brown/tawny coat. It has a small population size (likely < 1,000 individuals) and a restricted range, limited to south-central Ontario and south-central Quebec. Most records come from scattered protected areas, where mortality and rates of hybridization with Eastern Coyotes occurs less frequently than elsewhere in its range. Population expansion is unlikely, owing to competition with Eastern Coyote and increased mortality rates outside protected areas.

Canadian Occurrence: Ontario, Quebec

COSEWIC Status History: In 1999, the Eastern Grey Wolf (*Canis lupus lycaon*) was considered a subspecies of the Grey Wolf and was placed in the Data Deficient category. Status was re-examined (as Eastern Wolf, *Canis lupus lycaon*) and designated Special Concern in May 2001. New genetic analyses indicate that the Eastern Wolf is not a subspecies of Grey Wolf. In May 2015, a new wildlife species, Eastern Wolf (*Canis sp. cf. lycaon*) was designated Threatened and the original designation was deactivated.

* COSEWIC (Committee on the Status of Endangered Wildlife in Canada)

³ The use of “sp. cf. *lycaon*” in the scientific name means that the Eastern Wolf is recognized as a distinct species based on the best available data, while taking into account that the current taxonomic debate has not yet been completely resolved. The Species at Risk Public Registry has been updated and uses “*Canis sp. cf. lycaon*” as the scientific name for the Eastern Wolf, reflecting the name change by COSEWIC. However, to accurately reflect the legal name for this species under the *Species at Risk Act* (SARA; S.C. 2002, c. 29) at the time of publication of this management plan, the species’ scientific name remains “*Canis lupus lycaon*” in the title of this document.

2. Species Status Information

The conservation status of the Eastern Wolf is not yet ranked at the global Canadian, Ontario or Quebec level (NatureServe 2019).

The Eastern Wolf (*Canis lupus lycaon*) is listed as a species of special concern⁴ in Schedule 1 of the *Species at Risk Act* (SARA) (S.C. 2002, c. 29). In May 2015, COSEWIC designated the Eastern Wolf as a distinct species and uplisted it to a higher risk category, i.e., threatened. As a result of this reassessment and change in risk category, the species will be eligible for an amendment of its status under Schedule 1 of SARA. In January 2016, the Committee on the Status of Species at Risk in Ontario (COSSARO) reassessed the taxon formerly recognized as “Eastern Wolf (*Canis lupus lycaon*)” as “Algonquin Wolf (*Canis* sp.)” and reassessed it as “Threatened” (COSSARO 2016). In June 2016, the Ontario government listed the Algonquin Wolf (*Canis* sp.) as Threatened⁵ under the Ontario *Endangered Species Act, 2007* (S.O. 2007, c. 6). In Quebec, only the Grey Wolf (*Canis lupus*) is officially recognized by the provincial government and this species is not listed under the Quebec *Act respecting threatened or vulnerable species* (C.Q.L.R., c. E-12.01).

In this document, the Eastern Wolf will be considered a distinct species, in accordance with the most recent COSEWIC status report on the species (COSEWIC 2015). In the status report, the Eastern Wolf is defined by a combination of morphological⁶ characteristics, its ecological role and specific genetic characteristics (COSEWIC 2015).

3. Species information

3.1 Species History

Eastern Wolves have long been recognized as being morphologically distinct from Grey Wolves (Young and Goldman 1944; Kolenosky and Standfield 1975) as they are slightly smaller and have a lower body mass than Grey Wolves. Aboriginal traditional knowledge from the Mohawk First Nation of Akwesasne notes that more than one type of canid was recognized in the region before European contact, based on differences in body size, temperament and size of prey (Lickers, pers. comm. 2015, in COSEWIC 2015). The historical range of the Eastern Wolf likely covered a large part of northeastern North America (COSEWIC 2015; see section 3.3 *Species Population and Distribution*).

⁴ A wildlife species that could become threatened or endangered because of a combination of biological characteristics and identified threats.

⁵ A species that lives in the wild in Ontario, is not endangered, but is likely to become endangered if steps are not taken to address factors threatening it.

⁶ The size, shape and structure of an organism or of one of its parts.

There has been some debate in the literature as to whether the Eastern Wolf is the result of hybridization between the Coyote (*Canis latrans*)⁷ and the Grey Wolf (*Canis lupus*)⁸ either in the past (Lehman et al. 1991; Roy et al. 1994; Wayne and Vila 2003, cited in Kyle et al. 2006) or more recently (vonHoldt et al. 2016), or whether it is an ecotype or population of the Grey Wolf (Koblmüller et al. 2009), or a completely distinct wolf species (Wilson et al. 2000; Kyle et al. 2006; Rutledge et al. 2012). Recent genetic research (e.g., Wilson et al. 2000, 2003, 2009; Rutledge 2010a; Rutledge et al. 2010b, 2010c, 2012; vonHoldt et al. 2011; Benson et al. 2012; Rutledge et al. 2015) and the lack of evidence of hybridization between the Coyote and the Grey Wolf (Mech 2011) strongly suggest that the Eastern Wolf is not a subspecies of the Grey Wolf, nor a hybrid,⁹ but rather a distinct species.¹⁰ In keeping with the COSEWIC status report (2015), the latter classification was used in this management plan (see Appendix 1 of the status report for further details). According to this definition, the Gray Wolf–Eastern Wolf hybrid is called the “Great Lakes–Boreal Wolf” (*C. lupus* x *C. sp. cf. lycaon*), and “Eastern Coyote” (*C. latrans* x *C. sp. cf. lycaon*), which is sometimes also called “coywolf” (Way et al. 2010), is the name used for the Eastern Wolf–Coyote hybrid. The Eastern Coyote is widespread in southern Ontario and Quebec and in the Atlantic provinces (Way 2007; Way et al. 2010; NLDEC 2016).

However, on the basis of analyses that demonstrate a similar genetic signature (Wilson et al. 2000; Chambers et al. 2012; Rutledge et al. 2012), some researchers have also suggested that the Eastern Wolf may be conspecific¹¹ with the Red Wolf (*Canis rufus*), an endangered species that is present in the southeastern United States (Wilson et al. 2000; Grewal et al. 2004; Kyle et al. 2006). It remains possible that the Eastern Wolf and Red Wolf populations have the same historical origin (Grewal et al. 2004; Kyle et al. 2006, 2008; Rutledge et al. 2012; COSEWIC 2015). However, due to introgression¹² of Coyote, Domestic Dog (*Canis lupus familiaris*) or Grey Wolf genes during the Red Wolf captive breeding program (Hailer and Leonard 2008; vonHoldt et al. 2011; Wilson et al. 2012), it is likely that the Red Wolf, as now found in the southeastern United States, no longer meets the criteria used to define the Eastern Wolf (see section 3.3 *Species Population and Distribution*; COSEWIC 2015).

⁷ In this document, as in the COSEWIC status report (2015), coyote (*Canis latrans*) means the small canid native to areas west of the Mississippi River that moved northward and eastward at the time of European settlement of North America. The Coyote is believed to have arrived in Ontario around the early 1900s, in southeastern Ontario in 1919, and in Quebec (Outaouais region) in 1944 (Nowak 1979).

⁸ This theory corresponds to the “2-species hypothesis” (Grey Wolf and Coyote), whose hybridization produced a variety of hybrids, including the Eastern Wolf (COSEWIC 2015, Appendix 1).

⁹ See also Hohenlohe et al. (2017), who disagree with the hypothesis that the Eastern Wolf is the result of recent hybridization between the Grey Wolf and the Coyote.

¹⁰ This theory corresponds to the “3-species hypothesis” (Grey Wolf, Eastern Wolf and Coyote) (COSEWIC 2015, Appendix 1).

¹¹ Belonging to the same species.

¹² Introduction of genes of one species in the gene pool of another species, occurring when matings between the two produce fertile hybrids.

3.2 Species Description

The common names of Eastern Wolf include, but are not limited to, the Eastern Grey Wolf, Eastern Timber Wolf, Canadian Wolf and Algonquin Wolf. The body size and skull characteristics of Eastern Wolves are generally considered to be intermediate between those of the Grey Wolf and the Coyote (Kolenosky and Standfield 1975; Schmitz and Kolenosky 1985; Sears et al. 2003; Theberge and Theberge 2004; Rutledge et al. 2010c, 2010d; Benson et al. 2012). In Algonquin Provincial Park, the average weight of Eastern Wolf adult males was 30.3 kg (standard error of 0.6 kg, n=48) and the average weight of adult females was 23.9 kg (standard error of 0.6 kg, n=40; Theberge and Theberge 2004). In Quebec, Hénault and Jolicoeur (2003) report that the average weight of male wolves, including Eastern Wolves, Grey Wolves and most likely hybrids, ranged from 24.7 kg (Papineau-Labelle Wildlife Reserve) to 44.5 kg (La Mauricie National Park and surrounding area, including the Saint-Maurice Wildlife Reserve, the Mastigouche Wildlife Reserve and the controlled harvesting zone [ZEC] of Chapeau-de-Paille), and the average weight of females ranged from 21.7 kg (on the outskirts of the La Vérendrye Wildlife Reserve) to 28.2 kg (La Mauricie National Park and surrounding area). The coat colour of the Eastern Wolf is quite variable, but is usually described as tawny, with more reddish-brown and brown highlights than the Grey Wolf and the Great Lakes–Boreal Wolf (Young and Goldman 1944; Kolenosky and Standfield 1975). Eastern Wolves observed in the wild are also sometimes confused with Coyotes, which are similar in appearance, although Coyotes are generally considered to have a narrower muzzle, larger ears and proportionally smaller feet relative to their body size (OMNR 2005a). Eastern Coyotes can be difficult to differentiate morphologically from the Eastern Wolf, since their appearance is intermediate and may resemble both the Coyote and the Eastern Wolf (Benson et al. 2012). Given the significant morphological variability in the Eastern Wolf and its similarities with other canids (Rutledge et al. 2010c), it is often very difficult to identify the Eastern Wolf without an assignment test using genetic markers (Rutledge 2010a; COSEWIC 2015).

3.3 Species Population and Distribution

The analysis of the current populations and distribution of the Eastern Wolf is complicated by the confusion caused by its ability to hybridize with other species of the genus *Canis* (see section 3.1 *Species History*). Given the varying degrees of hybridization of Eastern Wolves with other canids, the recognition of individuals as members of the species *Canis* sp. cf. *lycaon* is based, according to the COSEWIC status report (2015), on the probability of genetic assignment to this group ($Q \geq 0.8$; Rutledge et al. 2010c; COSEWIC 2015). This threshold establishes that individuals for which the probability of genetic assignment is less than 0.8 are considered canid hybrids.¹³

¹³ Eastern Wolf in Quebec was identified by two similar methods. One of the methods is the same as that used in Algonquin Provincial Park (Rutledge and White 2013, 2014), whereas the analyses by Stronen et al. (2012), Rogic et al. (2014) and Tessier (unpub. data) used a small proportion of samples from the

The Eastern Wolf exploits a relatively narrow ecological niche, which differs from that of the Grey Wolf and the Eastern Coyote (COSEWIC 2015). Unlike the Eastern Coyote, it is found in minimally human-impacted regions and needs larger prey (e.g., White-tailed Deer [*Odocoileus virginianus*], Beaver [*Castor canadensis*]) to meet its energy requirements (Rutledge et al. 2010c). The Eastern Coyote has thus not replaced the Eastern Wolf in its ecological niche (Benson et al. 2017). In addition, the Eastern Wolf occurs more often in mixedwood forest regions, while the Grey Wolf is generally found in boreal forests, where there is a higher density of Moose (COSEWIC 2015). See section 3.4 *Needs of the Eastern Wolf* for further details on the ecological niche of the Eastern Wolf.

Although its historical range likely covered part of the United States (Wilson et al. 2000; Kyle et al. 2006; Rutledge et al. 2010d; COSEWIC 2015), the known range of the Eastern Wolf, as described here, is currently limited to Canada, namely Ontario and Quebec (COSEWIC 2015). Indeed, assuming that the Eastern Wolf and the Red Wolf are conspecific or were once conspecific, it is believed that the historical range of the Eastern Wolf included the entire temperate forest region of the eastern United States and southeastern Canada (Wilson et al. 2000; Rutledge et al. 2010d). However, it is also possible that the species' historical range was more limited and further to the north and did not include the range of the Red Wolf (Nowak 1995). The historical Canadian range is estimated at approximately 100,000 to 500,000 km² (COSEWIC 2015). In the early 19th century, logging, agricultural practices and other human activities resulted in large-scale landscape changes as well as an intensification of hunting and trapping of top-level predators.¹⁴ It is approximately at this time that the range of Eastern Wolves and their main prey is believed to have shifted to include areas further north, i.e., in Ontario and Quebec (Hall and Kelson 1959; Kyle et al. 2006; Wilson et al. 2009). The Eastern Coyote, a species tolerant to human disturbances, likely replaced the Eastern Wolf in certain parts of southern Canada over the last century, including southern Ontario and southern Quebec.

In the United States, data on distribution are incomplete or have not been examined for this taxon (NatureServe 2017). It is possible that the species' range may extend from eastern Minnesota to Wisconsin and Michigan, but the wolves containing Eastern Wolf genetic material found in these regions appear to be mainly Great Lakes–Boreal Wolves (Wheeldon and White 2009; Mech 2010a; Thiel and Wydeven 2011).

Algonquin Park area and, according to COSEWIC (2015), included as Eastern Wolves samples that would likely have been identified as admixed animals under the Rutledge and White method (2013; 2014).

¹⁴ Also known as apex predators or super predators, these organisms generally have no predators apart from other super predators, reside at the top of the food chain and are believed to have the capacity to self-regulate their population density (Wallach et al. 2015). These organisms often play an essential role in regulating their ecosystem (Estes et al. 2011; Ripple et al. 2014).

The extent of occurrence¹⁵ is estimated at 154,257 km² (based on data shown in Figure 1), while the area of occupancy¹⁶ is estimated at 31,821 km² (COSEWIC 2015; Tessier, unpub. data¹⁷). In Canada, the current distribution of the Eastern Wolf is thought to be restricted to the mixed forest zone of central Ontario and southwestern Quebec, i.e., the Great Lakes and St. Lawrence Forest Region (COSEWIC 2015). In Ontario, the Eastern Wolf is found in Algonquin Provincial Park and the surrounding townships, as well as in Killarney Provincial Park and near French River Provincial Park, in Queen Elizabeth II Wildlands Provincial Park and in Kawartha Highlands Provincial Park (Benson et al. 2012; COSEWIC 2015; Rutledge et al. 2016). Its distribution in Quebec includes the Papineau-Labelle Wildlife Reserve and surrounding area, Maganasipi ZEC, the Mont-Tremblant National Park and Rouge-Matawin Wildlife Reserve area, the Laurentides Wildlife Reserve and La Mauricie National Park and surrounding area (including in particular the Saint-Maurice Wildlife Reserve and the Mastigouche Wildlife Reserve; Potvin 1987; Villemure and Festa-Bianchet 2002; Villemure 2003; Villemure and Masse 2004; Rogic et al. 2014; Mainguy et al. 2017; Hénault 2019; Tessier, unpub. data). Outside the current extent of occurrence, research including samples of wolves from Duck Mountain Provincial Park and Riding Mountain National Park in Manitoba have established the presence of Eastern Wolf genetic material in that province (Wilson et al. 2000; Stronen et al. 2010; Rutledge et al. 2010b). Eastern Wolf genetic material has also been detected in wolves living as far west as Manitoba and Saskatchewan (Stronen et al. 2010; Stronen, pers. comm. 2011). However, it appears that most of the wolves in Manitoba that have *Canis* sp. cf. *lycaon* genetic material are Great Lakes–Boreal Wolves (Wheeldon 2009). Lastly, until recently, no microsatellite profiles characteristic of the Eastern Wolf had been found in canids in the Maritimes (Way et al. 2010; COSEWIC 2015), but Eastern Wolf genetic material has since been detected in an individual (*C. lupus* x *C. sp. cf. lycaon*) captured in New Brunswick (McAlpine et al. 2015).

¹⁵ The area included in a convex polygon encompassing the geographic distribution of all known populations of a species.

¹⁶ The index calculates the area within the “extent of occurrence” that is potentially occupied by the species. This index is based on the size of sites where the Eastern Wolf has been recorded. See the COSEWIC status report (2015) for more details.

¹⁷ The difference between the area presented here and the area shown in Table 2 of the COSEWIC status report (COSEWIC 2015) corresponds to the areas of the Mastigouche Wildlife Reserve (1,565 km²) and the Saint-Maurice Wildlife Reserve (784 km²), adjacent to La Mauricie National Park, where Eastern Wolves have been identified by genetic analysis (N. Tessier, unpub. data).




Figure 1. Distribution of the Eastern Wolf. The locations shown were taken from various sources, some of which use a methodology common to several publications¹⁸ (stars: data from COSEWIC [2015, orange stars], Mainguy et al. [2017, blue stars], ONHIC [2017, red stars], Hénault [2019, yellow stars] and Wolf Lake Anishinabeg Community and R. Van Schie [unpub. data, purple stars]), or other methods (circles: Rogic et al. 2014; Tessier, unpub. data). The number within the circles indicates the number of individuals identified, and the position of the circle indicates the approximate site (and not necessarily the exact location) where the individuals were found. The areas shown in dark green correspond to protected areas (MDDELCC 2017; OMNRF 2018a)¹⁹, the areas shown in light yellow are Quebec wildlife reserves, and the areas shown in light grey are ZECs. It should be noted that the Eastern Wolf locations shown here are not exhaustive, but reflect the best available information.

¹⁸ Including Rutledge (2010a), Rutledge et al. (2010b, 2010c), Rutledge and White (2013, 2014), WLFN (2018) and Hénault (2019).

¹⁹ The protected areas represented here include territories under various management regimes. There are several different types of protected areas in Ontario (e.g., Ontario provincial parks, conservation reserves, wilderness areas; OMNRF 2018b) and Quebec (e.g., Quebec provincial parks, wildlife refuges, wildlife habitats; MDDELCC, 2017), as well as federal protected areas (e.g., national parks of Canada, national wildlife areas). Consult the Ontario (OMNRF 2018b) and Quebec (MDDELCC 2017) government registries for a complete list of the types of protected areas in these provinces.

To estimate the minimum number of Eastern Wolves within the extent of occurrence, a method similar to that used in the COSEWIC status report (2015) was employed. For the purpose of this calculation, certain sites where the species has been detected (e.g. protected areas according to the governments of Ontario [OMNRF 2018b] and Quebec [MDDELCC, 2017], wildlife reserves and areas where hunting and trapping of wolves and Eastern coyotes are prohibited) were considered to support the Eastern Wolf. At this point, only these sites are taken into consideration in the calculation, as few records of Eastern Wolves come from outside of them, and those records generally correspond to single individuals (COSEWIC 2015). The estimate was obtained by assessing the abundance of wolves using the estimated density of wolves,²⁰ to which a correction was applied based on the approximate proportion of wolves that are Eastern Wolves.²¹ The proportion of mature individuals (i.e., two years and older) within the population was estimated at 46% (Patterson, unpub. data cited in COSEWIC 2015). In order to estimate the maximum number of Eastern Wolves within the extent of occurrence, the total area of the extent of occurrence (154,257 km²) was multiplied by the estimated density of wolves in Algonquin Provincial Park (3.0 individuals/100 km² x 69% = 2.07 individuals/100 km²).

It was thus estimated that the number of Eastern Wolves within the extent of occurrence was between 450 and 3,193 individuals, 205 to 1,466 of that number being mature individuals.²² The estimation of the upper value of the range was based on the assumption that Eastern Wolf density throughout the extent of occurrence is equivalent to that observed in Algonquin Provincial Park, which is unlikely, considering that the

²⁰ The estimated densities of wolves in Algonquin Provincial Park (3.0 individuals/100 km²; Rutledge et al. 2010e) were used for the sites located in Ontario, and the respective estimated densities in the Papineau-Labelle Wildlife Reserve (2.6 individuals/100 km²; Potvin 1986, 1987), in the Rouge-Matawin Wildlife Reserve and Mont-Tremblant National Park (1.6 individuals/100 km²; Hénault and Jolicoeur 2003), in La Mauricie National Park, in the Saint-Maurice Wildlife Reserve and the Mastigouche Wildlife Reserve (1.3 individuals/100 km²; Villemure 2003), and in the Laurentides Wildlife Reserve (0.44 individuals/100 km²; Jolicoeur 1998) were used for the sites in Quebec. This density was multiplied by the respective area of the sites, the largest of which include the Laurentides Wildlife Reserve (7,861 km²), Algonquin Provincial Park (7,571 km²) and the surrounding townships (6,340 km²), the Rouge-Matawin Wildlife Reserve and Mont-Tremblant National Park (3,165 km²), the La Mauricie National Park, Saint-Maurice Wildlife Reserve and Mastigouche Wildlife Reserve region (2,885 km²), and the Papineau-Labelle Wildlife Reserve (1,628 km²).

²¹ A correction was applied to the values obtained by multiplying them by an estimate of the proportion of wolves assigned to the Eastern Wolf species (69%; data from Rutledge et al. 2010c, in Algonquin Provincial Park; COSEWIC 2015). While this is the best estimate available, the use of this value likely leads to an overestimation of the number of Eastern Wolves at some sites (e.g., possibly at several sites in Quebec, including the Papineau-Labelle wildlife reserve [Hénault, 2019]).

²² The estimates mentioned in the COSEWIC status report (COSEWIC 2015) are from 516 to 2,620 individuals, of which 236 to 1,203 are mature individuals. The differences between the estimates of minimum numbers of individuals provided here and those indicated in Table 2 of the COSEWIC status report (COSEWIC 2015) are attributable in part to the fact that more specific densities are used here for the various sectors of Quebec (as previously mentioned), and in part to the inclusion of the areas of the Mastigouche Wildlife Reserve and of the Saint-Maurice Wildlife Reserve, adjacent to La Mauricie National Park, where Eastern Wolves have been identified by genetic analysis (N. Tessier, unpub. data). The maximum numbers differ slightly from COSEWIC (2015) estimates, due to the recalculation of the extent of occurrence using the data shown in Figure 1.

extent of occurrence is highly sensitive to the addition of peripheral data. In addition, given the threats to the Eastern Wolf outside protected areas and outside areas where hunting and trapping of wolves and Eastern coyotes are prohibited, the number of mature individuals is probably closer to the lower value of the range and is very likely below 1000 (COSEWIC 2015). These estimates may be refined as knowledge gaps are addressed and additional information is available.

The effective population size²³ of the Eastern Wolf population in and around Algonquin Provincial Park has been estimated at 24 to 122 (depending on the approach used), with a harmonic mean of 46 (Rutledge et al. 2016).

3.4 Needs of the Eastern Wolf

Wolves²⁴ live and hunt in social groups called “packs.” Although packs of 12 to 14 wolves have occasionally been reported (Theberge and Theberge 1998), a pack usually consists of 2 to 10 wolves (Pimlott et al. 1969; Potvin 1987; Villemure 2003; Loveless 2010).

Wolves use scent marking (i.e., the smell of feces, urine, anal glands) to demarcate territory and minimize encounters with neighbouring packs, although they do defend their territory if necessary.

Eastern Wolves use various types of habitats, but are generally found in extensive forested areas (COSEWIC 2015). Certain wolf habitat preferences may be explained by an association with prey (Mladenoff et al. 1997; McLoughlin et al. 2004; Desy 2007; Loveless 2010), but wolves may select their habitat with a view to facilitating their movements, increasing their hunting success, and avoiding intraspecific conflicts²⁵ as well as areas associated with human presence or busy roads. Estimated average territory sizes are 199 km² (85 to 324 km²) in the Papineau-Labelle Wildlife Reserve in Quebec (Potvin 1987), 645 km² (623 to 659 km²) in La Mauricie National Park and surrounding area in Quebec (Villemure 2003; data including hybrids), and 190 km² (49 to 330 km²) in Algonquin Provincial Park in Ontario (Loveless 2010).

Breeding females in Eastern Wolf packs produce litters of up to seven pups annually if resources permit, with an average of 4.6 (Mills et al. 2008). In the spring, the females give birth in dens, where the pups will also be raised (Jolicœur et al. 1998; Norris et al.

²³ The effective population size is a theoretical value, corresponding to the number of individuals in an ideal population, i.e., a population in which all individuals reproduce equally (Fisher 1930; Wright 1931), for which there is believed to be a rate of inbreeding or rate of genetic drift equivalent to that of the actual population. This value is used to estimate the risk of genetic decline in a population. The effective population size required for a population to retain its evolutionary potential sets a lower limit to viable population size for wildlife (Soulé 1987 cited in Franklin and Frankham 1998).

²⁴ Since very little information specific to Eastern Wolves exists at present and since the data for certain areas may relate to both Eastern Wolves and hybrids, some of the information on “wolves” is given merely as an example, except where the “Eastern Wolf” is specified.

²⁵ Occurring between members of the same species.

2002; Trapp 2004). Dens can include underground tunnels, generally excavated in well-drained cohesive soils (e.g., sandy soils, that may include silt or gravel), as well as crevasses, rock caves, spaces under tree roots, hollow logs or stumps, or abandoned beaver lodges (Joslin 1967; Pimlott 1967; Jolicoeur et al. 1998; Norris et al. 2002; Patterson, pers. comm. 2009; Benson et al. 2015). Dens are often selected near wetlands or water bodies, likely because of the importance of water to lactating females, which seek to avoid leaving their pups unattended (Mills et al. 2008; Benson et al. 2015). The pack uses one or more dens during the first 4 to 15 weeks of the pups' life (Jolicoeur et al. 1998; Benson et al. 2015). In Algonquin Provincial Park, Eastern Wolves commonly establish their den sites in conifer forests (Benson et al. 2015), including pine (*Pinus* spp.) stands (Norris et al. 2002), possibly owing to the sandy soils associated with these stands (Mech 1970; Ballard and Dau 1983). However, the study by Benson et al. (2015) shows that Eastern Wolves exhibited only marginal selection for conifer stands (33% of dens) in Algonquin Provincial Park. A description of the dens used by wolf packs in the Laurentides Wildlife Reserve and surrounding area also indicates that dens may be surrounded by several types of stands, including black spruce stands, balsam fir stands and scrubland (Jolicoeur et al. 1998). The dens are sometimes reused from year to year by the pack (Jolicoeur et al. 1998; Benson et al. 2015), but the rate of reuse in Algonquin Provincial Park is low (Benson et al. 2015), where the availability of dens is not believed to be a limiting factor (Norris et al. 2002; Argue et al. 2008).

During the rest of the summer, the pups spend most of their time in a series of rendezvous sites, which become the focal point of pack activities (Joslin 1967; Argue et al. 2008). Rendezvous sites are usually located near a permanent water source, often on the edges of lakes, ponds, streams or peatlands, and may be found in various open and forested habitats (Joslin 1967; Jolicoeur et al. 1998). In Algonquin Provincial Park, primarily conifer stands are selected (Benson et al. 2015). The location of rendezvous sites appears to depend on various factors, including kill sites and type of prey (Theberge and Theberge 2004; Benson et al. 2015). Between May and August, wolves that are not in the den or at a rendezvous site appear to be generally found near a permanent water source, such as a peatland, pond, lake or stream (on average at $91 \text{ m} \pm 27 \text{ m}$ from the water), possibly related to predation of beavers or ungulates coming to drink there (Theberge and Theberge 2004). Pup dispersal is a normal part of the species' life cycle and can occur as early as age 15 weeks (Mills et al. 2008). See section 3.4.1 *Limiting factors* regarding the establishment of territories by dispersing individuals.

The Eastern Wolf feeds on White-tailed Deer, Moose (*Alces alces*) and Beaver, as well as a wide variety of smaller animals (Forbes and Theberge 1996b; Potvin et al. 1988; Villemure 2003; OMNR 2005a; Kittle et al. 2007; Loveless 2010; Benson et al. 2017). Like other canids, the Eastern Wolf can cache food for later use. Moldowan and Kitching (2017) observed an Eastern Wolf (or possibly a hybrid) caching part of a White-tailed Deer calf carcass in a sphagnum bog in Algonquin Provincial Park, whose microhabitat could help preserve the hidden food.

According to COSEWIC (2015), yearlings and adults have a life expectancy of 6.2 years and pups have a life expectancy of 0.7 to 3.5 years.

3.4.1 Limiting Factors

Eastern Wolf populations are limited by prey availability: wolf densities are generally higher in areas with a higher density of prey (Messier and Crête 1985; Fuller et al. 2003). For example, the decline in the abundance of wolves in Algonquin Provincial Park between the early 1960s and the 1990s was attributed in part to a change in prey populations, specifically a lower number of deer in the park during the winter and summer months and a decrease in the number of beavers (Algonquin Wolf Advisory Group 2000; Quinn 2005). Wolves living in areas with a low prey density also have lower survival and reproduction rates (Messier 1985, 1987). Although wolves are considered generalist predators, whose main types of prey are determined by availability, accessibility and profitability²⁶ (Peterson and Ciucci 2003), the Eastern Wolf is generally found less frequently in areas where the White-tailed Deer is absent. Although predator-prey dynamics are important, they are not necessarily sufficient to explain the variations in Eastern Wolf abundance, particularly in the presence of human-caused mortality (Theberge and Theberge 2004).

Furthermore, Benson and Patterson (2013) have shown that packs of large canid species (Eastern Wolves, Gray Wolves and Eastern Coyotes) are territorial and that this reduces the possibility of Eastern Wolf population expansion outside of already occupied sites. It is therefore not very likely, for example, that dispersing Eastern Wolves will become established in a territory already occupied by a pack of Eastern Coyotes. They are more likely to join other canid packs and possibly hybridize with Eastern Coyotes (see *Introduced genetic material (IUCN Threat 8.3) – hybridization of the Eastern Wolf with the Eastern Coyote*).

Hybridization may in some cases be considered a natural evolutionary process, but it may also be exacerbated by human activity (Allendorf et al. 2001). For this reason, hybridization of the Eastern Wolf with the Grey Wolf is considered a limiting factor rather than a threat, since the two species were already evolving in sympatry²⁷ prior to European settlement of North America (Nowak 1995) and their hybridization does not appear to be intensified by human activity. Therefore, a natural hybridization zone exists between the areas occupied by the Eastern Wolf and those occupied by the Grey Wolf, where Great Lakes–Boreal Wolves occur, whose stability is ensured by the fact that the Grey Wolf and the Eastern Wolf have different habitat and prey needs (Kolenosky and Standfield 1975; Geffen et al. 2004; Wheeldon 2009). In a stable system such as that, hybridization can even benefit Eastern Wolves by increasing genetic diversity and thus the species' ability to respond to environmental change (such as climate change) (Hamilton and Miller 2015; Jackiw et al. 2015). However, hybridization of the Eastern Wolf with the Eastern Coyote is considered a threat because it is exacerbated by

²⁶ Profitability corresponds to the ratio between the energy obtained and the time required to catch a prey.

²⁷ Whose area of occupancy overlaps.

human activity (see *Introduced genetic material (IUCN Threat 8.3) – hybridization of the Eastern Wolf with the Coyote*).

Competition with Grey Wolves or Great Lakes–Boreal Wolves is likely another limiting factor for Eastern Wolf populations. However, there is little documented evidence of this.

Finally, it is also possible that intraspecific aggression is more frequent when Eastern Wolf densities increase (Rutledge et al. 2010e), and this may be a limiting factor intrinsically related to population density, as has been demonstrated for the Grey Wolf in Yellowstone Park (Cubaynes et al. 2014).

4. Threats

The threats to the Eastern Wolf may vary regionally and locally throughout its range in Canada. The information presented in Table 1 is an overall assessment of the threats to the species across Canada. Where information is known on the significance of a given threat at the regional or local scale, additional information is provided in the threats description section.

4.1 Threat Assessment

The threat assessment for the Eastern Wolf is based on the IUCN-CMP (World Conservation Union–Conservation Measures Partnership) unified threats classification system (Table 1). Threats are defined as the proximate activities or processes that have caused, are causing, or may cause in the future the destruction, degradation, and/or impairment of the entity being assessed (population, species, community, or ecosystem; in this case, the Eastern Wolf) in the area of interest (global, national, or subnational; in this case Canada). Limiting factors are not considered during this threat assessment process. For purposes of threat assessment, only present and future threats are considered. Historical threats, indirect or cumulative effects of the threats, or any other relevant information that would help understand the nature of the threats are presented in section 4.2 *Description of Threats*.

The predominant threats to the Eastern Wolf are hunting, trapping and poaching, as well as roads and hybridization with the Eastern Coyote.

Table 1. Threat Classification Table for the Eastern Wolf

Threat #	Threat Description	Impact^a	Scope^b	Severity^c	Timing^d	Detailed threats
1	Residential and commercial development	Low	Small	Extreme	High	Indirect threats (e.g., increased access and hunting and trapping pressure and increased hybridization) are dealt with under these other threats. It was assumed that new urban developments are causing the most severe impact; however, Eastern Wolves can use cottage or housing areas that remain relatively undeveloped.
1.1	Housing and urban areas	Low	Small	Extreme	High	–
1.2	Commercial and industrial areas	Negligible	Negligible	Extreme	High	–
1.3	Tourism and recreation areas	Low	Small	Serious-Moderate	High	–
2	Agriculture and aquaculture	Negligible	Negligible	Serious-Slight	Moderate	The scope of the threat posed by agriculture was assessed as negligible considering that a large proportion of the current area of occupancy of the Eastern Wolf is located in areas where this type of activity is absent or strictly regulated.
2.1	Annual and perennial non-timber crops	Negligible	Negligible	Serious-Slight	Moderate	Given the uncertainty concerning use by the Eastern Wolf of habitat outside protected areas, severity was defined by means of a range of values (Serious-Slight).
2.2	Wood and pulp plantations	Negligible	Negligible	Negligible	High	It seems that plantations do not pose a threat in Ontario, but will constitute a threat in Quebec.
2.3	Livestock farming and ranching	Negligible	Negligible	Unknown	High	Livestock farming may increase in Ontario. There is much uncertainty in this regard in Quebec. The threat assessed here corresponds to the direct effects of habitat alteration. The aspect of the threat associated with canids killed “in defence of property” is addressed under IUCN threat 5.1 - <i>hunting, trapping and poaching</i>).

Threat #	Threat Description	Impact^a	Scope^b	Severity^c	Timing^d	Detailed threats
3	Energy production and mining	Negligible	Negligible	Extreme	High	The scope of the threat posed by drilling and mining and quarrying was assessed as negligible considering that a large proportion of the current area of occupancy of the Eastern Wolf is located in areas where these activities are not permitted, and moreover it appears that few projects of this type will take place in the near future.
3.1	Oil and gas drilling	Negligible	Negligible	Negligible	Negligible	–
3.2	Mining and quarrying	Negligible	Negligible	Extreme	High	The scope may be greater in Ontario, near the lower limit of “small.”
4	Transportation and service corridors	Medium-Low	Pervasive	Moderate-Slight	High	–
4.1	Roads and railroads	Medium-Low	Pervasive	Moderate-Slight	High	In the assessment of this threat, mortality associated with roads was taken into account, as well as habitat loss. Indirect threats (e.g., increased access for hunting and trapping and increased hybridization) are dealt with under these other threats. The scope of the threat posed by roads was assessed as pervasive, since all types of roads were considered (including logging roads). With regard to severity, “moderate-slight” was used to denote uncertainty. Mortalities have been signaled in Quebec and in Ontario.
4.2	Utility and service lines	Negligible	Small	Negligible	High	The assessment of the scope of this threat (including for example power transmission lines and pipelines) includes only the new structures that might be built and associated rights-of-way. The scope is considered to be lower than 1% in Ontario and nearly 1% in Quebec.

Threat #	Threat Description	Impact^a	Scope^b	Severity^c	Timing^d	Detailed threats
5	Biological resource use	High-Medium	Large	Serious-Moderate	High	–
5.1	Hunting and collecting terrestrial animals	High-Medium	Large	Serious-Moderate	High	<p>In the assessment of this threat, trapping, hunting and poaching were taken into account, as well as incidental take. The scope “large” rather than “pervasive” is explained by the fact that a large proportion of the packs have territories located largely in areas where hunting and trapping are prohibited.</p> <p>For severity, the range “serious-moderate” was used to denote uncertainty.</p> <p>Ontario: when wolves leave areas where hunting and trapping are prohibited, the threats they face are much greater.</p> <p>Québec: severity depends on the different group and the proportion of animals that are outside areas where hunting and trapping are prohibited; may not be serious in some locations.</p>
5.3	Logging and wood harvesting	Not a threat	Large	Neutral or potentially beneficial	High	<p>COSEWIC has assessed logging and wood harvesting and determined that it was not a threat. An impact could occur in the event of the cutting of stands surrounding dens or rendezvous sites, but the cutting of other stands could be of some benefit associated with an increase in ungulate prey density.</p>
6	Human intrusions and disturbances	Negligible	Pervasive	Negligible	High	–
6.1	Recreational activities	Negligible	Pervasive	Negligible	High	<p>The recreational activities considered include, but are not limited to: snowmobiling, use of ATVs, boating, hiking and ecotourism activities associated with wolves.</p>

Threat #	Threat Description	Impact ^a	Scope ^b	Severity ^c	Timing ^d	Detailed threats
7	Natural system modifications	Negligible	Restricted-Small	Negligible	High	–
7.1	Fire and fire suppression	Negligible	Restricted-Small	Negligible	High	Forest fires are a natural process that can, however, restrict the habitat available for the Eastern Wolf.
7.2	Dams and water management/use	Negligible	Negligible	Extreme	High	The implementation of new major dam construction projects in the area of occupancy of the Eastern Wolf is not anticipated. The scope was therefore considered negligible. The severity of the threat has been assessed as “extreme” given that large areas could become unsuitable for the Eastern Wolf in the event of flooding of habitat due to possible future major dam construction.
8	Invasive and other problematic species and genes	Medium	Pervasive	Moderate	High	–
8.1	Invasive non-native/alien species	Unknown	Pervasive	Unknown	High	This threat deals with the diseases, parasites and threats associated with domestic animals (mainly the domestic dog); the hybridization of the Eastern Wolf with the Eastern Coyote is dealt with in point 8.3 – <i>Introduced genetic material</i> . The severity of this threat is unknown, in both Quebec and Ontario. The scope is pervasive in Ontario and unknown in Quebec.
8.3	Introduced genetic material	Medium	Pervasive	Moderate	High	In the assessment of this threat, the hybridization of the Eastern Wolf with the Eastern Coyote was taken into account. The assessment of the threat takes into consideration the fact that a large proportion of the packs have territories located largely in areas where hunting and trapping are prohibited and, consequently, where the introgression of Eastern Coyote genes is more limited (see details in the description of the threat in section 4.2). Severity may be more serious in the area of Algonquin Provincial Park, but “moderate” when all populations are considered.

Threat #	Threat Description	Impact ^a	Scope ^b	Severity ^c	Timing ^d	Detailed threats
9	Pollution	Negligible	Small	Negligible	High	–
9.4	Garbage and solid waste	Negligible	Small	Negligible	High	The severity of the threat associated with waste collection and landfill sites was considered negligible given the positive and negative effects on the species. As a matter of fact, landfill sites can encourage movements out of the territory, thereby increasing the risk of human-related mortality or interpack aggression (Patterson, pers. comm. 2017), but can also be a food source.

^a **Impact** – The degree to which a species is observed, inferred, or suspected to be directly or indirectly threatened in the area of interest. The impact of each threat is based on Severity and Scope rating and considers only present and future threats. Threat impact reflects a reduction of a species population or decline/degradation of the area of an ecosystem. The median rate of population reduction or area decline for each combination of scope and severity corresponds to the following classes of threat impact: Very High (75% declines), High (40%), Medium (15%), and Low (3%). Unknown: used when impact cannot be determined (e.g., if values for either scope or severity are unknown); Not Calculated: impact not calculated as threat is outside the assessment timeframe (e.g., timing is insignificant/negligible or low as threat is only considered to be in the past); Negligible: when scope or severity is negligible; Not a Threat: when severity is scored as neutral or potential benefit.

^b **Scope** – Scope – Proportion of the species that can reasonably be expected to be affected by the threat within 10 years. Usually measured as a proportion of the species' population in the area of interest. (Pervasive = 71–100%; Large = 31–70%; Restricted = 11–30%; Small = 1–10%; Negligible < 1%).

^c **Severity** – Within the scope, the level of damage to the species from the threat that can reasonably be expected to be affected by the threat within a 10-year or 3-generation timeframe. Usually measured as the degree of reduction of the species' population. (Extreme = 71–100%; Serious = 31–70%; Moderate = 11–30%; Slight = 1–10%; Negligible < 1%; Neutral or Potential Benefit ≥ 0%).

^d **Timing** – High = continuing; Moderate = only in the future (could happen in the short term [< 10 years or 3 generations]) or now suspended (could come back in the short term); Low = only in the future (could happen in the long term) or now suspended (could come back in the long term); Insignificant/Negligible = only in the past and unlikely to return, or no direct effect but limiting.

4.2 Description of Threats

The threats to the Eastern Wolf are described below in descending order of threat impact by category (main threats, threats of low or negligible impact, and potential threats). Under each category, the numbering of the threats corresponds to the numbering in Table 1, and they are therefore not necessarily presented in order of relative impact.

4.2.1 Main threats

Roads and railroads (IUCN Threat 4.1)

Transportation corridors in themselves do not necessarily negatively impact wolves, particularly narrow logging roads or roads with a low traffic volume, especially those at a greater distance from primary roads (Popp and Donovan 2016). In fact, wolves can use certain types of secondary and tertiary roads to facilitate their movements and improve their predation success (Noss 1990; Whittington et al. 2011; Lesmerises et al. 2012). Studies have shown that wolves can persist in areas where habitat is fragmented, provided they are not subject to harvesting and provided these fragmented habitats maintain the characteristics necessary to the survival of wolves in this area (Mech 1989; Fuller et al. 1992; Berg and Benson 1999; Merrill 2000; Fritts et al. 2003; Holloway 2009). The increased access that roads provide to humans (including hunters and trappers; see *Hunting and collecting terrestrial animals (IUCN Threat 5.1) – hunting, trapping and poaching* section) is an indicator of the increase in the hunting and trapping harvest (Benson et al. 2012, 2014). Furthermore, in some areas collisions with vehicles are a significant source of Eastern Wolf mortality, including in and around the Papineau-Labelle Wildlife Reserve (Potvin 1987) and Algonquin Provincial Park (Friends of Algonquin Park 2015). Roads can also cause a loss of suitable habitat (e.g., a reduction in habitats containing sufficient prey) and give Eastern Coyotes better access to the prey species that live on the forest edge. Higher road density also increases the probability of hybridization between Eastern Wolves and Eastern Coyotes (Benson et al. 2012).

It has been suggested that maintenance of road density to less than 0.3 to 0.7 km of roads per km² was necessary to support wolves, with a low proportion of these roads being highways (roughly below 0.02 km/km²) (Wydeven et al. 1998; Rateaud et al. 2001). Areas in southern Ontario and southern Quebec where the road network is very dense can significantly impede natural expansion of the Eastern Wolf in its historical range (COSEWIC 2015). The road density in southern Ontario is generally greater than 0.6 km/km² (Buss and deAlmeida 1997) and has been increasing over the past several decades (COSEWIC 2015).

Hunting and collecting terrestrial animals (IUCN Threat 5.1) – hunting, trapping and poaching

Wolves are regulated species of furbearing mammals in both Ontario (*Fish and Wildlife Conservation Act* [S.O. 1997, c. 41]) and Quebec (*Act Respecting the Conservation and Development of Wildlife* [C.Q.L.R., c. C-61.1]).

In some regions of Canada, Eastern Wolves are or have been hunted or trapped for both commercial (e.g., by licensed trappers) and recreational purposes. Bounties were once offered for killing canids in certain areas, some of which still exist (Hénault and Jolicœur 2003; Hénault, pers. comm. 2017). The trapping harvest largely exceeds the hunting harvest. For example, based on a mail-in survey conducted in southern Quebec, 97% of the reported harvest of wolves came from trapping (Jolicœur *et al.* 2000). In addition to direct losses of individuals, harvesting activities can cause indirect impacts. Rutledge *et al.* (2010e) showed that a decrease in human-caused mortality, following the wolf harvesting ban in Algonquin Provincial Park and the surrounding townships, was accompanied by a restoration of the social structure of the packs and a decrease in the number of unrelated animals in the pack. Conversely, high harvesting rates are related to an increase in hybridization of the Eastern Wolf with the Coyote (Rutledge *et al.* 2011; Benson *et al.* 2014; see *Introduced genetic material [IUCN Threat 8.3] – hybridization of the Eastern Wolf with the Coyote*), which suggests that the presence of areas where harvesting is prohibited promotes the persistence of packs containing few or no hybrid individuals. The annual survival of Eastern Wolves is much higher in areas where hunting and trapping have been reduced or banned outright, compared to areas where hunting and trapping are permitted (Benson *et al.* 2014). Hunting could also have negative physiological effects (e.g., increased stress, which can contribute to the development of pathologies) in the remaining wolves in the pack, as illustrated in the Grey Wolf (Bryan *et al.* 2015; Molnar *et al.* 2015). In order to enhance the understanding of the effects of trapping on the Eastern Wolf population in Canada, additional information is provided in Appendix A for Quebec and Ontario.

Hunting and trapping of Eastern wolves and coyotes are prohibited in certain protected areas located within the area of occupancy, including the national parks of Canada (e.g., La Mauricie National Park), Quebec national parks (e.g., Mont-Tremblant National Park), and some Ontario provincial parks (e.g., Algonquin, Queen Elizabeth II Wildlands, and Killarney provincial parks [OMNRF 2016]). Hunting and trapping are also banned in certain other areas. Appendix C presents areas closed to wolf and coyote hunting and trapping in Ontario as of September 2016. Although a large proportion of Eastern Wolf pack territories are located in protected areas or in areas where hunting and trapping are banned, there are a number of packs whose territories are located in whole or in part on lands where harvesting activities are permitted (e.g., wildlife reserves in Quebec, ZECs or private lands near protected areas; see, for example, Potvin 1987; Theberge and Theberge 2004).

Eastern Wolves can also be incidentally harvested (legally or illegally) in association with big game hunting (e.g., during the hunting seasons for White-tailed Deer, Moose and Black Bear [*Ursus americanus*]) (Hénault and Jolicœur 2003; OMNR 2005a;

Patterson, pers. comm. 2012 in COSEWIC 2015), or be killed out of hatred or fear of wolves (Hénault and Jolicoeur 2003; Theberge and Theberge 2004; Bath 2006). A certain quantity of Eastern Wolves have been harvested illegally in Ontario in locations where harvesting of coyotes and wolves is banned (Rutledge et al. 2010e). In addition, in areas where hunting of wolf is or was prohibited, but not that of Eastern Coyote, Eastern Wolves are and were likely killed because of confusion in differentiating them from Eastern Coyotes (OMNR 2005a). Moreover, wolves have been hunted and taken for sustenance and ceremonial purposes, as well as in the interest of public safety or in defence of property (e.g., livestock and domestic animals) (OMNR 2005a; *Endangered Species Act*, 2007). The number of Eastern Wolves taken annually for these purposes is currently unknown.

Introduced genetic material (IUCN Threat 8.3) – hybridization of the Eastern Wolf with the Eastern Coyote

Coyotes quickly colonized eastern North America in the 20th century, including the historical area of occupancy of the Eastern Wolf, in part owing to human-induced changes to the landscape, such as the spread of agriculture (Kays et al. 2010). Coyotes reached southern Ontario in 1919 (Nowak 1979) and Quebec in 1944 (Naughton 2012). Intensive hybridization has been documented between Eastern Wolves and Coyotes, resulting in the establishment of a large population of Eastern Coyotes in northeastern North America (Way et al. 2010). It has been demonstrated that hybridization, accompanied by introgression, constitutes a significant threat to several species of the genus *Canis* on a global scale, including the Red Wolf (Wayne and Jenks 1991; Adams et al. 2003) and the Ethiopian Wolf (*Canis simensis*; Gottelli et al. 1994) as well as the Dingo (*Canis lupus dingo*; Elledge et al. 2008). This phenomenon appears to be even more frequent in the case of species that were historically allopatric²⁸ such as the Eastern Wolf and the Coyote (Stronen et al. 2012b). Hybridization is a significant threat to the long-term maintenance of the genetic identity of the Eastern Wolf, particularly in regions where its habitat is fragmented by human activities, which favours the presence of Eastern Coyotes, or when Eastern Wolf population densities are low owing to high mortality rates (Kays et al. 2010; COSEWIC 2015). Indeed, human-caused mortality of Eastern Wolves (see *Hunting and collecting terrestrial animals (IUCN Threat 5.1) – hunting, trapping and poaching*) is identified as one of the main causes of hybridization between Eastern Coyotes and Eastern Wolves in the area of Algonquin Provincial Park (Rutledge et al. 2011), in response to a lack of conspecific mates and disruption of Eastern Wolf pack structure (Rutledge et al. 2010c, 2010e). Gene introgression from Eastern Coyotes to Eastern Wolves appears, however, to be more limited in protected areas, such as Algonquin Provincial Park (Rutledge et al. 2011; Benson et al. 2012; Heppenheimer et al. 2018), because the environmental conditions and resource management regimes in such areas (e.g., regulations banning wolf harvesting in certain protected areas) help make conditions less conducive to hybridization (OMNR 2005a; Rutledge et al. 2010e).

²⁸ Whose area of occupancy does not overlap.

4.2.2 Threats of low or negligible impact

Residential and commercial development (IUCN Threat 1)

Residential and commercial development that results in changes at the landscape scale (e.g., residential and cottage construction, shopping centre construction, development of golf courses or ski trails) can reduce to varying degrees the area of suitable habitat available to the Eastern Wolf as well as to its main prey, while increasing the likelihood of encounters between wolves and humans (Mech 1996; Boitani 2003; Quinn 2005). Where there is some level of human-related wolf mortality, it is suggested that a population density of less than 4 to 8 humans per km² is required for the continued existence of the wolf population (Wydeven et al. 1998; Rateaud et al. 2001). Residential and commercial development can also cause declines in prey populations or changes in prey migration patterns, which can have a negative impact on the viability of a wolf population. It should be noted that residential and commercial development is typically accompanied by new road construction and can be associated with an increase in harvesting around the periphery of development. The effects of these threats are assessed in the *Hunting and collecting terrestrial animals (IUCN Threat 5.1) – hunting, trapping and poaching* and *Roads and railroads (IUCN Threat 4.1)* sections. Large protected areas such as Algonquin Provincial Park are rare in the range of the Eastern Wolf, and smaller areas, such as La Mauricie National Park, rarely support more than one wolf pack (Villemure 2003). Ongoing development continues to reduce suitable Eastern Wolf habitat and can increase the likelihood of hybridization with Eastern Coyotes, which more easily become established in disturbed areas (Lehman et al. 1991; Roy et al. 1994; see section *Introduced genetic material (IUCN Threat 8.3) – hybridization of the Eastern Wolf with the Eastern Coyote*). In addition to effects at the landscape scale, residential and commercial developments could cause adverse effects at the local scale, since Eastern Wolves, like Grey Wolves, likely avoid human infrastructure in the selection of breeding sites (dens, rendezvous sites) (Sazatornil et al. 2016).

Agriculture (IUCN Threat 2)

The conversion of suitable Eastern Wolf habitat to agricultural land results in a loss of forested habitat, which is an important component of the species' habitat. Rateaud et al. (2001) have shown that, in southern Quebec, when prey availability is not a limiting factor, wolves (Eastern Wolves, Grey Wolves and hybrids) live and persist in habitats that have an average forest cover of 82% or greater, but may be present irregularly when the cover is close to 60%. The threat caused by agriculture is also attributable to an increase in wolf mortality rates owing to the attitudes of some farmers towards wolves (Stronen et al. 2007; Mech 2010b; Way and Bruskotter 2012; COSEWIC 2015; this aspect is addressed in the section *Hunting and collecting terrestrial animals (IUCN Threat 5.1) – hunting, trapping and poaching*). The indirect threat associated with the increase in Eastern Coyote densities in agricultural areas is discussed in the section *Introduced genetic material (IUCN Threat 8.3) – hybridization of the Eastern Wolf with the Eastern Coyote*.

Recreational activities (IUCN Threat 6.1)

Recreational activities involving human intrusions into wolf habitat (e.g., wilderness camping) could have an impact on the Eastern Wolf packs affected. Wolves display avoidance behaviour to humans and tend to relocate newborn pups following a disturbance of their den or rendezvous site (Frame et al. 2007; Argue et al. 2008). This threat is not well documented. If we include all the types of recreational activities that take place in the range of the Eastern Wolf (e.g., snowmobiling, ATV use, boating, hiking), its scope is pervasive. However, the severity of the threat was assessed as negligible since the current known area of occupancy of the Eastern Wolf is mainly located in parks or wildlife reserves where these activities are well regulated.

4.2.3 Potential threats

Invasive/non-native alien species (IUCN Threat 8.1) – diseases, parasites and domestic animals

Although the severity of this threat is unknown (Brand et al. 1995), diseases and parasites can be a concern for small, threatened populations (Boitani 2003). Eastern Wolves are susceptible to a certain number of viral diseases, including rabies, canine distemper, canine parvovirus and canine hepatitis (Theberge et al. 1994; Theberge and Theberge 1998). They can also suffer significant mortality caused by mange, an ectoparasite²⁹ (Kreeger 2003), which could result in high energetic costs, as observed in the Grey Wolf (Cross et al. 2016). Eastern Wolves could also be exposed to various vector-borne diseases, such as anaplasmosis, ehrlichiosis, heartworm or Lyme's disease, as in the case of Grey Wolves in Wisconsin (Jara et al. 2016). Research on wolves in and around Algonquin Provincial Park suggests that diseases and parasites pose a low degree of threat to this population (Theberge et al. 1994; Theberge and Theberge 1998; Kreeger 2003). However, the occurrence of diseases and parasites among Eastern Wolves is believed to be more significant outside the large protected areas owing to increased contact with domestic dogs, Eastern Coyotes and other animals which can act as “reservoirs” for parasites and diseases.

Habitat shifting and alteration (IUCN Threat 11.1) – climate change

The severity of the threat that climate change poses to the Eastern Wolf has not yet been assessed. However, models predict a reduction in Moose densities in southern Ontario, including in the Algonquin Provincial Park area (Murray et al. 2006; Rempel 2011), as well as a possible increase in White-tailed Deer densities in Ontario and Quebec (Thompson et al. 1998; Murray et al. 2006). Although this increase in White-tailed Deer density would be beneficial to the Eastern Wolf, the anticipated reduction in snow cover will likely lead to reduced hunting success for the Eastern Wolf, which has an advantage over ungulates in heavy snow cover (DelGiudice et al. 2002;

²⁹ Parasite that lives on the outer surface of its host.

Crête and Larivière 2003). Climate change could also adversely affect the length of time that cached food can be preserved (Sutton et al. 2016).

5. Management Objective

The management objective is to achieve and maintain a viable³⁰ Eastern Wolf population within the species' current range in Canada.

To meet this management objective, it is necessary, at a minimum, to maintain the Eastern Wolf density in the Algonquin Provincial Park area at its current level, which is estimated at approximately 2.1 individuals per 100 km² ^[31] (Rutledge et al. 2010c, 2010e; COSEWIC 2015). This area is important for the conservation of this species in Canada and the status of the Eastern Wolf is better documented in the park than in the rest of the species' Canadian range. Eastern Wolf distribution and abundance elsewhere in Ontario and Quebec has not been studied much. Until more detailed information is available as a baseline, a precautionary approach is important to maintain the species' presence in known occupied sites. Conservation of the species at these sites contributes significantly to the resilience³² and redundancy³³ of the Canadian population. It also allows for maintenance of regional representativeness within the Canadian distribution. Lastly, connectivity between occupied sites and other regions of suitable habitat is required in order to facilitate dispersal of individuals and maintain genetic cohesion of the species in Canada. However, the dispersal routes used by the Eastern Wolf are not well documented and additional research is required to assess the species' connectivity needs.

An effective population size³⁴ of at least 500 mature individuals is generally believed to be needed to sustain the genetic diversity required to ensure the viability of a population such as that of the Eastern Wolf in Canada (Franklin and Frankham 1998; Rutledge et al. 2016)³⁵. The effective size of the Algonquin Provincial Park Eastern Wolf population is estimated at between 24 and 122 individuals (Rutledge et al. 2016; see section 3.3 *Species Population and Distribution*). Even considering the Eastern Wolves present outside the study area of Rutledge et al. (2016), including in Quebec, it is unlikely that the effective population size is currently at the 500-individual threshold. Moreover, Rutledge et al. (2016) estimate that a total population size of 2500 to 4545 Eastern

³⁰ A population that is sufficiently abundant and well adapted to its environment that it can persist in the long term (in the face of demographic, genetic and environmental stochasticity and natural disasters), without the need to manage it or to continually invest resources.

³¹ The density figure of 3.0 individuals/100 km² from Rutledge et al. (2010e) was modified by multiplying it by an estimate of the proportion of these wolves assigned to the "Eastern Wolf" species (69%; data from Rutledge et al. 2010c; COSEWIC 2015).

³² Resilience is the ability of a population to recover from a disturbance. Resilience is influenced by population size, level of genetic diversity, as well as characteristics of the species and its habitat.

³³ Redundancy is the presence of multiple populations of the species to guard against catastrophic loss.

³⁴ See section 3.3 for a definition.

³⁵ The effective population size of 500 mature individuals is referred to here for discussion purposes and should not be interpreted as the target for this management plan.

Wolves would be needed to reach an effective population size of 500 individuals, which is much higher than the current estimate of 450-2578 individuals, the upper figure of which is considered unlikely (see section 3.3 *Species Population and Distribution*). It is unclear whether the Eastern Wolf's current area of occupancy and the connectivity between occupied areas are sufficient to maintain a viable population.

In the assessment of the success of the plan, the management objective may be revisited or further specified in light of new information on the taxonomy, abundance or distribution of the species.

6. Broad Strategies and Conservation Measures

6.1 Actions Already Completed or Currently Underway

The following actions have been completed or are currently underway to contribute to the conservation of the Eastern Wolf in Canada.

- Eastern Wolves in and around Algonquin Provincial Park, in Ontario, have been the subject of research and monitoring since the 1960s. As a result, there is an exhaustive collection of data on the ecology, population dynamics and genetics of the wolves in this area (Forbes and Theberge 1992, 1995, 1996a, 1996b; Theberge et al. 1994, 2006; Vucetich and Paquet 2000; Norris et al. 2002; Grewal et al. 2004; Theberge and Theberge 1998, 2004; Mills 2006; Argue et al. 2008; Mills et al. 2008; Patterson and Murray 2008; Rutledge et al. 2010c, 2010e, 2011, 2015, 2016; Benson et al. 2012, 2013, 2014, 2015; Benson and Patterson 2013; Heppenheimer et al. 2018).
- Ecological data have been collected concerning wolves in Ontario outside Algonquin Provincial Park. These data have contributed greatly to our understanding of the distribution, behaviour, ecological role and genetics of the Eastern Wolf as well as of the factors that influence hybridization (Schmitz and Kolenosky 1985; Sears et al. 2003; Wheeldon 2009; Wheeldon and White 2009; Wilson et al. 2009; Holloway 2009; Loveless 2010; Rutledge 2010a; Rutledge et al. 2010b 2010d, 2016; Benson et al. 2012; Otis et al. 2017; Heppenheimer et al. 2018).
- In 2001, a ban on the harvesting of wolves was enacted in 40 townships surrounding Algonquin Provincial Park and in three townships located within the park (one of which overlaps the park boundary and is included in the 40 townships referred to above). This ban was extended in 2004 to include Eastern Coyotes and other coyote species owing to the difficulty distinguishing them morphologically from Eastern Wolves. The regulations governing harvesting were also amended in other areas of central and northern Ontario in order to better manage and conserve wolf populations. In 2016, the ban on the harvesting of wolves, Eastern Coyotes and other coyotes was extended to three additional areas, namely the townships in and around Kawartha Highlands,

Queen Elizabeth II Wildlands and Killarney provincial parks (OMNRF 2016; Appendix C).

- Under a phase-out policy on trapping, trapping of wolves will be eliminated from one-third of the remaining Ontario provincial parks where it is currently permitted (COSEWIC 2015).
- Forest management on public lands in Ontario takes the habitat needs of the Eastern Wolf into account, either directly by restricting logging operations near known dens and rendezvous sites, or indirectly by creating habitat for its prey (White-tailed Deer, Moose and Beaver). The Ontario Ministry of Natural Resources and Forestry has published the *Forest Management Guide for Conserving Biodiversity at the Stand and Site Scales* (OMNR 2010a) and the *Forest Management Guide for Great Lakes-St. Lawrence Landscapes* (OMNR 2010b), which provide information about creating and conserving habitat.
- Although few data have been published on the relationships of the Eastern Wolf with its habitat, research has been conducted in Algonquin Provincial Park with a view to determining the effects of the management of forests and other landscape components on the habitat use and hunting practices of wolves (Loveless 2010). Factors having an impact on territory size and competition for resources between Eastern Wolf packs have also been studied (Arseneau 2010). The influence of the landscape on hybridization dynamics was also the subject of a large-scale study in certain parts of Quebec and Manitoba (Stronen et al. 2010) and, on a smaller scale, within and near Algonquin Provincial Park by researchers from Trent University and from the Ontario Ministry of Natural Resources (Benson et al. 2012).
- In partnership with the University of Sherbrooke, La Mauricie National Park conducted a study between 2000 and 2003 with the goal of gaining a better understanding of the ecology of wolves (including the Eastern Wolf) in the park and surrounding area (Villemure 2003; Villemure and Festa-Bianchet 2004). These studies prompted Parks Canada to develop a conservation strategy for wolves in and outside La Mauricie National Park (Villemure and Masse 2005).
- Research has also been conducted by the University of Montreal on the extent of hybridization and the distribution of the Eastern Wolf in Canada (Stronen et al. 2012a, 2012b; Rogic et al. 2014), in partnership with Parks Canada, Environment and Climate Change Canada, the Société des établissements de plein air du Québec and the Quebec Department of Forests, Wildlife and Parks. This research has made it possible to document the presence of Eastern Wolves in Mont-Tremblant National Park and surrounding area and in La Mauricie National Park and surrounding area.
- The Quebec Department of Forests, Wildlife and Parks released a report on the genetic identification and distribution of large wild canids in Quebec (Mainguy et al. 2017) using samples from 438 large canids, 13 of which were identified as Eastern Wolves.
- Research based on the genetic identification of canids from samples taken from pelts harvested by trappers has also been conducted in the Outaouais and Laurentides regions of Quebec (Hénault and Rutledge 2017; Hénault 2019).

- The Wolf Lake Anishinabeg community is conducting a project combining traditional knowledge and scientific research on the wolf, including surveys (DNA, telemetry tracking, population studies; the Kebaowek First Nation contributes to these surveys), harmonization of forest planning designed to maintain mature forests and steps to expand protected areas in the Maganasipi River watershed in Quebec.
- The Mohawk community of Kahnawà:ke is conducting a project involving genetic analyses of canids, potentially including the Eastern Wolf.
- Ontario Parks has instituted an educational program in Algonquin Provincial Park aimed at informing the general public about wolf ecology and at changing the public perception of this species, including a communication program on the unique aspects of the Eastern Wolf, as well as the ecology of wolves and their interactions with humans. In addition, this program helps provide information on the presence of wolves along the Highway 60 corridor in Algonquin Provincial Park (Manseau et al. 2003). Over 168,500 people participated in the Wolf Howl program in Algonquin Provincial Park between 1960 and 2016 (LeGros pers. comm. 2017), and this activity has been designated a “Canadian Signature Experience” by Destination Canada (formerly the Canadian Tourism Commission) (Steinberg, pers. comm. 2017).
- The wolf is the emblematic animal of Mont-Tremblant National Park, and an education program as well as a management plan for habituated wolves (wolves that have become tolerant or accustomed to humans or human activities) have been developed (Tennier 2008). A program of auditory surveys was carried out over several years, but was discontinued because of a lack of conclusive results (Tennier, pers. comm. 2017).
- In 2015, the Société des établissements de plein air du Québec, in collaboration with the Quebec Department of Forests, Wildlife and Parks and the Université du Québec à Rimouski, initiated a study on the canids of Mont-Tremblant National Park, including the Eastern Wolf. The study should enhance the understanding of the genetic identity of canids and their use of the area.
- Since the “Algonquin Wolf” was designated as Threatened, the Ontario Ministry of Natural Resources and Forestry has been working on a recovery strategy for this taxon.

The following actions have been taken or are underway in order to contribute to the conservation of wolves in Ontario, Quebec, Manitoba, Saskatchewan and the Maritimes. Consequently, they are likely to benefit the Eastern Wolf or to help further knowledge of the species.

- Several ecological studies have been conducted on wolf populations in Quebec and their distribution (e.g., Messier 1984, 1985, 1987; Messier and Crête 1985; Potvin 1987; Potvin et al. 1988; Jolicoeur 1998; Jolicoeur and Hénault 2002, 2010; Larivière et al. 2000; Rateaud et al. 2001; Villemure and Festa-Bianchet 2002; Villemure 2003; Villemure and Masse 2004; Houle 2008; Houle et al. 2010; Lesmerises 2012). Once current and future genetic research has determined the

distribution of Eastern Wolves in Quebec, it will be possible to interpret and use the results of these studies in an Eastern Wolf management context.

- The Quebec Department of Forests, Wildlife and Parks is continuing to collect data on large canids captured in traps in order to document their morphologic and genetic profile.
- The Quebec Department of Forests, Wildlife and Parks conducts biennial assessments of harvesting pressure on wolves (all species combined) by determining the number of wolf pelts sold or traded in various parts of the province and compares them against wolf population estimates (Jolicoeur and Hénault, pers. comm. 2010).
- The Nionwentsïo Office (Huron-Wendat Nation) gathered contemporary knowledge about wolves from Huron-Wendat trappers (Bureau du Nionwentsïo 2016).
- The Ontario Ministry of Natural Resources and Forestry released a Strategy for Wolf Conservation in Ontario (OMNR 2005b).
- Research has been underway since 2010 in northern Ontario on the ecological relationships between wolves and moose (Vander Vennen et al. 2016) and between wolves and caribou (Patterson, comm. pers. 2009) and on the factors that affect wolf density (Kittle et al. 2015).
- Genetic research conducted in Manitoba has provided information on wolves in Duck Mountain Provincial Park and Riding Mountain National Park (Stronen 2009; Stronen et al. 2010, 2012a, 2012b) as well as on the migratory patterns of wolves to and from this province (Crichton, pers. comm. 2010).
- Genetic and isotopic analyses were performed on a canid captured near Caraquet, New Brunswick, which turned out to be a Great Lakes–Boreal Wolf (McAlpine et al. 2015).
- The analysis of tissue samples provided by Paul Paquet (Saskatchewan) to the University of Montreal laboratory made it possible to validate information on the distribution of the genetic material of wolves in Saskatchewan (Stronen, pers. comm. 2011).
- Parks Canada studied the attitudes and perceptions of residents, hunters and trappers in the area of La Mauricie National Park toward wolves (Parks Canada 2007) and developed and instituted an educational program at the park on the importance of wolves aimed at modifying public perceptions of them (Bath 2006; Leith 2007; SOM Inc. 2007; TNS Inc. 2007a, 2007b). Education and awareness efforts are under way.
- The attitudes of farmers toward wolves in the area around Riding Mountain National Park in Manitoba and the factors that can influence these attitudes have also been studied (Stronen et al. 2007).
- The group Midwest Wolf Stewards has met annually since the late 1980s to discuss wolf conservation in the Great Lakes region. Meeting participants include representatives of provincial and federal organizations, state agencies and non-governmental organizations as well as First Nations interested in the management of wolves in the Great Lakes region.
- The Quebec Department of Forests, Wildlife and Parks has published and implemented management plans for Moose (since 1994; Lefort and Massé 2015)

and for White-tailed Deer (since 1996; Huot and Lebel 2012) in order to manage cervid populations, which are an important component of the Eastern Wolf's diet.

6.2 Broad Strategies

The broad strategies of this management plan are described below. They are not presented in order of priority. The priority is identified in relation to the conservation measures (see section 6.3 *Conservation Measures*).

1. Eliminate or reduce the main threats to the species and its habitat in Canada.
2. Increase the awareness, education and engagement of key stakeholders in Eastern Wolf conservation and promote research initiatives.
3. Conduct surveys, clarify certain demographic parameters and monitor the distribution and population of the Eastern Wolf.
4. Address knowledge gaps that need to be filled to manage the Eastern Wolf (e.g., taxonomy, habitat and threats).

6.3 Conservation Measures

The conservation measures, the associated priority and a proposed implementation timetable for applying the broad strategies are presented in Table 2.

Table 2. Conservation Measures and Implementation Schedule

Conservation and Management Measures	Priority ^a	Threats or Concerns ^b	Timeline
1. Broad strategy: Eliminate or reduce the main threats to the species and its habitat in Canada			
1.1 Institute or continue to implement management measures that reduce the likelihood of introgression by the Eastern Coyote or by other large canid species in areas in which the Eastern Wolf occurs.	High	3	2029
1.2 Where necessary and as applicable, establish buffer zones around areas occupied by the Eastern Wolf and develop specific management measures that promote conservation of the species.	High	2, 3	2029
1.3 Identify dispersal corridors and potential areas of colonization in the range of the Eastern Wolf in order to plan and implement measures that reduce the human footprint (e.g., trapping and hunting, roads, agriculture) to thresholds acceptable for the species.	High	1, 5	2024
1.4 Encourage the creation, conservation and stewardship of healthy forests or forest ecosystems in the species' range that will contribute to ensuring the natural predator-prey dynamics of the Eastern Wolf.	High	1, 3, 4, 5, 6, 9	2029
1.5 Consider the needs of the Eastern Wolf in the management plans and policies that apply to public lands, environmental assessments, and land use planning initiatives (e.g., forestry, mines, agriculture, energy) in areas in which the species occurs.	High	1, 2, 4, 5, 6, 9	Ongoing
1.6 Promote and support the application of existing acts and regulations that help to reduce threats to the Eastern Wolf in areas in which the species occurs.	High	1, 2, 3, 4, 5, 6	Ongoing
1.7 Where trapping of large canids is authorized, promote the use of humane trapping techniques.	Medium	2	Ongoing
1.8 Develop and apply best management practices to reduce the number of vehicle-wolf collisions.	Medium	1	Ongoing
1.9 In recreational/tourism areas, plan and implement activities designed to minimize disturbance of the Eastern Wolf.	Medium	6	Ongoing
1.10 Develop and apply management measures that target White-tailed Deer, Moose and Beaver harvesting rates in order to maintain the natural predator-prey dynamics of the Eastern Wolf.	Medium	9	Ongoing
1.11 Encourage the development, or improvement, and application of acts, regulations or policies where deemed necessary.	Medium	1, 2, 3, 4, 5, 6	2024

Conservation and Management Measures	Priority ^a	Threats or Concerns ^b	Timeline
2. Broad strategy: Increase awareness, education and engagement of key stakeholders in Eastern Wolf conservation and promote research initiatives			
2.1 Study the attitudes and perceptions of key stakeholders (e.g., farmers, trappers, hunters) with respect to the wolf in the range of the Eastern Wolf, and develop education and awareness programs aimed at increasing their tolerance.	High	2, 4, 5, 6	2029
2.2 Educate and raise the awareness of communities that coexist with the Eastern Wolf (e.g., farmers, trappers, hunters, municipalities) about the status of the species and practices compatible with persistence of the species.	High	2, 4, 5, 6	Ongoing
2.3 Maintain or develop initiatives designed to provide key stakeholders with information on the presence of the wolf at the local scale (e.g., citizen science natural resources monitoring program)	High	10	2029
2.4 Institute or continue initiatives that promote the engagement and cooperation of governments, First Nations and key stakeholders (e.g., non-governmental organizations, private land owners, forest companies, trappers) in Eastern Wolf conservation efforts (e.g., at the landscape scale).	High	1, 2, 3, 4, 5, 6, 7	2024
2.5 Promote and support research and knowledge transfer initiatives, including traditional ecological knowledge related to the Eastern Wolf.	High	10	Ongoing
2.6 Raise public awareness about wolves and their habitat, in order to change negative attitudes and behaviour toward wolves (e.g., through the educational programs offered in the protected areas).	Medium	2, 5	Ongoing
3. Broad strategy: Conduct surveys, clarify certain demographic parameters and monitor the distribution and population of the Eastern Wolf			
3.1 Develop and promote the use of standardized protocols (e.g., data collection and processing, assignment tests) and databases.	High	10	Ongoing
3.2 Using standardized genetic assignment tests, conduct surveys in the range of the Eastern Wolf and adjacent areas, particularly in previously unsurveyed suitable habitat.	High	10	2024
3.3 Estimate local densities of Eastern Wolf populations at known sites.	High	10	2024
3.4 Maintain a program for monitoring the Eastern Wolf population in order to identify population trends in Algonquin Provincial Park, and initiate such monitoring programs at other important occupied sites within the species' range.	High	10	2029
3.5 Obtain a more precise estimate of the Eastern Wolf population in Canada and of the effective population size throughout the species' range.	High	10	2029

Conservation and Management Measures	Priority ^a	Threats or Concerns ^b	Timeline
3.6 Establish a system for monitoring distribution based on the area of occupancy of the Eastern Wolf.	Medium	10	2024
3.7 Locate the areas of overlap between the Eastern Wolf and the “Eastern Coyote” (<i>C. latrans</i> x <i>C. sp. cf. lycaon</i>) and the “Great Lakes–Boreal Wolf” (<i>C. lupus</i> x <i>C. sp. cf. lycaon</i>) in order to target areas in which efforts to reduce hybridization should be focused.	Medium	10	2029
4. Broad strategy: Address knowledge gaps that need to be filled to manage the Eastern Wolf (e.g., taxonomy, habitat and threats)			
4.1 Estimate the number of Eastern Wolves harvested by trapping and hunting activities and the impact of the harvest rate on the Eastern Wolf population.	High	2, 10	2024
4.2 Assess the role played by occupied sites other than Algonquin Provincial Park in terms of the conservation of naturally regulated Eastern Wolf populations, and determine whether buffer zones with a ban on wolf harvesting around these areas are necessary to ensure adequate conservation of Eastern Wolf packs.	High	10	2024
4.3 Study the factors that promote or reduce Eastern Wolf hybridization (e.g., habitat fragmentation, human-related mortality).	High	3, 10	2029
4.4 Determine the ecological conditions and thresholds that promote persistence of the Eastern Wolf (e.g., competition, predation, availability of prey, availability of dens and rendezvous sites, human density).	High	9, 10	2029
4.5 Determine the most appropriate spatial unit for Eastern Wolf management (e.g., Canadian range, local population range).	High	10	2029
4.6 Obtain and compare genetic samples from Saskatchewan, Manitoba, Ontario, Quebec, New Brunswick and the United States in order to determine the genetic identity and relationships of the different species of the <i>Canis</i> genus and more accurately determine the range of the Eastern Wolf.	Medium	3, 10	2024
4.7 Determine whether the Eastern Wolf could be reliably identified by precise morphological characteristics.	Medium	10	2024
4.8 Conduct studies on habitat selection (e.g., determine and quantify the biophysical characteristics required for conservation of a pack) at different spatial scales (e.g., landscape, territory) for the Eastern Wolf.	Medium	10	2029
4.9 Obtain more detailed data on the aspects of Eastern Wolf biology for which knowledge is insufficient (e.g., survival rate, generation time, social structure, adaptability).	Medium	10	2029

Conservation and Management Measures	Priority ^a	Threats or Concerns ^b	Timeline
4.10 Estimate the number of incidental kills of Eastern Wolf (e.g., during hunting for Moose or White-tailed Deer).	Medium	2, 10	2024
4.11 Study the effect of roadkill on Eastern Wolf populations in the known area of occupancy.	Medium	1, 10	2029
4.12 Study the effect of landscape fragmentation (e.g., roads, agriculture, residential and commercial development) on the Eastern Wolf and its requirements in terms of connectivity.	Medium	1, 4, 5, 10	2029
4.13 Compile cases of wolf mortality within the Eastern Wolf's range that are associated with protecting human life or property, and make this information available in order to be able to identify and implement appropriate reduction and mitigation measures, where required.	Low	2	2029
4.14 Obtain more detailed data on the impact and current extent of threats posed by parasites and diseases on Eastern Wolf populations in Canada. If parasites and diseases constitute a significant threat for maintenance of Eastern Wolf populations, study the methods of transmission of parasites and diseases (e.g., by populations of other wildlife species or by domestic dogs) and develop techniques for mitigating their effects.	Low	7, 10	2029
4.15 Conduct the necessary studies in order to assess the anticipated direct and indirect effects of climate change or any other threats that could arise in the future on Eastern Wolf populations (e.g., habitat shifting and alteration, parasitic diseases, higher trophic level interactions).	Low	8, 10	2029

^a **“Priority”** reflects the degree to which the measure contributes directly to the conservation of the species or is an essential precursor to a measure that contributes to the conservation of the species. High priority measures are considered those most likely to have an immediate and/or direct influence on attaining the management objective for the species. Medium priority measures may have a less immediate or less direct influence on reaching the management objective, but are still important for the management of the population. Low priority conservation measures will likely have an indirect or gradual influence on reaching the management objective, but are considered important contributions to the knowledge base and/or public involvement and acceptance of the species.

^b **Threats** – 1) roads and railroads, 2) hunting and collecting terrestrial animals – hunting, trapping and poaching, 3) introduced genetic material – hybridization of the Eastern Wolf with the Eastern Coyote, 4) residential and commercial development, 5) agriculture, 6) recreational activities, 7) invasive/non-native alien species – diseases, parasites and domestic animals, 8) habitat shifting and alteration – climate change.
Concerns: 9) predator-prey dynamics (limiting factor), 10) knowledge gaps.

6.4 Narrative to Support Conservation Measures and Implementation Schedule

A number of specific characteristics of the Eastern Wolf must be taken into account in preparing a management strategy for the species. First, the current Canadian range of the Eastern Wolf is small compared to its historical range. Its expansion is limited by urbanization to the south within Ontario and Quebec and by the presence of the Grey Wolf or Great Lakes–Boreal Wolf to the north. The range of the Eastern Wolf also appears to be fragmented by areas of unsuitable habitat or by areas in which management measures are not compatible with the maintenance of Eastern Wolf packs. Second, the Eastern Wolf is a top predator, i.e., it is typically not a prey species of any other species and is therefore at the top trophic level of its ecosystem. Like most mammals that are top predators, Eastern Wolves occur in small densities in large areas of habitat and have a relatively long generation time. The management strategy for the species must therefore also be adapted to the characteristics of this type of predator, for example by promoting the conservation of large tracts of forest enabling them to meet their ecological needs, which includes ensuring the availability of their preferred prey. Like many top predators, the Eastern Wolf may also raise public safety concerns or fear on the part of the public.

In order to apply appropriate conservation measures, it is important to more accurately determine the current and historical range of the Eastern Wolf in Canada, along with the demographic data (e.g., wolf density). This will make it possible to implement the necessary monitoring to identify population and distribution trends. The use of standard sampling protocols and standardized databases will be of prime importance for clarifying uncertainties related to the status of Eastern Wolf populations and the species' Canadian range and for determining to what extent the Canadian population is resilient and redundant.

Given the knowledge gaps on Eastern Wolf population and distribution, which are due in part to uncertainties regarding the extent of hybridization with other species and hybrids of the genus *Canis* and to a lack of data, further studies will be required to resolve the taxonomic uncertainties and to manage the technical challenges associated with the need to identify the species by molecular analysis. Given these uncertainties, it is likely that some conservation and management measures will need to be broadly applied to all large canids found within the Eastern Wolf range, at least temporarily.

Knowledge gaps relating to the impact, scope, severity and timing of certain threats, including hunting, trapping and poaching, hybridization with Eastern Coyotes and road mortality, also need to be addressed. Certain characteristics of Eastern Wolf habitat and ecology also need to be further defined, including the ecological thresholds promoting viability of the population and the species' connectivity requirements.

Given the threat to the Eastern Wolf posed by hybridization with Eastern Coyotes, maintaining a viable Eastern Wolf population in Canada will require the elimination or reduction of several threats, including those associated with roads, hunting, trapping

and poaching, which have an effect on hybridization with Coyotes (see section 4.2 *Threats Description*). At the same time, in Ontario and Quebec, integrated, large-scale management of species and wild hybrids of the genus *Canis* will be required to promote stable hybrid systems (see Bohling 2016). The threats that cause increased mortality or loss of suitable habitat will also have to be reduced or eliminated in order to ensure the viability of the population. More extensive use of certain conservation measures in the area of occupancy of the Eastern Wolf aimed at reducing or eliminating the primary threats (e.g., implement a ban on the harvesting of Eastern Wolves and Eastern Coyotes in some areas – see section 6.1 *Actions already completed or underway*) could thus contribute to the achievement of the management objective. Ensuring the well-being of Eastern Wolves must also be considered in territories where it can be harvested by trapping; the use of humane trapping techniques (AIHTS 1999; Proulx et al. 2012, 2015) should be advocated to minimize animal pain and suffering and would benefit all canids. Ensuring connectivity between occupied areas containing suitable unoccupied habitats in the surrounding area is also important, as it contributes to the natural expansion and genetic cohesion of the Eastern Wolf population in Canada.

Finally, the involvement of governments, First Nations and a wide range of stakeholders (e.g., non-governmental organizations, universities, private landowners, forest companies) will be critical to the implementation of many conservation measures. Transfer of knowledge (e.g., detailed genetic profile) to affected parties will be particularly important since many knowledge acquisition measures are set out in the implementation schedule (Table 2). The introduction of communications programs designed to improve the public's perception of wolves and to modify negative behaviours and attitudes toward wolves also appears to be very important, given that social acceptance is critical to reducing certain threats, such as the killing of Eastern Wolves out of fear or hatred.

7. Measuring Progress

Every five years, success in the implementation of this management plan will be measured against the following performance indicators:

1. The Canadian Eastern Wolf population is assessed as viable;
2. The extent of the species' range in Canada is maintained.

8. References

- Adams, J.R., B.T. Kelly, and L.P. Waits. 2003. Using faecal DNA sampling and GIS to monitor hybridization between red wolves (*Canis rufus*) and Coyotes (*Canis latrans*). *Molecular Ecology* 12:2175–2186.
- AIHTS. 1999. Agreement on International Humane Trapping Standards between the European Union, Canada and the Russian Federation. Website: <http://fur.ca/fur-trapping/humane-trapping-standards-and-animal-welfare/> [accessed January 2017].
- Algonquin Wolf Advisory Group. 2000. The Wolves of Algonquin Provincial Park: A Report to the Minister of Natural Resources. 26 p. + appendix.
- Allendorf, F.W., R.F., Leary, P. Spruell and J.K. Wenburg. 2001. The problems with hybrids: setting conservation guidelines. *Trends in Ecology and Evolution* 16(11):613-622.
- Argue, A.M., K.J. Mills, and B.R. Patterson. 2008. Behavioural response of eastern wolves (*Canis lycaon*) to disturbance at homesites and its effects on pup survival. *Canadian Journal of Zoology* 86:400-406.
- Arseneau, J. 2010. Causes and consequences of group dominance in eastern wolves (*Canis lycaon*). Master's thesis, Trent University, Peterborough, Ontario, Canada. 53 p.
- Ballard, W.B., and J.R. Dau. 1983. Characteristics of gray wolf, *Canis lupus*, den and rendezvous sites in southcentral Alaska. *Canadian Field-Naturalist* 97:299–302.
- Bath, A. J. 2006. Human dimensions in wolf conservation: understanding trappers' perceptions toward wolves in and around La Mauricie National Park of Canada. Report submitted to the Quebec Service Centre, Parks Canada. 96 p.
- Benson, J.F. Pers. comm. 2011. E-mail communication with Kari Van Allen. December 2011. Trent University, Peterborough, Ontario.
- Benson, J.F., B.R. Patterson, and T.J. Wheeldon. 2012. Spatial genetic and morphologic structure of wolves and coyotes in relation to environmental heterogeneity in a *Canis* hybrid zone. *Molecular Ecology* 21:5934-5954.
- Benson, J.F., and B.R. Patterson. 2013. Inter-specific territoriality in a *Canis* hybrid zone: spatial segregation between wolves, coyotes, and hybrids. *Oecologia* 173:1539-1550.
- Benson, J.F., K.J. Mills, K.M. Loveless, and B.R. Patterson. 2013. Genetic and environmental influences on pup mortality risk for wolves and coyotes within a *Canis* hybrid zone. *Biological Conservation* 166:133-141.

- Benson, J.F., B.R. Patterson, and P.J. Mahoney. 2014. A protected area influences genotype-specific survival and the structure of a *Canis* hybrid zone. *Ecology* 95:254-264.
- Benson, J.F., K.J. Mills, and B.R. Patterson. 2015. Resource selection by wolves at dens and rendezvous sites in Algonquin Park, Canada. *Biological Conservation* 182:223-232.
- Benson, J.F., K.M. Loveless, L.Y. Rutledge and B.R. Patterson. 2017. Ungulate predation and ecological roles of wolves and coyotes in eastern North America. *Ecological Applications* 27:718-733.
- Berg, W., and S. Benson. 1999. Updated wolf population estimate for Minnesota, 1997-1998. Minnesota Department of Natural Resources. Grand Rapids, Minnesota. 14 p.
- Bohling, J.H. 2016. Strategies to address the conservation threats posed by hybridization and genetic introgression. *Biological Conservation* 203:321-327.
- Boitani, L. 2003. Wolf conservation and recovery, *in* L.D. Mech and L. Boitani (eds.). *Wolves: behavior, ecology and conservation*, The University of Chicago Press, Chicago, Illinois. Pp. 317-340.
- Brand, C.J., M.J. Pybus, W.B. Ballard, and R.O. Peterson. 1995. Infectious and parasitic diseases of the gray wolf and their potential effects on wolf populations in North America, *in* L.N. Carbyn, S.H. Fritts and D.R. Seip (eds.). *Ecology and conservation of wolves in a changing world*, Canadian Circumpolar Institute, Edmonton, Alberta. Pp. 419-429.
- Bryan, H.M., J.E.G. Smits, L. Koren, P.C. Paquet, K.E. Wynne-Edwards, and M. Musiani. 2015. Heavily hunted wolves have higher stress and reproductive steroids than wolves with lower hunting pressure. *Functional Ecology* 29(3):347-356.
- Bureau du Nionwentsïo. 2016. Connaissances contemporaines sur le loup de l'Est recueillies auprès de piégeurs hurons-wendat titulaires de lots de piégeage. Report submitted to Environment and Climate Change Canada. 5 p.
- Buss, M., and M. deAlmeida. 1997. A review of wolf and coyote status and policy in Ontario. Ontario Ministry of Natural Resources. 88 p.
- Chambers, S.M., S.R. Fain, B. Fazio, and M. Amaral. 2012. An account of the taxonomy of North American wolves from morphological and genetic analyses. *North American Fauna* 77:1-67.
- COSEWIC (Committee on the Status of Endangered Wildlife in Canada). 2015. COSEWIC assessment and status report on the Eastern Wolf (*Canis lycaon*) in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa. xii + 67 p.

- COSSARO (Committee on the Status of Species at Risk in Ontario). 2016. Ontario Species at Risk Evaluation Report for Algonquin Wolf (*Canis* sp.), an evolutionarily significant and distinct hybrid with *Canis lycaon*, *C. latrans*, and *C. lupus* ancestry. Website: <https://www.ontario.ca/page/ontario-species-risk-evaluation-report-algonquin-wolf-canis-sp-evolutionarily-significant-and> [accessed November 2016].
- Crête, M., and S. Larivière. 2003. Estimating the costs of locomotion in snow for coyotes. *Canadian Journal of Zoology* 81:1808-1814.
- Crichton, V. Pers. comm. 2010. Communication with Angela McConnell. Conservation Manitoba, Government of Manitoba, Winnipeg, Manitoba.
- Cross, P.C., E.S. Almberg, C.G. Haase, P.J. Hudson, S.K. Maloney, M.C. Metz, A.J. Munn, P. Nugent, O. Putzeys, D.R. Stahler, A.C. Stewart, and D.W. Smith. 2016. Energetic costs of range in wolves estimated from infrared thermography. *Ecology* 97(8):1938-1948.
- Cubaynes, S., D.R. MacNulty, K.A. Quimby, D.W. Smith, and T. Coulson. 2014. Density-dependent intraspecific aggression regulates survival in northern Yellowstone wolves (*Canis lupus*). *Journal of Animal Ecology* 83(6):1344-1356.
- DelGiudice, G.D., M.R. Riggs, P. Joly, and W. Pan. 2002. Winter severity, survival, and cause-specific mortality of female white-tailed deer in north-central Minnesota. *Journal of Wildlife Management* 66:698-717.
- Desy, G.E. 2007. Predicting fine-scale distributions of gray wolves: is habitat an effective surrogate for prey availability? Master's thesis, University of Guelph, Guelph, Ontario, Canada. 72 p.
- Elledge, A.E., L.R. Allen, B.-L. Carlsson, A.N. Wilton, and L.K.-P. Leung. 2008. An evaluation of genetic analyses, skull morphology and visual appearance of assessing dingo purity. *Wildlife Research* 35(8):812-820.
- Estes, J.A., J. Terborgh, J.S. Brashares, M.E. Power, J. Berger, W.J. Bond, S.R. Carpenter, T.E. Essington, R.D. Holt, J.B.C. Jackson, R.J. Marquis, L. Oksanen, T. Oksanen, R.T. Paine, E.K. Pikitch, W.J. Ripple, S.A. Sandin, M. Scheffer, T.W. Schoener, J.B. Shurin, A.R.E. Sinclair, M.E. Soulé, R. Virtanen, and D.A. Wardle. 2011. Trophic downgrading of planet earth. *Science* 333: 301-306.
- Fisher, R.A. 1930. The genetical theory of natural selection. Clarendon Press, Oxford. 272 p.
- Forbes, G.J., and J.B. Theberge. 1992. Importance of scavenging on moose by wolves in Algonquin Park, Ontario. *Alces* 28:235-241.

- Forbes, G.J., and J.B. Theberge. 1995. Influences of a migratory deer herd on wolf movements and mortality in and around Algonquin Park, Ontario, *in* L. Carbyn, S. Fritts and D. Seip (eds.). Ecology and Conservation of Wolves in a Changing World. Canadian Circumpolar Institute, University of Alberta, Edmonton, Alberta. Pp. 303-314.
- Forbes, G.J., and J.B. Theberge. 1996a. Cross-boundary management of Algonquin Park wolves. *Conservation Biology* 10:1091-1097.
- Forbes, G.J., and J.B. Theberge. 1996b. Response to prey variation by wolves in north-central Ontario. *Canadian Journal of Zoology* 74:1511-1520.
- Frame, P., D. Cluff, and D. Hik. 2007. Response of wolves to experimental disturbance at homesites. *Journal of Wildlife Management* 71:313-320.
- Franklin, J.R., and R. Frankham. 1998. How large must populations be to retain evolutionary potential? *Animal Conservation* 1:69-73.
- Friends of Algonquin Park. 2015. Eastern Wolf, Wolf Research in Algonquin Provincial Park. The Science Behind Algonquin's Animals. Website: <http://www.sbaa.ca/projects.asp?cn=314> [accessed October 2011].
- Fritts, S.H., R.O. Stephenson, R.D. Hayes, and L. Boitani. 2003. Wolves and humans, *in* L.D. Mech and L. Boitani (eds.). Wolves: behavior, ecology and conservation, The University of Chicago Press, Chicago, Illinois. Pp. 289-316.
- Fuller, T.K. 1989. Population dynamics of wolves in North-Central Minnesota. *Wildlife Monographs* 105:1-41.
- Fuller, T.K., W.E. Berg, G.L. Radde, M.S. Lenarz, and G.B. Joselyn. 1992. A history and current estimate of wolf distribution and numbers in Minnesota. *Wildlife Society Bulletin* 20:42-55.
- Fuller, T.K., L.D. Mech, and J.F. Cochrane. 2003. Wolf Population Dynamics, *in* L.D. Mech and L. Boitani (eds.). Wolves: behavior, ecology and conservation, The University of Chicago Press, Chicago, Illinois. Pp. 161-191.
- Geffen, E., M.J. Anderson and R.K. Wayne. 2004. Climate and habitat barriers to dispersal in the highly mobile grey wolf. *Molecular Ecology* 13:2841-2490.
- Gottelli, D., C. Sillero-Zubiri, G.D. Applebaum, M.S. Roy, D.J. Girman, J. Garcia-Moreno, E.A. Ostrander, and R.K. Wayne. 1994. Molecular genetics of the most endangered canid: the Ethiopian wolf *Canis simensis*. *Molecular Ecology* 3(4):301-312.
- Grewal, S.K., P.J. Wilson, T.K. Kung, K. Shami, M.T. Theberge, J.B. Theberge, and B.N. White. 2004. A genetic assessment of the Eastern Wolf (*Canis lycaon*) in Algonquin Park. *Journal of Mammalogy* 85:625-632.

- Hailer, F., and J.A. Leonard. 2008. Hybridisation among three native North American *Canis* species in a region of natural sympatry. PLoS ONE 3(10): e3333.
- Hall, R.E., and K.R. Kelson. 1959. The Mammals of North America, Vol. 2. The Ronald Press Company, New York, New York.
- Hamilton, J.A., and J.M. Miller. 2015. Adaptive introgression as a resource for management and genetic conservation in a changing climate. Conservation Biology 30(1):33-41.
- Hénault, M., and H. Jolicoeur. 2003. Les loups au Québec: meutes et mystères. Société de la Faune et des Parcs du Québec. Quebec City, Quebec. 129 p.
- Hénault, M. 2015. Unpub. data. Information contained in the Quebec government's review of the Eastern Wolf Status Report.
- Hénault, M. Pers. comm. 2017. Comments submitted to the Species at Risk Public Registry during the 60-day public comment period.
- Hénault, M., and L. Rutledge. 2017. Évaluation de la présence du loup de l'Est (*Canis* sp. cf. *lycaon*) dans la région de l'Outaouais. Report for Environment and Climate Change Canada, Canadian Wildlife Service –Quebec Region. 24 p.
- Hénault, M. 2019. Évaluation de la présence du loup de l'Est dans la réserve faunique Papineau-Labelle et sa périphérie. Report for Environment and Climate Change Canada, Canadian Wildlife Service – Quebec Region. 11 p. + appendices.
- Heppenheimer, E., R.J. Harrigan, L.Y. Rutledge, K.-P. Koepfli, A.L. DeCandia, K.E. Brzeski, J.F. Benson, T. Wheeldon 8,9, B.R. Patterson, R. Kays, P.A. Hohenlohe, and B.M. von Holdt. 2018. Population genomic analysis of North American Eastern Wolves (*Canis lycaon*) supports their conservation priority status. Genes 9(606):1-18.
- Hohenlohe, P.A., L.Y. Rutledge, L.P. Waits, K.R. Andrews, J.R. Adams, J.W. Hinton, R.M. Nowak, B.R. Patterson, A.P. Wydeven, P.A. Wilson, and B.N. White. 2017. Comment on "Whole-genome sequence analysis shows two endemic species of North American wolf are admixtures of the coyote and gray wolf." Science Advances 3:e1602250.
- Holloway, J. 2009. Size Dependent Resource Use of a Hybrid Wolf (*C. lycaon* × *C. lupus*) Population in Northeast Ontario. Master's thesis, Trent University, Peterborough, Ontario, Canada. 73 p.
- Houle, M. 2008. Effets cumulés des activités forestières sur la sélection d'habitat du loup gris (*Canis lupus*) en forêt boréale aménagée. Master's thesis, Laval University, Quebec City, Quebec, Canada. 42 p.

- Houle, M., D. Fortin, C. Dussault, R. Courtois, and J.P. Ouellet. 2010. Cumulative effects of forestry on habitat use by gray wolf (*Canis lupus*) in the boreal forest. *Landscape Ecology* 25(3):419-433.
- Huot, M., and F. Lebel. 2012. Plan de gestion du cerf de Virginie au Québec 2010-2017. Ministère des Ressources naturelles et de la Faune – Secteur Faune Québec, Direction générale de l'expertise sur la faune et ses habitats, Québec (Québec). 578 p.
- IUCN (International Union for Conservation of Nature). 2016. Protected Areas Categories. Website: <https://www.iucn.org/theme/protected-areas/about/protected-areas-categories> [accessed November 2016].
- Jackiw, R.N., G. Mandil, and H.A. Hager. 2015. A framework to guide the conservation of species hybrids based on ethical and ecological considerations. *Conservation Practice and Policy* 29(4):1040-1051.
- Jara, R.F., A.P. Wydeven, and M.D. Samuel. 2016. Gray Wolf Exposure to Emerging Vector-Borne Diseases in Wisconsin with Comparison to Domestic Dogs and Humans. *PLoS ONE* 11(11):e0165836.
- Jolicoeur, H. 1998. Le loup du massif du Lac-Jacques-Cartier. Ministère de l'Environnement et de la Faune, Direction de la faune et des habitats, Direction de la conservation du patrimoine écologique. Quebec City, Quebec. 132 p.
- Jolicoeur, H., R. Lemieux, J.-P. Ducruc, and C. Fortin. 1998. Caractérisation des tanières de loup dans le massif du lac Jacques-Cartier. Ministère de l'Environnement et de la Faune, Direction de la faune et des habitats, Direction de la conservation du patrimoine écologique. Quebec City, Quebec. 42 p.
- Jolicoeur, H., R. Lafond, N. Scaringella, W. Grenier, and R. Morin. 2000. Résultats d'une enquête postale maison effectuée en 1997 auprès des piégeurs et des chasseurs de loups et de coyotes du sud du Québec. Société de la faune et des parcs, Direction du développement de la faune (Québec). 58 p.
- Jolicoeur, H., and M. Hénault. 2002. Répartition géographique du loup et du coyote au sud du 52^e parallèle et estimation de la population de loups au Québec. Société de la faune et des parcs du Québec. 41 p + appendix.
- Jolicoeur, H., and M. Hénault. 2010. Current status and management of Wolves in Québec. Wolf Stewards Meeting, Dorset, Ontario.
- Jolicoeur, H., and M. Hénault. Pers. comm. 2010. Communication with Sylvain Giguère. Quebec Department of Natural Resources and Wildlife, Quebec City, Quebec, Canada.

- Joslin, P.W.B. 1967. Movements and homesites of timber wolves in Algonquin Park. *American Zoologist* 7: 279-288.
- Kays, R., A. Curtis, and J.J. Kirchman. 2010. Rapid adaptive evolution of northeastern coyotes via hybridization with wolves. *Biology Letters* 6:89-93.
- Kittle, A.M., J. Fryxell, G. Desy, and J. Hamr. 2007. The scale-dependent impact of wolf predation risk on resource selection by three sympatric ungulates. *Oecologia* 157:163-175.
- Kittle, A.M., M. Anderson, T. Avgar, J.A. Baker, G.S. Brown, J. Hagens, E. Iwachewski, S. Moffatt, A. Mosser, B.R. Patterson, D.E.B. Reid, A.R. Rodgers, J. Shuter, G.M. Street, I.D. Thompson, L.M. Vander Vennen, and J.M. Fryxell. 2015. Wolves adapt territory size, not pack size to local habitat quality. *Journal of Animal Ecology* 84:1177–1186.
- Koblmüller, S., M. Nord, R.K. Wayne, and J.A. Leonard. 2009. Origin and status of the Great Lakes wolf. *Molecular Ecology* 18:2313-2326.
- Kolenosky, G.B. 1972. Wolf predation on wintering deer in east-central Ontario. *Journal of Wildlife Management* 36:357-369.
- Kolenosky, G.B., and R. Standfield. 1975. Morphological and ecological variation among gray wolves (*Canis lupus*) of Ontario, Canada, in M.W. Fox (ed.). *The wild canids: their systematics, behavioural ecology, and evolution*, Van Nostrand Reinhold, New York, New York. Pp. 62-72.
- Kreeger, T.J. 2003. The internal wolf: Physiology, Pathology, and Pharmacology, in L.D. Mech and L. Boitani (eds.). *Wolves: behavior, ecology and conservation*, The University of Chicago Press, Chicago, Illinois. Pp. 192-217.
- Kyle, C.J., A.R. Johnson, B.R. Patterson, P.J. Wilson, K. Shami, S.K. Grewal, and B.N. White. 2006. Genetic nature of eastern wolves: Past, present and future. *Conservation Genetics* 7:273-287.
- Kyle, C.J., A.R. Johnson, B.R. Patterson, P.J. Wilson, and B.N. White. 2008. The conspecific nature of eastern and red wolves: conservation and management implications. *Conservation Genetics* 9:699-701.
- Larivière, S., H. Jolicoeur and M. Crête. 1998. Densités et tendance démographique du loup (*Canis lupus*) dans les réserves fauniques du Québec entre 1983 et 1997. Ministère de l'Environnement et de la Faune, Direction de la faune et des habitats, Quebec City, Quebec. 33 p.
- Larivière, S., H. Jolicoeur, and M. Crête. 2000. Status and conservation of the gray wolf (*Canis lupus*) in wildlife reserves in Quebec. *Biological Conservation* 94:143-151.

- Lefort, S., and S. Massé. 2015. Plan de gestion de l'original au Québec 2012-2019. Ministère des Forêts, de la Faune et des Parcs – Secteur de la faune et des parcs, Direction générale de l'expertise sur la faune et ses habitats et Direction générale du développement de la faune, Quebec City, Quebec. 443 p.
- LeGros, D. Pers. comm. 2017. Email communication. Naturalist with the Natural Heritage Education program, Ontario Ministry of Natural Resources and Forestry, Ontario Parks, Algonquin Provincial Park, Ontario, Canada.
- Lehman, N.E., A. Eisenhawer, K. Hansen, L.D. Mech, R.O. Peterson, J.P. Gogan, and R.K. Wayne. 1991. Introgression of Coyote mitochondrial DNA into sympatric North American gray wolf populations. *Evolution* 45:104-119.
- Leith, M.A. 2007. Évaluation sommative des activités éducatives sur le loup de l'Est pour le parc national de la Mauricie. Submitted to the Service d'interprétation et de la mise en valeur du parc national de la Mauricie, Quebec Service Centre, Parks Canada. 42 p.
- Lesmerises, F. 2012. Impacts des perturbations anthropiques sur la sélection d'habitat et la distribution spatiale du loup gris (*Canis lupus*) en forêt boréale. Master's thesis, Université du Québec à Rimouski, Rimouski, Quebec, Canada. 76 p.
- Lesmerises, F., C. Dussault, and M-H. St-Laurent. 2012. Wolf habitat selection is shaped by human activities in a highly managed boreal forest. *Forest Ecology and Management* 276:125-131.
- Loveless, K. 2010. Foraging strategies of eastern wolves in relation to migratory prey and hybridization. Master's thesis, Trent University, Peterborough, Ontario. 81 p.
- Mainguy, J., M. Hénault, H. Jolicœur, and E. Dalpé-Charron. 2017. Identification génétique et répartition spatiale des grands canidés sauvages au Québec. Ministère des Forêts, de la Faune et des Parcs, Direction de l'expertise sur la faune terrestre, l'herpétofaune et l'avifaune and Direction de la gestion de la faune de Lanaudière et des Laurentides, Québec. 82 p.
- Manseau, M., S. Czetwertynski, R. Lemieux, A. Demers, and H. Jolicoeur. 2003. Impact des appels de loups faits dans le cadre d'activités écotouristiques sur comportement de deux meutes de loups dans le massif du lac Jacques-Cartier. *Le Naturaliste Canadien* 127(1):43-54.
- McAlpine, D.F., D. X. Soto, L.Y. Rutledge, T.J. Wheeldon, B.N. White, J.P. Goltz, and J. Kennedy. 2015. Recent Occurrences of Wild-origin Wolves (*Canis* spp.) in Canada South of the St. Lawrence River Revealed by Stable Isotope and Genetic Analysis. *The Canadian-Field Naturalist* 129:386-394.
- McLoughlin, P.D., L.R. Walton, H.D. Cluff, P.C. Paquet, and M.A. Ramsay. 2004. Hierarchical habitat selection by tundra wolves. *Journal of Mammalogy* 85:576-580.

- MDDELCC (Ministère du Développement durable, de l'Environnement et de la Lutte contre les changements climatiques). 2017. Registre des aires protégées. Website: http://www.mddelcc.gouv.qc.ca/biodiversite/aires_protegees/registre/
- Mech, L.D. 1970. The wolf: ecology and behavior of an endangered species. Natural History Press, New York, New York. 389 p.
- Mech, L.D. 1977. Productivity, mortality, and population trends of wolves in northeastern Minnesota. *Journal of Mammalogy* 58:559-574.
- Mech, L.D. 1989. Wolf population survival in an area of high road density. *American Midland Naturalist* 121:387-389.
- Mech, L.D. 1996. A new era for carnivore conservation. *Wildlife Society Bulletin* 24:397-401.
- Mech, L.D. 2010a. What is the taxonomic identity of Minnesota wolves? *Canadian Journal of Zoology* 88:129-138.
- Mech, L.D. 2010b. Considerations for developing wolf harvest regulations in the contiguous United States. *Journal of Wildlife Management* 74:1421-1424.
- Mech, L.D. 2011. Non-genetic data supporting genetic evidence for the eastern wolf. *Northeastern Naturalist* 18:521-526.
- Merrill, S.B. 2000. Road densities and wolf, *Canis lupus*, habitat suitability: an exception. *Canadian Field-Naturalist* 114:312-313.
- Messier, F. 1984. Moose-wolf dynamics and the natural regulation of moose populations. Master's thesis, University of British Columbia, Vancouver, British Columbia. 143 p.
- Messier, F. 1985. Social organization, spatial distribution and population density of wolves in relation to moose density. *Canadian Journal of Zoology* 63:1068-1077.
- Messier, F., and M. Crête. 1985. Moose-wolf dynamics and the natural regulation of moose populations. *Oecologia (Berlin)* 65:503-512.
- Messier, F. 1987. Physical condition and blood physiology of wolves in relation to moose density. *Canadian Journal of Zoology* 65:91-95.
- MFFP (Ministère des Forêts, de la Faune et des Parcs). Unpub. data. Statistiques de piégeage 2011-2012. Ministère des Forêts, de la Faune et des Parcs.
- Mills, K.J. 2006. Wolf (*Canis lycaon*) pup survival, dispersal, and movements in Algonquin Provincial Park, Ontario. Master's thesis, Trent University, Peterborough, Ontario, Canada. 67 p.

- Mills, K.J., B.R. Patterson, and D.L. Murray. 2008. Direct estimation of early survival and movements in Eastern Wolf pups. *Journal of Wildlife Management* 72:949-954.
- Mladenoff, D.J., R.G. Haight, T.A. Sickley, and A.P. Wydeven. 1997. Causes and implications of species restoration in altered ecosystems: A spatial landscape projection of wolf population recovery. *Bioscience* 47:21-31.
- Moldowan, P.D., and H. Kitching. 2017. Observation of an Eastern Wolf (*Canis* sp. cf. *lycaon*) Food Caching in a Sphagnum Bog in Algonquin Provincial Park, Ontario. *Canadian Field-Naturalist* 130(4):351-354.
- Molnar, B., J. Fattebert, R. Palme, P. Ciucci, B. Betschart, D.W. Smith, and P.-A. Diehl. 2015. Environmental and Intrinsic Correlates of Stress in Free-Ranging Wolves. *PLoS ONE* 10(9): e0137378.
- Murray, D.L., E.W. Cox, W.B. Ballard, H.A. Whitlaw, M.S. Lenarz, T.W. Custer, T. Barnett, and T.K. Fuller. 2006. Pathogens, nutritional deficiency, and climate influences on a declining moose population. *Wildlife Monographs* 166: 1-20.
- NatureServe. 2017. NatureServe Explorer: An online encyclopedia of life [web application]. Version 7.1. Website: <http://www.natureserve.org/explorer> [accessed January 2017].
- NatureServe. 2019. NatureServe Explorer: An online encyclopedia of life [web application]. Version 7.1. Website: <http://www.natureserve.org/explorer> [accessed October 2019].
- Naughton, D. 2012. *The Natural History of Canadian Mammals*. University of Toronto Press, Toronto, Ontario. 784 p.
- NLDEC. 2016. Living with Coyotes in Newfoundland and Labrador. Newfoundland and Labrador Department of Environment and Conservation. Website: http://www.env.gov.nl.ca/env/wildlife/all_species/coyotes.html [accessed January 2017].
- Norris, D.R., M.T. Theberge, and J.B. Theberge. 2002. Forest composition around wolf (*Canis lupus*) dens in eastern Algonquin Provincial Park, Ontario. *Canadian Journal of Zoology* 80:866-872.
- Noss, T.D. 1990. The Ecological Effects of Roads. *Wild Earth Guardians*. Website: <http://www.wildlandscpr.org/ecological-effects-roads> [accessed January 2012].
- Nowak, R.M. 1979. North American Quaternary Canis. Monograph No. 6, Museum of Natural History, University of Kansas, Lawrence, Kansas. 154 p.
- Nowak, R.M. 1995. Another look at wolf taxonomy, *in* L.N. Carbyn, S.H. Fritts and D.R. Seip (eds.). *Ecology and conservation of wolves in a changing world*, Canadian Circumpolar Institute, Edmonton, Alberta. Pp. 375-397.

- OMNR (Ontario Ministry of Natural Resources). 2005a. Backgrounder on wolf conservation in Ontario. Ontario Ministry of Natural Resources. Website: www.web2.mnr.gov.on.ca/mnr/ebr/wolves/backgrounder.pdf [accessed January 2012].
- OMNR (Ontario Ministry of Natural Resources). 2005b. Strategy for Wolf Conservation in Ontario. 8 p.
- OMNR (Ontario Ministry of Natural Resources). 2010a. Forest Management Guide for Conserving Biodiversity at the Stand and Site Scales. Queen's Printer for Ontario, Toronto, Ontario. 211 p.
- OMNR (Ontario Ministry of Natural Resources). 2010b. Forest Management Guide for Great Lakes – St. Lawrence Landscapes. Queen's Printer for Ontario, Toronto, Ontario. 57 p.
- OMNRF (Ontario Ministry of Natural Resources and Forestry). 2016. Hunting Notices and Updates. Website: <https://www.ontario.ca/page/hunting-notice-and-updates#section-4> [accessed November 2016].
- OMNRF (Ontario Ministry of Natural Resources and Forestry). 2018a. Land Information Ontario. Website: <https://www.ontario.ca/page/land-information-ontario>.
- OMNRF (Ontario Ministry of Natural Resources and Forestry). 2018b. Ontario's parks and protected areas. Website: <https://www.ontario.ca/page/ontarios-parks-and-protected-areas>.
- OMNRF (Ontario Ministry of Natural Resources and Forestry). Unpub. data. Trapper harvest data for wolf and coyote in Ontario – seasons 2004-2005 to 2015-2016.
- ONHIC (Ontario Natural Heritage Information Centre). 2017. Data on Eastern Wolf/Algonquin Wolf.
- Otis, J.-A., D. Thornton, L. Rutledge, and D.L. Murray. 2017. Ecological niche differentiation across a wolf-coyote hybrid zone in eastern North America. *Diversity and Distributions* 23:529-539.
- Packard, J.M. 2003. Wolf behavior: reproductive, social, and intelligent, *in* L.D. Mech and L. Boitani (eds.). *Wolves – behavior, ecology, and conservation*. The University of Chicago Press, Chicago, Illinois. Pp. 35-65.
- Parks Canada. 2007. Étude sur les attitudes, perceptions et connaissances des résidents de la région limitrophe du parc national du Canada de la Mauricie à l'égard du loup de l'Est. Rapport final, March 2007.
- Patterson, B.R., and D.L. Murray. 2008. Flawed population viability analysis can result in misleading population assessment: a case study for wolves in Algonquin Park, Canada. *Biological Conservation* 141:669-680.

- Patterson, B.R. Pers. comm. 2009. Communication with John Benson and Karen Loveless. Ontario Ministry of Natural Resources, Wildlife Research and Development Section, Peterborough, Ontario, Canada.
- Patterson, B.R. 2009. Unpub. data. Provided by the Ontario Ministry of Natural Resources, Wildlife Research and Development Section, Peterborough, Ontario, Canada.
- Patterson, B.R. Pers. comm. 2012. E-mail correspondence with L. Rutledge concerning the draft status report on the Eastern Wolf. Ontario Ministry of Natural Resources, Wildlife Research and Development Section, Peterborough, Ontario, Canada.
- Peterson, R.O., and P. Ciucci. 2003. The wolf as a carnivore, *in* L.D. Mech and L. Boitani (eds.). *Wolves: behavior, ecology and conservation*. The University of Chicago Press, Chicago, Illinois. Pp. 104-130.
- Pimlott, D.H. 1967. Wolf predation and ungulate populations. *American Zoologist* 7:267-268.
- Pimlott, D.H., J.A. Shanon, and G.B. Kolenosky. 1969. The ecology of the timber wolf in Algonquin Provincial Park. Fish and Wildlife Report No. 87. Ontario Ministry of Natural Resources, Peterborough, Ontario. 92 p.
- Popp, J.N., and V.M. Donovan. 2016. Fine-scale tertiary-road features influence wildlife use: a case study of two major North American predators. *Animal Biology* 66:229-238.
- Potvin, F. 1986. Écologie du loup dans la réserve de Papineau-Labelle. Ministère du Loisir, de la Chasse et de la Pêche du Québec, Direction générale de la faune. Quebec City, Quebec. 103 p.
- Potvin, F. 1987. Wolf movements and population dynamics in Papineau-Labelle reserve in Quebec. *Canadian Journal of Zoology* 66:1266-1273.
- Potvin, F., H. Jolicoeur, and J. Huot. 1988. Wolf diet and prey selectivity during two periods for deer in Quebec: decline versus expansion. *Canadian Journal of Zoology* 66:1274-1279.
- Proulx, G., M.R. Cattet, and R.A. Powell. 2012. Humane and efficient capture and handling methods for carnivores, *in* L. Boitani and R.A. Powell (eds.). *Carnivore ecology and conservation: a handbook of techniques*. Oxford University Press, London, UK. p. 70-129
- Proulx, G., D. Rodtks, M.W. Barrett, M. Cattet, D. Dekker, E. Moffatt, and R.A. Powell. 2015. Humaneness and Selectivity of Killing Neck Snares Used to Capture Canids in Canada: A Review. *Canadian Wildlife Biology and Management* 4(1):55-65.

- Quinn, N.W.S. 2005. Reconstructing changes in abundance of White-tailed Deer, *Odocoileus virginianus*, Moose, *Alces alces*, and Beaver, *Castor canadensis*, in Algonquin Park, Ontario, 1860-2004. *Canadian Field-Naturalist* 119:330-342.
- Rateaud, W., H. Jolicoeur, and P. Etcheverry. 2001. Habitat du loup dans le sud-ouest du Québec: occupation actuelle et modèles prédictifs. *Société de la faune et des parcs du Québec*. 56 p.
- Rempel, R.S. 2011. Effects of climate change on moose populations: exploring the response horizon through biometric and systems models. *Ecological Modelling* 222:3355-3365.
- Ripple, W.J., and R.L. Beschta. 2004. Wolves and the ecology of fear: Can predation risk structure ecosystems? *Bioscience* 54:755-766.
- Ripple, W.J., J.A. Estes, R.L. Beschta, C.C. Wilmers, E.G. Ritchie, M. Hebblewhite, J. Berger, B. Elmhagen, M. Letnic, M.P. Nelson, O.J. Schmitz, D.W. Smith, A.D. Wallach, and A.J. Wirsing. 2014. Status and Ecological Effects of the World's Largest Carnivores. *Science* 343:1241484.
- Rogic, A., N. Tessier, and F.-J. Lapointe. 2014. Identification of canids found within Parc national du Mont-Tremblant and its surroundings using microsatellite markers. University of Montreal and Ministère des Forêts, de la Faune et des Parcs. Report submitted to the head of the Service de la conservation et de l'éducation du Parc national du Mont-Tremblant. Montreal, Quebec. 30 p.
- Roy, M.S., E. Geffen, D. Smith, E.A. Ostrander, and R.K. Wayne. 1994. Patterns of differentiation and hybridization in North American wolflike canids, revealed by analysis of microsatellite loci. *Molecular Biology and Evolution* 11:553-570.
- Rutledge, L.Y. 2010a. Evolutionary origins, social structure, and hybridisation of the eastern wolf (*Canis lycaon*). PhD thesis, Trent University, Peterborough, Ontario, Canada. 187 p.
- Rutledge, L.Y., B.R. Patterson, and B.N. White. 2010b. Analysis of *Canis* mitochondrial DNA demonstrates high concordance between the control region and ATPase genes. *BMC Evolutionary Biology* 10:215.
- Rutledge, L.Y., C.J. Garroway, K.M. Loveless, and B.R. Patterson. 2010c. Genetic differentiation of eastern wolves in Algonquin Park despite bridging gene flow between coyotes and grey wolves. *Heredity* 105:520-531.
- Rutledge, L.Y., K.I. Bos, R.J. Pearce, and B.N. White. 2010d. Genetic and morphometric analysis of sixteenth century *Canis* skull fragments: implications for historical eastern and grey wolf distribution in North America. *Conservation Genetics* 11:1273-1281.

- Rutledge, L.Y., B.R. Patterson, K.J. Mills, K.M. Loveless, D.M. Murray, and B.N. White. 2010e. Protection from harvesting restores the natural social structure of Eastern Wolf packs. *Biological Conservation* 143:332-339.
- Rutledge, L.Y., B.N. White, J.R. Row, and B.R. Patterson. 2011. Intense harvesting of eastern wolves facilitated hybridisation with coyotes. *Ecology and Evolution* 2(1):19-33.
- Rutledge, L.Y., P.J. Wilson, C.F.C. Klütsch, B.R. Patterson, and B.N. White. 2012. Conservation genomics in perspective: a holistic approach to understanding *Canis* evolution in North America. *Biological Conservation* 155:186-192.
- Rutledge, L.Y., and B.N. White. 2013. Genetic characterization of *Canis* in Québec. Short Form Contract No. 111720000. Final Report. 10 p.
- Rutledge, L.Y., and B.N. White. 2014. Genetic characterization of *Canis* in Québec. Short Form Contract No. 111720000. Final Report Addendum. 14 p.
- Rutledge, L.Y., S. Devillard, J.Q. Boone, P.A. Hohenlohe, and B.N. White. 2015. RAD sequencing and genomic simulations resolve hybrid origins within North American *Canis*. *Biology Letters* 11(7): 20150303.
- Rutledge, L.Y., G. Desy, J.-M. Fryxell, K. Middel, B.N. White, and B.R. Patterson. 2016. Patchy distribution and low effective population size raise concern for an at-risk top predator. *Diversity and Distributions* 2016:1-11.
- Sazatornil, V., A. Rodríguez, M. Klaczek, M. Ahmadi, F. Álvares, S. Arthur, J.C. Blanco, B.L. Borg, D. Cluff, Y. Cortés, E.J. García, E. Geffen, B. Habib, Y. Iliopoulos, M. Kabi, M. Krol, L. Llaneza, F. Marucco, J.K. Oakleaf, D.K. Person, H. Potočník, N. Ražen, H. Rio-Maior, H. Sand, D. Unger, P. Wabakken, and J.V. López-Bao. 2016. The role of human-related risk in breeding site selection by wolves. *Biological Conservation* 201:103-110.
- Schmitz, O.J., and G.B. Kolenosky. 1985. Wolves and coyotes in Ontario – morphological relationships and origins. *Canadian Journal of Zoology* 63:1130-1137.
- Sears, H.J., J.B. Theberge, M.T. Theberge, I. Thornton, and G.D. Campbell. 2003. Landscape influence on *Canis* morphological and ecological variation in a coyote-wolf *C. lupus x latrans* hybrid zone, southeastern Ontario. *Canadian Field-Naturalist* 117:589-600.

- Sebbane, A., R. Courtois, and H. Jolicoeur. 2008. Changements de comportement du caribou de Charlevoix entre 1978 et 2001, en fonction des modifications de l'habitat. Quebec Department of Natural Resources and Wildlife, Direction de l'expertise sur les faune et ses habitats, Québec (Quebec). 52 p. + annex.
- SOM Inc. 2007. Étude auprès des visiteurs du parc national de la Mauricie à propos de leurs perceptions, attitudes, comportements et connaissances par rapport au loup. Report submitted to the Quebec Service Centre, Parks Canada. 41 p.
- Steinberg, B. Pers. comm. 2017. Email communication. Coordinator of the Natural Heritage Education program, Ontario Ministry of Natural Resources and Forestry, Ontario Parks, Ontario, Canada.
- Stronen, A.V., R.K. Brook, P.C. Paquet, and S. Mclachlan. 2007. Farmer attitudes toward wolves: implications for the role of predators in managing disease. *Biological Conservation* 135:1-10.
- Stronen, A.V. 2009. Dispersal in a plain landscape: wolves in southwestern Manitoba, Canada. PhD thesis, University of New Brunswick, Fredericton, New Brunswick, Canada. 243 p.
- Stronen, A.V., G.J. Forbes, T. Sallows, G. Goulet, M. Musiani, and P.C. Paquet. 2010. Wolf body mass, skull morphology and the mitochondrial DNA haplotypes in the Riding Mountain National Park region of Manitoba, Canada. *Canadian Journal of Zoology* 88:496-507.
- Stronen, A.V. Pers. comm. 2011. E-mail communication with Nathalie Tessier. Department of Biological Sciences, University of Montreal, Montreal, Quebec.
- Stronen, A.V., G.J. Forbes, P.C. Paquet, G. Goulet, T. Sallows, and M. Musiani. 2012a. Dispersal in a plain landscape: short-distance genetic differentiation in southwestern Manitoba wolves, Canada. *Conservation Genetics* 13(2):359-371.
- Stronen, A.V., N. Tessier, H. Jolicoeur, P.C. Paquet, M. Hénault, M. Villemure, B.R. Patterson, T. Sallows, G. Goulet, and F.-J. Lapointe. 2012b. Canid hybridization: contemporary evolution in human-modified landscapes. *Ecology and Evolution* 2:2128-2140.
- Sutton, A.O., D. Strickland, and D.R. Norris. 2016. Food storage in a changing world: implications of climate change for food-caching species. *Climate Change Responses* 3:12.
- Tennier, H. 2008. Lignes directrices pour la prévention et la gestion des loups familiers au parc national du Mont-Tremblant, parc national du Mont-Tremblant, Société des établissements de plein air du Québec. Lac Supérieur, Quebec. 53 p.

- Tennier, H. Pers. comm. 2017. Comments submitted to the Species at Risk Public Registry during the 60-day public comment period. Parc national du Mont-Tremblant, Société des établissements de plein air du Québec, Lac Supérieur, Quebec, Canada.
- Tessier, N. 2015. Unpub. data. Quebec Department of Forests, Wildlife and Parks (MFFP).
- Theberge, J.B., G.J. Forbes, I.K. Barker, and T. Bollinger. 1994. Rabies in wolves of the Great Lakes Region. *Journal of Wildlife Diseases* 30:563-566.
- Theberge, J.B., and M.T. Theberge. 1998. Wolf country: Eleven years tracking the Algonquin wolves. McClelland & Stewart Inc., Toronto, Ontario. 306 p.
- Theberge, J.B., and M.T. Theberge. 2004. The wolves of Algonquin Park: a 12-year ecological study. Publication No. 56 of the Department of Geography, University of Waterloo, Waterloo, Ontario. 163 p.
- Theberge, J.B., M.T. Theberge, J.A. Vucetich, and P.C. Paquet. 2006. Pitfalls of applying adaptive management to a wolf population in Algonquin Provincial Park, Ontario. *Environmental Management* 37:451-460.
- Thiel, R.P., and A.P. Wydeven. 2011. Eastern Wolf (*Canis lycaon*) Status Assessment Report –Covering East-Central North America. Report submitted to the United States Fish and Wildlife Service. Tomah and Park Falls, Wisconsin. 81 p.
- Thompson, I.D., M.D. Fannigan, B.M. Wotton, and R. Suffling. 1998. The effects of climate change on landscape diversity: an example in Ontario Forests. *Environmental Monitoring and Assessment* 49:213-233.
- TNS Inc. 2007a. Étude sur les attitudes, perceptions et connaissances des résidents de la région limitrophe du parc national de la Mauricie à l'égard du loup de l'Est. Report submitted to the Quebec Service Centre, Parks Canada. 88 p.
- TNS Inc. 2007b. Étude sur l'évaluation des dimensions humaines des chasseurs de la région limitrophe du parc national de la Mauricie à l'égard de la conservation du loup de l'Est. Report submitted to the Quebec Service Centre, Parks Canada. 81 p.
- Trapp, J.R. 2004. Wolf den site selection and characteristics in the northern Rocky Mountains: A multi-scale analysis. Master's thesis, Prescott College, Prescott, Arizona. 63 p.
- Vander Vennen, L.M., B.R. Patterson, A.R. Rodgers, S. Moffatt, M.L. Anderson, and J.M. Fryxell. 2016. Diel movement patterns influence daily variation in wolf kill rates on moose. *Functional Ecology* 30:1568-1573.

- Villemure, M., and M. Festa-Bianchet. 2002. Écologie du loup au parc national de La Mauricie. Report prepared for Parks Canada. Department of Biology, University of Sherbrooke. Sherbrooke, Quebec. 130 p.
- Villemure, M. 2003. Écologie et conservation du loup dans la région du parc national de la Mauricie. Master's thesis, University of Sherbrooke, Sherbrooke, Quebec, Canada. 103 p.
- Villemure, M., and D. Masse. 2004. La conservation du loup (*Canis lupus*) au parc national de la Mauricie. *Le Naturaliste Canadien* 128:35-42.
- Villemure, M., and M. Festa-Bianchet. 2004. Écologie du loup au parc national du Canada de la Mauricie. Report prepared for Parks Canada. Department of Biology, University of Sherbrooke. Sherbrooke, Quebec. 118 p.
- Villemure, M., and D. Masse. 2005. Stratégie de conservation du loup dans la région du parc national de la Mauricie. Prepared for the Resource Conservation Service, La Mauricie National Park. 64 p.
- vonHoldt, B.M., J.P. Pollinger, D.A. Earl, J.C. Knowles, A.R. Boyko, H. Parker, E. Geffen, M. Pilot, W. Jedrzejewski, B. Jedrzejewska, V. Sidorovich, C. Greco, E. Randi, M. Musiani, R. Kays, C.D. Bustamante, E.A. Ostrander, J. Novembre, and R.K. Wayne. 2011. A genome-wide perspective on the evolutionary history of enigmatic wolf-like canids. *Genome Research* 21:1294-1305.
- vonHoldt, B.M., J.A. Cahill, Z. Fan, I. Gronau, J. Robinson, J.P. Pollinger, B. Shapiro, J. Wall, and R.K. Wayne. 2016. Whole-genome sequence analysis shows that two endemic species of North American wolf are admixtures of the coyote and gray wolf. *Science Advances* 2:e1501714.
- Vucetich, J., and P. Paquet. 2000. The demographic population viability of Algonquin wolves. Report prepared for the Algonquin Park Wolf Advisory Committee. Houghton, Michigan, and Meacham, Saskatchewan. 18 p. + appendix.
- Wallach, A.D., I. Izhaki, J.D. Toms, W.J. Ripple, and U. Shanas. 2015. What is an apex predator? *Oikos* 124:1453-1461.
- Way, J.G., L. Rutledge, T. Wheeldon, and B.N. White. 2010. Genetic Characterization of Eastern "Coyotes" in Eastern Massachusetts. *Northeastern Naturalist* 17(2):189-204.
- Way, J.G., and J.T. Bruskotter. 2012. Additional considerations for gray wolf management after their removal from Endangered Species Act protections. *Journal of Wildlife Management* 76:457-461.
- Wayne, R.K., and S.M. Jenks. 1991. Mitochondrial DNA analysis imply extensive hybridization of the endangered red wolf *Canis rufus*. *Nature* 351:565–568.

- Wayne, R.K., and C. Vila. 2003. Molecular genetics studies of wolves, *in* L.D. Mech and L. Boitani (eds.). *Wolves: behavior, ecology and conservation*. The University of Chicago Press, Chicago, Illinois. Pp. 218-238.
- Wheeldon, T. 2009. Genetic characterization of *Canis* populations in the Western Great Lakes regions. Master's thesis, Trent University, Peterborough, Ontario. 113 p.
- Wheeldon, T., and B.N. White. 2009. Genetic analysis of historic western Great Lakes region wolf samples reveals early *Canis lupus/lycaon* hybridization. *Biology Letters* 5:101-104.
- Whittington, J., M. Hebblewhite, N.J. DeCesare, L. Neufeld, M. Bradley, J. Wilmshurst, and M. Musiani. 2011. Caribou encounters with wolves increase near roads and trails: a time-to-event approach. *Journal of Applied Ecology* 48:1535-1542.
- Wilmers, C.W., R.L. Crabtree, D.W. Smith, K.M. Murphy, and W.M. Getz. 2003a. Trophic facilitation by introduced top predators: grey wolf subsidies to scavengers in YNP. *Journal of Animal Ecology* 72:909-916.
- Wilmers, C.W., D.R. Stahler, R.L. Crabtree, D.W. Smith, and W.M. Getz. 2003b. Resource dispersion and consumer dominance: scavenging at wolf- and hunter-killed carcasses in Greater Yellowstone, USA. *Ecology Letters* 6:996-1003.
- Wilson, P.J., S.K. Grewal, I.D. Lawford, J.N.M. Heal, A.G. Granacki, D. Pennock, J.B. Theberge, M.T. Theberge, D.R. Voigt, W. Waddell, R.E. Chambers, P.C. Paquet, G. Goulet, D. Cluff, and B.N. White. 2000. DNA profiles of the eastern Canadian wolf and the red wolf provide evidence for a common evolutionary history independent of the gray wolf. *Canadian Journal of Zoology* 78:2156-2166.
- Wilson, P.J., S.K. Grewal, T. McFadden, R.C. Chambers, and B.N. White. 2003. Mitochondrial DNA extracted from eastern North American wolves killed in the 1800s is not of gray wolf origin. *Canadian Journal of Zoology* 81:936-940.
- Wilson, P.J., S.K. Grewal, F.F. Mallory, and B.N. White. 2009. Genetic characterization of hybrid wolves across Ontario. *Journal of Heredity* 100 (suppl. 1):S80-S89.
- Wilson, P.J., L.Y. Rutledge, T.J. Wheeldon, B.R. Patterson, and B.N. White. 2012. Y-chromosome evidence supports widespread signatures of three-species *Canis* hybridisation in eastern North America. *Ecology and Evolution* 2(9):2325-2332.
- Wolf Lake Anishinabeg Community and Rosanne Van Schie (Trent University, Faculty of Forestry, Wildlife Ecology Research Lab). Unpub. data. Data collected as part of projects that include genetic analysis of samples of large canids.
- Wright, S. 1931. Evolution in Mendelian populations. *Genetics* 16:97-159.

- Wydeven, A.P., T.K. Fuller, W. Weber, and K. MacDonald. 1998. The potential for wolf recovery in the northeastern United States via dispersal from southeastern Canada. *Wildlife Society Bulletin* 26: 776-784.
- Young, S.P., and E.A. Goldman. 1944. *The Wolves of North America*. American Wildlife Institute, Washington, D.C. 636 p.

Appendix A: Additional Information on the Effects of Trapping on the Eastern Wolf Population in Canada

In Quebec, an analysis of the data on wolves harvested in furbearer management units indicates that 172 wolves on average were harvested each year between 2005-2006 and 2014-2015 (MFFP, unpub. data). During the 1990-1997 period, the estimated trapping harvest rates³⁶ in seven wildlife reserves in Quebec located, at least partly, in the area of occurrence of the Eastern Wolf, ranged from 1.2% (± 0.3 ; Mastigouche Wildlife Reserve) to 74.4% (± 23.8 ; Papineau-Labelle Wildlife Reserve) (Larivière et al. 1998). Furthermore, in the Papineau-Labelle Wildlife Reserve and surrounding area, where documented mortality from 1980 to 1984 was entirely of human origin and 83% of this mortality was attributable to harvesting (permitted in the area bordering the wildlife reserve), the annual survival rate of radio-collared wolves was 64% at a time when commercial trapping was banned in the reserve (Potvin 1987).³⁷ In La Mauricie National Park and surrounding area, where significant parts of wolf territories extend outside the park, the annual mortality rate of radio-collared wolves was 36% between 2000 and 2003, with 90% of these mortalities caused by trapping. Adult mortality totalled two deaths for every 13 wolf-years (15%) compared to eight pups for every nine wolf-years (89%), with most of these resulting from captures outside the park (Villemure 2003). Pups are often more vulnerable to trapping mortality than adults (Mech 1977; Fuller 1989; Jolicœur 1998), and juvenile Eastern Wolves are believed to be particularly vulnerable given that they typically disperse much earlier than what is reported in other wolf populations (Packard 2003; Mills et al. 2008). Since the early 1990s, there has been a downward trend in the number of fur pelts sold in Quebec (Hénault and Jolicoeur 2003; MFFP unpub. data). However, the number of pelts sold is not necessarily a reliable indicator of the number of individuals taken. Given that the Eastern Wolf may be confused with the Eastern Coyote or the Grey Wolf and that the Eastern Wolf is not yet officially recognized by Quebec authorities, it is impossible to determine the proportion of Eastern Wolves in the harvest of wolves or Eastern Coyotes.

In Ontario, according to the numbers reported by trappers, the average number of wolves (all species combined) taken annually by trappers between the 2004-2005 and 2015-2016 seasons was 514, while the average number of coyotes taken annually was 2,598 (OMNRF, unpub. data). In Algonquin Provincial Park and the surrounding townships, where wolf trapping and hunting have been banned since 2001, the proportion of human-caused mortality has fallen from 67% before the ban (1989-1999) to 16% after the ban (2002-2007). The survival rate and population density have not increased, however, owing to an increase in natural mortality rates; nonetheless, the proportion of packs containing unrelated individuals declined significantly, from 80% to 6% (Rutledge et al. 2010e).

³⁶ Percent trapping mortality, calculated from data on reported fur sales and regional estimates of wolf populations (Hénault and Jolicoeur 2003).

³⁷ Potvin (1987) reported that no difference had been observed in annual survival rates between age classes (pups, yearlings and adults (> 24 months)).

Appendix B: Effects on the Environment and Other Species

A strategic environmental assessment (SEA) is conducted on all SARA recovery planning documents, in accordance with the [*Cabinet Directive on the Environmental Assessment of Policy, Plan and Program Proposals*](#).³⁸ The purpose of a SEA is to incorporate environmental considerations into the development of public policies, plans, and program proposals to support environmentally sound decision-making and to evaluate whether the outcomes of a recovery planning document could affect any component of the environment or any of the [*Federal Sustainable Development Strategy*](#)'s³⁹ (FSDS) goals and targets.

Conservation planning is intended to benefit species at risk and biodiversity in general. However, it is recognized that the implementation of management plans may inadvertently lead to environmental effects beyond the intended benefits. The planning process based on national guidelines directly incorporates consideration of all environmental effects, with a particular focus on possible impacts upon non-target species or habitats. The results of the SEA are incorporated directly into the management plan itself, but are also summarized below.

Eastern Wolf habitat conservation measures will probably have a beneficial effect on a number of other species that use similar habitats, particularly species that require extensive forested areas, including the Grey Wolf and the Great Lakes–Boreal Wolf. Conservation measures are also likely to benefit species that feed on the prey of the Eastern Wolf (Pimlott et al. 1969; Kolenosky 1972; Wilmers et al. 2003a, 2003b; Ripple and Beschta 2004). Table 3 provides a non-exhaustive list of species that may benefit from conservation measures targeting the Eastern Wolf.

³⁸ www.canada.ca/en/environmental-assessment-agency/programs/strategic-environmental-assessment/cabinet-directive-environmental-assessment-policy-plan-program-proposals.html

³⁹ www.fsds-sfdd.ca/index.html#/en/goals/


Table 3. Species that may benefit from Eastern Wolf conservation and management measures in the areas where the Eastern Wolf is present

Common Name	Scientific Name	SARA Status
Blue Jay	<i>Cyanocitta cristata</i>	Not at Risk
Common Raven	<i>Corvus corax</i>	Not at Risk
Grey Jay	<i>Perisoreus canadensis</i>	Not at Risk
Bald Eagle	<i>Haliaeetus leucocephalus</i>	Not at Risk
Grey Wolf	<i>Canis lupus</i>	Not at Risk
American Marten	<i>Martes americana</i>	Not at Risk
Fisher	<i>Martes pennanti</i>	Not at Risk
Red Fox	<i>Vulpes vulpes</i>	Not at Risk
Wood Turtle	<i>Glyptemys insculpta</i>	Threatened
Blanding's Turtle	<i>Emydoidea blandingii</i>	Threatened

Although some of the proposed conservation measures have benefits for the environment in general and have a positive impact on other sympatric native species, there could be consequences for species whose requirements differ from those of the Eastern Wolf. For example, the relationship between the Eastern Wolf and its main prey, White-tailed Deer, Moose and Beaver, could be modified. Research indicates that Eastern Wolves attack young and old individuals of White-tailed Deer (Pimlott et al. 1969) and Moose (Loveless 2010) populations to a disproportionate degree, supporting the theory that predation by wolves can be compensatory, reducing the number of old or sick individuals within the population, while healthy breeding animals have a higher survival rate. However, during periods of prey scarcity, this relationship may no longer be compensatory (Potvin et al. 1988; Delguidice et al. 2002). In areas where Moose have been hunted by humans, Eastern Wolves have increased their predation of Moose calves and juveniles (Loveless 2010). In addition, canids present in the extent of occurrence of the Eastern Wolf attack Woodland Caribou (*Rangifer tarandus caribou*), such as in the Laurentides Wildlife Reserve (Sebbane et al. 2008), but it has not yet been confirmed whether the canids are Eastern Wolves, Grey Wolves or Great Lakes–Boreal Wolves. A study on the relationship between wolves and caribou in northern Ontario was undertaken in 2010; the preliminary data suggest that the Grey Wolf is the main predator of Woodland Caribou in this area (Benson, pers. comm. 2011) and that the impact of the Eastern Wolf on Woodland Caribou populations would therefore be very limited. Moreover, certain conservation measures could also have an effect on Eastern Coyote populations present in the area of occurrence of the Eastern Wolf. Some measures leading to a reduction in canid mortality rates could benefit them, while others (e.g., focussed on habitat management) could limit their expansion.

Consequently, it is important that Eastern Wolf habitat management activities be planned from an ecosystem perspective and with input from responsible authorities through the development, of multi-species plans, ecosystem-based recovery strategies and area management plans that take into account the needs of the various species, including other species at risk. Many stewardship and habitat improvement activities undertaken for the benefit of the Eastern Wolf will be implemented through ecosystem-based management programs that take the needs of species at risk into account.

Appendix C: Season Closures for Hunting and Trapping Wolf and Coyote in Ontario, September 2016



Source: <http://apps.mnr.gov.on.ca/public/files/er/wolf-coyote-season-map.pdf>