



STOCK STATUS UPDATE OF AMERICAN LOBSTER (*HOMARUS AMERICANUS*) IN LOBSTER FISHING AREA 41 (4X + 5ZE) FOR 2020

Context

The status of American Lobster (*Homarus americanus*) in Lobster Fishing Area (LFA) 41 was last assessed in the fall of 2017 (DFO 2018; Cook et al. 2017) with annual updates in the following years. This update applies the suite of indicators from the 2017 assessment to determine the stock status up to the end of the 2020 season. The Northeast Fisheries Science Centre (NEFSC) surveys were not conducted in 2020, and the at-sea observer companies were restricted, or limited, in their operations due to concerns with the COVID-19 global pandemic. The survey data were updated where possible. Indicators for Lobster in LFA 41 are consistent with the Fisheries and Oceans Canada (DFO) precautionary approach and allow for the evaluation and monitoring of the offshore Lobster fishery.

This Science Response Report results from the Science Response Process of October 22, 2020, on the Stock Status Update of American Lobster in Lobster Fishing Area (LFA) 41.

Background

Commercial Lobster fishing in LFA 41 (Figure 1) occurs offshore, from the 50 nautical mile line (92 km) to the upper continental slope. While LFA 41 extends to the easterly boundary of the 4V Northwest Atlantic Fisheries Organization (NAFO) line, the fishery is limited to NAFO Divisions 4X and 5Zc.




Figure 1. Map showing LFA 41 offshore subareas for primary indicators (4X—Crowell Basin, Southwest Browns, and Southeast Browns, and 5Z—Georges Basin and Georges Bank).

The LFA 41 fishery operates under the Offshore Lobster and Jonah Crab Integrated Fisheries Management Plan (DFO 2020). It is the only Lobster fishery in Canada that is managed with a

Maritimes Region

Total Allowable Catch (TAC). The minimum legal size is 82.5 mm Carapace Length (CL), and there is a prohibition on landing berried and/or v-notched females. The fishery operates year-round. Currently, there is no trap limit.

The annual TAC (720 t) was established in 1985 based on historical landings. Annual landings from 2002 to 2020 are presented in Figure 2. In recent years, the TAC has been managed under a three-year management cycle that allows for quota overruns and carry-forward of uncaught quota. At the end of the third year of a cycle, no more than three times the annual quotas (i.e., no more than 2,160 t) may be landed.




Figure 2. Landings (t) for Lobster Fishing Area 41 from 2002–2020 against a Total Allowable Catch of 720 t. Horizontal red line denotes the TAC. Note: Red bar (hash marks) for 2020 landings indicates incomplete data.

Analysis and Response

Indicators of Stock Status

The status of Lobster in LFA 41 is assessed using two indicators of stock health: survey commercial biomass and reproductive potential. The reference points defining the Healthy, Cautious, and Critical zones—the Upper Stock Reference (USR) and the Limit Reference Point (LRP)—are based on the survey biomass. Both indicators use fishery-independent data available from four multispecies surveys, two conducted by DFO and two conducted by NEFSC. The NEFSC surveys were not conducted in 2020 due to concerns with the COVID-19 global pandemic; therefore, the data from these surveys are not updated. Information is instead provided using the 2019 data. One year of missing survey data will have little to no impact on the median because it is a 3-year running median. The DFO Summer Research Vessel Survey (RV41) covers the offshore portions on the Scotian Shelf, and the DFO Spring Research Vessel Survey (GB) covers the offshore portions on Georges Bank. The NEFSC surveys cover the Gulf of Maine and Georges Bank in the spring (NSpr41) and autumn (Naut41).

Primary Indicators and Stock Status

Commercial Biomass from Research Vessel Surveys

Lobster biomass is measured by four multispecies surveys from which commercial biomass indices are used to determine overall stock health. The commercial biomass is calculated for each survey, and a 3-year running median is used to assess stock status relative to reference indicators. The Limit Reference Indicator (LRI) for each index is defined as the median of the five lowest non-zero biomasses in the time series. The Upper Stock Indicator (USI) is defined as 40% of the median of the higher productivity period (i.e., 2000–2015). For the stock to be considered in the Healthy Zone, the commercial biomass indices for at least three of the four surveys must be above their respective USIs (Figure 3). Currently, all four surveys are above their respective USIs. Therefore, the stock is in the Healthy Zone, and it has been since 2002.




Figure 3. Commercial biomass time series along with the 3-year running median (red line), compared to LRI (solid blue line) and USI (dot-dash grey line). Top row: left—RV41, right—GB. Bottom row: left—NSpr41, right—Naut41. Note: Different scales are used on both x-axis and y-axis.

Reproductive Potential

Reproductive potential consists of an integrated index combining female abundance-at-size, fecundity-at-size, and size-at-maturity (Cook et al. 2017). It represents an estimate of total eggs produced within the stock area and can also be viewed as a surrogate for Spawning Stock Biomass (SSB). An Upper Boundary (UB) and Lower Boundary (LB) have been set (where sufficient data are available) to help gauge the significance of changes in egg production relative to long-term medians. Reproductive potential is above the long-term median and the respective UBs in all survey indices. Estimates of reproductive potential are among the highest values on record (Figure 4). An increase in overall abundance was the main driver of the increase in reproductive potential despite a decrease in median size of the Lobsters, as was observed in the at-sea samples and documented during the 2017 stock assessment.




Figure 4. Reproductive potential in millions of eggs estimated from the 4 surveys covering LFA 41 along with the 3-year running median (solid red line). Lower bounds are represented by solid blue lines and upper bounds by dot-dash grey lines. No bounds are identified for the Georges Bank DFO survey, and only upper bounds are identified for the Summer Research Vessel Survey due to the brevity of the time series. Top row: left—RV41, right—GB. Bottom row: left—NSpr41, right—Naut41. Note: Different scales are used on both x-axis and y-axis.

Bycatch

At-sea observer data are aggregated by 3-year time blocks to represent average annual bycatch estimates in LFA 41 (Table 1). Bycatch amounts for crabs (*Cancer* sp.), Cusk, Atlantic Cod, and White Hake have decreased consistently since 2011. Non-retained Lobster catch consists of undersized, berried, v-notched, and potentially cull (one or zero claws), soft, and jumbos (≥ 140 mm CL). The target for number of observed trips is 6 per season for LFA 41. The total number of trips, observed trips, and the percentage of observer coverage are reported in Table 2.

Table 1. The average annual total weight (kg) across 3-year blocks, for the top 6 non-Lobster bycatch and the non-retained Lobster in Lobster Fishing Area 41. These 7 groups represent > 99% of the total bycatch by weight.

Bycatch/non-retained	Annual Average (kg)		
	2011–2013	2014–2016	2017–2019
AMERICAN LOBSTER	132,607	158,221	205,468
CANCER SP.*	24,073	5,622	2,154
CUSK	11,892	5,840	3,122
COD(ATLANTIC)	4,778	4,133	2,778
WHITE HAKE	5,588	4,854	1,603
RED HAKE	1,430	662	1,632
SEA RAVEN	1,107	299	169
HADDOCK	182	150	430

CANCER SP.* is Jonah crab and rock crab combined to account for any misidentification.

Table 2. Number of observed trips per year from 2009 to 2020 for Lobster Fishing Area 41.

Year	Total Number of Trips	Observed Trips	% of Observer Coverage
2009	78	4	5.13
2010	76	3	3.95
2011	51	3	5.88
2012	32	5	15.63
2013	36	6	16.67
2014	35	6	17.14
2015	34	4	11.76
2016	36	6	16.67
2017	34	4	11.76
2018	34	7	20.59
2019	43	5	11.63
2020	34	2	5.88

Conclusions

The primary indicators of stock status for Lobster in LFA 41 show the stock is in the Healthy Zone, with all four multispecies-survey-commercial-biomass indices above their respective USIs. Reproductive potential estimates were also above the upper boundaries where defined. Despite not having a removal reference or estimates of removal rates, the TAC of 720 t poses minimal risk to the stock status falling into the Cautious Zone, as the stock has proven its resilience to this level of removal.

Contributors

Name	Affiliation
Victoria Howse (Lead)	DFO Science, Maritimes Region
Adam Cook	DFO Science, Maritimes Region
Cheryl Denton	DFO Science, Maritimes Region
Rabindra Singh	DFO Science, Maritimes Region
Brad Hubley	DFO Science, Maritimes Region
Kyle Gillespie	DFO Science, Maritimes Region
Verna Docherty	DFO Fisheries Management, Maritimes Region
Brady Stevenson	DFO Fisheries Management, Maritimes Region

Approved by

Alain Vézina
Regional Director of Science, DFO Maritimes Region
Dartmouth, Nova Scotia
Ph. 902-426-3490

Date: November 19, 2020

Sources of information

- Cook, A.M., Cassista Da-Ros, M., and Denton, C. 2017. [Framework Assessment of the Offshore American Lobster \(*Homarus americanus*\) in Lobster Fishing Area \(LFA\) 41](#). DFO Can. Sci. Advis. Sec. Res. Doc. 2017/065. viii + 186 p.
- DFO. 2018. [Assessment of Lobster \(*Homarus americanus*\) in Lobster Fishing Area 41 \(4X + 5Z\) for 2016](#). DFO Can. Sci. Advis. Sec. Sci. Advis. Rep. 2018/004.
- DFO. 2020. [Offshore lobster and Jonah crab – Maritimes Region. Integrated Fisheries Management plans](#).

This Report is Available from the:

Center for Science Advice (CSA)
Maritimes Region
Fisheries and Oceans Canada
PO Box 1006, 1 Challenger Drive
Dartmouth, Nova Scotia
Canada B2Y 4A2

Telephone: 902-426-7070

Fax: 902-426-5435

E-Mail: MaritimesRAP.XMAR@dfo-mpo.gc.ca

Internet address: www.dfo-mpo.gc.ca/csas-sccs/

ISSN 1919-3769

ISBN 978-0-660-38728-4 N° cat. Fs70-7/2021-017E-PDF

© Her Majesty the Queen in Right of Canada, 2021



Correct Citation for this Publication:

DFO. 2021. Stock Status Update for American Lobster (*Homarus americanus*) in Lobster Fishing Area 41 (4X + 5Ze) for 2020. DFO Can. Sci. Advis. Sec. Sci. Resp. 2021/017.

Aussi disponible en français :

MPO. 2021. Mise à jour sur l'état du stock de homard d'Amérique (*Homarus americanus*) de la zone de pêche du homard 41 (4X + 5Ze) en 2020. Secr. can. de consult. sci. du MPO. Rép. des Sci. 2021/017.