

EXTREME HEAT *and* HUMAN HEALTH: Information for Pharmacists and Pharmacist Technicians

HEAT-RELATED Illnesses

As the number of extreme heat events in Canada increases each year due to climate change, pharmacists and pharmacy technicians can expect an increase in heat-related illnesses across the country.

Heat exposure can result in direct illnesses, including heat exhaustion and heat stroke.

Heat exhaustion is the most common, milder form of heat illness with symptoms such as heavy sweating, fatigue, weakness, dizziness, headache, diarrhea, malaise, nausea/vomiting after prolonged exposure to heat.

Heat stroke is a medical emergency. It is a severe form of heat illness that has overlapping symptoms with heat exhaustion and it needs to be treated in hospital.



	Heat EXHAUSTION Symptoms	Heat STROKE Symptoms
Temperature	Core temperature elevated but < 40°C (104°F)	Core temperature elevated but ≥ 40°C (104°F)
Neurological	<ul style="list-style-type: none">Anxiety and confusionDizziness and light-headednessHeadache	Mental status changes: <ul style="list-style-type: none">Delirium/hallucinationsAtaxia (lack of coordination indicating neurological dysfunction)Confusion, irritability, emotional instability, aggressiveness, seizuresLoss of consciousness
Cardiac	<ul style="list-style-type: none">Tachycardia	<ul style="list-style-type: none">Cardiac arrhythmias and tachycardia
Skin	<ul style="list-style-type: none">Cutaneous flushing (hot, red skin)Sweating present	Classic Heat Stroke: <ul style="list-style-type: none">Hot, red, dry skinTypically affects sedentary, vulnerable people Exertional Heat Stroke: <ul style="list-style-type: none">Profuse sweatingTypically associated with high physical activity
Other	<ul style="list-style-type: none">Nausea, vomitingHypotension	<ul style="list-style-type: none">ShockTachypnea (rapid breathing)
Outcome	Untreated and with ongoing heat exposure, heat exhaustion can worsen and become heat stroke.	In later stages, pulmonary edema, hepatic failure, renal failure, rhabdomyolysis (muscle fibre breakdown), and death.


AT-RISK Populations

Many of the risk factors that can increase the susceptibility to heat-related illness and death are interrelated. Some pre-existing conditions are exacerbated by exposure to heat and increase a person's vulnerability to heat-related illness.

Populations *disproportionately* affected by heat include

- Older adults and children
- Those who are pregnant
- Athletes & people who are active in heat (such as outdoor workers)
- People with cardiovascular illnesses (including hypertension), obesity, diabetes, respiratory illnesses, renal disease, neurological disease, and mental health disorders including addictions

- People who are malnourished or dehydrated
- People who have a history of heat illness or an active sunburn
- People living alone & people who are socially isolated
- People with lower socioeconomic status



HEAT RISK FACTORS

Associated with Medications

Various drugs and medications are known to interfere with the body's thermoregulatory mechanisms, predisposing the person to heat illness.

Note: There is no straightforward relationship between the consumption of a drug and a resultant hyperthermic state. Heat stroke may occur in those who are not on any medications. Heat stroke episodes can be due to a physiological condition that predisposes a patient to heat hypersensitivity (e.g., poor hydration), and may not be associated with any drug(s) in question. Secondly, there are various drugs that can induce a hyperthermic state in the absence of extreme environmental heat or exercise.

Any medication with the potential to affect the following can put someone at a higher risk of heat illness:



- The hypothalamus, and the body temperature set-point
- Heat perception, leading to behavioural change (avoidance)
- Cardiac output
- Peripheral vasodilation
- Sweat rate
- Renal function
- Body hydration
- Electrolyte status

Some examples of drug classes include:



- Antipsychotics/ Neuroleptics
- Antidepressants
- Lithium
- Anti-epileptics
- Anti-Alzheimer's agents (Cholinesterase inhibitors)
- Anti-Parkinson's agents
- Diuretics
- Nitrate vasodilators
- ACE inhibitors
- Calcium channel blockers & beta blockers

Prevention of HEAT-RELATED Illnesses

- Drink water before having the sensation of thirst, drink MORE when it is hot
- Lower consumption of alcohol, if possible
- Do not increase consumption of caffeine
- Wear light-coloured, loose-fitting, breathable clothing and wear a hat
- Reduce strenuous activity (e.g., reduce duration & intensity of activity)
- Be active during cooler parts of day or move activities to a cooler area
- Spend more time in air conditioned spaces, use cool showers/baths as alternative if air conditioning is not available
- Mist the skin with cool water to cool down when hot
- Seek medical attention, as needed

Recommendations for Pharmacists and Technicians during EXTREME HEAT EVENTS

- **RECOGNIZE** early signs of heat illnesses and direct patients to appropriate medical care.
- **DISCUSS** safe handling of medications sensitive to heat.
- **IDENTIFY** patients more at risk during extreme heat events.
- **IDENTIFY** medication history or prescribed medications that may exacerbate heat-related conditions.
- **PROVIDE HEALTH CANADA HEAT HEALTH INFORMATION** to patients, especially those prescribed medications that may increase their vulnerability to heat related illnesses. Resources can be found online and ordered at Canada.ca



Extreme Heat Events Guidelines: Technical Guide for Healthcare Workers
www.canada.ca/en/health-canada/services/environmental-workplace-health/reports-publications/climate-change-health/extreme-heat-events-guidelines-technical-guide-health-care-workers.html