

Management Plan for the Horned Grebe (*Podiceps auritus*), Western population, in Canada

Horned Grebe, Western population



2022



Government
of Canada

Gouvernement
du Canada

Canada

Recommended citation:

Environment and Climate Change Canada. 2022. Management Plan for the Horned Grebe (*Podiceps auritus*), Western population, in Canada. *Species at Risk Act* Management Plan Series. Environment and Climate Change Canada, Ottawa. v + 49 pp.

Official version

The official version of the recovery documents is the one published in PDF. All hyperlinks were valid as of date of publication.

Non-official version

The non-official version of the recovery documents is published in HTML format and all hyperlinks were valid as of date of publication.

For copies of the management plan, or for additional information on species at risk, including the Committee on the Status of Endangered Wildlife in Canada (COSEWIC) Status Reports, residence descriptions, action plans, and other related recovery documents, please visit the [Species at Risk \(SAR\) Public Registry](#)¹.

Cover illustration: A pair of Horned Grebes © iStock.com/pum_eva

Également disponible en français sous le titre
« Plan de gestion du Grèbe esclavon (*Podiceps auritus*), population de l'ouest, au Canada »

© His Majesty the King in Right of Canada, represented by the Minister of Environment and Climate Change, 2022. All rights reserved.

ISBN 978-0-660-45554-9

Catalogue no. En3-5/124-2022E-PDF

Content (excluding the illustrations) may be used without permission, with appropriate credit to the source.

¹ www.canada.ca/en/environment-climate-change/services/species-risk-public-registry.html

Preface

The federal, provincial, and territorial government signatories under the [Accord for the Protection of Species at Risk \(1996\)](#)² agreed to establish complementary legislation and programs that provide for effective protection of species at risk throughout Canada. Under the *Species at Risk Act* (S.C. 2002, c.29) (SARA), the federal competent ministers are responsible for the preparation of management plans for listed species of special concern and are required to report on progress within five years after the publication of the final document on the SAR Public Registry.

The Minister of Environment and Climate Change and Minister responsible for the Parks Canada Agency is the competent minister under SARA for the Horned Grebe, Western population, and has prepared this management plan, as per section 65 of SARA. To the extent possible, it has been prepared in cooperation with the governments of Yukon, Northwest Territories, Nunavut, British Columbia, Alberta, Saskatchewan, Manitoba, Ontario, the Gwich'in Renewable Resources Board, the Sahtú Renewable Resources Board, the Wek'èezhii Renewable Resources Board and the Wildlife Management Advisory Council as per section 66(1) of SARA.

Success in the conservation of this species depends on the commitment and cooperation of many different constituencies that will be involved in implementing the directions set out in this plan and will not be achieved by Environment and Climate Change Canada, the Parks Canada Agency, or any other jurisdiction alone. All Canadians are invited to join in supporting and implementing this plan for the benefit of the Horned Grebe, Western population, and Canadian society as a whole.

Implementation of this management plan is subject to appropriations, priorities, and budgetary constraints of the participating jurisdictions and organizations.

² www.canada.ca/en/environment-climate-change/services/species-risk-act-accord-funding.html#2

Acknowledgments

This management plan was prepared by Benoit Laliberté (Environment and Climate Change Canada, Canadian Wildlife Service [ECCC-CWS] – National Capital Region), with the support of Marc-André Cyr (ECCC-CWS – National Capital Region).

This document was drafted with the help of: Christian Artuso (ECCC-CWS – National Capital Region), André Breault (ECCC-CWS – Pacific Region), John Brett (ECCC-CWS – Ontario Region), Kaytlin Cooper (Gwich'in Renewable Resources Board), Katherine Conkin (Government of Saskatchewan), Kiel Drake (Bird Studies Canada – Saskatchewan), Annie Larivière Clermont (ECCC-CWS – National Capital Region), Ann McKellar (ECCC-CWS – Prairie Region), Cynthia Paszkowski (University of Alberta) and Cindy Wood (ECCC-CWS – Northern Region).

Numerous people reviewed the document and provided helpful insight:

M. Archambault (ECCC-CWS – Ontario Region), Angela Barakat (ECCC-CWS – National Capital Region), Stephanie Behrens (Government of Northwest Territories), Suzanne Carrière (Government of Northwest Territories), Diane Casimir (Parks Canada), Ian Cruickshank (Parks Canada), Jim Forbes (British Columbia Ministry of Agriculture), Aimee Guile (Wek'èezhii Renewable Resources Board), Eric Gross (ECCC-CWS – Pacific Region), A. Heatherington (British Columbia Ministry of Environment and Climate Change), David Johns (Alberta Environment and Parks), Cindy Kemper (Alberta Environment and Parks), T. Kohler (Department of National Defence), Shannon Landels (Parks Canada), Colin McDonald (Sahtú Renewable Resources Board), David Moore (ECCC-CWS – Ontario Region), Rosemin Nathoo (Wildlife Management Advisory Council), T. O'Dell (British Columbia Ministry of Agriculture), Cynthia Pekarik (ECCC-CWS – National Capital Region), S. Piquette (Department of National Defense), David Prescott (Alberta Environment and Parks), K. Risto (Department of National Defense), Pam Sinclair (ECCC-CWS – Northern Region), Megan Stanley (ECCC-CWS National Capital Region), J. Steciw (British Columbia Ministry of Environment and Climate Change), Karen Stefanyk (British Columbia Ministry of Environment and Climate Change), Julie Thomas (Government of Yukon), L. Vickers (British Columbia Ministry of Agriculture) and Joanna Wilson (Government of Northwest Territories).

Executive Summary

The Horned Grebe (*Podiceps auritus*) is a waterbird species found in Eurasia and North America. There are two populations in North America: the Western population and a small isolated population in the east (on the Magdalen Islands, Quebec). The Western population, which represents the bulk of the breeding population in Canada and in North America, is the subject of this Management Plan.

The Horned Grebe, Western population, was assessed as Special Concern by the Committee on the Status of Endangered Wildlife in Canada (COSEWIC) in 2009 and was listed in Schedule 1 of Canada's *Species at Risk Act* in 2017. The IUCN Red List³ increased the global rank of the Horned Grebe from Least Concern to Vulnerable in 2015 because of ongoing population declines in North America and Europe. In North America, the Horned Grebe is protected in Canada under the *Migratory Birds Convention Act* and in the United States under the *Migratory Bird Treaty Act*.

The Western population of the Horned Grebe is estimated at between 200,000 and 500,000 individuals. Approximately 92% of the North American breeding range of this population is located in Canada, and most birds are presumed to winter in the United States. While the COSEWIC status report of this population suggested a long-term decline of the continental population on the wintering grounds, a new analysis of the Christmas Bird Count (CBC) data using a hierarchical model⁴ now suggests that the population has been relatively stable since the 1970s and might have been increasing in recent years. However, this continental trend masks regional variations. For example, the number of Horned Grebe wintering in Alaska, and on large inland lakes and reservoirs appears to have increased, while it has been decreasing in British Columbia and in many areas along the east coast. On the breeding grounds, Breeding Bird Survey (BBS) data suggest the species has experienced long-term declines in Canada since 1970, with the most important declines occurring in British Columbia and Saskatchewan. However, the BBS has significant drawbacks when it comes to assessing population trends of wetland species, and lacks sufficient coverage in the boreal forest to properly assess population trends over the entire breeding range of the Horned Grebe, Western population. Additional research and monitoring is required to assess breeding population trends and to understand the connectivity between breeding and wintering grounds.

Because the Horned Grebe, Western population has such a widespread distribution, it faces numerous threats. Wetland loss and degradation occurs throughout the breeding range at the hand of various activities, but is particularly problematic in the Canadian Prairies where a large number of small, shallow wetlands are vulnerable to conversion to agriculture and contamination by pesticides. Additionally, droughts in the Prairies (a likely result of climate change) will further reduce the availability of breeding sites and exacerbate the conversion of wetlands to cropland. As a waterbird, Horned Grebes are

³ International Union for the Conservation of Nature Red List of Threatened Species

⁴ See Link et al. 2006, Sauer and Link 2011 and Soykan et al. 2016

vulnerable to fisheries bycatch and oil spills. Additional threats include eutrophication of nesting sites, diseases (Type E botulism) and collisions with power lines and wind turbines.

The management objective for the Horned Grebe, Western population, is to maintain, over the next 30 years (2022-2052), population levels at or above the average population levels of the past 30 years (1987-2017), and to maintain the population's current distribution in Canada.

The broad strategies and conservation measures identified to achieve these objectives are: i) conserving and restoring Horned Grebe breeding habitat in the Prairies through stewardship programs and best-management practices on privately owned land, ii) conserving and restoring Horned Grebe breeding habitat in the boreal forest through wetland conservation policies and best-management guidelines for natural resources industries, iii) addressing key knowledge gaps regarding threats other than habitat loss, particularly the impact of pesticides and the magnitude of mortality associated with fisheries bycatch, oil spills, diseases and collisions with power lines and wind turbines, iv) understanding the connectivity between breeding and wintering grounds, v) establishing a monitoring program suited to this (and other) wetland species.

Table of contents

Preface.....	i
Acknowledgments	ii
Executive Summary	iii
1. COSEWIC Species Assessment Information.....	1
2. Species Status Information	1
3. Species Information	3
3.1. Species Description	3
3.2. Species Population and Distribution.....	3
3.3. Needs of the Horned Grebe, Western Population	14
4. Threats.....	16
4.1. Threat Assessment	16
4.2. Description of Threats	19
5. Management Objective	28
6. Broad Strategies and Conservation Measures.....	29
6.1. Actions Already Completed or Currently Underway	29
6.2. Broad Strategies	31
6.3. Conservation Measures	32
6.4. Narrative to Support Conservation Measures and Implementation Schedule ..	33
7. Measuring Progress	34
8. References.....	35
Appendix A: Map of Bird Conservation Regions	48
Appendix B: Effects on the Environment and Other Species	49

1. COSEWIC* Species Assessment Information

Date of Assessment: April 2009

Common Name (population): Horned Grebe – Western population

Scientific Name: *Podiceps auritus*

COSEWIC Status: Special Concern

Reason for Designation: Approximately 92% of the North American breeding range of this species is in Canada and is occupied by this population. It has experienced both long-term and short-term declines and there is no evidence to suggest that this trend will be reversed in the near future. Threats include degradation of wetland breeding habitat, droughts, increasing populations of nest predators (mostly in the Prairies), and oil spills on their wintering grounds in the Pacific and Atlantic Oceans.

Canadian Occurrence: Yukon Territory, Northwest Territories, Nunavut, British Columbia, Alberta, Saskatchewan, Manitoba, Ontario

COSEWIC Status History: Designated Special Concern in April 2009. Assessment based on a new status report.

* COSEWIC (Committee on the Status of Endangered Wildlife in Canada)

2. Species Status Information

The North American Horned Grebe (*Podiceps auritus cornutus*) population is divided into two distinct populations: the Western population, which comprises the bulk of the individuals across the continent, and a small long-standing breeding population (approximately 15 adults) on the Magdalen Islands in Quebec. The Western population is the subject of this Management Plan.

The Horned Grebe, Western population, is found breeding in Canada from the Yukon to the extreme northwestern part of Ontario, including British Columbia, Alberta, Saskatchewan, Manitoba, the Northwest Territories and the southern part of Nunavut. Approximately 92% of the Western population's breeding range is in Canada, but extends to Alaska, in the northwest, and to Montana and North Dakota, to the south (Stedman 2020).

The Horned Grebe's global conservation status rank is G5 (Secure). It is also considered Secure in Canada (N5B, N5B and N5M; NatureServe 2018). The statuses in each province and territory are presented in Table 1. The IUCN Red List increased the rank of the global Horned Grebe population from Least Concern to Vulnerable in 2015

because of ongoing population declines in North America and Europe (BirdLife International 2018).

Table 1. NatureServe Status^a for the Horned Grebe and its populations in Canada

Region	Horned Grebe species' status	Magdalen Islands population's Status (Pop. 1) ^b	Western population's Status (Pop. 2) ^b
Global	G5	G5TNR	G5TNR
Canada	N5B, N5N, N5M	NNR	NNR
Alberta	S3B	-	SNR
British Columbia	S4B, SNRN	-	SNR
Labrador	SU	-	-
Manitoba	S3B	-	SNR
New Brunswick	S4N, S4M	-	-
Newfoundland Island	SNA	-	-
Northwest Territories	S3B	-	SNR
Nova Scotia	S4N	-	-
Nunavut	SUB, SUM	-	SNR
Ontario	S1B, S4N	-	SNR
Prince Edward Island	SNA	-	-
Quebec	S1	SNR	-
Saskatchewan	S5B	-	SNR
Yukon Territory	S4B	-	SNR

^a The NatureServe status (rank) is made up of a letter which reflects the spatial level for which the status has been granted (G = global, N = national and S = provincial, state or territorial level). The numbers which follow it refer to the following statuses: 1- critically imperiled; 2- imperiled; 3-vulnerable to extirpation or disappearance; 4- apparently secure; 5- demonstrably widespread, abundant and secure. A breeding code is used when a breeding population and a non-breeding population are found within the same province or territory: B = breeding, N = non-breeding, M = migratory. Finally, the code NR signifies that the status has not yet been assessed. Two ranking values next to each other (e.g. S4S5N) show a range of uncertainty regarding the status of the species for the region.

^b NatureServe acknowledges the presence of two distinct populations (Population 1 being the Magdalen Islands population and Population 2 being the Western population), but a complete assessment was only conducted at the species (global) level in Canada.

In Canada, the Western population was designated as Special Concern by COSEWIC and was listed as such under Schedule 1 of the *Species at Risk Act* (S.C. 2002, c.29) in 2017, while the Magdalen Islands population has been listed as Endangered since 2011. The Horned Grebe is also a migratory bird protected in Canada under the *Migratory Birds Convention Act*, 1994 and in the United States under the *Migratory Bird Treaty Act*. It is designated as a priority species in eleven Bird Conservation Regions^{5,6}

⁵ Bird Conservation Regions or BCR are bird ecoregions developed by the North American Bird Conservation Initiative (NABCI 2019; see Map of BCRs in Appendix A).

⁶ Northwestern Interior Forest, Pacific & Yukon Region (BCR 4), Northern Pacific Rainforest (BCR 5), Boreal Taiga Plains, Prairie & Northern Region (BCR 6), Taiga Shield and Hudson Plain, Prairie & Northern Region (BCR 7), Boreal Softwood Shield, Prairie & Northern Region and Ontario subregions (BCR 8-PNR and BCR 8-ON), Great Basin, Pacific & Yukon Region (BCR 9), Northern Rockies, Pacific & Yukon Region (BCR 10), Prairie Potholes (Prairie & Northern Region (BCR 11), Boreal Hardwood Transition, Ontario and Manitoba Region (BCR 12) and Lower Great Lakes/St. Lawrence Plain (BCR 13).

across Canada. Finally, at the national level, it is listed as a Tier 2 species in Canada's Waterbird Conservation Plan (Environment Canada 2003).

At the provincial and territorial levels, the Horned Grebe has been listed as Special Concern in Ontario under the *Endangered Species Act* since 2009, and it is designated as Special Concern in New Brunswick under the *Species at Risk Act* (O.C. 2013-143). In Alberta, it has the general status of Sensitive, but this does not provide legal protection. In British Columbia, the species is listed on the Yellow List (species that are apparently secure and not at risk of extinction), but this listing does not provide legal protection either.

In Quebec, the Magdalen Islands population is listed as Threatened under the *Loi sur les espèces menacées ou vulnérables* (L.R.Q. c. E-12.01).

3. Species Information

3.1. Species Description

The Horned Grebe is a waterbird weighing between 300 and 570 g (Stedman 2020). Its breeding plumage is characterized by a distinctive patch of bright yellow feathers which extends into tufts behind the eye. Its eyes are red, its neck and flanks are chestnut-red and the back is black. Males and females are similar in colouration. In winter, the plumage is black (back) and white (belly), with the white cheeks contrasting with a black crown.

3.2. Species Population and Distribution

The Horned Grebe has a holarctic⁷ distribution: it is found both in North America and Eurasia and it is represented by a different subspecies in each hemisphere. While the global population is estimated at 239,000–583,000 individuals (Wetlands International, 2012). The Western population of the Horned Grebe is estimated at between 200,000 and 500,000 individuals, while the European subspecies, also known as the Slavonian Grebe (*P. a. auritus*), is estimated at only 12,800-18,400 mature individuals (BirdLife International 2017). Table 2 presents a non-exhaustive list of translations of “Horned Grebe” into a few different Indigenous languages.

⁷ of, relating to, or being the biogeographic region including the northern parts of the Old and the New Worlds and comprising the Nearctic and Palearctic regions or subregions (Merriam-Webster, 2019)

Table 2. Non-exhaustive list of Indigenous names of the Horned Grebe

Name	Language/Origin	Translation	Source
zhingibis	Anishinaabemowin (Ojibwe)	grebe, a hell-diver, [Podiceps spp.; grebe]	Ojibwe People's Dictionary 2021
sihkihp	Nêhiyawêwin (Plains Cree)	grebe	Cree online dictionary 2021
nōtáh	Chipewyan (Dene)	Horned Grebe or Eared Grebe	South Slave Divisional Education Council 2012
tagwaatsik	Teet'it Gwich'in First Nation	Horned Grebe	K. Cooper pers. comm. 2020
s-xwátísh	She shashishalhem	grebe	FirstVoices 2021a
yuuyuučkinł	čišaaʔath	grebe	FirstVoices 2021b
miçuk	Ktunaxa	grebe	FirstVoices 2021c
too dzèè'	Kwadacha Tsek'ene	Eared Grebe or Horned Grebe	FirstVoices 2021d
hiixuudaada	Hlgaagilda Xaayda Kil	Western Grebe but may include Red-necked Grebe and Horned Grebe	FirstVoices 2021e
surilitchiaq	Inuvialuktun	Horned Grebe	K. Cooper pers. comm. 2020

3.2.1 Breeding distribution, abundance and trends

Horned Grebes of the Western population breed south of the treeline from Alaska through the Northwest Territories, and from eastern British Columbia to the northern United States and east to the Ontario border (Sinclair et al. 2003, Stedman 2020). Approximately 92% of the Horned Grebe Western population's breeding range is in Canada (COSEWIC 2009, Figure 1).



Figure 1. Distribution of the Horned Grebe in North America.

In Canada, Horned Grebes breed at different densities across their range. The highest densities (1.5 to 3.3 pairs per km²) occur in the Prairies, particularly in southern Saskatchewan, Alberta and, to a lesser extent, Manitoba (Sugden 1977, COSEWIC 2009; Figure 2). A significant portion of the Horned Grebe's breeding range is also located in the boreal forest of western Canada and Alaska. Although the Horned Grebe is widespread in the boreal forest, it is less common and breeds at lower densities than in the Prairie Parkland Region (Semenchuk 2007, COSEWIC 2009, Mitchell 2018).




Figure 2. Horned Grebe distribution and relative abundance during the breeding season (7 June to 27 July), based on eBird⁸ data from 2005 to 2020 (Fink *et al.* 2020). Light coloring indicate lower relative abundance, while dark coloring indicate higher relative abundance.

In Alberta, Horned Grebes are most often found in the Grassland and Parkland Natural Region and only found occasionally in the Boreal Forest, Foothills and Rocky Mountain Natural Regions (Semenchuk 2007). The second Atlas of the Breeding Birds of Alberta also suggests there has been a range contraction in the northwestern part of the province (Semenchuk 2007).

In Saskatchewan, the Horned Grebe is a common breeder in the Prairie Parkland Region, but less common and localized in the boreal and subarctic regions (Smith 1996, K. Drake pers. comm. 2019).

⁸ eBird is an online database of bird observations submitted through checklists.

In Manitoba, the probability of observing this species is the highest in the Prairie Pothole Region, especially south and west of Riding Mountain National Park (Mitchell 2018). Detections within the boreal biome were few and far between, with the exception of a cluster of sites around The Pas, Churchill and Nueltin Lake (Mitchell 2018).

They are considered common in the southern Yukon Territory (Sinclair et al. 2003) and widespread in the Northwest Territories. Although relatively high breeding densities were recorded near Yellowknife (2.2 pairs/km²), this is probably not representative of the rest of the Northwest Territories, where breeding density is probably less than 0.1 bird/km² overall (Stotts 1988, Fournier and Hines 1999; Figure 2). They breed in small numbers in northern Manitoba up to the border with Nunavut (Mitchell 2018), which suggest that breeding in the southernmost portions of Nunavut is possible. Recent extralimital nesting records from Charlton and Danby islands, in James Bay, Nunavut (Fink *et al.* 2020) suggest that nesting may occur in this region periodically.

In British Columbia, the species breeds in sparse clusters east of the Coast Mountains, with the largest clusters located in the Peace River lowlands, the Cariboo plateau, and the Thompson-Nicola plateau (Howie 2015).

In Ontario, the Horned Grebe is an irregular, even rare breeder in the extreme northwestern parts of the province, near the border with Manitoba. During the surveys for the second Atlas of the Breeding Birds of Ontario, breeding evidence was reported in three locations (Opasquia Provincial Park, Pikangikum Lake and the Rainy River sewage lagoons) (Hoar 2007).

3.2.1.1 Breeding Bird Survey (BBS)

The BBS is designed to detect breeding bird species through standardized roadside surveys conducted primarily by volunteers. Surveys are run along routes that comprise 50 stops, spaced 0.8 km apart. Participants record the total number of individuals of all bird species heard or visually observed during a 3-minute observation period (Government of Canada 2021).

The BBS it is not well-suited to survey wetland species like the Horned Grebe, because BBS routes under-sample wetlands in general, and they are conducted in June, a time of the year when Horned Grebes are incubating and less visible and active vocally. Its coverage is heavily skewed toward the Prairie portion of the breeding range, with relatively few routes in the boreal part of Horned Grebe's distribution. Yet, it provides some information on the species' population trend in Canada.

According to BBS data, the Horned Grebe population has been declining in Canada since 1970 (Figure 3). The probability that this decline is at least 30% is 22%, while the probability that this declines is at least 50% is 2.9 % (Smith et al. 2020).




Figure 3. BBS 10-year rolling trends. Based on trend in 2019: 22% probability of a 30% decrease and 2.9% probability of a 50% decrease (Smith et al. 2020).

BBS data indicate long- and short-term declines in the Canadian Horned Grebe Western population (Table 3, Figure 4). From 1970 to 2019, there was an average annual trend of -1.71% (95% CI -4.56, 0.67), and an overall decline of -57.0% (95% CI -89.9, 38.7). Regionally, trends were most negative in British Columbia (cumulative decline: -81.3%, 95% CI -97.9, -16.0), Manitoba (-67.2%, 95% CI -90.6, 6.1) and Saskatchewan (-58.3%, 95% CI -81.2, -7.89).

Over the most recent three generations (2006-2019), the national rate of decline is -1.11% per year (95% CI -6.05, 4.54), equivalent to -13.5% cumulatively (95% CI -55.5, 78.0). At a regional scale, trends are most negative in British Columbia (cumulative decline -46.8%, 95% CI -98.1, 44.5) and Manitoba (-34.2%, 95% CI -76.3, 41.6), and most positive in Yukon (0.2%, 95% CI -57.0, 173.6) and Alberta (29.1%, 95% CI -29.2, 160.0; Table 3, Figure 4).

Table 3. Short-term (2006-2019) and long-term (1970-2019) population trends for Horned Grebe in Canada, based on Breeding Bird Survey data; bolded trends have 95% credible intervals that do not cross zero and are highly likely to represent a substantial rate of change (Smith et al. 2020).

Region	Annual % Rate of Change (95% lower/upper credible intervals)	Cumulative % Change (95% lower/upper credible intervals)	Probability of decline >30%	Number of routes	Reliability
Short-term (2006-2019)					
Canada	-1.11 (-6.05, 4.54)	-13.5 (-55.5, 78.0)	0.26	161	Low
Alberta	1.99 (-2.62, 7.63)	29.1 (-29.2, 160.0)	0.02	69	Low
British Columbia	-4.73 (-14.59, 2.87)	-46.8 (-87.1, 44.5)	0.69	10	Low
Manitoba	-3.17 (-10.49, 2.71)	-34.2 (-76.3, 41.6)	0.56	18	Low
Northwest Territories	-0.52 (-9.48, 9.88)	-6.6 (-72.6, 240.4)	0.29	4	Low
Saskatchewan	-1.21 (-5.87, 3.56)	-14.6 (-54.5, 57.5)	0.26	47	Low
Yukon	0.01 (-6.28, 8.05)	0.2 (-57.0, 173.6)	0.21	11	Low
Long-term (1970-2019)					
Canada	-1.71 (-4.56, 0.67)	-57.0 (-89.9, 38.7)	0.80	193	Medium
Alberta	-0.66 (-2.35, 1.30)	-27.7 (-68.8, 87.9)	0.47	77	Medium
British Columbia	-3.36 (-7.56, -0.36)	-81.3 (-97.9, -16.0)	0.96	11	Low
Manitoba	-2.25 (-4.70, 0.12)	-67.2 (-90.6, 6.1)	0.90	18	Medium
Northwest Territories	-0.98 (-4.82, 4.12)	-38.3 (-91.1, 621.9)	0.54	5	Low
Saskatchewan	-1.77 (-3.35, -0.17)	-58.3 (-81.2, -7.89)	0.90	67	High
Yukon	-1.23 (-4.58, 2.49)	-45.6 (-90.0, 233.7)	0.61	12	Low




Figure 4. Annual rates of population change estimated over three generations (2006-2019) from Breeding Bird Survey trends at the scale of Bird Conservation Regions within provinces, territories, and states (Smith et al. 2020).

3.2.2 Wintering distribution, abundance and trends

The Horned Grebe, Western population, winters across a vast area at various densities (Figure 5; Meehan et al. 2020). In western North America, it winters in coastal estuaries and bays from Alaska to California, as well as inland on large water bodies and rivers, particularly in Idaho, Oregon and California (Stedman 2020; Figure 5). The highest densities are observed in southern British Columbia, Washington State and in Alaska. In eastern North America, it winters from the Atlantic coast of Nova Scotia and New Brunswick to the Gulf of Mexico coast. It is also found inland on the Great Lakes to central Maryland, eastern Virginia, eastern North Carolina, central South Carolina, northern Georgia, throughout Florida except southern Florida peninsula, Tennessee, and from north-central Arkansas, northern and southeastern Louisiana west through southeastern Oklahoma and northeastern Texas (Stedman 2020; Figure 5).

Based on the Christmas Bird Counts (CBC; Meehan et al. 2020), an average of 48% of Horned Grebes winter on the west coast of the continent, while 35% winter on the east coast (including Florida). Twelve percent (12%) of birds counted were reported in states located on the Gulf of Mexico, and 5% on inland waterbodies of south and southeastern states.




Figure 5. Map of relative abundance for the Horned Grebe, based on CBC data 1966-2019 (Meehan et al. 2020).

The CBC documents winter bird populations through annual surveys within fixed 24 km diameter count circles (Birds Canada 2020). Being a land-based survey, the CBC coverage of coastal waters is limited to the line-of-sight of the observers, so it is not optimal to survey Horned Grebes where they winter further offshore. Given that very few Horned Grebes overwinter in Canada, but 92% of the North American population breeds in Canada, the North American CBC results are considered to provide the best insight into trends of Canadian birds.

Based on the overall CBC trends at the continental scale (results from Canada and the USA combined; Table 4), the population trend has increased by 0.53%/year (95% CI - 0.50, 1.81) between 1970 and 2019. However, the continental trend masks that populations wintering in different areas are fluctuating at different rates and directions. For example, in the western portion of the wintering range, the number of Horned Grebe wintering in Alaska appears to have increased, while it has been decreasing in British Columbia. In the eastern portion of the wintering range, numbers appears to have declined in a number of coastal states (e.g. Florida, Maryland, New Jersey, New York and Virginia; Table 4).

Most 95% credible intervals presented in Table 4 cross zero, so there is a high uncertainty about the actual trend of the population in many regions. The trend in British Columbia, however, does not cross zero. Studies conducted in British Columbia examining coastal waterbird trends documented a statistically significant decrease in the Salish Sea of -2.60% per year (1999 – 2011; Crewe *et al.* 2012), and a non-significant increase in the same region over a longer temporal period (1999 – 2019; Ethier *et al.* 2020). There appears to have been a slight shift in wintering distribution since the 1970s and the Horned Grebe could winter more frequently inland than previously (Figure 6; Meehan *et al.* 2020, Stedman 2020), although the relative abundance of this species is generally much lower inland than along the coast.

Table 4. Long-term (1970-2019) and short-term (2009-2019) population trends for Horned Grebe in North America, based on CBC data (Meehan *et al.* 2020). Bolded trends have 95% credible intervals that do not cross zero and are highly likely to represent a substantial rate of change.

Geographic area	Long-term trend (1970-2019)			Short-term trend (2009-2019)			Abundance Index ^a (2019)
	%/year	Lower CI	Upper CI	%/year	Lower CI	Upper CI	
Continental (Canada and USA combined)	0.53	-0.50	1.81	1.36	-1.78	5.12	35.42
Canada	-1.58	-2.31	-0.81	-1.74	-4.91	1.00	3.02
British Columbia	-1.66	-2.41	-0.85	-1.90	-5.39	1.10	2.73
Nova Scotia	-1.57	-2.61	-0.48	-1.80	-4.34	0.00	0.10
New Brunswick	0.49	-1.76	2.74	0.37	-4.09	4.35	0.08
Ontario	1.71	0.83	2.80	4.87	-4.40	12.19	0.04

Geographic area	Long-term trend (1970-2019)			Short-term trend (2009-2019)			Abundance Index ^a (2019)
	%/year	Lower CI	Upper CI	%/year	Lower CI	Upper CI	
United States	0.96	-0.28	2.24	1.74	-1.70	5.76	32.30
Alabama	-0.07	-1.24	1.15	-0.06	-3.81	4.02	1.24
Alaska	1.40	-0.35	3.16	1.95	-3.42	9.07	10.45
Arizona	2.63	-0.66	6.67	7.49	-9.16	30.67	< 0.01
Arkansas	0.77	-0.81	2.33	0.69	-7.18	11.55	0.24
California	-0.18	-1.41	0.72	-0.58	-5.08	4.30	0.60
Colorado	-0.97	-5.45	2.25	-0.85	-8.05	9.64	0.03
Connecticut	-2.37	-3.20	-1.54	-2.52	-6.36	0.46	0.01
Delaware	-2.20	-3.38	-1.02	-1.90	-5.28	3.65	0.01
Florida	-2.08	-2.81	-1.29	-2.30	-5.90	0.16	0.63
Georgia	2.88	1.70	4.03	1.61	-5.01	7.39	0.35
Idaho	5.15	2.92	7.52	4.06	-4.23	8.83	1.91
Illinois	4.51	2.73	6.78	-1.70	-13.81	12.89	0.03
Indiana	3.97	2.50	5.38	1.62	-5.34	7.22	0.08
Kansas	5.71	1.77	10.22	3.54	-13.03	14.97	0.03
Kentucky	6.90	5.43	8.43	6.77	2.06	11.75	0.58
Louisiana	-0.93	-2.21	0.35	-7.55	-17.13	-1.07	0.08
Maine	-1.47	-2.39	0.11	-1.31	-3.83	1.05	0.63
Maryland	-1.95	-2.68	-1.21	14.15	6.29	22.83	0.06
Massachusetts	0.41	-0.26	1.07	-6.54	-11.64	-1.69	0.05
Michigan	0.89	-0.31	2.05	-9.27	-16.14	-2.25	0.04
Minnesota	-4.40	-9.78	0.26	-8.51	-39.72	1.40	< 0.01
Mississippi	-0.27	-1.38	0.89	-10.03	-17.33	-2.85	0.13
Missouri	3.70	1.52	5.94	2.56	-9.76	9.40	1.83
Montana	7.89	5.36	10.90	4.08	-9.70	11.29	0.08
Nevada	0.93	-3.14	5.41	0.97	-4.80	7.15	0.10
New Jersey	-3.35	-4.36	-0.52	-1.36	-6.67	6.44	0.01
New Mexico	-2.93	-6.52	0.66	-9.01	-32.05	17.31	< 0.01
New York	-1.18	-1.76	-0.61	-2.99	-7.77	1.05	0.07
North Carolina	-0.02	-0.77	0.71	5.93	-0.16	12.26	0.51
Ohio	1.68	0.73	2.62	4.54	-3.42	12.86	0.02
Oklahoma	3.51	1.43	5.29	10.50	1.69	27.57	0.62
Oregon	0.36	-0.45	1.16	-0.67	-5.33	2.08	0.15
Pennsylvania	-0.25	-1.62	1.07	9.18	-3.89	23.98	0.03
Rhode Island	-0.64	-1.91	0.67	-0.49	-3.03	2.83	0.10
South Carolina	0.95	-0.28	2.52	3.43	-1.88	9.07	0.45
Tennessee	1.46	-0.01	2.86	2.04	-1.27	5.66	0.48
Texas	-1.17	-2.37	0.21	-4.26	-10.36	1.66	0.23
Utah	23.32	6.63	44.10	14.59	-10.31	40.98	0.33
Vermont	-1.20	-2.62	0.24	-1.41	-6.18	2.14	0.03
Virginia	-2.12	-2.91	-1.22	6.58	0.72	13.74	0.11
Washington	0.60	-2.77	1.80	1.01	-1.54	3.29	3.72
West Virginia	-0.38	-2.33	1.62	-5.01	-19.75	13.58	0.01
Wisconsin	3.19	0.86	5.89	0.14	-15.81	18.63	< 0.01

^a Abundance index allows for comparison of relative abundance between different geographic areas. These are derived from CBC observations and corrected for different parameters, such the effect of observer effort (Meehan et al. 2020)




Figure 6. Map of Horned Grebe wintering population trends between 1966 and 2019, based on CBC data (Meehan et al. 2020). Regions are delimited by administrative and Bird Conservation Region boundaries. Dots represent CBC circles where Horned Grebes were detected and black areas had no Horned Grebe data.

3.2.3 Migration

After breeding, Horned Grebes move to larger lakes to undergo molt (Stout and Cooke 2003). Adult Horned Grebes molt wing-feathers simultaneously, which results in a flightless period of a few weeks, usually between July and October (Stedman 2020). Molting locations are largely unknown. Most birds molt away from breeding areas (Stout and Cooke 2003), but a few adults will molt on the breeding grounds (André Breault, CWS-Pacific Region pers. comm.).

During migration, Horned Grebes stop on large lakes, rivers and wetlands. In Ontario, large numbers of birds can be observed on the Great Lakes, with peaks around mid-October and mid-April (highest count in April: 3,000; Kirk 2014).

More research connecting breeding and wintering grounds is required and very few bands have been recovered for this species (Figure 7).




Figure 7. Horned Grebe banding and band recovery locations (from Dunn et al. 2009). Squares represent band recovery locations for each individual.

3.3. Needs of the Horned Grebe, Western Population

3.3.1 Breeding season

The Horned Grebe breeds in small (generally 0.5 to 2 ha, but ranging from 0.24 to 18.2 ha), shallow (at least 20 cm deep, but on average 40 cm), and usually fishless, perennial wetlands, but they can also nest on larger lakes with shallow edges and sufficient emergent vegetation. Breeding sites usually contain at least 40% open water with beds of emergent vegetation, such as sedges (*Carex* spp.), rushes (*Juncus* spp.) and cattails (*Typha* spp.) (Faaborg 1976, Kuczynski et al. 2012, Routhier 2012, Stedman 2020).

Horned Grebes are territorial and are usually solitary nesters (Palmer 1962), but occasionally, more than one pair and even loose colonies will occur on larger waterbodies with highly productive habitats (Fjeldså 1973, Sugden 1977, Stedman 2020). Horned Grebes are also opportunistic in their selection of a breeding site and will readily nest in human-created habitat (Fournier and Hines 1999, Hoar 2007, Kuczynski et al. 2012).

Horned Grebes generally breed in their first year and the generation time⁹ is 4 years (COSEWIC 2009). During the breeding season, Horned Grebes will feed mainly on aquatic and some airborne arthropods (Stedman 2020). The chicks are semi-precocial¹⁰. They leave the nest hours after hatching and are looked after by the adults, which feed and carry them on their back up to 14 days after hatching (Stedman 2020).

⁹ Average age of parents in the population

¹⁰ Hatched with eyes open, covered with down, and capable of leaving the nest soon after hatching (they can walk and often swim), but stay at the nest and are fed by parents (Ehrlich et al. 1988)

3.3.2 Migration and wintering periods

Horned Grebes migrate mostly at night (Stedman 2020) and will stop on large lakes, rivers and wetlands. During severe winters or storms, Horned Grebes might become stranded and be forced to dry land. They can also land in areas they mistakenly believe to be water bodies (e.g. mining tailing ponds, solar farms).

In winter, they are mostly found at sea, near coastlines and in bays along the Atlantic, Pacific and Gulf of Mexico coasts (del Hoyo et al. 1992, Stedman 2020), generally south of 38° N, where average January temperature is warmer than -1° C (Kirk 2014). They will sometimes winter on lakes (Godfrey 1986). On their wintering grounds, diet shifts to fish, crustaceans (especially amphipods and crayfish – at least in North America) and polychaetes¹¹ (Stedman 2020).

¹¹ A class of annelid worms, generally marine.

4. Threats

4.1. Threat Assessment

The Horned Grebe, Western population's threat assessment uses the IUCN-CMP (International Union for the Conservation of Nature –Conservation Measures Partnership) unified threat classification system. Threats are defined as the proximate activities or processes that have caused, are causing, or may cause in the future the destruction, degradation, and/or impairment of the entity being assessed (population, species, community, or ecosystem) in the area of interest (global, national, or subnational). Limiting factors are not considered during this assessment process. Historical threats, indirect or cumulative effects of the threats, or any other relevant information that would help understand the nature of the threats are presented in detail in section 4.2 Description of Threats.

Table 5. Threat Assessment Calculator Summary

Threat #	Threat description	Impact ^a	Scope ^b (next 10 Yrs)	Severity ^c (10 Yrs or 3 Gen.)	Timing ^d
1	Residential & commercial development	Negligible	Negligible (<1%)	Slight (1-10%)	High (Continuing)
1.1	Housing & urban areas	Negligible	Negligible (<1%)	Slight (1-10%)	High (Continuing)
2	Agriculture & aquaculture	Medium	Large (31-70%)	Moderate (11-30%)	High (Continuing)
2.1	Annual & perennial non-timber crops	Medium	Large (31-70%)	Moderate (11-30%)	High (Continuing)
2.3	Livestock farming & ranching	Negligible	Negligible (<1%)	Negligible (<1%)	High (Continuing)
3	Energy production & mining	Low	Pervasive – Large (31%-100%)	Slight (1-10%)	High (Continuing)
3.1	Oil & gas drilling	Low	Small (1-10%)	Slight (1-10%)	High (Continuing)
3.2	Mining & quarrying	Negligible	Negligible (<1%)	Negligible (<1%)	High (Continuing)
3.3	Renewable energy	Low	Pervasive – Large (31%-100%)	Slight (1-10%)	High (Continuing)

Threat #	Threat description	Impact ^a	Scope ^b (next 10 Yrs)	Severity ^c (10 Yrs or 3 Gen.)	Timing ^d
4	Transportation & service corridors	Low	Pervasive – Large (31%-100%)	Slight (1-10%)	High (Continuing)
4.1	Roads & railroads	Not a Threat	Negligible (<1%)	Neutral or Potential Benefit	High (Continuing)
4.2	Utility & service lines	Low	Pervasive – Large (31%-100%)	Slight (1-10%)	High (Continuing)
5	Biological resource use	Low	Large (31-70%)	Slight (1-10%)	High (Continuing)
5.3	Logging & wood harvesting	Negligible	Negligible (<1%)	Negligible (<1%)	High (Continuing)
5.4	Fishing & harvesting aquatic resources	Low	Large (31-70%)	Slight (1-10%)	High (Continuing)
6	Human intrusions & disturbance	Negligible	Negligible (<1%)	Slight (1-10%)	High (Continuing)
6.1	Recreational activities	Negligible	Negligible (<1%)	Slight (1-10%)	High (Continuing)
7	Natural system modifications	Low	Large (31-70%)	Slight (1-10%)	High (Continuing)
7.1	Fire & fire suppression	Unknown	Unknown	Unknown	High (Continuing)
7.3	Other ecosystem modifications	Low	Large (31-70%)	Slight (1-10%)	High (Continuing)
8	Invasive & problematic species, pathogens & genes	Negligible	Restricted (11-30%)	Negligible (<1%)	High (Continuing)
8.2	Problematic native plants and animals	Negligible	Restricted (11-30%)	Negligible (<1%)	High (Continuing)
8.4	Pathogens & microbes	Negligible	Small (1-10%)	Negligible (<1%)	Moderate (Possibly in the short term < 10 yrs)

Threat #	Threat description	Impact ^a	Scope ^b (next 10 Yrs)	Severity ^c (10 Yrs or 3 Gen.)	Timing ^d
9	Pollution	Medium - Low	Large (31-70%)	Moderate - Slight (1-30%)	High (Continuing)
9.2	Industrial & military effluents	Low	Restricted - Small (1-30%)	Moderate - Slight (1-30%)	High (Continuing)
9.3	Agricultural & forestry effluents	Medium - Low	Large (31-70%)	Moderate - Slight (1-30%)	High (Continuing)
11	Climate change	Medium - Low	Large (31-70%)	Moderate - Slight (1-30%)	High (Continuing)
11.4	Changes in precipitation & hydrological regimes	Medium - Low	Large (31-70%)	Moderate - Slight (1-30%)	Moderate (Possibly in the short term < 10 yrs)
11.5	Severe / Extreme Weather Events	Negligible	Negligible (<1%)	Negligible (<1%)	High (Continuing)

^a **Impact** – The degree to which a species is observed, inferred, or suspected to be directly or indirectly threatened in the area of interest. The impact of each threat is based on Severity and Scope rating and considers only present and future threats. Threat impact reflects a reduction of a species population or decline/degradation of the area of an ecosystem. The median rate of population reduction or area decline for each combination of scope and severity corresponds to the following classes of threat impact: Very High (75% declines), High (40%), Medium (15%), and Low (3%). Unknown: used when impact cannot be determined (e.g. if values for either scope or severity are unknown); Not Calculated: impact not calculated as threat is outside the assessment timeframe (e.g., timing is insignificant/negligible or low as threat is only considered to be in the past); Negligible: when scope or severity is negligible; Not a Threat: when severity is scored as neutral or potential benefit.

^b **Scope** – Proportion of the species that can reasonably be expected to be affected by the threat within 10 years. Usually measured as a proportion of the species' population in the area of interest. (Pervasive = 71–100%; Large = 31–70%; Restricted = 11–30%; Small = 1–10%; Negligible < 1%).

^c **Severity** – Within the scope, the level of damage to the species from the threat that can reasonably be expected to be affected by the threat within a 10-year or three-generation timeframe. Usually measured as the degree of reduction of the species' population. (Extreme = 71–100%; Serious = 31–70%; Moderate = 11–30%; Slight = 1–10%; Negligible < 1%; Neutral or Potential Benefit ≥ 0%).

^d **Timing** – High = continuing; Moderate = only in the future (could happen in the short term [< 10 years or 3 generations]) or now suspended (could come back in the short term); Low = only in the future (could happen in the long term) or now suspended (could come back in the long term); Insignificant/Negligible = only in the past and unlikely to return, or no direct effect but limiting.

4.2. Description of Threats

The Horned Grebe, Western population has a widespread distribution in North America, so it is vulnerable to numerous threats. Wetlands in the Prairies have been severely impacted by the conversion of grassland to cropland and wetland drainage, which has resulted in the permanent loss of small wetlands used for breeding. This habitat loss is further exacerbated in the short-term and medium-term due to potential droughts. In agricultural landscapes, there are also concerns that pesticides might affect small wetlands by contaminating their food chains. In the boreal forest, natural resources industries (oil & gas, mining and forestry) can have negative impacts on wetlands, such as removal of riparian vegetation, changes in water flows and changes in nutrient dynamic.

Direct causes of mortality include diseases (i.e. avian botulism), fisheries bycatch, dry landings, collisions with power lines and wind turbines. On the wintering grounds, oil spills can affect large numbers of grebes.

Considering all these threats, the overall threat impact for the Horned Grebe was scored as “High” (see Master et al. 2012 for a detailed explanation of the methodology).

1.1 Housing & urban areas (negligible)

New residential developments near and around lakes can negatively impact grebes nesting habitat causing permanent habitat loss, habitat modification (e.g. removal of vegetation) and disturbance. In the Canadian prairies, new developments (including new roads, farm infrastructures, housing development and extraction activities) represented a small fraction (<6%) of the total area of wetlands lost between 1985 and 2001 (Watmough and Schmoll 2007). Most ponds where this species breeds are not usually the preferred waterbodies for housing development due to their size and depth, so this threat was considered negligible. Also, a large portion of the Horned Grebe’s breeding distribution is located in areas of relatively low human density, so this threat does not appear as a major driver of habitat loss compared to other activities. However, during dry years in the Prairies, ponds dry up and are at risk of encroachment, particularly near settlements.

In Europe, Horned Grebes will abandon lakes with many summer houses and continuous human activity on the water during summer (Summers et al. 2009, J. Fjeldså, pers. communication *in* Stedman 2020). Kuczynski et al. (2012) also observed that Horned Grebes avoided ponds with human structures located inside ponds (e.g. docks, machinery) in Alberta.

2.1 Annual & perennial non-timber crops (medium)

In the Prairies, wetlands have been impacted severely by conversion of grassland to cropland and wetland drainage (Sugden and Beyersbergen 1984, COSEWIC 2009). The conversion of wetlands to agricultural land is deeply rooted in the landscape of the

Prairies: up to 70% of the wetlands have disappeared in some areas since European settlement (Canadian Wetland Inventory 2008). Currently, more than 90% of wetlands show signs of alteration while the number of remaining unaltered wetlands in the landscape is decreasing (Bartzen et al. 2010). Kuczynski et al. (2012) suggested that small pothole-type wetlands might have disappeared faster from the Prairie landscape than larger, natural wetlands, which might be too large to support Horned Grebes.

Between 1985 and 2001, wetland area in the Prairie Habitat Joint Venture (PHJV) decreased by 5% (228,500 ha of wetlands lost). 77% of wetlands lost were <0.25 ha in area. Wetland basin of >0.26 ha and <1 ha in size accounted for 19% of all complete losses, whereas basins >1 ha accounted for 4% of losses (Watmough and Schmoll 2007). During this period, the most common cover types replacing lost wetland area were annual cultivation crops (62%) and perennial grass (21%; Watmough and Schmoll 2007).

Despite ongoing habitat conservation and restoration efforts, it is estimated that approximately 152,000 acres of wetlands will be lost over the next 10 years in the PHJV area (PHJV 2014a) and temporary and seasonal ponds are the most vulnerable to conversion to agriculture (Bartzen et al. 2010).

Considering the high relative abundance of Horned Grebes in the Prairie Potholes Region and their preference for small (generally <2 ha), shallow wetlands, the threat of habitat loss and modification through agriculture is considered one of the major threats to Horned Grebes.

2.3 Livestock farming & ranching (negligible)

Livestock can impact wetlands in a number of ways. They use wetlands as a source of drinking water, defecating or trampling the surrounding vegetation. Horned Grebes require ponds with emergent and riparian vegetation for nesting (Kuczynski et al. 2012), so by trampling vegetation, livestock might make small ponds unsuitable for the species. Also, by removing riparian vegetation, livestock might increase nutrient and sediment loading into small wetlands (see section 7.3 Other ecosystem modifications).

Although ill-managed livestock can have a negative impact on Horned Grebe habitat, proper and environmentally sound livestock management offers potential benefits for this species. Good practices include, for example, fencing wetlands and maintaining riparian vegetation (Cox and Cullington 2009). Once these practices are adopted, the small ponds used by Horned Grebes are more likely to be retained and protected. Additionally, dugouts and farm ponds are often created to maintain a steady supply of water. The Horned Grebe will readily adopt human-made ponds (Fournier and Hines 1999, Kuczynski et al. 2012), and so dugouts and farm ponds, if properly vegetated and naturalized, can also become potential habitat.

For these reasons, livestock management was scored as negligible and could even become beneficial if environmentally sound livestock management practices and best management practices for dugouts and farm ponds are adopted.

3.1 Oil & gas drilling (low)

Oil & gas development is widespread across the western boreal forest and involves the construction of roads, pipelines, seismic lines, well pads and other infrastructures, all of which can lead to modification and disappearance of wetlands. The degree of overlap between wetlands and these activities is unknown, but is assumed to be considerable given the extent of area affected by anthropogenic activities and the abundance of wetlands in the western boreal forest (PHJV 2014b). Most of the western boreal forest is publicly owned, so industry best-management practices and wetland protection policies should provide a minimal degree of protection to wetlands, although small, shallow and fishless ponds, might, by their nature, be overlooked. Overall, this threat was scored as low given the geographic scale at which it occurs, the population size of the Horned Grebe and the existence of best management practices and wetland conservation policies meant to protect wetlands against destruction by development projects. However, the cumulative effect of industries (forestry, oil & gas, mining & quarrying, transportation corridors, etc.) in the western boreal forest might have a higher global impact on habitat loss (Webster et al. 2015).

3.2 Mining & quarrying (negligible)

Similarly to oil & gas and forestry operations, mining and quarrying activities can lead to habitat loss in the boreal zone (Webster et al. 2015). The modification and drainage of small wetlands can be particularly problematic for the Horned Grebe. Mining and quarrying activities are, however, more localized and not as widespread as other industrial activities and as such was scored as negligible. However, the cumulative effect of industries (forestry, oil & gas, mining & quarrying, transportation corridors, etc.) in the western boreal forest might have a higher global impact on habitat loss (Webster et al. 2015).

3.3 Renewable energy (low)

Grebes in general are vulnerable to collisions with wind turbines due to their low manoeuvrability and low flight altitude (Furness et al. 2013). In Canada, there has been at least one report of a Horned Grebe killed by collision with a wind turbine in Alberta (Bird Studies Canada, Canadian Wind Energy Association, Environment and Climate Change Canada and Ontario Ministry of Natural Resources and Forestry 2018).

Offshore wind turbines can also affect bird populations in other ways than collisions. In Europe, many species, including the Slavonian Grebe, avoid areas with offshore wind turbines (Furness et al. 2013). However, offshore wind turbine development is limited in North America at the moment.

Grebes can also land in solar farms, which they mistake for water bodies (Kagan et al. 2004). This is called “dry landing”, since grebes are unable to easily take flight from land. Once stranded, they die from starvation or predation (Kagan et al. 2004). There is not enough information to assess the magnitude of this impact on Horned Grebe. Because grebes are vulnerable to collisions with wind turbines and dry landing in solar farms and the number of Horned Grebes killed each year is likely underestimated (i.e. not all carcasses are found or identifiable at the species’ level), this threat was scored as “low”.

4.1 Roads & railroads (not a threat)

Transportation corridors, which include roads and railroads, are generally considered as a source of habitat loss and fragmentation. Road building combined with other natural resources developments (e.g. oil & gas, mining and logging) can have a large cumulative impact on wetlands in the boreal region (Webster et al. 2015).

However, the Horned Grebe is known to colonize and successfully raise broods on borrow pits created from soil removal in road construction (Fournier and Hines 1999, Kuczynski et al. 2012). In a study conducted in Alberta (Kuczynski et al. 2012), Horned Grebes occupied 36% of constructed ponds and 74.5%-81.3% of these produced chicks. The borrow pits that were occupied were usually 0.6 ha to 2 ha with more emergent (73%) and riparian vegetation (80%) covering the pond periphery and exempt of beavers. Hence, initial habitat loss during construction might be partially or completely compensated by the creation of new habitat which meets their needs.

For this reason, roads are considered to have a neutral or potentially beneficial impact if proper borrow pit restoration guidelines are adopted. These guidelines should focus on creating ponds that are large and deep enough for Horned Grebes (even during drought years), and include a revegetation program.

4.2 Utility & service lines (low)

Grebes are one of the most vulnerable bird groups to collisions with power lines (Bevanger 1998, APLIC 2012, Rioux et al. 2013), but there is no estimate of Horned Grebe mortality due to these collisions in Canada. Bevanger (1998) conducted a review of literature of 16 investigations of bird collisions with power lines (1972-1993) and reported a total 303 casualties of unspecified grebe species. In the early 1980s, Malcolm (1982) reported important grebe mortality due to collisions with a power line near a wetland in Montana. No Horned Grebe was reported, but Eared Grebes represented 29% of the 3,218 bird mortality detected. However, Horned Grebes have been found dead under powerlines (C. Kemper pers. comm. 2020) and the estimated mortality is likely underestimated (i.e. not all carcasses are found or identifiable at the species’ level).

Because grebes are vulnerable to collisions with power lines and the number of Horned Grebes killed each year is likely underestimated (i.e. not all carcasses are found or identifiable at the species' level), this threat was scored as "low".

5.3 Logging & wood harvesting (negligible)

Similarly to oil & gas and mining operations, forestry operations are widespread across the western boreal forest and overlap greatly with the Horned Grebe's distribution. Forestry operations, which involves building roads, can cause the modification or destruction of wetlands. Most of the western boreal forest is publicly owned, so industry best-management practices and wetland protection policies should provide a minimal degree of protection to wetlands, although small, shallow and fishless ponds, might, by their nature, be overlooked. Although this threat can lead to habitat loss, it was scored as negligible given the geographic scale at which it occurs, the population size of the Horned Grebe in the western boreal forest and the existence of best management practices and wetland conservation policies meant to protect wetlands against destruction by the industry. However, the cumulative effect of industries (forestry, oil & gas, mining & quarrying, transportation corridors, etc.) in the western boreal forest might have a higher global impact on habitat loss (Webster et al. 2015).

5.4 Fishing & harvesting aquatic resources (low)

Because of their diet specialization during migration and on their wintering grounds (i.e. fish), grebes are vulnerable to getting caught and drowning in fishing nets (Riske 1976, Piersma 1988, Ulfvens 1989, Harrison and Robins 1992). The COSEWIC report (2009) mentions that grebes are killed annually on the Great Lakes during both spring and fall migration and that grebe species are caught in gillnet fisheries in California (Mills et al. 2005). Bartonek (1965) estimated that 3,000 grebes (not identified at the species level) and loons were netted annually on the southern part of Lake Winnipegosis, Manitoba. In Europe, the order of magnitude of Slavonian Grebes reported as bycatch¹² in the Baltic and North Seas was in the "tens" of individuals (Zydelis et al. 2009). This represents the mortality of approximately 1% (total wintering population of approximately 1,850), so the number of birds potentially affected in North America could be much higher than what has been reported in Europe, but evidence is lacking because bycatch mortality is not systematically reported.

On the Pacific coast, it is estimated that gill-net fishing is responsible for an annual bycatch of 12,085 (range 1,129–24,002) seabirds, but Horned Grebes were not reported as being taken (Smith and Morgan 2005, Ellis et al. 2013).

Additionally, fishery regulations and reduced fishing efforts have resulted in an overall decline of bycatch mortality of seabirds on the eastern (Regular et al. 2013) and western coasts of North America (Zydelis et al. 2013), which most likely reduced the impact of this threat on Horned Grebe. Although detailed data on Horned Grebe bycatch

¹² Caught accidentally

are lacking, this impact was scored as “low” because there is evidence that the species is vulnerable and individuals likely get caught on a yearly basis.

6.1 Recreational activities (negligible)

In Europe, watercraft disturbance on grebes (including Slavonian/Horned Grebes) has been documented to impact reproductive success either by flushing birds from their nest or causing nest destruction through wave action (Ruddock and Whitfield 2007). This threat is probably limited on their Canadian breeding grounds due to the nature of their preferred habitat (i.e. fishless, small, shallow wetlands) which do not present a lot of interest to recreational boaters or anglers. Disturbance by watercraft is more likely to occur on their staging and wintering areas, particularly after the breeding season when they are molting and are flightless (Stedman 2020). Specific information on the intensity of this disturbance in North America is lacking, but this threat is generally considered marginal and was scored as “negligible”.

7.1 Fire & fire suppression (unknown)

Forest fires can have a profound impact on the boreal forest (e.g. habitat destruction, changes in vegetation, run-offs of sediments, changes in nutrient cycles and hydrological processes). With climate change, frequency and intensity of forest fires is predicted to increase (Amiro et al. 2003). Specific impacts of forest fires on Horned Grebes have not been studied, but the effects are likely cumulative, causing habitat loss and a decrease of the reproductive success. Additional research should provide insight on forest fire’s impact on waterbirds and wetlands (particularly small, shallow wetlands). For this reason, this threat was scored as unknown.

7.3 Other ecosystem modifications (low)

Horned Grebes are exclusively aquatic birds and rely heavily on waterbodies to feed, breed and molt. Any activity that could potentially affect water quality is likely to impact the Horned Grebe. Such activities include agriculture (e.g. fertilizer run-offs), livestock farming and ranching, oil & gas, mining & quarrying, logging & wood harvesting, and the construction of related infrastructure, such as roads, camps, pipelines, well pads, and cut lines. These activities are widespread throughout much of the Horned Grebe’s distribution, and although their cumulative impact is unknown, they can increase nutrient loading (Bayley et al. 2012) by removing riparian vegetation, changing water flows and changing nutrient dynamic.

Horned Grebes use primarily eutrophic environments, so nutrient loading can be beneficial to a certain point by making small ponds more productive and increasing the abundance of macroinvertebrates which grebes eat. However, excessive eutrophication can be detrimental by decreasing water quality (Sánchez-Carrillo et al. 1999, Scheffer et al. 2001). An increase of water turbidity and aquatic plant growth could reduce Horned Grebes’ ability to forage and a decrease the amount of open water, making small ponds unsuitable for breeding.

Because Horned Grebes use small ponds and marshes for breeding, these are generally considered vulnerable to alteration of hydrologic regime and sediment loading due to their shallow depth (Scheffer et al. 2001, Bayley et al. 2012). The scope of this threat was estimated to be large because many activities might impact breeding ponds throughout the species' range, but the severity over the next 3 generations was considered small overall. Hence, this threat was scored as low.

8.2 Problematic native plants & animals (negligible)

Raccoons (*Procyon lotor*) are known nest predators and have greatly expanded their range northwards over the course of the last century. They are now widespread in the Canadian Prairies and in parts of the boreal forest (Larivière 2004, Latham 2008). Changes to habitat, such as conversion to cropland, is a contributing factor to a reduced nest success due to predation (Sovada et al. 2001, Watmough and Schmoll 2007), so it is possible that Horned Grebes are facing an increased predation rate from raccoons.

8.4 Pathogens & Microbes (negligible)

Avian botulism is the most significant pathogens threat to waterfowl and shorebirds (Rocke and Bollinger 2007). Horned Grebes are particularly vulnerable to Type E botulism, which mostly affects fish-eating birds in the Great Lakes in Canada. Type E botulism outbreaks in the Great Lakes were documented for the first time in the United States in 1963 (Rocke and Bollinger 2007) and in Ontario in 1998 (Campbell and Barker 1999, CWHC 2000). During some outbreaks, more than 10,000 fatalities have been reported, with Horned Grebe sometimes in the top 5 species affected (Chipault et al. 2015). Kirk (2014) also reports that “between 2004 and 2013, the USGS National Wildlife Health Center received reports of 660 known (2,304 estimated) dead Horned Grebes recovered on Great Lakes shores and associated with confirmed or suspected botulism type E mortality events in Wisconsin, Michigan, Pennsylvania, New York, and Ontario (J. Chipault, pers. comm. 2014)”.

The species could also be vulnerable to Type C avian botulism (Smith 1977 *cited in* Rocke and Bollinger 2007), but to a lesser extent than Type E. In Canada, Type C mostly affects waterfowl in the Prairies (CWHC 2000). In the database maintained by the Canadian Wildlife Health Cooperative (2019), there are 9 records of botulism Type E in Horned Grebe and 4 records of an “unknown” botulism strain, which could be either Type C or E.

Invasive non-native species which could be intermediate hosts are a contributing factor to mortality associated with avian botulism. Zebra mussels (*Dreissena polymorpha*) and round gobies (*Gobius sp.*) are contributing factors to outbreaks of Type E botulism in Great Lakes fish-eating and mussel-eating birds (CWHC 2000).

Overall, the impact of pathogens & microbes on the Horned Grebe is considered negligible.

9.2 Industrial & military effluents (low)

The increase of natural resources extraction activities leads to a higher risk of contaminant spills occurring in the environment, which could impact Horned Grebes. However, such incidents are likely to affect a small number of birds, since they are usually solitary nesters.

Tailings ponds¹³ are located across internationally important migratory corridors and pose a significant mortality risk for birds, including mass mortality events (Timoney and Ronconi 2010). In 2008, approximately 1,600 ducks were found dead after landing on tailing ponds located on Syncrude Canada's Aurora North tar sands mine (R. v. Syncrude Canada Ltd. 2010). In 2011, a standardized monitoring program called the Oil Sands Bird Contact Monitoring Program (OSBCMP) was created to estimate the number of birds landing in tailing ponds of oil sands mines in Alberta (Ronconi 2011). Horned Grebes were reported as one of the species affected by tailing ponds and the number of Horned Grebes that landed on tailing ponds varied between year (2011-2018) from one to 263 (145 on average). However, the overall number of Horned Grebes that were oiled and died was relatively low. It varied from one individual to a high of nine individuals in 2015 (St. Clair et al. 2012, St. Clair et al. 2013, Owl Moon Environmental Inc 2015, Owl Moon Environmental Inc 2016, Owl Moon Environmental Inc 2017, Hatfield Consultants 2018, Canadian Natural Resources Limited, Canadian Natural Albian Sands, Fort Hills Energy Corporation, Imperial Oil Canada Ltd., Suncor Energy Inc. and Syncrude Canada Ltd. 2019). However, data regarding the non-lethal effects on birds that survive after leaving tailing ponds are lacking.

Because of their higher trophic position in the food chain, grebes are vulnerable to bioaccumulation of contaminants. DDE (dichlorodiphenyldichloroethylene) and PCB (polychlorinated biphenyl) have been reported in Horned Grebe egg shells in Manitoba (Forsyth et al. 1994) and high levels of dioxins and furans were detected in Horned Grebe livers downstream of a pulp and paper plant in British Columbia (Vermeer et al. 1993).

Horned Grebes are also vulnerable to oil pollution on their wintering grounds and have been reported as casualties in many incidents. In the southern USA, 12.3% of 34,717 oiled birds killed in eight oil spills were Horned Grebes (del Hoyo et al. 1992). In 1976, an oil spill in Chesapeake Bay was responsible for the death of more than 4,000 Horned Grebes (Roland et al. 1977). On the Pacific coast, Horned Grebes are also regularly affected by oil spills: 78 oiled Horned Grebes were collected (both alive and dead) in California after the Cosco Busan oil spill (California Department of Fish and Game, 2008), 12 were collected after the Selendang Ayu oil spill in Alaska (Industrial Economics Inc. 2015) and 16 were collected during an oiling episode in the winter of 1997-98 in central California (Hampton *et al.* 2003). Following the Deepwater Horizon Spill along the Gulf Coast, a total of 4 unidentified grebes (2 oiled and 2 not visibly oiled) died (USFWS, 2011).

¹³ In mining facilities, tailings are stored in artificial ponds. Tailings are a mixture of water and other byproducts of the extraction processes.

Overall, the number of Horned Grebe impacted by the oil spills described above is likely underestimated, as individuals may die offshore and may not be recovered. Even birds that do come ashore can die outside the search areas, are difficult to capture until they are dead, and once dead can easily be missed or misidentified by searchers (Ford et al. 2009). The above figure only reflects the direct, short-term impact of the oil spills and does not take into account long term and cumulative effects. For example, the Horned Grebe wintering population had not recovered from the Exxon Valdez spills in Alaska even years after the event (Day et al. 1997, McKnight et al. 2008).

The large wintering area of this species in North America partially protects this population from catastrophic losses due to isolated, localized oil spills (Stedman 2020). For these reasons, the overall impact of Industrial & military effluents was scored as “low”, but mortality from oil spill incidents requires additional and continent-wide monitoring.

9.3 Agricultural & forestry effluents (medium-low)

There are two major types of run-offs that can impact Horned Grebes: fertilizers and pesticides. The former is covered under section 7.3 Other ecosystem modifications. Pesticides have been documented to have a negative impact on invertebrates (Stehle and Schulz 2015) which can eventually impact grebes' productivity.

In an agricultural landscape, pesticides are likely to contaminate some wetlands (through surface runoffs, leaching, spray-drift and wind erosion). The presence of pesticides, such as atrazine and glyphosate (two herbicides), has been documented in Prairie Pothole wetlands (Donald et al. 2001, Anderson et al. 2012, McMurry et al. 2016, Eveltier and Skopec 2018). Although the impact of these herbicides on the Horned Grebe requires further research, they are generally considered as a threat to biodiversity, including birds (Mineau and Palmer 2013).

Among different pesticides used in agriculture, neonicotinoids have received more attention in recent years. First introduced in the 1990s, neonicotinoids are now the most widely used insecticide in the world (Douglas and Tooker 2015). Neonicotinoids are persistent insecticides that have the propensity to integrate water systems and can have negative impacts on aquatic invertebrates (Mineau and Palmer 2013, Anderson et al. 2015, Morrissey et al. 2015). This class of pesticide is widely used in the Prairies (Main et al. 2014), and although its impact on Horned Grebe has not been studied yet, it might contribute to a reduction of invertebrate prey availability, as well as contributing to sub-lethal effects on the species, such as a decrease in reproductive output (Main et al. 2014). Since information regarding how Horned Grebe are affected by pesticides is lacking, there is some uncertainty regarding the overall impact of this threat and it was scored as “low to medium”.

11.4 Changes in precipitation & hydrological regimes (medium-low)

Climate change is a complex phenomenon that is generally expected to lead to a warmer climate and changes in precipitation patterns. A climate change scenario where the combined effects of changes in temperature and precipitation lead to higher evapotranspiration would reduce the persistence of shallow wetlands in the Prairie Potholes Region (Millett et al. 2005, Werner et al. 2010), thus posing a significant threat to Horned Grebes. In fact, Horned Grebes already respond to wet/dry cycles in the Prairie Potholes Region, where the May pond index is positively correlated with the BBS Annual Index (Figure 4).

There is still considerable uncertainty regarding impacts of climate change on wetlands in western Canada, but some cumulative impacts are foreseeable. A warmer and drier climate could increase the frequency and intensity of forest fires, cause the melting of the permafrost and the dessication of wetlands (Cheskey et al. 2011). Additionally, droughts are a contributing factor to shallow wetlands conversion to cropland (Bartzen et al. 2010). This impact was scored as “low to medium” because of the uncertainty about the climate trend over the next 10 years, but recognizing that it can have a significant impact on habitat.

11.5 Severe / Extreme Weather Events (negligible)

Severe weather conditions can force Horned Grebes to land in areas where they are not able to take off again (e.g. dry landing). Stranded birds are frequently reported during migration and severe winters, but some “mass” dry landing episodes can occur following particularly severe storms, as shown in three documented cases in which 68, 75 and 124 individuals were forced to dry land (Hodgdon 1979, Bell 1980, Eaton 1983). Storms will also create waves that can flood nests or kill adults (i.e. hail). Storm frequency (e.g. episodes of hail and tornadoes) are predicted to increase, although information is lacking to predict if this will have a significant impact on the population. Based on current information, this threat was considered to be negligible.

5. Management Objective

The management objective for the Horned Grebe, Western population, is to maintain, over the next 30 years (2022-2052), population levels at or above the average population levels of the past 30 years (1987-2017), and to maintain the population’s current distribution in Canada.

The Horned Grebe, Western population, was designated as Special Concern because of population declines and the many threats it faces, particularly habitat loss and degradation. However, the latest CBC analysis suggests a relatively stable population trend at the continental scale since the 1970s. Nonetheless, this global trend masks the fact that populations wintering in different areas are fluctuating at different rates and directions: while the abundance of birds wintering along the West Coast has increased

over the years, it declined along the East Coast until the mid-2000s and increased subsequently. The combined net effect is a relatively stable trend, with a slight increase in recent years.

On the breeding grounds, BBS data suggest that the Canada-wide decline is ongoing, and that it is steeper in the Prairie Potholes Region. However, this decline might be related to wet/dry cycles, which might reduce the number of small, shallow ponds available for breeding in dry years (or increase their availability in wet years). Also, BBS has, by its design, significant limitations to evaluate population trends of wetland species. This is particularly true in the northern parts of the Horned Grebe's breeding range where BBS coverage is limited.

Considering i) that the CBC continental trend masks different regional trends, such as the decline and subsequent increase that occurred along the East coast, and ii) that Horned Grebe abundance in the Prairie Potholes Region is affected by climate conditions and the availability of ponds; the population objective was established using population level over the past 30 years to provide a more robust baseline (i.e. which encompasses potential cycles) against which to assess progress. At the moment, and considering the significant limitations of the BBS to establish trends for the species in the boreal forest, the CBC appears to be the most reliable source of information available against which to assess progress.

Although the CBC data suggests a stable long-term continental trend, the species remains vulnerable to a number of threats, such as habitat loss and degradation, contamination of wetland food chains by pesticides, mortality by fisheries bycatch, oil spills and collisions with power lines and wind turbines. Habitat loss is mostly due to conversion to agriculture in the Prairie Potholes Region and, to a lesser extent, to natural resources development in the boreal forest. Climate change could exacerbate habitat loss, particularly in the Prairies where, in dry years, semi-permanent wetlands are more vulnerable to being converted to agricultural lands. Additional monitoring is required to assess population trends in the boreal forest, and additional research is required to understand links between breeding and wintering grounds.

6. Broad Strategies and Conservation Measures

6.1. Actions Already Completed or Currently Underway

- A recovery strategy (2013) and an action plan (2015) were developed for the Horned Grebe, Magdalen Islands population. One of the action plan's approach is to support actions targeting the maintenance of the Western population to help ensure it remains abundant and thereby increases the probability of exchanges with the Magdalen Islands population.
- The Horned Grebe, Western population, is surveyed by the major monitoring programs in North America such as the BBS, the CBC and the WBPBS.

Although these programs provide long-term trends, they are not specifically designed for secretive marsh birds. Routhier et al. 2012 suggested that playbacks could be used in the WBPHS ground survey component to increase grebe detection.

- Between 1998 and 2000, 144 adults and 51 young Horned Grebes were banded in the Yellowknife area as part of a Master's thesis by Bonnie Stout (see Stout and Cook 2003).
- The British Columbia Coastal Waterbird Survey has been conducted since 1999 and provides information on trends and abundance of wintering Horned Grebe along British Columbia's coastline.
- In 2007 and 2008, the Canadian Wildlife Service – Northern Region proceeded to the banding of Horned Grebes in the Yellowknife area. 55 adults and 18 young were captured and all banded with a uniquely numbered metal band. Some were also banded with leg color bands (C. Wood, pers. comm. 2020). In 2017, an additional 4 adults were banded (metal band only).
- The Prairie Marsh Monitoring Program, which ran from 2008 to 2012, is a survey dedicated to marsh birds and employs playback call (although the Horned Grebe call was not used; Bird Studies Canada 2018).
- A 2010 report to Parks Canada on the status and distribution of birds and mammals in the Southern Gulf Islands identified Horned Grebe as a priority species for monitoring in the area (Davidson 2010).
- In 2014, the Ontario government released a provincial management plan for the Horned Grebe that identified threats, population objectives and conservation measures for the species (Kirk 2014).
- In 2015, the Ontario government released a Government response statement on the Horned Grebe Management Plan.
- The Horned Grebe, including both Western and Magdalen Island populations, has been identified as a priority species in 11 BCR strategies which established population objectives and conservation measures.
- The Prairie Habitat Joint Venture developed habitat objectives, and conservation programs and partnerships for the Prairies Parklands (PHJV 2014a) and the Western Boreal Forest Regions (PHJV 2014b). These plans establish programs and partnerships that address the threat of habitat loss and degradation for waterfowl species, which will also benefit the Horned Grebe, Western population, across its Canadian breeding range.
- Parks Canada multi-species action plans identify recovery measures specific to species at risk in Parks Canada places. For a list of current multi-species action plans including Horned Grebe, Western population, refer to the documents published for the species on the [Species at Risk Public Registry](#).
- The Canadian Wildlife Health Cooperative actively monitors mortality by diseases in a wide range of bird species including the Horned Grebe.
- Research on habitat use and selection has been conducted at the University of Alberta and data have been gathered in the Aspen Parkland (Moenting et al. unpublished 2009) and the Peace River Parkland (Kuczinski et al. 2009).

- Bird landings and mortality on liquid impoundment facilities in the oil sand region of Alberta are monitored since 2011 through the Canada-Alberta oil sands environmental monitoring and the Oil Sands Bird Contact Monitoring Program.
- Several wetland policies and guidance documents have been published or are underway at the provincial and territorial levels, which might contribute to protecting Horned Grebe habitat:
 - Alberta wetland policy (Alberta Government 2013)
<https://open.alberta.ca/publications/9781460112878>
 - Wetland Ways: Interim Guidelines for Wetland Protection and Conservation in British Columbia (Cox and Cullington 2009)
<https://www2.gov.bc.ca/gov/content/environment/air-land-water/water/water-planning-strategies/wetlands-in-bc>
 - Managing Saskatchewan Wetlands, A Landowner's Guide (Huel 2000)
<http://www.saskh2o.ca/PDF/managingsaskatchewanwetlands.pdf>
 - The Manitoba Water Strategy (Manitoba Conservation 2003)
<https://www.gov.mb.ca/sd/waterstewardship/waterstrategy/pdf/water-strategy.pdf>
 - The Yukon Territory is developing a wetland policy (2018-2019)
<https://online.engageyukon.ca/project/yukon-wetlands>

6.2. Broad Strategies

The broad strategies of this management plan are as follows:

- Habitat conservation and stewardship
- Population monitoring and surveys
- Research

6.3. Conservation Measures

Table 6. Conservation Measures and Implementation Schedule

Conservation Measure	Priority ^e	Threats or Concerns Addressed	Timeline
6.3.1 Broad Strategy: Habitat conservation and stewardship			
a) Empower private landowners, through stewardship programs, to conserve and restore seasonal and semi-permanent wetlands, particularly in the Prairie Potholes Region.	High	IUCN Threat 2.1	2022-2032
b) Adopt environmentally responsible best practice guidelines for crop and livestock management that integrate conservation of seasonal and semi-permanent wetlands.	Medium	IUCN Threat 2.1 and 2.3 and 9.3	2022-2027
c) Support the adoption, implementation and enforcement of best-management practices and wetland conservation policies, particularly for industries (e.g. oil & gas, mining and forestry) in the western boreal forest.	High	IUCN Threat 2.1 and 9.3	2022-2032
d) Develop and adopt breeding habitat restoration guidelines (e.g. borrow pits, farm dugouts and stock ponds), particularly drought resilient habitat.	High	IUCN Threat 3.1 and 4.1	2022-2027
6.3.2 Broad Strategy: Population monitoring and surveys			
a) Establish a long-term monitoring program of wetland bird species to track abundance and habitat use throughout the Horned Grebe's range.	High	All	2022-2027
b) Establish a reporting program that compiles incidents, species and number of individuals affected by oil spills, fisheries bycatch, diseases, dry landing and collisions with power lines and wind turbines.	Low	IUCN Threat 3.3, 4.2, 5.4, 8.4, 9.2 and 11.5	2022-2032
6.3.3 Broad Strategy: Research			
a) Conduct research to understand connectivity between breeding, molting, staging and wintering grounds.	High	All	2022-2032
b) Conduct research to understand impacts of pesticides on wetland bird species.	Medium	IUCN Threat 9.3	2022-2032

^e "Priority" reflects the degree to which the measure contributes directly to the conservation of the species or is an essential precursor to a measure that contributes to the conservation of the species. High priority measures are considered those most likely to have an immediate and/or direct influence on attaining the management objective for the species. Medium priority measures may have a less immediate or less direct influence on reaching the management objective, but are still important for the management of the population. Low priority conservation measures will likely have an indirect or gradual influence on reaching the management objective, but are considered important contributions to the knowledge base and/or public involvement and acceptance of the species.

6.4. Narrative to Support Conservation Measures and Implementation Schedule

The conservation measures for the Horned Grebe, Western population, were developed to address all threats this species is facing, while putting more emphasis on the most important threats and addressing important knowledge gaps.

The Prairie Potholes Region has been identified as a focal region for conservation measures because it is an area of relatively high breeding densities where the most important threats occur (i.e. conversion of breeding habitat to agriculture, vulnerability of breeding habitat due to drier climate and pollution by pesticides).

The most imminent threat to Horned Grebe in the Prairie Potholes Region is the degradation and conversion of wetlands on privately owned lands. Small, shallow wetlands are particularly vulnerable to conversion to agriculture, especially during dry years. The involvement of private land owners is thus crucial to implementing this management plan. Outreach and education regarding the importance of seasonal and semi-permanent wetlands as well as stewardship programs will support and empower private landowners to conserve, restore and create wetlands on their property. Farm dugout and water stock ponds can be used as nesting ponds if they are properly managed. Horned Grebes will readily breed in human-created habitats, such as borrow pits, farm dugout and water stock ponds. Restoration guidelines promoting habitat features preferred by the Horned Grebe (and other aquatic animals), such as maintenance of riparian and emergent vegetation, should be adopted. Also, best management practices for livestock management will also support the protection of small wetlands by maintaining riparian vegetation and limiting disturbance and destruction by livestock. Examples of best management practices for wetlands are listed in section 6.1 *Actions already completed or currently underway*.

Habitat loss is also driven by natural resource development projects in the western boreal forest, where industrial activities (e.g. oil & gas, mining and forestry) largely overlap with the Horned Grebe's distribution. Wetland conservation on the public lands should be guided by larger frameworks which are typically of provincial jurisdiction. Wetland policies should be implemented in collaboration with relevant policy makers and include guidelines and best management practices regarding the conservation and the restoration of wetlands. Examples of best management practices for wetlands are listed in section 6.1 *Actions already completed or currently underway*.

Information about connectivity between breeding and wintering grounds is essential to adopt more targeted conservation measures since populations wintering in different areas are fluctuating at different rates and directions. For example, the reasons behind the decline and subsequent increase of Horned Grebes wintering along the East Coast remain largely unknown. Understanding where these individuals breed would help reduce threats throughout their life cycle.

Because the BBS has significant shortcomings as a tool to assess breeding population size and trends in Horned Grebes, a monitoring program targeting Horned Grebes and other marsh-bird species is recommended to properly assess population trends. This program should be implemented to cover, as much as possible, the entire breeding range of the species.

Finally, a number of secondary threats affecting waterbirds (e.g. oil spills, fisheries bycatch, diseases, collisions with power lines and wind turbines, and dry landing) have been identified, but in all these cases, data are currently collected on a case-by-case basis and seldom compiled at national or continental scales. Hence, the overall understanding of the impact of these threats is incomplete and potentially underestimated. Concerted and integrated monitoring programs are required to monitor these threats in the future.

7. Measuring Progress

The performance indicators presented below provide a way to measure progress towards achieving the management objective and monitoring the implementation of the management plan.

- The indicator of progress for a population that is equal or above the average population level of the past 30 years (1987-2017) is the population trend and abundance index:
 - o The population trend and abundance index of Horned Grebes will be inferred using a combination of available data sources, particularly the CBC, but also the BBS, the WBPHS and other surveys targeting wetland species once they are developed. The population objective was established using population level over the past 30 years to provide a more robust baseline (i.e. one which encompasses potential cycles) against which to assess progress. At the moment, and considering the significant limitations of the BBS to establish trends for the species, the CBC appears to be the most reliable source of information available against which to assess progress. However, a breeding ground monitoring program and more information on connectivity between breeding and wintering grounds are required to better understand the different trends observed on the wintering grounds.

- The indicator of progress for maintaining of the current distribution of the species in Canada (based on 2007-2017 records) is the population's distribution:
 - o The distribution of Horned Grebes in Canada will be measured using a combination of available data sources, including the eBird database, provincial breeding bird atlases, the BBS and other surveys targeting wetland species once they are developed.

8. References

Alberta Government. 2013. Alberta Wetland Policy. Ministry of Environment and Parks. Environment and Sustainable Resource Development. 25p. [online]. Available at <https://open.alberta.ca/publications/9781460112878>. (Accessed in December 2019).

Amiro, B.D., M.D. Flannigan, B.J. Stocks, J.B Todd and B.M. Wotton. 2003. Boreal forest fires: an increasing issue in a changing climate. Paper submitted to the XII World Forestry Congress, 2003 [online]. Available at <http://www.fao.org/3/XII/0207-B3.htm>. (Accessed in May 2019).

Anderson, A.-M., G. Byrtus, J. Thompson, D. Humphries, D. Hill and M. Bilyk. 2012. Baseline pesticide data for semi-permanent wetlands in the Aspen Parkland of Alberta. Prepared for Alberta Environment Water Research User Group, Alberta Environment Ecosystem User Group and Alberta North American Waterfowl Management Plan Partnership. 104 p.

Anderson, J.C., C. Dubetz, V.P. Palace. 2015. Neonicotinoids in the Canadian aquatic environment: A Literature review on current use products with a focus on fate, exposure, and biological effects. *Science of the Total Environment*. 505: 409-422.

Avian Power Line Interaction Committee (APLIC). 2012. Reducing avian collisions with power lines: the state of the art in 2012. Edison Electric Institute and APLIC. Washington, D.C.

Bartonek, J.C. 1965. Mortality of diving ducks on Lake Winnipegosis through commercial fishing. *The Canadian Field-Naturalist*. 79:15-20.

Bartzen, A. B., K.W. Dufour, R.G. Clark and F. D. Caswell. Trends in agricultural impact and recovery of wetlands in prairie Canada. *Ecological Applications* 20(2): 525-538.

Bayley, S.E., A.S. Wong and J.E. Thompson. 2013. Effects of agricultural encroachment and drought on wetlands and shallow lakes in the Boreal Transition Zone of Canada. *Wetlands* 33:17-28

Bell, R. K. 1980. Horned Grebes forced down by ice storm. *Redstart* 47(4): 142–144.

BirdLife International. 2018. *Podiceps auritus*. In: The IUCN Red List of Threatened Species 2018 [online]. Available at www.iucnredlist.org. (Accessed January 2019).

BirdLife International, 2017. European birds of conservation concern: populations, trends and national responsibilities. Cambridge, UK: BirdLife International.

Bird Studies Canada. 2018. The Prairie Marsh Monitoring Program [online]. <https://www.bsc-eoc.org/volunteer/ppmmp/>. (Accessed in March 2019).

Bird Studies Canada, Canadian Wind Energy Association, Environment and Climate Change Canada and Ontario Ministry of Natural Resources and Forestry. 2018. Wind Energy Bird and Bat Monitoring Database. Summary of the Findings from Post-construction Monitoring Reports [online].

https://www.birdscanada.org/resources/wind/2018_Database_Summary_Report.pdf. (Accessed in October 2019).

Campbell, D.G. and Barker, I.K. 1999. Botulism Type E in fish-eating birds, Lake Erie and Lake Huron. CWHC Newsletter. 6(3)7-8.

Canadian Wildlife Health Cooperative (CWHC). 2019. Database query in May 2019. Canadian Wildlife Health Cooperative National Office. Saskatoon, SK.

Canadian Wildlife Health Cooperative (CWHC). 2000. Type E Botulism in Birds [online]. Available at http://www2.cwhc-rcsf.ca/wildlife_health_topics/botulism/botulisme_org.php. (Accessed in February 2019).

Canadian Natural Resources Limited, Canadian Natural Albian Sands, Fort Hills Energy Corporation, Imperial Oil Canada Ltd., Suncor Energy Inc. and Syncrude Canada Ltd. 2019. Oil sands bird contact monitoring program. 2018 Annual report. Submitted to the Alberta Energy Regulator. 241 p.

California Department of Fish and Game. 2008. Natural Resource Damage Assessment for the Cosco Busan Oil Spill [online]. Available at nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=17518 (Accessed in March 2019).

Calvert, A. M., C. A. Bishop, R. D. Elliot, E. A. Krebs, T. M. Kydd, C. S. Machtans, and G. J. Robertson. 2013. A synthesis of human-related avian mortality in Canada. *Avian Conservation and Ecology* 8(2): 11.

Cheskey, E., J. Wells, S. Casey-Lefkowitz. 2011. Birds at Risk. The Importance of Canada's Boreal Wetlands and Waterways. Natural Resources Defense Council, Boreal Songbird Initiative and Nature Canada. 32p.

Chipault, J.G., C.L. White, D.S. Blehert, S.K. Jennings, S.M. Strom. 2015. Avian botulism type E in waterbirds of Lake Michigan, 2010-2013. *Journal of Great Lakes Research*. 41(2): 659-664.

Committee on the Status of Endangered Wildlife in Canada (COSEWIC). 2009. COSEWIC assessment and status report on the Horned Grebe *Podiceps auritus*, Western population and Magdalen Islands population, in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa. vii + 42 pp.
(https://www.sararegistry.gc.ca/virtual_sara/files/cosewic/sr_horned_grebe_0809_e.pdf)

Cox, R.K. and J. Cullington. 2009. Wetland Ways: Interim guidelines for wetland protection and conservation in British Columbia, Chapter 4: Grazing. Wetland Stewardship Partnership. 17p [online]. Available at <https://www2.gov.bc.ca/gov/content/environment/air-land-water/water/water-planning-strategies/wetlands-in-bc> (Accessed in August 2019).

Cree online dictionary. 2021. Online Cree Dictionary [online]. Available at <http://www.creedictionary.com/mobile/> (Accessed in September 2021).

Crewe, T., K. Barry, P. Davidson and D. Lepage. 2012. Coastal waterbird population trends in the Strait of Georgia 1999–2011: Results from the first 12 years of the British Columbia Coastal Waterbird Survey. *British Columbia Birds*. Vol. 22 (2012).

Davidson, P., R.W. Butler, A. Couturier, S. Marquez and D. Lepage. 2010. Status and distribution of birds and mammals in the Southern Gulf Islands, British Columbia. Bird Studies Canada, Pacific Wildlife Foundation and Parks Canada. Unpublished report. 120 pp.

Day, R.H., S.M. Murphy, J.A. Wiens, G.D. Hayward, E.J. Harner and L.N. Smith. 1997. Effects of the Exxon Valdez oil spill on habitat use by birds in Prince William Sound, Alaska. *Ecological Applications*, 7(2): 593-613.

Del Hoyo, J., A. Elliot and J. Sargatal. 1992. Handbook of the Birds of the World; Ostrich to Ducks. Lynx Edicions, Barcelona. 696 p.

Donald, B.D., N.P. Gurprasad, L. Quinnet-Abbott and K. Cash. 2001. Diffuse geographic distribution of herbicides in northern prairie wetlands. *Environmentak Toxicology and Chemistry*. 20(2): 273-279.

Douglas, M.R. and J.F. Tooker. 2015. Large-scale deployment of seed treatments has driven rapid increase in use of neonicotinoid insecticides and preemptive pest management in U.S. field crops. *Environmental Science & Technology* 49: 5088-5097.

Drought Research Initiative. 2011. The 1999-2005 Canadian Prairies drought: Science, impacts and lessons. R. Stewart and R. Lawford (Eds). [online]. Available at <http://www.meteo.mcgill.ca/dri/errata.php> (Accessed in November 2019).

Dunn, E.H., A.D. Brewer, A.W. Diamond, E.J. Woodsworth and B.T. Collins. 2009. Canadian Atlas of Bird Banding, Volume 3: Raptors and Waterbirds, 1921-1995. Special Publication, Canadian Wildlife Service, Environment Canada. 202 pp.

Eaton, S. W. 1983. Horned Grebes downed by ice storm. *American Birds* 37(5): 836–837.

Ehrlich, P.R., D.S. Dobkin, D. Wheye. 1988. Precocial and Altricial Young [online]. Available at

https://web.stanford.edu/group/stanfordbirds/text/essays/Precocial_and_Altricial.html (Accessed in December 2019).

Ellis, J.I., S.I. Wilhelm, A. Hedd, G.S. Fraser, G.J. Robertson, J.-F. Rail, M. Fowler and K.H. Morgan. 2013. Mortality of migratory birds from marine commercial fisheries and offshore oil and gas production in Canada. *Avian Conservation and Ecology* (82): 4.

Environment and Climate Change Canada, 2017. North American Breeding Bird Survey - Canadian Trends Website, Data-version 2015. Environment Canada, Gatineau, Quebec, K1A 0H3.

Environnement Canada. 2003. Wings Over Water, Canada's Waterbird Conservation Plan. Environnement Canada. 27p.

Ethier, D.M., P.J.A. Davidson, G. Sorenson, C. Jardine, D. Lepage, K. Barry, K. Devitt, D.W. Bradley. 2020. Twenty years of coastal waterbird trends suggest regional patterns of environmental pressure in British Columbia, Canada. Manuscript submitted for publication.

Evelsler, V. and M. Skopec. 2018. Pesticides, Including Neonicotinoids, in Drained Wetlands of Iowa's Prairie Pothole Region. *Wetlands*. 38: 221-232.

Faaborg, J. 1976. Habitat selection and territorial behavior of the small grebes of North Dakota. *Wilson Bulletin* 88(3): 390–399.

Ferguson, R. S. and S. G. Sealy. 1983. Breeding ecology of the Horned Grebe, *Podiceps auritus*, in southwestern Manitoba. *Canadian Field-Naturalist*. 97:401-408.

Fink, D., T. Auer, A. Johnston, M. Strimas-Mackey, O. Robinson, S. Ligocki, W. Hochachka, C. Wood, I. Davies, M. Iliff, L. Seitz. 2020. eBird Status and Trends, Data Version: 2019; Released: 2020. Cornell Lab of Ornithology, Ithaca, New York. <https://doi.org/10.2173/ebirdst.2019>

FirstVoices. 2021a [online]. Available at

<https://www.firstvoices.com/explore/FV/sections/Data/Salish/she%20shashishalhem/she%20shashishalhem/learn/words/3ec2f46c-5d29-43e7-aae2-96166727a8e7> (Accessed in September 2021).

FirstVoices. 2021b. Available at

<https://www.firstvoices.com/explore/FV/sections/Data/c%CC%95i%C5%A1aa%CA%94at%E1%B8%A5/Nuu%C4%8Daan%CC%93u%C9%AB/c%CC%95i%C5%A1aa%CA%94at%E1%B8%A5/learn/words/5160652f-a92f-44b6-b3ed-9b28496e651c> (Accessed in September 2021).

FirstVoices. 2021c [online]. Available at <https://www.firstvoices.com/explore/FV/sections/Data/Ktunaxa/Ktunaxa/Ktunaxa/learn/words/20c19d3f-87a7-4740-aaa4-6a41de768d2c> (Accessed in September 2021).

FirstVoices. 2021d [online]. Available at <https://www.firstvoices.com/explore/FV/sections/Data/Athabasca/Kwadacha%20Tsek'ene/Kwadacha%20Tsek'ene/learn/words/1c1cf8e5-2dc9-4abe-bd33-49299874718c> (Accessed in September 2021).

FirstVoices. 2021e [online]. Available at <https://www.firstvoices.com/explore/FV/sections/Data/Haida/Hlg%CC%B1aagilda%20X%CC%B1aayda%20Kil/Hlg%CC%B1aagilda%20X%CC%B1aayda%20Kil/learn/words/af7db0ce-3ff7-4656-8125-3a7625b18dc8> (Accessed in September 2021). Godfrey, W. E. 1986. The Birds of Canada. Revised edition. National Museum of Natural Sciences, National Museums of Canada, Ottawa. 650 p.

Fjeldså, J. 1973. Antagonistic and heterosexual behaviour of the Horned Grebe *Podiceps auritus*. *Sterna* 12(3):161–217.

Ford, R.G., J.L. Casey, W.A. Williams. 2009. Acute seabird and waterfowl mortality resulting from the M/V Cosco Busan oil spill, November 7, 2007. Final Report. Prepared for: California Department of Fish and Game, Office of Spill Prevention and Response. R.G.Ford Consulting Company, Portland, OR. 54 pp.

Forsyth, D. J., P. A. Martin, K. D. De Smet, M. E. Riske. 1994. Organochlorine contaminants and eggshell thinning in grebes from prairie Canada. *Environmental Pollution* 85:51-58.

Fournier M. A. and J. E. Hines. 1999. Breeding ecology of the Horned Grebe *Podiceps auritus* in subarctic wetlands. Occasional Paper No. 99. Canadian Wildlife Service. 32 p.

Government of Canada. 2021 [online]. Breeding Bird Survey instructions. Available at <https://www.canada.ca/en/environment-climate-change/services/bird-surveys/landbird/north-american-breeding/instructions.html> (Accessed in September 2021).

Graf, M.D. 2009. Literature review on the Restoration of Alberta's Boreal Wetlands affected by oil, gas and in situ oil sands development. Prepared for Ducks Unlimited Canada. 53 p.

Hallmann, C.A., R.P.B. Foppen, C.A.M. van Turnhout, H. de Kroon and E. Jongejans. 2014. Declines in insectivorous birds are associated with high neonicotinoid concentrations. *Nature* 511: 341–343.

Hampton, S.F., R.G. Ford, H.R. Carter, C. Abraham and D. Humple. 2003. Chronic oiling and seabird mortality from the sunken vessel S.S. Jacob Luckenbach in central California. *Marine Ornithology*. 31: 35-41.

Harrison, N. and M. Robins. 1992. The threat from nets to seabirds. *RSPB Conservation Review* 6: 51-56.

Hatfield Consultants. 2018. Oil Sands Bird Contact Monitoring Program. 2017 Regional Report. Prepared for Oil Sands Bird Technical Team, Alberta. 268 p.

Hoar, T. 2007. Horned Grebe, pp. 144-145 in Cadman, M.D., D.A. Sutherland, G.G. Beck, D. Lepage, and A.R. Couturier, eds. *Atlas of the Breeding Birds of Ontario, 2001-2005*. Birds Studies Canada, Environment Canada, Ontario Field Ornithologists, Ontario Ministry of Natural Resources, and Ontario Nature, Toronto, xxii + 706 pp.

Hodgdon, K.Y. 1979. Operation Horned Grebe. *North American Bird Bander* 4(3): 110.

Howie, R. 2015. Horned Grebe in Davidson, P.J.A., R.J. Cannings, A.R. Couturier, D. Lepage, and C.M. Di Corrado (eds.). 2015. *The Atlas of the Breeding Birds of British Columbia, 2008-2012*. Bird Studies Canada, Delta, B.C [online]. Available at <https://www.birdatlas.bc.ca/accounts/speciesaccount.jsp?sp=HOGGR&lang=en> (Accessed in March 2019).

Huel, D. 2000. *Managing Saskatchewan Wetlands, A Landowner's Guide*. Saskatchewan Wetland Conservation Corporation. 68p. [online]. Available at <http://www.saskh2o.ca/PDF/managingsaskatchewanwetlands.pdf> (Accessed in December 2019).

Industrial Economics, Inc. 2015. *Natural Resource Damage Assessment Plan for the M/V Selendang Ayu Oil Spill*. Final Draft. Cambridge, MA. 62 pp.

Johnson, W.C., Werner, B., Guntenspergen, G.R., Voldseth, R.A., Millett, B., Naugle, D.E., Tulbure, M., Carroll, R.W.H., Tracy, J. and C. Olawsky. 2010. Prairie Wetland Complexes as Landscape Functional Units in a Changing Climate. *BioScience* 60(2): 128-140.

Johnson, W.C., Millett, B.V., Gilmanov, T., Voldseth, R.A., Guntenspergen, G.R. and D.E. Naugle. 2005. Vulnerability of Northern Prairie Wetlands to Climate Change. *BioScience* 55(10): 863-872.

Kirk, D. A. 2014. *Management Plan for the Horned Grebe (Podiceps auritus) in Ontario*. Ontario Management Plan Series. Prepared for the Ontario Ministry of Natural Resources and Forestry, Peterborough, Ontario. viii + 32pp.

Kuczynski, E. C., Paszkowski, C. A. and B. A. Gingras. 2012. Horned Grebe habitat use of constructed wetlands in Alberta, Canada. *Journal of Wildlife Management* 76(8): 1694–1702.

Langor, D.W., E.K. Cameron, C.J.K. MacQuarrie, A. McClay, B. Peter, M. Pybus, T. Ramsfield, K. Ryall, T. Scarr, D. Yemshanov, I. DeMarchant, R. Footitt and G.R. Pohl. 2014. Non-native species in Canada's boreal zone: diversity, impacts and risk. Canadian Science Publishing. *Environ. Rev.* 22: 372-420.

Latham, A.D.M. 2008. Evidence of Raccoon, *Procyon lotor*, Range Extension in Northern Alberta. *The Canadian Field-Naturalist.* 122(2): 176-178.

Larivière, S. 2004. Range expansion of raccoons in the Canadian prairies: review of hypotheses. *Wildlife Society Bulletin* 32(3):955-963.

Link, W.A., J.R. Sauer and D. K. Niven. 2006. A hierarchical model for regional analysis of population change using Christmas Bird Count data, with application to the American Black Duck. *The Condor* 108:13-24.

Main, A.R., Headley, J. V., Peru, K. M., Michel, N. L., Cessna, A. J. and C. A. Morrissey. 2014. Widespread use and frequent detection of neonicotinoid insecticides in wetlands of Canada's Prairie Pothole Region. *PLoS ONE* 9(3): e92821.

Malcolm, J. M. 1982. Bird collisions with a power transmission line and their relation to botulism at a Montana wetland. *Wildlife Society Bulletin* 10:297–304.

Manitoba Conservation. 2003. The Manitoba Water Strategy. 10 p. [online]. Available at <http://digitalcollection.gov.mb.ca/awweb/pdfopener?smd=1&did=10676&md=1#:~:text=The%20objective%20of%20Manitoba%27s%20water,to%20existing%20and%20future%20generations> (Accessed in December 2019).

Master, L. L., D. Faber-Langendoen, R. Bittman, G. A. Hammerson, B. Heidel, L. Ramsay, K. Snow, A. Teucher, and A. Tomaino. 2012. NatureServe Conservation Status Assessments: Factors for Evaluating Species and Ecosystem Risk. NatureServe, Arlington, VA.

McKellar, Ann E. 2015. A comparison of the BBS and BPOP for assessing population trends of grebes in prairie Canada. Canadian Wildlife Service. Unpublished report.

McKnight, A., K.M. Sullivan, D.B. Irons, S.W. Stephensen and S. Howlin. Prince William Sound Marine Bird Surveys, Synthesis and Restoration, Exxon Valdez Oil Spill Restoration Project Final Report. 136 pp.

McMurry, S. J.B. Belden, L.M. Smith, S.A. Morrison, D.W. Daniel, B.R. Euliss, N.H. Euliss, B.J. Kensinger and B.A. Tangen. 2016. Land use effects on pesticides in sediments of prairie potholewetlands in North and South Dakota. *Science of Total Environment*. 565: 682-689.

Meehan, T. D., G. S. LeBaron, K. Dale, N. L. Michel, A. Krump, and C. W. Wilsey. 2020. Abundance trends of birds wintering in the USA and Canada from Audubon Christmas Bird Counts, 1966–2019, version 3.0. National Audubon Society, New York, New York, USA. <https://www.audubon.org/conservation/where-have-all-birds-gone>.

Merriam-Webster, 2019. Holarctic [online]. Available at <https://www.merriam-webster.com/dictionary/Holarctic>. (Accessed December 2019).

Mills, K.L., W.J. Sydeman and P.J. Hodum. 2005. Marine Bird Conservation Plan, Chapter 6. Marine Ecology Division, PRBO Conservation Science, CA [online]. Available at http://www.prbo.org/cms/docs/marine/CCS%20Plan_Chpt%206_web.pdf (Accessed in May 2019).

Mineau, P. and C. Palmer. 2013. The impact of the nation's most widely used insecticides on birds. American Bird Conservancy. 96 pp.

Mitchell, L. 2018. Horned Grebe *in* Artuso, C., A. R. Couturier, K. D. De Smet, R. F. Koes, D. Lepage, J. McCracken, R. D. Mooi, and P. Taylor (eds.). *The Atlas of the Breeding Birds of Manitoba, 2010-2014*. Bird Studies Canada. Winnipeg, Manitoba

Moenting, A. L.E. Hamilton, R.M. Corrigan and C.A. Paskowski. 2009. Species richness patterns of aquatic birds in the Buffalo Lake Moraine, Alberta, Canada in relation to pond size, water level, and land-use. Department of Biological Sciences, University of Alberta, Unpublished paper.

NABCI. 2019. Bird conservation regions of Canada [online]. Available at <http://nabci.net/foundation-for-conservation/bird-conservation-regions-of-canada/> (Accessed in December 2019).

National Audubon Society. 2019. The Christmas Bird Count Historical Results [online]. Available at <http://www.christmasbirdcount.org> (Accessed in March 2019).

NatureServe. 2018. NatureServe Explorer: An online encyclopedia of life, version 7.1 [online]. NatureServe, Arlington, Virginia. Available at: <http://explorer.natureserve.org>. (Accessed in February 2019).

Niemuth, N.D. and J. W. Solberg. 2003. Response of Waterbirds to Number of Wetlands in the Prairie Pothole Region of North Dakota, U.S.A. *Waterbirds* 26 (2): 233-238.

Ojibwe People's Dictionary. 2021. The Ojibwe People's Dictionary [online]. Available at <https://ojibwe.lib.umn.edu/main-entry/zhingibis-na> (Accessed in September 2021).

Owl Moon Environmental Inc. 2015. Oil sands bird contact monitoring program. 2014 Annual report. Prepared for Alberta Energy Regulator and Alberta Environment and Sustainable Resource Development. 321 p.

Owl Moon Environmental Inc. 2016. Oil sands bird contact monitoring program. 2015 Annual report. Prepared for Alberta Energy Regulator and Alberta Environment and Sustainable Resource Development. 214 p.

Owl Moon Environmental Inc. 2017. Oil sands bird contact monitoring program. 2016 Annual report. Prepared for Alberta Energy Regulator and Alberta Environment and Sustainable Resource Development. 262 p.

Palmer, R. S. 1962. Handbook of North American Birds, Vol. 1 (Loons through Flamingos). Yale University Press, New Haven. 567 p.

Piersma, T. 1988. Body size, nutrient reserves and diet of Red-necked and Slavonian Grebes *Podiceps grisegena* and *P. auritus* on Lake IJsselmeer, the Netherlands. *Bird Study* 35:13-24.

Prairie Habitat Joint Venture (PHJV). 2014a. Prairie Habitat Joint Venture Implementation Plan 2013-2020: The Prairie Parklands. Report of the Prairie Habitat Joint Venture. Environment Canada, Edmonton, AB.

Prairie Habitat Joint Venture (PHJV). 2014b. Prairie Habitat Joint Venture Implementation Plan 2013-2020: The western boreal forest. Report of the Prairie Habitat Joint Venture. Environment Canada, Edmonton, AB.

Regular, P. W. Montevecchi, A. Hedd, G. Robertson and S. Wilhelm. 2013. Canadian fishery closures provide a large-scale test of the impact of gillnet bycatch on seabird populations. *Biology Letters* 9: 20130088. <http://dx.doi.org/10.1098/rsbl.2013.0088>

Rioux, S., J.-P. L. Savard, and A. A. Gerick. 2013. Avian mortalities due to transmission line collisions: a review of current estimates and field methods with an emphasis on applications to the Canadian electric network. *Avian Conservation and Ecology* 8(2): 7.

Riske, M. E. 1976. Environmental impact upon grebes breeding in Alberta and British Columbia. PhD Thesis, University of Calgary, Calgary, Alberta. 482 p.

Rocke, T.E. and T.K. Bollinger. 2007. Avian botulism in Infectious diseases of wild birds. Thomas, N. J., Hunter, D.B. and C.T. Atkinson (eds). Blackwell Publishing. p. 377-416.

- Roland, J. V., G. E. Moore, and M. A. Bellanca. 1977. The Chesapeake Bay oil spill-February 2, 1976: A case history. *International Oil Spill Conference Proceedings* Vol. 1977(1): 523-527.
- Routhier, D.D. 2012. Spatiotemporal variation in occupancy and productivity of grebes in prairie Canada: estimation and conservation applications. Master's Thesis, University of Saskatchewan, Saskatoon. 113 p.
- Routhier, D.D., K.W. Dufour, M.T. Bidwell, and R.G. Clark. 2014. Surveying populations of breeding grebes in prairie parkland Canada: Estimation problems and conservation applications. *Wildlife Society Bulletin* 38:14–17.
- Roy, C. 2019. Predicted trends of Horned Grebe populations in distinct wintering areas using Christmas Bird Count data. Unpublished analysis.
- Roy, C., S.G. Cumming, E. McIntire, S.M. Slattery. 2018. Population synchrony of dabbling ducks in western North America. *Faculté de foresterie, de géographie et de géomatique and Centre d'étude de la forêt, Université Laval*. Unpublished paper.
- R. v. Syncrude Canada Ltd. 2010. ABPC 229 [online]. <http://www.canlii.org/en/ab/abpc/doc/2010/2010abpc229/2010abpc229.pdf> (Accessed 29 August 2016)
- Sánchez-Carrillo S. D. Angeler, M. Alvarez-Cobelas and R. Sánchez-Andres. 1999. Freshwater Wetland Eutrophication (Chap. 9). In Ansari et al. (Eds) *Eutrophication: Causes, Consequences and Control* (Vol. 1). Springer. 394 p.
- Sauer, J. R. and W. A. Link. 2011. Analysis of the North American Breeding Bird Survey using hierarchical models. *The Auk* 128(1): 87-98.
- Sauer, J. R., D. K. Niven, J. E. Hines, D. J. Ziolkowski, Jr, K. L. Pardieck, J. E. Fallon, and W. A. Link. 2017. *The North American Breeding Bird Survey, Results and Analysis 1966 - 2015*. Version 2.07.2017 USGS Patuxent Wildlife Research Center, Laurel, MD.
- Scheffer, M. S. Carpenter, J.A. Foley, C. Folke and B. Walker. 2001. Catastrophic shifts in ecosystems. *Nature*. 413: 591-596.
- Semenchuk, G. P. 2007. Horned Grebe in *The Atlas of Breeding Birds of Alberta: a second look*. Federation of Alberta Naturalists. Edmonton, Alberta. vii + 626 p.
- Sinclair, P.H., W.A. Nixon, C.D. Eckert, and N.L. Hughes. 2003. *Birds of the Yukon Territory*. UBC Press, Vancouver, British Columbia, Canada.
- Smith, A.C., Hudson, M-A.R. Aponte, V.I., and Francis, C.M. 2020. North American Breeding Bird Survey - Canadian Trends Website, Data-version 2019. Environment and Climate Change Canada, Gatineau, Quebec, K1A 0H3.

Smith, A. R. 1996. Atlas of Saskatchewan Birds. Saskatchewan Natural History Society (Nature Saskatchewan), Special publication No. 22. 456 p.

Smith, J.L. and K.H. Morgan. 2005. An assessment of seabird bycatch in longline and net fisheries in British Columbia. Canadian Wildlife Service, Pacific and Yukon Region. Technical report series No. 401.

South Slave Divisional Education Council. 2012. Dÿne Dédliné Yatié Denínu Kuç Yatié Chipewyan Dictionary. National Library of Canada Publication Data. 369 p.

Sovada, M. A., R. M. Anthony, and B. D. J. Batt. 2001. Predation on waterfowl in arctic tundra breeding areas: A review. Wildlife Society Bulletin 29:6-15.

Soykan, C.U., J. Sauer, J. G. Schuetz, G. S. LeBaron, K. Dale and G. M. Langham. Population trends for North American winter birds based on hierarchical models. Ecosphere 7(5).

St. Clair, C.C., Habib T., Loots, S., Ball, J. and McCallum, C. 2012. 2011 Annual Report of the Regional Bird Monitoring Program for the Oil Sands Region. Department of Biological Sciences, University of Alberta. 167 p.

St. Clair, C.C., Habib T., Loots, S., Ball, J. and McCallum, C. 2013. 2012 Report of the Regional Bird Monitoring Program for the Oil Sands. Department of Biological Sciences, University of Alberta. 60p.

Stedman, S.J. 2020. Horned Grebe (*Podiceps auritus*), version 2.0. The Birds of North America (P.G. Rodewald, ed.) [online]. Cornell Lab of Ornithology, Ithaca, NY. Retrieved from The Birds of North America Online. Available at <https://birdsna.org/Species-Account/bna/species/horgre/introduction> (Accessed in March 2019).

Stehle, S. and R. Schulz. 2015. Agricultural insecticides threaten surface waters at the global scale. Proceeding of the National Academy of Sciences of the U.S.A. 112(18): 5750-5755.

Stout, B.E. and F. Cooke. 2003. Timing and location of wing molt in Horned, Red-necked and Western Grebes in North America. Waterbirds 26:88-93.

Stotts, V.D. 1988. Observation of miscellaneous wildlife during 1980-82 waterfowl breeding ground surveys in Northwest Territories. Draft paper prepared for submission to the Canadian Field Naturalist. Copy on file with CWS, Yellowknife, Northwest Territories.

Summers, R. W., R.A. Mavor and M.H. Hancock. 2009. Correlates of breeding success of Horned Grebe in Scotland. *Waterbirds* 32(2): 265-275.

Sugden, L. G. 1977. Horned Grebe breeding habitat in Saskatchewan parklands. *Canadian Field-Naturalist* 91(4):372-376.

Sugden, L. G., and G. W. Beyersbergen. 1984. Farming intensity on waterfowl breeding grounds in Saskatchewan parklands. *Wildl. Soc. Bull.* 12:22-26.

Timoney, K.P. and R.A. Ronconi 2010. Annual bird mortality in the bitumen tailings ponds in northeastern Alberta, Canada. *Wilson Journal of Ornithology*. 122(3): 569-576.

Ulfvens, J. 1989. Clutch size, productivity and population changes in a population of the Horned Grebe *Podiceps auritus* in an exposed habitat. *Ornis Fennica* 66(2):75-77.

US Fish Wildlife Service (USFWS). 2011. Deepwater Horizon Bird Impact Data from the DOI-ERDC NRDA Database 12 May 2011 [online]. Available at www.fws.gov/home/dhoilspill/pdfs/Bird%20Data%20Species%20Spreadsheet%2005122011.pdf (Accessed in February 2019).

Van der Valk, A. G. 2005. Water-level fluctuations in North American prairie wetlands. *Hydrobiologia* 539:171-188.

Venier, L.A., I.D. Thompson, R. Fleming, J. Malcolm, I. Aubin, J.A. Trofymow, D. Langor, R. Sturrock, C. Patry, R.O. Outerbridge, S.B. Holmes, S. Haeussler, L. De Grandpré, H.Y.H. Chen, E. Bayne, A. Arsenault, and J.P. Brandt. 2014. Effects of natural resource development on the terrestrial biodiversity of Canadian boreal forests. *Environ. Rev.* 22: 457-490.

Vermeer, K., W. J. Cretney, J. E. Elliott, R. J. Norstrom, P. E. Whitehead. 1993. Elevated polychlorinated dibenzodioxin and dibenzofuran concentrations in grebes, ducks, and their prey near Port Alberni, British Columbia, Canada. *Marine Pollution Bulletin* 26: 431-435.

Watmough, M.D. and M.J. Schmoll. 2007. Environment Canada's Prairie & Northern Region Habitat Monitoring Program Phase II: Recent habitat trends in the Prairie Habitat Joint Venture. Technical Report Series No. 493. Environment Canada, Canadian Wildlife Service, Edmonton, Alberta Canada. 135 pp.


Webster, K.L., Beall, F.D., Creed, I.F. and D.P. Kreuzweiser. 2015. Impacts and prognosis of natural resource development on water and wetlands in Canada's boreal zone. *Environmental Review* 23: 78–131.

Žydelis, R., J. Bellebaum, H. Österblom, M. Vetemaa, B. Schirmeister, A. Stipiece, M. Dagys, M. van Eerden and S. Garthe. 2009. Bycatch in gillnet fisheries – An overlooked threat to waterbird populations. *Biological Conservation*. 142: 1269-1281.

Žydelis, R., C. Small and G. French. 2013. The incidental catch of seabirds in gillnet fisheries: a global review. *Biological conservation* 162: 67–88.

Appendix A: Map of Bird Conservation Regions

Terrestrial Bird Conservation Regions



- | | | | |
|---|---------------------------------------|---|---|
| 1. Aleutian/Bering Sea Islands | 18. Shortgrass Prairie | 35. Chihuahuan Desert | 52. Planicie Costera Y Lomerios Humedos Del Golfo De Mexico |
| 2. Western Alaska | 19. Central Mixed Grass Prairie | 36. Tamaulipan Brushlands | 53. Sierra Madre Del Sur |
| 3. Arctic Plains And Mountains | 20. Edwards Plateau | 37. Gulf Coastal Prairie | 54. Sierra Norte De Puebla-Oaxaca |
| 4. Northwestern Interior Forest | 21. Ooko/And Prairies | 38. Islas Marianas | 55. Planicie Noroccidental De Yucatan |
| 5. Northern Pacific Rainforest | 22. Eastern Tallgrass Prairie | 39. Sierras De Baja California | 56. Planicie De La Peninsula De Yucatan |
| 6. Boreal Taiga Plains | 23. Prairie Hardwood Transition | 40. Desierto De Baja California | 57. Isla Cozumel |
| 7. Taiga Shield And Hudson Plains | 24. Central Hardwoods | 41. Islas Del Golfo De California | 58. Altos De Chiapas |
| 8. Boreal Softwood Shield | 25. West Gulf Coastal Plain/Ouachitas | 42. Sierras Y Planicies Del Cabo | 59. Depresiones Intermontanas |
| 9. Great Basin | 26. Mississippi Alluvial Valley | 43. Planicie Costera, Lomerios Y Canones De Occidente | 60. Sierra Madre De Chiapas |
| 10. Northern Rockies | 27. Southeastern Coastal Plain | 44. Marismas Nacionales | 61. Planicie Costera Del Soconusco |
| 11. Prairie Potholes | 28. Appalachian Mountains | 45. Planicie Costera Y Lomerios Del Pacifico Sur | 62. Archipelago De Revillagigedo |
| 12. Boreal Hardwood Transition | 29. Piedmont | 46. Sur Del Altiplano Mexicano | 63. Isla Guadalupe |
| 13. Lower Great Lakes/ St. Lawrence Plain | 30. New England/Mid-Atlantic Coast | 47. Faja Neovolcanica Transversal | 64. Arrecife Alacranes |
| 14. Atlantic Northern Forest | 31. Peninsular Florida | 48. Sierra Madre Oriental | 65. Los Tuxtlas |
| 15. Sierra Nevada | 32. Coastal California | 49. Planicie Costera Y Lomerios Secos Del Golfo De Mexico | 66. Pantanos De Centla-Laguna De Terminos |
| 16. Southern Rockies/Colorado Plateau | 33. Sonoran And Mojave Deserts | 50. Cuenca Del Rio Balsas | |
| 17. Badlands And Prairies | 34. Sierra Madre Occidental | 51. Valle Tehuacan-Cuicatlan | |



Map © National GIS Laboratory, Bird Studies Canada, 2014

0 1,000 km

Appendix B: Effects on the Environment and Other Species

A strategic environmental assessment (SEA) is conducted on all SARA recovery planning documents, in accordance with the [Cabinet Directive on the Environmental Assessment of Policy, Plan and Program Proposals](#)¹⁴. The purpose of a SEA is to incorporate environmental considerations into the development of public policies, plans, and program proposals to support environmentally sound decision-making and to evaluate whether the outcomes of a recovery planning document could affect any component of the environment or any of the [Federal Sustainable Development Strategy](#)'s¹⁵ (FSDS) goals and targets.

Conservation planning is intended to benefit species at risk and biodiversity in general. However, it is recognized that implementation of management plans may also inadvertently lead to environmental effects beyond the intended benefits. The planning process based on national guidelines directly incorporates consideration of all environmental effects, with a particular focus on possible impacts upon non-target species or habitats. The results of the SEA are incorporated directly into the management plan itself, but are also summarized below in this statement.

The Horned Grebe is a waterbird species nesting on small ponds and wetlands of the prairie and boreal ecoregions upon which many other species depend for nesting and feeding. Conservation measures aimed at conserving and restoring ecosystems are expected to alleviate threats for other SARA-listed wetlands species, such as the Olive-sided Flycatcher (*Contopus cooperi*), the Rusty Blackbird (*Euphagus carolinus*) and the Yellow Rail (*Coturnicops noveboracensis*), Western Tiger Salamander (*Ambystoma mavortium*), Western Toad (*Anaxyrus boreas*), Northern Leopard Frog (*Lithobates pipiens*), Great Plains Toad (*Anaxyrus cognatus*), non-inclusively. On western wintering grounds, mitigating stresses related to fisheries bycatch and contamination is expected to benefit seabird species such as the Marbled Murrelet (*Brachyramphus marmoratus*), the Pink-footed Shearwater (*Puffinus creatopus*) and the Short-tailed Albatross (*Phoebastria albatrus*).

Although it is possible that this management plan may negatively influence other species, it is concluded that it is unlikely to produce significant negative effects, given the non-intrusive nature of the proposed actions and the abundant populations of potentially affected species.

¹⁴ www.canada.ca/en/environmental-assessment-agency/programs/strategic-environmental-assessment/cabinet-directive-environmental-assessment-policy-plan-program-proposals.html

¹⁵ www.fsds-sfdd.ca/index.html#/en/goals/