

COMMITTEE ON THE
STATUS OF ENDANGERED
WILDLIFE IN CANADA

OTTAWA, ONT. K1A 0H3
(819) 997-4991

COMITÉ SUR LE STATUT
DES ESPÈCES MENACÉES
DE DISPARITION AU
CANADA

OTTAWA (ONT.) K1A 0H3
(819) 997-4991

**STATUS REPORT ON THE PYGMY SPERM WHALE
*KOGIA BREVICEPS***

IN CANADA

3611008I

QL
88
573
Vol. 10

BY

**ROBIN W. BAIRD
DAWN NELSON
JON LIEN**



AND

DAVID W. NAGORSEN


**STATUS ASSIGNED IN 1994
NOT AT RISK**

**REASON: THE PYGMY SPERM WHALE IS AN UNCOMMON SPECIES IN
CANADIAN WATERS, BUT THERE ARE NO KNOWN CURRENT
THREATS TO THIS SPECIES IN CANADA.**

**OCCURRENCE: COASTAL WATERS OF NEW BRUNSWICK, NOVA SCOTIA AND
BRITISH COLUMBIA [ATLANTIC AND PACIFIC OCEANS].**

COSEWIC - A committee of representatives from
Federal, provincial and private agencies which
assigns national status to species at risk in
Canada.

CSEMDC - Un comité de représentants d'organismes
fédéraux, provinciaux et privés qui attribue un
statut national aux espèces canadiennes en péril.



Committee
on the Status
of Endangered
Wildlife
in Canada

Comité sur le
statut des espèces
menacées
de disparition
au Canada

JUNE 1990

OH3 (819)
Ottawa, Ont. K1A 0S2 (613) 997-4991

NOTES

1. This report is a working document used by COSEWIC in assigning status according to criteria listed below. It is released in its original form in the interest of making scientific information available to the public.
2. Reports are the property of COSEWIC and the author. They may not be presented as the work of any other person or agency. Anyone wishing to quote or cite information contained in status reports may do so provided that both the author and COSEWIC are credited. Reports may be cited as in the following example:

Bredin, E. J. 1989. Status report on the Northern Prairie Skink, Eumeces septentrionalis, in Canada. Committee on the Status of Endangered Wildlife in Canada. 48 pp.
3. Additional copies of this report may be obtained at nominal cost from Canadian Nature Federation, 453 Sussex Drive, Ottawa, Ontario, K1N 6Z4.

DEFINITIONS

- SPECIES:** "Species" means any species, subspecies, or geographically separate population.
- VULNERABLE SPECIES:** Any indigenous species of fauna or flora that is particularly at risk because of low or declining numbers, occurrence at the fringe of its range or in restricted areas, or for some other reason, but is not a threatened species.
- THREATENED SPECIES:** Any indigenous species of fauna or flora that is likely to become endangered in Canada if the factors affecting its vulnerability do not become reversed.
- ENDANGERED SPECIES:** Any indigenous species of fauna or flora that is threatened with imminent extinction or extirpation throughout all or a significant portion of its Canadian range.
- EXTIRPATED SPECIES:** Any indigenous species of fauna or flora no longer known to exist in the wild in Canada but occurring elsewhere.
- EXTINCT SPECIES:** Any species of fauna or flora formerly indigenous to Canada but no longer known to exist anywhere.

**STATUS REPORT ON THE PYGMY SPERM WHALE
*Kogia breviceps***

IN CANADA

3611008I

BY

**ROBIN W. BAIRD
MARINE MAMMAL RESEARCH GROUP
BOX 6244
VICTORIA, BRITISH COLUMBIA
V8P 5L5**



**DAWN NELSON
WHALE RESEARCH GROUP
MEMORIAL UNIVERSITY OF NEWFOUNDLAND
230 MT. SCIO ROAD
ST. JOHN'S, NEWFOUNDLAND
A1C 5S7**

**JON LIEN
OCEAN SCIENCES CENTRE AND DEPARTMENT OF PSYCHOLOGY
MEMORIAL UNIVERSITY OF NEWFOUNDLAND
ST. JOHN'S, NEWFOUNDLAND
A1C 5S7**

AND

**DAVID W. NAGORSEN
ROYAL BRITISH COLUMBIA MUSEUM
675 BELLEVILLE STREET
VICTORIA, BRITISH COLUMBIA
V8V 1X4**

**STATUS ASSIGNED IN 1994
NOT AT RISK**

Status of the Pygmy Sperm Whale, *Kogia breviceps*, in Canada

ROBIN W. BAIRD^{1,2}, DAWN NELSON³, JON LIEN⁴ and DAVID W. NAGORSEN⁵

¹ Marine Mammal Research Group, Box 6244, Victoria, British Columbia V8P 5L5

² Department of Biological Sciences, Simon Fraser University, Burnaby, British Columbia V5A 1S6

³ Whale Research Group, 230 Mt. Scio Road, Memorial University of Newfoundland, St. John's, Newfoundland A1C 5S7. Current address: 16 Maples Road, Raymond, ME 04071

⁴ Ocean Sciences Centre and Department of Psychology, Memorial University of Newfoundland, St. John's, Newfoundland A1C 5S7

⁵ Royal British Columbia Museum, 675 Belleville Street, Victoria, British Columbia V8V 1X4

Robin W. Baird, Dawn Nelson, Jon Lien and David W. Nagorsen. 1993. Status of the Pygmy Sperm Whale, *Kogia breviceps*, in Canada. Report to the Committee on the Status of Endangered Wildlife in Canada (COSEWIC), Canadian Wildlife Service, Ottawa, Ontario.

This report reviews the general biology, world-wide status and management of the Pygmy Sperm Whale, *Kogia breviceps*, with special reference to its status in Canadian waters. Pygmy Sperm Whales appear to be uncommon in Canadian waters; there are several unconfirmed sighting records of this species off British Columbia and only four stranding records off the Canadian east coast. Little is known about its biology or world-wide status, and although it is taken in small numbers both directly and incidentally in fisheries, no serious threats to its status are apparent. No COSEWIC designation is required regarding its status in Canadian waters.

Le présent rapport résume la biologie générale, le statut international et la gestion du Cachalot Pygmée, *Kogia breviceps*, et s'attache plus particulièrement à son statut dans les eaux canadiennes. Le Cachalot Pygmée semble être une espèce inhabituelle dans les eaux canadiennes; il existe plusieurs mentions visuelles de cachalots au large de la Colombie-Britannique, mais aucune n'a été confirmée. En outre, on ne dispose que de quatre mentions de spécimens échoués le long de la côte est du Canada. On ne connaît que peu de détails au sujet de sa biologie ou de son statut international. Bien que le Cachalot Pygmée soit capturé en petit nombre, que ce soit directement ou comme prise fortuite par les pêcheurs, il ne semble y avoir aucune menace grave à son statut. Son statut dans les eaux canadiennes ne nécessite donc aucune désignation par le CSEMDC.

Taxonomy

Class	Mammalia
Order Cetacea	
Family	Physeteridae
Scientific Name	<i>Kogia breviceps</i> (de Blainville, 1838)
Common Name	Pygmy Sperm Whale
Other Names:	
Latin America	Cachalote Pigméo
Japanese	Komakkō Kujira
Russian	Karlikovyy Kashalot

Key Words: Pygmy Sperm Whale, Cachalot Pygmée, *Kogia breviceps*, Canada, North Atlantic, British Columbia, status, Cetacea.

Little is known about the Pygmy Sperm Whale, *Kogia breviceps* (de Blainville, 1838) [Figure 1]. We summarize the current state of knowledge of the species, with special reference to its status and management in Canadian waters, by request of the Fish and Marine Mammal Subcommittee of the Committee on the Status of Endangered Wildlife in Canada (COSEWIC). There are two species within the genus *Kogia*: *Kogia breviceps*, and the Dwarf Sperm Whale, *Kogia simus*. Before 1966, however, most authors recognized one species, *Kogia breviceps*, within the genus (Yamada 1954; Handley 1966), resulting in considerable confusion as to which species is actually referred to in early publications. This taxonomic uncertainty can be attributed to the similarity of the two species and the scarcity of specimens. Both taxa are small (less than 3.8 m), have a small, underslung shark-like mouth which is set well back from the tip of the snout. They are dark bluish-gray dorsally and shade from a lighter gray laterally to a dull white or pink on the belly (Leatherwood and Reeves 1983; Nagorsen 1985). Both have a bulbous head and, like the other member of the Family Physeteridae, the Sperm Whale (*Physeter macrocephalus*), they have a spermaceti organ (Handley 1966). The tail stock is elongated and laterally compressed, and the flukes are notched and concave along the rear margin. Stranded specimens of both species are occasionally confused with sharks, due to the shark-like mouth and the presence of a lightly pigmented bracket-shaped mark on each side of the head between the eye and flipper, sometimes referred to as "false gills" (Figure 1).

The two species can be discriminated by a variety of external and cranial characters (see Table 1). Many authors note that they can be distinguished by the position of the dorsal fin on the back, set well posterior to the midpoint of the back for the Pygmy Sperm Whale and with the

anterior insertion of the fin near the midpoint of the back for the Dwarf Sperm Whale, but Ross (1979) cautioned against using this character in isolation, as there is considerable individual variation in both species. Pygmy Sperm Whales reach a greater length than Dwarf Sperm Whales, but the maximum length attained by *Kogia breviceps* is unclear. Caldwell et al. (1971) reported an individual of 4.25 m, but Ross (1979) suggested that the length may have been estimated, rather than measured. According to Ross (1979) the next-largest recorded specimen was 3.5 m in length. Leatherwood et al. (1988) noted that Pygmy Sperm Whales may grow to a maximum length of at least 3.7 m, and Eliason and Houck (1986) reported an animal 3.82 m in length from the records of the Smithsonian Institution. The taxonomic position of the genus *Kogia* within the Odontoceti is unclear (eg. see Rice and Wolman 1990).

Distribution

The Pygmy Sperm Whale is found virtually world-wide in tropical and warm-temperate seas. In the western Pacific it has been reported from Japan in the north to New Zealand and Australia in the south (Omura and Takahashi 1981; Baker 1983; Brabyn 1991), and in the eastern Pacific from Washington State in the north to Peru and Chile in the south (Allen 1941; Scheffer and Slipp 1948; Hubbs 1951; Brownell 1969; Waerebeek et al. 1987). Pike and Giovando (1963) stated that Pygmy Sperm Whales were known to occur in the offshore waters of British Columbia, but we could find no documented records from the Canadian west coast. Four unconfirmed sighting reports are listed as *Kogia breviceps* from the B.C. coast (Baird unpublished data), but positive confirmation awaits photographs or a specimen. Three records are confirmed from the adjacent waters of Washington State (Scheffer and Slipp 1948; Everitt

et al. 1979; Osborne et al. 1988) and it is likely that this species will eventually be documented in B.C. *Kogia simus* has been recorded in British Columbia (Nagorsen and Stewart 1983).

In the western Atlantic, Pygmy Sperm Whales have been recorded from Canada southward along the U.S. east coast, throughout the Gulf of Mexico, and as far south as Brazil, Uruguay and Argentina (Piers 1923; Allen 1941; Gunter et al. 1955; Carvalho 1967; Hysmith et al. 1976; Geise and Borobia 1987). Four stranding records have been reported on the Canadian east coast (Figure 2; Table 2). Three of these were from Canadian waters, and one from the French Island of Miquelon (Piers 1923; Sergeant et al. 1970; Nelson et al. 1991; McAlpine and Murison in prep.). These are the most northern records in the western Atlantic. In the eastern Atlantic this species has been recorded from Ireland, the Netherlands and France south to South Africa (Allen 1941; Fraser 1974; Maul and Sergeant 1977; Teixeira 1979; Ross 1984). Pygmy Sperm Whales are found in the Indian Ocean (Leatherwood and Reeves 1989; Chantrapornsyl et al. 1991) but do not appear to have been recorded in the Mediterranean Sea (Baccetti et al. 1991).

Protection

International

The Pygmy Sperm Whale is listed under Appendix II of the Convention on International Trade in Endangered Species of Wild Fauna and Flora 1973 (CITES) (see Birnie 1982). Such a listing allows for the regulation of international trade between members and non-members of the convention by requiring export permits from the country of origin. However, there appears to be no current international trade in Pygmy Sperm Whale products. The

International Whaling Commission (IWC) regulates the taking of whales in accordance with the current Schedule provisions, but whether this Commission's mandate covers the Pygmy Sperm Whale is unclear, as members of the Commission are divided as to whether "whale" refers to all cetaceans, or to only some species (Klinowska 1987, 1991).

National

Canada: Until they were repealed in 1993, the Cetacean Protection Regulations of the Fisheries Act of Canada of 1867 protected all cetacean species from "hunting". "Hunting" was defined as "to chase, shoot at, harpoon, take, kill, attempt to take or kill, or to harass cetaceans in any manner", and could only be undertaken under licence. Aboriginal "hunting" however could be undertaken without licences. The Cetacean Protection Regulations were replaced with the Marine Mammal Regulations of the Fisheries Act in early 1993. These regulations appear to provide no more or no less protection, by stating only that "no person should disturb a marine mammal except when ... under the authority of these Regulations". No provisions exist for regulation of incidental catches in fishing operations. Canada is not currently a member of the International Whaling Commission, having withdrawn in 1982 (IWC 1982).

United States: All cetaceans are protected through the Marine Mammal Protection Act of 1972, as well as through the Packwood-Magnuson Amendment of the Fisheries and Conservation Act and the Pelly Amendment of the Fisherman's Protective Act.

Population Size(s) and Trends

In the recent IUCN Cetacean Red Data Book, Klinowska (1991) noted that there is insufficient information to accurately classify the world status of this species. No information is available on population sizes or trends, nor on stock identity. In fact, very few sightings of either species of *Kogia* have ever been reported, and knowledge of both species comes largely from stranded animals. In some areas, Pygmy Sperm Whales are among the most frequently recorded stranded species. For example, Odell (1991) noted that they are the second most frequent to strand in the southeastern United States; off the Hawaiian Islands the Pygmy Sperm Whale is the fourth most frequently stranded species (Nitta 1991). In his analysis of strandings in New Zealand, Brabyn (1991) noted that although they do not represent the largest number of individuals (as mass strandings are infrequent), Pygmy Sperm Whales are the most frequently recorded species. Many authors have interpreted these frequent strandings as evidence that this species is fairly common. Recent numerous aerial sightings of *Kogia* (not discriminated to species) in the northern Gulf of Mexico seem to support this supposition (Jefferson et al. 1992).

Only four strandings have been reported off the Canadian east coast and it appears that Pygmy Sperm Whales become more common in the south, with a greater number of records reported off the northeast U.S. coast (Early and McKenzie 1991).

Habitat

Pygmy Sperm Whales generally inhabit offshore waters in warm temperate and tropical areas. Brabyn (1991) suggested that the Mahia

Peninsula area of the north island of New Zealand is a calving area for this species, based on a high proportion of mother/calf strandings. However, information on the habitat type is not presented. Klages et al. (1989) noted that both species in the genus feed on prey typical of the continental slope, although *Kogia simus* feeds more inshore than *Kogia breviceps*.

General Biology

Reproduction

Very little is known about the reproductive biology of this species. Most estimates of reproductive parameters are based on the examination of a small number of stranded animals, and should be considered preliminary. Ross (1979) suggested that mating and calving occur from autumn through spring, and noted that while there was insufficient data to accurately estimate gestation, two alternative methods suggest a duration between 7 and 11 months. Individuals have been recorded as being simultaneously pregnant and accompanied by a calf, indicating that some females may breed annually (Ross 1979; Price et al. 1984; Eliason and Houck 1986). Ross (1979) reported that length at birth averages about 1.2 m, sexual maturity is attained at about 2.7-2.8 m for females and 2.7 to 3.0 m for males, and physical maturity is reached at lengths of 3.0-3.3 m for both sexes. The sex ratio of stranded animals varies. Brabyn (1991) found that of 79 Pygmy Sperm Whales that stranded in New Zealand, there were more than twice as many females as males. Off South Africa, the sex ratio of 16 adult and sub-adult animals was 1:1, while the sex ratio for 15 fetuses and calves was heavily biased towards males (12 males: 3 females) [Ross 1979]. Such disparate figures likely arise from the small sample sizes, and clearly more information is needed.

Ross (1979, 1984) and Eliason and Houck (1986) discussed sectioning teeth for age determination, but no information is available to calibrate the deposition rate of layers. Ross (1984) noted that one sexually mature female which stranded with a calf had only 3.5 growth layers in the dentine. Assuming that one layer is deposited annually, this would imply the female was only about two years old when she first conceived (Ross 1984). No information on longevity is available.

Movements

Strandings off South Africa and the U.S. southeast coast occur throughout the year, possibly suggesting a lack of seasonal movements (Ross 1979; Leatherwood and Reeves 1983). Off South Australia however, Kemper and Ling's (1991) analysis of stranding records indicates the presence of Pygmy Sperm Whales only during April through October. In the eastern North Pacific the majority of records are concentrated during the fall and winter (Eliason and Houck 1986), and along the coast of Europe, Fraser (1974) suggested that this species follows the North Atlantic current extension of the Gulf Stream in search of food. There is also evidence for a possible seasonal movement of *Kogia breviceps* near the west coast of New Caledonia, as strandings there occur mainly between June and December (Sylvestre 1988).

Behaviour

Behavioural observations are based on a few sightings at sea and on instances where individuals have been kept in captivity after stranding alive (eg. see Sylvestre 1983). Pygmy Sperm Whales are seen singly or in groups of up to about six individuals (Allen 1941; Vidal et al. 1987; Leatherwood et al.

1988), and do not appear to regularly associate with other species of cetaceans or with seabirds (Au and Pitman 1988). Surface behaviour is typically undemonstrative; individuals typically rise slowly to the surface, produce an inconspicuous blow, and dive without showing the flukes. They are occasionally observed lying still at the water's surface with the top of the head and back exposed. Allen (1941) noted that animals at the surface appeared to be very easy to approach closely, even after being harpooned. However, as with several other species of cetaceans, Pygmy Sperm Whales may release a cloud of reddish brown feces into the water when startled (Leatherwood et al. 1988). The function of this behaviour is unclear, although Scott and Cordaro (1987) observed a Dwarf Sperm Whale mother and calf pair exhibiting this behaviour while trapped inside a tuna purse-seine net, and then apparently hiding within the cloud when approached by dolphins which were also trapped.

Pygmy Sperm Whales appear to feed primarily on cephalopods, as well as crustaceans and fish. Cephalopods recorded from stomach contents of this species include *Abrealia* sp., *Abrealiopsis* sp., *Ancistrocheirus* sp., *Galiteuthis* sp., *Histioteuthis* sp., *Loligo vulgaris*, *Lycoteuthis diadema*, *Moroteuthis* sp., *Octopoteuthis cyeletron*, *Ommastrephes* sp., *Onychoteuthis boreali-japonicus*, *Phasmatopsis* sp., *Pygropsis* sp., *Pyroteuthis* sp., *Sepioteuthis australis*, *Taningia* sp., *Taonius pavo*, *Todarodes* sp., *Tuethowenia pellucida*, and *Vampyroteuthis* sp. (Hale 1947; Raun et al. 1970; Ross 1979; Eliason and Houck 1986; Klages et al. 1989). Crustaceans recovered include *Aristaeomorpha foliacea*, *Carcinides maenas*, *Gnathophausia ingens*, *Goneplax angulata*, *Hymenodora* sp., *Pandalopsis* sp., *Pandalus* sp., *Pasiphaea pacifica*, and *Penaeus californiensis* (Allen 1941; Hale 1947; Scheffer and Slipp 1948; Raun et al. 1970;

Ross 1979; Vidal et al. 1987). Fish recorded include *Lampanyctus* sp., *Maurolicus muelleri*, *Photichthys argenteus*, *Pyrosoma* sp., *Rexea solandri*, *Scopelopsis multipunctatus*, and *Symbolophorus* sp. (Ross 1979, 1984). Raun et al. (1970) also noted the seaweed *Sargassum* from the stomach of a stranded animal, but suggest that this may have been ingested accidentally. An isotope analysis of muscle tissue from a stranded animal off Miquelon suggested that the animal had been feeding at the same trophic level as Sperm Whales, ie. it had probably been feeding on squid in offshore waters (Nelson et al. 1991). This species likely uses echolocation to find prey, as echolocation-type clicks have been recorded from live-stranded animals (Caldwell and Caldwell 1991).

Limiting Factors

Data on natural mortality is scarce. There are no reported observations of Killer Whales (*Orcinus orca*) attacking Pygmy Sperm Whales (Jefferson et al. 1991), but the species has been recorded from the stomach contents of Killer Whales from both the Caribbean and the Indian Ocean (Perrin 1982). Long (1991) noted an apparent attack by a White Shark (*Carcharodon carcharias*) on a Pygmy Sperm Whale off California. Information on other causes of natural mortality is sparse. Virtually all recorded strandings of this species are of single animals or of cow/calf pairs. In New Zealand, Robson (1984) reported one case of three dead individuals found together on a beach, in association with seven stranded False Killer Whales (*Pseudorca crassidens*), and Brabyn (1991) described a stranding of four individuals. Robson (1984) attributed many of the single strandings of this species to heavy infestations of parasites in the head and inner ear. Parasites recorded from this species include the cestode *Phyllobothrium delphini*, and the nematodes *Anisakis*

physeteris, *Anisakis typica*, *Anisakis simplex*, *Phocanema kogiae*, *Stenurus* sp., *Terranova* sp., and possibly *Crassicauda* sp. (Zam et al. 1971; Ross 1979; Vidal et al. 1987). Buck (1984) noted the presence of the bacteria *Enterobacter agglomerans*, *Enterobacter cloacae*, *Pseudomonas cepacia*, *Pseudomonas maltophilia*, *Bacillus*, *Flavobacterium*, and the yeasts *Rhodotorula pallida*, *Rhodotorula rubra*, *Torulopsis*, and *Aureobasidium*, but the role of such pathogens in natural mortality is unknown. Severe vascular disease, including thrombus formation, has been noted in one animal (Roberts et al. 1964, cited in Sweeney and Ridgway 1975).

There appear to be no known major threats (past or present) to this species (Klinowska 1991). Small numbers of Pygmy Sperm Whales have been taken in fisheries, both directly and incidentally. In previous years *Kogia* have been taken by shore-based whaling operations off Japan, but it is unclear if both species were included (Yamada 1954). Edmondson (1948) noted one animal which was speared, and another accidentally hooked and landed while fishing with a hand line off Hawaii. Van Waerebeek et al. (1987) found a specimen in a dump in Peru where the remains of fish and dolphins taken deliberately for human consumption were discarded. One animal was illegally killed in Australia in 1989 (Australia 1991). Whalers in the Timor Sea, Indonesia, have hunted *Kogia* (Weber 1923), although there is no evidence that they have been taken in recent years (Barnes 1991). *Kogia*, probably including *Kogia breviceps*, have been and probably still are taken occasionally in the Lesser Antilles (Caldwell and Caldwell 1975; Reeves 1988). Vidal et al. (1990) noted that Pygmy Sperm Whales were recorded caught in gillnet fisheries in Mexican or Central American waters, but presented no details on exact locations or numbers. This species is taken in directed fisheries and

incidentally in gillnet and seine fisheries in the Philippines (Aragones et al. 1991; Dolar et al. 1991). Animals have also been caught incidentally in gillnets in the central North Pacific and off Sri Lanka (Omura et al. 1984; Leatherwood and Reeves 1989). Ingestion of foreign objects (Jones 1984; Ross 1979, 1984) and boat collisions (Sylvestre 1988) may occasionally result in mortality. Little has been reported on levels of environmental contaminants in this species (Cockcroft et al. 1991).

Evaluation

Based on existing information the Pygmy Sperm Whale appears to be only a rare visitor to Canadian waters. No serious threats to its status in Canadian waters are apparent. As such, no COSEWIC status designation is required.

Acknowledgements

We thank the World Wildlife Fund (Canada) for financial assistance for the preparation of this report, Robert Campbell and COSEWIC for providing assistance and support, Sherry Smrstik of the National Marine Mammal Laboratory for providing references, and Tamara Guenther and two anonymous reviewers (Randall Reeves and David Gaskin) for helpful comments on the manuscript. We would also like to thank Donald McAlpine and Laurie Murison for allowing us to cite their manuscript in preparation.

Literature Cited

Allen, G.M. 1941. Pygmy sperm whale in the Atlantic. Field Museum of Natural History Zoology Series 27:17-36.

- Aragones, L.V., S. Leatherwood, M.L. Dolar, and C.L. Hill. 1991. Species composition of Philippine marine mammal fauna. Page 2 in Abstracts of the Ninth Biennial Conference on the Biology of Marine Mammals, December 5-9, 1991, Chicago, IL.
- Au, D.W., and R.L. Pitman. 1988. Seabird relationships with tropical tunas and dolphins. Pages 174 - 212 in *Seabirds and other marine vertebrates*. Edited by J. Burger. Columbia University Press, New York.
- Australia. 1991. Progress report on cetacean research, May 1989 to May 1990. Reports of the International Whaling Commission 41:223-229.
- Baccetti, N., F. Cancelli, and T. Renieri. 1991. First record of *Kogia simus* (Cetacea, Physeteridae) from the Mediterranean Sea. *Mammalia* 55(1):152-154.
- Baker, A.N. 1983. Whales and dolphins of New Zealand and Australia: an identification guide. Victoria University Press, Wellington.
- Barnes, R.H. 1991. Indigenous whaling and porpoise hunting in Indonesia. UNEP Marine Mammal Technical Report 3:99-106.
- Birnie, P. 1982. Legal measures for the prevention of "pirate" whaling. International Union for Conservation of Nature and Natural Resources (IUCN) Environmental Policy and Law Paper 19:1-163.
- Brabyn, M.W. 1991. An analysis of the New Zealand whale stranding record. Department of Conservation Science and Research Series 29.
- Brownell, R.L. Jr. 1969. Pygmy sperm whale in the Gulf of California. *Journal of Mammalogy* 50(2):356-357.
- Buck, J.D. 1984. Microbiological observations on two stranded live whales. *Journal of Wildlife Diseases* 20(2):148-150.
- Caldwell, D.K., and M.C. Caldwell. 1975. Dolphin and small whale fisheries of

the Caribbean and West Indies: occurrence, history, and catch statistics - with special reference to the Lesser Antillean Island of St. Vincent. *Journal of the Fisheries Research Board of Canada* 32:1105-1110.

Caldwell, D.K., H. Neuhauser, M.C. Caldwell, and H.W. Coolidge. 1971. Recent records of marine mammals from the coasts of Georgia and South Carolina. *Cetology* 5:1-12.

Caldwell, M.C., and D.K. Caldwell. 1991. A note describing sounds recorded from two cetacean species, *Kogia breviceps* and *Mesoplodon europaeus*, stranded in northeastern Florida. Pages 151-154 in *Marine mammal strandings in the United States*. Edited by J.E. Reynolds and D.K. Odell. NOAA Technical Report NMFS 98.

Carvalho, C.T. 1967. Pygmy sperm whale (*Kogia breviceps*) in the Brazil. *Mammalia* 31:168-169.

Chantrapornsyl, S., C.C. Kinze, S. Leatherwood, and W.P. Prematunga. 1991. Notes on the genus *Kogia* in the northern Indian Ocean. UNEP Marine Mammal Technical Report 3:79-88.

Cockcroft, V.G., G.J.B. Ross, A.D. Connell, B.D. Gardner, and A.C. Butler. 1991. Occurrence of organochlorines in stranded cetaceans and seals from the east coast of southern Africa. UNEP Marine Mammal Technical Report 3:271-276.

Dolar, M.L.L., S. Leatherwood, L.V. Aragonés, and C.L. Hill. 1991. Directed fisheries for whales and dolphins in Palaway, the Southern Visayas and the Mindanao Sea, Philippines. Page 18 in *Ninth Biennial Conference on the Biology of Marine Mammals*, December 5-9, 1991, Chicago, IL.

- Early, G., and T.P. McKenzie. 1991. The northeast regional marine mammal stranding network. Pages 63-68 in Marine mammal strandings in the United States. Edited by J.E. Reynolds and D.K. Odell. NOAA Technical Report NMFS 98.
- Edmondson, C.H. 1948. Records of *Kogia breviceps* from the Hawaiian islands. Journal of Mammalogy 29(1):76-77.
- Eliason, J.J., and W.J. Houck. 1986. Notes on the biology of a gravid pygmy sperm whale (*Kogia breviceps*) from California. Cetology 51:1-5.
- Everitt, R.D., C.H. Fiscus, and R.L. DeLong. 1979. Marine mammals of northern Puget Sound and the Strait of Juan de Fuca - A report on investigations November 1, 1977 - October 31, 1978. NOAA Technical Memorandum ERL MESA-41.
- Fraser, F.C. 1974. Report on cetacea stranded on the British coasts from 1948 to 1966. British Museum (Natural History) 14.
- Geise, L., and M. Borobia. 1987. New Brazilian records for *Kogia*, *Pontoporia*, *Grampus*, and *Sotalia* (Cetacea, Physeteridae, Platanistidae, and Delphinidae). Journal of Mammalogy 68(4):873-875.
- Gunter, G., C.L. Hubbs, and M.A. Beal. 1955. Records of *Kogia breviceps* from Texas, with remarks on movements and distribution. Journal of Mammalogy 36(2):263-270.
- Hale, H.M. 1947. The pigmy sperm whale (*Kogia breviceps*, Blainville) on South Australian coasts. Records of the South Australian Museum 8:531-546.
- Handley, C.O. 1966. A synopsis of the genus *Kogia* (pygmy sperm whales). Pages 62-69 in Whales, dolphins, and porpoises. Edited by K.S. Norris. University of California Press, Berkeley.
- Hysmith, B.T., R.L. Colura, J.C. Kana. 1976. Beaching of a live pygmy sperm

whale (*Kogia breviceps*) at Port O'Connor, Texas. Southwestern Naturalist 21(3):409.

Hubbs, C.L. 1951. Eastern Pacific records and general distribution of the pygmy sperm whale. Journal of Mammalogy 32(4):403-410.

IWC. 1982. International Whaling Commission Report 1980-1981. Reports of the International Whaling Commission 32:3-7.

Jefferson, T.A., P.J. Stacey, and R.W. Baird. 1991. A review of killer whale interactions with other marine mammals: predation to co-existence. Mammal Review 21:151-180.

Jefferson, T.A., S. Leatherwood, L.K.M. Shoda, and R.L. Pitman. 1992. Marine mammals of the Gulf of Mexico: a field guide for aerial and shipboard observers. Texas A&M University Printing Center, College Station, Texas.

Jones, S.C. 1984. Letter to the editor. Whalewatcher 18(4):22-23.

Katona, S.K., V. Rough, and D.T. Richardson. 1983. A field guide to the whales, porpoises and seals of the Gulf of Maine and eastern Canada. Third Edition. Charles Scribner's Sons, New York.

Kemper, C.M., and J.K. Ling. 1991. Whale strandings in South Australia (1881-1989). Transactions of the Royal Society of South Australia 115(1):37-52.

Klages, N., V.G. Cockcroft, and P.B. Best. 1989. Abstract. Stomach contents of pygmy *Kogia breviceps* and dwarf *Kogia simus* sperm whales stranded on South African beaches. Page 35 in Abstracts of the Eighth Biennial Conference on the Biology of Marine Mammals, December 7-11, 1989, Pacific Grove, CA.

Klinowska, M. 1987. The status of marine mammals in the southern North Sea. Pages 75- 95 in Reasons for Concern - Proceedings of the 2nd North Sea

- Seminar '86. *Edited by G. Peet.* Amsterdam.
- Klinowska, M. 1991. Dolphins, porpoises and whales of the world - the IUCN Red Data Book. IUCN, Gland, Switzerland.
- Leatherwood, S., and R.R. Reeves. 1983. The Sierra Club handbook of whales and dolphins. Sierra Club Books, San Francisco.
- Leatherwood, S., and R.R. Reeves. *Editors.* 1989. Marine mammal research and conservation in Sri Lanka 1985-1986. UNEP Marine Mammal Technical Report 1.
- Leatherwood, S., R.R. Reeves, W.F. Perrin, and W.E. Evans. 1988. Whales, dolphins, and porpoises of the eastern North Pacific and adjacent Arctic waters - A guide to their identification. Dover Publications, Inc., New York.
- Long, D.J. 1991. Apparent predation by a white shark *Carcharodon carcharias* on a pygmy sperm whale *Kogia breviceps*. Fishery Bulletin, U.S. 89:538-540.
- Maul, G.E., and D.E. Sergeant. 1977. New cetacean records from Madeira. *Bocagiana* 43:1-8.
- Nagorsen, D. 1985. *Kogia simus*. *Mammalian Species* 239:1-6.
- Nagorsen, D.W., and G.E. Stewart. 1983. A dwarf sperm whale (*Kogia simus*) from the Pacific coast of Canada. *Journal of Mammalogy* 64(3):505-506.
- Nelson, D., A. Desbrosse, J. Lien, P. Ostrom, and R. Seton. 1991. A new stranding record of the pygmy sperm whale, *Kogia breviceps*, in waters off eastern Canada. *Canadian Field-Naturalist* 105(3):407-408.
- Nitta, E.T. 1991. The marine mammal stranding network for Hawaii, an overview. Pages 55-62 in *Marine mammal strandings in the United States*. Edited by J.E. Reynolds and D.K. Odell. NOAA Technical Report

NMFS 98.

Odell, D.K. 1991. A review of the southeastern United States marine mammal stranding network: 1978-1987. Pages 19-23 in Marine mammal strandings in the United States. Edited by J.E. Reynolds and D.K. Odell. NOAA Technical Report NMFS 98.

Omura, H., and Y. Takahashi. 1981. A pygmy sperm whale stranded at Tokaimura, Ibaragi, Japan. Scientific Reports of the Whales Research Institute 33:119-124.

Omura, H., M. Shirakihara, and H. Ito. 1984. A pygmy sperm whale accidentally taken by drift net in the North Pacific. Scientific Reports of the Whales Research Institute 35:183-193.

Osborne, R., J. Calambokidis, and E.M. Dorsey. 1988. A guide to marine mammals of greater Puget Sound. Island Publishers, Anacortes, Washington.

Perrin, W.F. Editor. 1982. Report of the workshop on identity, structure and vital rates of killer whale populations, Cambridge, England, June 23-25, 1981. Reports of the International Whaling Commission 32:617-632.

Piers, H. 1923. Accidental occurrence of the pygmy sperm whale (*Kogia breviceps*) on the coast of Nova Scotia; an extension of its known range, with remarks on the probability of the former presence in these waters of the true sperm whale (*Physeter macrocephalus*). Proceedings and Transactions of the Nova Scotian Institute of Science 15:95-114.

Pike, G.C., and L. Giovando. 1963. Whales and dolphins of the west coast of Canada. Fisheries Research Board of Canada Circular No. 68.

Price, M.C., G.J. MacDonald, S.L. Graham, and E.J. Kirk. 1984. Treatment of a strandling whale (*Kogia breviceps*). New Zealand Veterinary Journal

31(3):31-33.

Raun, G.G., H.D. Hoese, and F. Moseley. 1970. Pygmy sperm whales, genus *Kogia*, on the Texas coast. Texas Journal of Science 21(3):269-274.

Reeves, R.R. 1988. Exploitation of cetaceans in St. Lucia, Lesser Antilles, January 1987. Reports of the International Whaling Commission 38:445-447.

Rice, D.W., and A.A. Wolman. 1990. The stomach of *Kogia breviceps*. Journal of Mammalogy 71(2):237-242.

Robson, P.D. 1984. Strandings: ways to save whales. The Science Press Ltd., Johannesburg.

Ross, G.J.B. 1979. Records of pygmy and dwarf sperm whales, genus *Kogia*, from southern Africa, with biological notes and some comparisons.

Annals of the Cape Provincial Museums (Natural History) 11(14):259-327.

Ross, G.J.B. 1984. The smaller cetaceans of the south east coast of southern Africa. Annals of the Cape Provincial Museums (Natural History) 15(2):173-410.

Scheffer, V.B., and J.W. Slipp. 1948. The whales and dolphins of Washington State with a key to the cetaceans of the west coast of North America. American Midland Naturalist 39(2):257-337.

Scott, M.D., and J.G. Cordaro. 1987. Behavioral observations of the dwarf sperm whale, *Kogia simus*. Marine Mammal Science 3(4):353-354.

Sergeant, D.E., A.W. Mansfield, and B. Beck. 1970. Inshore records of cetacea for eastern Canada, 1949-68. Journal of the Fisheries Research Board of Canada 27(11):1903-1915.

Sweeney, J.C., and S.H. Ridgway. 1975. Common diseases of small cetaceans. Journal of the American Veterinary Medical Association 167(7):533-540.

- Sylvestre, J.P. 1983. Review of *Kogia* specimens (Physeteridae, Kogiinae) kept alive in captivity. *Investigations on Cetacea* 15:201-219.
- Sylvestre, J.P. 1988. On a specimen of pygmy sperm whale *Kogia breviceps* (Blainville 1838) from New-Caledonia. *Aquatic Mammals* 14(2):76-77.
- Teixeira, A.M.A.P. 1979. Marine mammals of the Portuguese coast. *Zeitschrift für Saugetierkunde* 44(4):221-238.
- Vidal, O., L.T. Findley, P.J. Turk, and R.E. Boyer. 1987. Recent records of pygmy sperm whales in the Gulf of California, Mexico. *Marine Mammal Science* 3(4):354-356.
- Vidal, O., K. van Waerebeek, L.T. Findley, and G. Alvarez-Manilla. 1990. *Abstract. Cetaceans and gillnet fisheries in Mexico, Central America and the Caribbean: a preliminary review. Page 51 in Abstracts of the Symposium on Mortality of Cetaceans in Passive Fishing Nets and Traps, October 20-21, 1990, La Jolla, CA.*
- Waerebeek, K. van, J.C. Reyes, and B.A. Luscombe. 1987. A second record of the pygmy sperm whale *Kogia breviceps* (de Blainville, 1838) (Cetacea, Physeteridae) from the Peruvian coast. *Zeitschrift für Saugetierkunde* 52:192-194.
- Weber, M. 1923. Die cetaceen der Siboga-Expedition. *Siboga-Expeditie Monograph* 58:1-38.
- Yamada, M. 1954. Some remarks on the pygmy sperm whale, *Kogia*. *Scientific Reports of the Whales Research Institute* 9:37-58.
- Zam, S.G., D.K. Caldwell, and M.C. Caldwell. 1971. Some endoparasites from small odontocete cetaceans collected in Florida and Georgia. *Cetology* 2:1-11.

Table 1. Distinguishable characteristics of *Kogia breviceps* and *Kogia simus* (Ross 1979; Leatherwood et al. 1988).

Characters	<i>Kogia breviceps</i>	<i>Kogia simus</i>
Maximum length ¹	3.8 m	2.7 m
Throat creases	absent	usually present
Dorsal fin height	usually less than 5% total body length	usually greater than 5% total body length
Dorsal fin position ¹	posterior to mid-point of back	near midpoint of back
Relative snout length	longer	shorter
Teeth in upper jaw	absent	up to 3 pairs
Teeth in lower jaw	10-16 pairs	8-13 pairs

¹ See text for further details.

Table 2. Records of *Kogia breviceps* from the Canadian east coast. All records are of single animals, found dead. No confirmed records from the Canadian west coast have been reported.


Date	Location	Source
17 January 1920	Halifax Harbour, NS	Piers 1923 ¹
29 January 1969	Sable Island, NS	Sergeant et al. 1970
8 October 1990	Isle de Miquelon	Nelson et al. 1991
8 December 1992	Saint John, NB	McAlpine and Murison, in prep

¹Katona et al. (1983) report a dead animal found under the ice in Halifax Harbour, NS, in the winter of 1970, but this appears to be a misprint, referring to the animal found under similar circumstances in 1920 (S. Katona, College of the Atlantic, personal communication).

Figure Captions

Figure 1. Pygmy Sperm Whale. Illustration by Dawn Nelson.

Figure 2. Records of *Kogia breviceps* off the Canadian east coast. See Table 2 for details of records.



Wm 1992

Fig 1

