DEATHS AND HOSPITALIZATIONS^{1,2}

Suicide rates are approx.

higher

among men compared to women




Suicide is the


SECOND

leading cause of death

among youth and young adults (15-34 years)







SUICIDE-RELATED BEHAVIOURS3,4

THOUGHTS

of suicide in their **LIFETIME**



had thoughts of suicide in the **PAST YEAR**

PLANS

suicide in their **LIFETIME**



had planned suicide in the **PAST YEAR**

ATTEMPTS

suicide in their LIFETIME



had attempted suicide in the **PAST YEAR**

Published data may underestimate the total number of reported deaths, attempts, plans and thoughts of suicide, due to the stigma and other factors.

ELP IS AVAILABLE

IF YOU OR SOMEONE YOU KNOW IS IN IMMEDIATE DANGER, PLEASE CALL 9-1-1.

Help is available 24/7 for suicide prevention and mental health. Here are some resources:

- Talk Suicide Canada: 1-833-456-4566 (or text 45645 from 4pm to midnight ET)
- For Quebec residents: 1-866-APPELLE (277-3553)
- Kids Help Phone: 1-800-668-6868 or text CONNECT to 686868
- Hope for Wellness Helpline for Indigenous peoples: 1-855-242-3310
- Trans Lifeline: 1-877-330-6366
- Wellness Together Canada
- Preventing suicide: Warning signs and how to help

LIKE US:

<u>@HealthyCdns</u>

FOLLOW US:

@GovCanHealth

^{4.} Statistics Canada. Canadian Community Health Survey—Annual component. 2020





, 2023 | Cat.: HP35-115/2022E-PDF | ISBN: 978-0-660-46509-8 | Pub.: 220621

 $Statistics\ Canadian\ Vital\ Statistics — Death\ Database.\ 2017-2019.\ Released\ in\ January,\ 2022.$

Canadian Institute of Health Information (CIHI). Discharge Abstract Database (DAD), National Ambulatory Care Reporting System (NACRS) and Ontario Mental Health Reporting System (OMHRS).

es are for the population in Canada aged 15 years and o