

The Attenuation of Light in Lake Winnipeg Waters

G.J. Brunskill, D.W.Schindler
S.E.M. Elliott and P.Campbell

Western Region
Fisheries and Marine Service
Department of Fisheries and the Environment
Winnipeg, Manitoba R3T 2N6

September 1979

**Fisheries & Marine Service
Manuscript Report No.1522**

Fisheries and Marine Service

Manuscript Reports

These reports contain scientific and technical information that represents an important contribution to existing knowledge but which for some reason may not be appropriate for primary scientific (i.e. *Journal*) publication. They differ from Technical Reports in terms of subject scope and potential audience: Manuscript Reports deal primarily with national or regional problems and distribution is generally restricted to institutions or individuals located in particular regions of Canada. No restriction is placed on subject matter and the series reflects the broad interests and policies of the Fisheries and Marine Service, namely, fisheries management, technology and development, ocean sciences and aquatic environments relevant to Canada.

Manuscript Reports may be cited as full publications. The correct citation appears above the abstract of each report. Each report will be abstracted by *Aquatic Sciences and Fisheries Abstracts* and will be indexed annually in the Service's index to scientific and technical publications.

Numbers 1-900 in this series were issued as Manuscript Reports (Biological Series) of the Biological Board of Canada, and subsequent to 1937 when the name of the Board was changed by Act of Parliament, as Manuscript Reports (Biological Series) of the Fisheries Research Board of Canada. Numbers 901-1425 were issued as Manuscript Reports of the Fisheries Research Board of Canada. The series name was changed with report number 1426.

Details on the availability of Manuscript Reports in hard copy may be obtained from the issuing establishment indicated on the front cover.

Service des pêches et des sciences de la mer

Manuscrits

Ces rapports contiennent des renseignements scientifiques et techniques qui constituent une contribution importante aux connaissances actuelles mais qui, pour une raison ou pour une autre, ne semblent pas appropriés pour la publication dans un journal scientifique. Ils se distinguent des Rapports techniques par la portée du sujet et le lecteur visé; en effet, ils s'attachent principalement à des problèmes d'ordre national ou régional et la distribution en est généralement limitée aux organismes et aux personnes de régions particulières du Canada. Il n'y a aucune restriction quant au sujet; de fait, la série reflète la vaste gamme des intérêts et des politiques du Service des pêches et de la mer, notamment gestion des pêches; techniques et développement, sciences océaniques et environnements aquatiques, au Canada.

Les Manuscrits peuvent être considérés comme des publications complètes. Le titre exact paraît au haut du résumé de chaque rapport, qui sera publié dans la revue *Aquatic Sciences and Fisheries Abstracts* et qui figuera dans l'index annuel des publications scientifiques et techniques du Service.

Les numéros de 1 à 900 de cette série ont été publiés à titre de manuscrits (Série biologique) de l'Office de biologie du Canada, et après le changement de la désignation de cet organisme par décret du Parlement, en 1937, ont été classés en tant que manuscrits (Série biologique) de l'Office des recherches sur les pêcheries du Canada. Les numéros allant de 901 à 1425 ont été publiés à titre de manuscrits de l'Office des recherches sur les pêcheries du Canada. Le nom de la série a été changé à partir du rapport numéro 1426.

La page couverture porte le nom de l'établissement auteur où l'on peut se procurer les rapports sous couverture cartonnée.



3 5555 00015104 5

Fisheries and Marine Service

Manuscript Report 1522

September 1979

THE ATTENUATION OF LIGHT IN LAKE WINNIPEG WATERS

by

G. J. Brunskill, D. W. Schindler

S. E. M. Elliott, and P. Campbell

Western Region

Fisheries and Marine Service

Department of Fisheries and the Environment

Winnipeg, Manitoba R3T 2N6

This is the 15th Manuscript Report

from the Western Region, Winnipeg

N. H. R. C. LIBRARY

BIBLIOTHEQUE DU C.N.R.C.

© Minister of Supply and Services Canada 1979

Cat. no. Fs 97-4/1522

ISSN 0701-7618

TABLE OF CONTENTS

	Page
ABSTRACT/RESUME	v
INTRODUCTION	1
METHODS	1
RESULTS	1
DISCUSSION	2
ACKNOWLEDGMENTS	4
REFERENCES	4

LIST OF TABLES

<u>Table</u>	<u>Page</u>
1 Secchi disc visibility and visual estimates of color at half the depth of secchi visibility, for stations on Lake Winnipeg in 1969	5
2 Measurements of the percentage of light ($\%T_\alpha$) transmitted from a battery-powered light source to a photosensitive cell over a variable, horizontal path length in the water, and computed horizontal extinction coefficients (K_w) based on this $\%T_\alpha$ data	11
3 Computed values for vertical extinction coefficients, based on measurements of the percentage of surface light reaching several depths in the water column	18
4 Percentage transmission ($\%T_\lambda$) of light (from a bank of fluorescent tubes) at various wavelengths (λ_j) through a 0.1 meter water column, vertical extinction coefficients ($K_{SR\alpha}$) for each wavelength over the 0.1 meter water column, and computed vertical extinction coefficients (\bar{K}_{SR}) and percent transmission (\bar{T}) for $\lambda = 400$ to 1000 μm through a 1.0 meter water column	20
5 Suspended sediment (SS) total dissolved solids (TDS), and total solids (TS) for Lake Winnipeg and Red River stations. Station locations can be found in Fig. 3	23
6 Daily solar radiation at Winnipeg and The Pas for days of our research on Lake Winnipeg	30
7 Total monthly solar radiation expressed as g cal cm^{-2} at Winnipeg and The Pas for 1969 to 1975	33
8 Regression equations for turbidity related parameters and <i>in situ</i> vertical extinction coefficients ($K_w(\text{IS})$) for solar radiation in Lake Winnipeg waters	34

LIST OF TABLES (cont'd)

<u>Table</u>	<u>Page</u>
9 Measured and estimated values for vertical extinction coefficients (K_w) of the transmission of incident light into Lake Winnipeg waters at selected stations	35
LIST OF FIGURES	
<u>Figure</u>	<u>Page</u>
1 The deck tank used on the wheelhouse deck of the Bradbury for light measurements and primary production work	38
2 Comparison of the distribution of light energy of the sun (on a cloudy day) with the light bank of fluorescent tubes as shown in Fig. 1	39
3 Location of stations where radiation measurements were made on Lake Winnipeg	40
4 Isopleths of the depths (meters) of secchi disc visibility in Lake Winnipeg during 4-12 June 1969	41
5 Isopleths of the depths (meters) of secchi disc visibility in Lake Winnipeg during 9-17 July 1969	42
6 Isopleths of the depths (meters) of secchi disc visibility in Lake Winnipeg during 24 July-1 August 1969	43
7 Isopleths of the depths (meters) of secchi disc visibility in Lake Winnipeg during 2-10 September 1969	44
8 Isopleths of the depths (meters) of secchi disc visibility in Lake Winnipeg during 3-13 October 1969	45
9 Isopleths of the depths (meters) of secchi disc visibility in Lake Winnipeg during 26-31 October 1969	46
10 Isopleths of horizontal (beam transmittance) extinction coefficients (K_α , in m^{-1}) for Lake Winnipeg during 9-17 July 1969	47
11 Isopleths of horizontal (beam transmittance) extinction coefficients (K_α , in m^{-1}) for Lake Winnipeg during 24 July-1 August 1969	48
12 Isopleths of horizontal (beam transmittance) extinction coefficients (K_α , in m^{-1}) for Lake Winnipeg during 2-10 September 1969	49
13 Isopleths of horizontal (beam transmittance) extinction coefficients (K_α , in m^{-1}) for Lake Winnipeg during 3-13 October 1969	50
14 Spectral extinction of radiation at 10 cm water depth in the fluorescent-lighted deck tank (Fig. 1) at Station 35 in the center of the north basin of Lake Winnipeg (Fig. 3)	50

LIST OF FIGURES (cont'd)

<u>Figure</u>	<u>Page</u>
on 8 October 1969	51
15 Spectral extinction of radiation at 10 cm water depth in the fluorescent-lighted deck tank (Fig. 1) at Station 58 in the northern part of the south basin (Fig. 3) on 9 September 1969	52
16 Spectral extinction of radiation at 10 cm water depth in the fluorescent-lighted deck tank (Fig. 1) at Station 51 in the mouth of Berens River (Fig. 3) on	53
17 Isopleths of concentrations of suspended sediments (g m^{-3}) in Lake Winnipeg during 9-17 July 1969	54
18 Isopleths of concentrations of suspended sediments (g m^{-3}) in Lake Winnipeg during 24 July-1 August 1969	55
19 Isopleths of concentrations of suspended sediments (g m^{-3}) in Lake Winnipeg during 2-10 September 1969	56
20 Isopleths of concentrations of suspended sediments (g m^{-3}) in Lake Winnipeg during 3-13 October 1969	57
21 Isopleths of concentrations of suspended sediments (g m^{-3}) in Lake Winnipeg during 26-31 October 1969	58
22 Distribution of daily radiation during the days of our research on Lake Winnipeg in 1969 and winter 1970	59
23 The relationship between the depth of secchi disc visibility (m) and vertical extinction coefficients ($K_w(\text{IS})$) for Lake Winnipeg in 1969	67
24 The relationship between vertical extinction coefficients measured in the deck tank (Fig. 1) open to solar radiation ($K_w(\text{DT})$) and an <i>in situ</i> measurement of vertical extinction coefficients ($K_w(\text{IS})$) in Lake Winnipeg waters in 1969	68
25 The relationship between horizontal (beam transmittance) extinction coefficients (K_α) and $K_w(\text{IS})$ in Lake Winnipeg waters in 1969	69
26 The relationship between horizontal (beam transmittance) extinction coefficient (with the transmissometer on an attenuated circuit) ($K_\alpha(A)$) and $K_w(\text{IS})$ in Lake Winnipeg waters in 1969	70
27 The relationship between vertical extinction coefficients ($K_w(\text{IS})$) and the concentration of suspended sediments (g m^{-3}) in Lake Winnipeg waters in 1969	71
28 Isopleths of predicted vertical extinction coefficient (\hat{K}_w , in m^{-1}) in Lake Winnipeg during 4-12 June 1969	72

LIST OF FIGURES (cont'd)

<u>Figure</u>	<u>Page</u>
29 Isopleths of predicted vertical extinction coefficient (\hat{K}_w , in m^{-1}) in Lake Winnipeg during 9-17 July 1969	73
30 Isopleths of predicted vertical extinction coefficient (\hat{K}_w , in m^{-1}) in Lake Winnipeg during 2-10 September 1969	74
31 Isopleths of predicted vertical extinction coefficient (\hat{K}_w , in m^{-1}) in Lake Winnipeg during 24 July-1 August 1969	75
32 Isopleths of predicted vertical extinction coefficient (\hat{K}_w , in m^{-1}) in Lake Winnipeg during 3-13 October 1969	76
33 The relationship between ($K_\alpha - K_w(\text{IS})$) as an estimator of scattering, and the concentration of suspended sediments in g m^{-3} . □ = South Basin, X = North Basin	77
34 The relationships between suspended sediment concentrations, $K_w(\text{IS})$, and the percentage of $K_w(\text{IS})$ due to suspended sediment	78
35 The relationships between chlorophyll α , phytoplankton biomass, $K_w(\text{IS})$, and the percentage of $K_w(\text{IS})$ due to chlorophyll α to phytoplankton biomass	79

ABSTRACT

Brunskill, G. J., D. W. Schindler, S. E. M. Elliott, and P. Campbell. 1979. The attenuation of light in Lake Winnipeg waters. Can. Fish. Mar. Serv. MS Rep. 1522: v + 79 p.

Lake Winnipeg water is usually turbid, and little light penetrates into the lake. Secchi disc visibility varied from 0.1 to 1.0 m in the south basin, and 0.5 to 2.6 m in the north basin. Horizontal extinction coefficients calculated from transmissometer (beam transmittance) data were high ($12\text{-}46 \text{ m}^{-1}$) in the south basin, and lower ($0.6\text{-}3 \text{ m}^{-1}$) in the central part of the north basin. Vertical extinction coefficients were calculated from percentages of surface light reaching the sensor of the submersible photometer cell, and these values were in the range $1\text{-}5 \text{ m}^{-1}$ in the south basin and narrows, and $0.5\text{-}1 \text{ m}^{-1}$ in the center of the north basin. All of these estimates of light transmission through lake water were highly correlated amongst themselves, and with concentrations of suspended sediments. Concentrations of algal biomass, chlorophyll α , and pheophytin were less strongly correlated with light transmission parameters. An attempt was made to estimate the relative proportion of the extinction of light due to scattering by suspended sediments. This was done by subtracting the vertical extinction coefficient from the horizontal (beam transmittance) extinction coefficient.

A sixteen channel spectroradiometer allowed computation of extinction coefficients for individual wavebands. High extinction coefficients were found for blue and green light in turbid waters, with maximum transmission of yellow and red light. Clearer waters of the central north basin had relatively uniform extinction coefficients over the 4,100 to 10,000 Å spectrum.

Key words: beam transmittance; vertical extinction coefficients; limnology; suspended sediments.

RESUME

Brunskill, G. J., D. W. Schindler, S. E. M. Elliott, and P. Campbell. 1979. The attenuation of light in Lake Winnipeg waters. Can. Fish. Mar. Serv. MS Rep. 1522: v + 79 p.

Les eaux du lac Winnipeg sont habituellement troubles, et peu de lumière y pénètre. La visibilité mesurée avec un disque de Secchi varie de 0.1 à 1 m dans le bassin du sud, et de 0.5 à 2.6 m dans le bassin du nord. Les coefficients d'extinction horizontale calculés à partir des données fournies par un transmissomètre à faisceaux dirigés étaient élevés ($12\text{-}46 \text{ m}^{-1}$) dans le bassin du sud et plus bas ($0.6\text{-}3 \text{ m}^{-1}$) dans la partie centrale du bassin du nord. On a calculé les coefficients d'extinction verticale à partir des pourcentages de la lumière superficielle qui atteignent le capteur de la cellule photométrique submersible et obtenu des chiffres de $1\text{-}5 \text{ m}^{-1}$ dans le bassin et les chenaux du sud, et $0.5\text{-}1 \text{ m}^{-1}$ dans le centre du bassin du nord. Toutes ces estimations de la transmission de la lumière à travers l'eau du lac montraient une très forte corrélation entre elles et avec les concentrations de sédiments en suspension. Les concentrations de la biomasse des algues, de la chlorophylle α et de la phaeophytine montraient une corrélation moins forte avec les paramètres de la transmission de la lumière. On a tenté d'estimer la proportion relative de l'extinction de la lumière qui est due à la dispersion par les sédiments en suspension, en soustrayant le coefficient d'extinction verticale du coefficient d'extinction horizontale (faisceaux dirigés).

Un spectroradiomètre à 16 canaux a permis de calculer les coefficients d'extinction pour des gammes d'ondes précises. Les coefficients se sont révélés hauts pour la lumière bleue et verte dans les eaux troubles, la transmission étant maximale pour les gammes du jaune et de rouge. Les eaux plus claires du bassin central du nord présentaient des coefficients d'extinction relativement uniformes situés au-dessus du spectre 4,000-10,000 Å.

Mots-clés: faisceaux dirigés; absorptivité, verticale; limnologie; sédiments en suspension.

INTRODUCTION

Franklin (1823) and Back (1936) noted the great turbidity of Lake Winnipeg waters while trying to avoid running their canoes onto rocks in the shallow water. Bajkov (1930) found that less turbid parts of Lake Winnipeg had larger abundances of zoobenthos, fish, and plankton. Brunskill (1973) and Brunskill, Schindler et al. (1979) showed that Lake Winnipeg phytoplankton had nutrient supplies in excess of their requirements for growth, and that extreme light attenuation by these turbid waters likely limits the growth of photosynthetic organisms, especially in the south basin of the lake. In this paper, we give an analysis of Lake Winnipeg light attenuation data collected in 1969.

Three factors (suspended inorganic sediments, phytoplankton, and dissolved organic matter in the water) likely control the depth of penetration of light into the lake water. The lake is shallow ($\bar{z} = 12 \text{ m}$) and large in surface area ($A_0 = 23,750 \text{ km}^2$). Strong winds over such a large and shallow water body frequently cause seiches, storm surges and large waves (Einarsson and Lowe 1968) which cause considerable resuspension of sediments. The Red River supplies $3-11 \times 10^6$ tonnes of sediments per year to the south basin of the lake (Brunskill 1973; Brunskill and Graham 1979). During calm weather in summer, dense algal blooms of *Alphanizomenon flos-aquae* and *Anabaena spp.* occur in the upper 0.5-1.0 m of the water column (Bajkov 1934; Brunskill, Schindler et al. 1979). In many bays and river mouths along the east shore of the lake, darkly-stained, humic acid-rich waters from Precambrian Shield watersheds are found. Watershed characteristics, sediment supply, hydrology, morphometry, nutrient chemistry and phytoplankton biology of Lake Winnipeg are given elsewhere (Brunskill, Elliott et al. 1979; Brunskill and Graham 1979; Brunskill, Schindler et al. 1979).

METHODS

Secchi disc visibility was measured *in situ* or in deck tanks (Fig. 1) with a 25 cm diameter secchi disc which was painted in black and white quadrants. The estimated precision for visibility was $\pm 5 \text{ cm}$ in most cases. Visual colour of the disc was also recorded at half secchi depth. Percentage light transmittance of lake water at selected stations was also estimated with a Whitney submersible photometer Model 8A by setting the meter to 100% with the sensor cell above water, and then lowering the cell to 10 to 300 cm water depth *in situ* or in deck tank water (Fig. 1). Vertical extinction coefficients (K_ω) were estimated from the slope of the regression line between $(\ln \% \text{ surface light})$ and water depth, ignoring the depth interval 0-10 cm, or were calculated from the attenuation of irradiation measured by this unit according to the equation of Vollenweider (1969):

$$K_\omega = \frac{\varepsilon^\lambda}{V} = \frac{1}{Z_2 - Z_1} (\ln i_1 - \ln i_2) \quad (\text{Eq. 1})$$

where $K_\omega = \frac{\varepsilon^\lambda}{V}$ = light extinction coefficient,

Z_1, Z_2 = depth intervals, below 10 cm meter depth

i_1, i_2 = light intensities at depths Z_1 and Z_2 , respectively

An estimate of horizontal extinction of light (beam transmittance) was obtained from the use of a Hydroproducts Model 412 Transmissometer. This instrument was calibrated to read 86% transmittance in air at 1 meter light path and 98-100% transmittance in a laboratory tank of distilled water. The distance between the light source and photocell was varied according to the turbidity of the water: in the north basin this distance was usually 50 cm, and in the south basin, 10 to 25 cm. The percentage transmission of light from the light source to the photocell ($\%T$) was used to calculate a horizontal extinction coefficient (K_ω) from the equation:

$$K_\omega = \ln \frac{100}{\%T} d^{-1} \quad (\text{Eq. 2})$$

where d = path length in meters.

Vertical extinction coefficients for light of wavelength (λ) from 4000 Å to 10,000 Å, and for specific wavebands, were obtained from an Agro-products Spectroradiometer. The sensor head of this apparatus was firmly fixed 10 cm below water level in a large tank (1.2 m deep, 0.7 m diameter). The residence time of water in this tank was approximately 20 minutes. This tank was mounted in a deck table over which a tight fitting lid closed (Fig. 1). The lid had a bank of eight fluorescent 80-watt tubes, approximately 1 m long, backed with aluminum foil, which was used as a constant light source for the Spectroradiometer sensor. This light source spectrum is compared to solar radiation on a cloudy day in Fig. 2. The plastic tank was painted black to exclude light from entering from the side walls of the tank. Extinction coefficients for 400 to 1,000 mμ, and for individual wavebands were calculated according to Eq. 1. Calibration of this spectroradiometer against a known light source in a dark room gave 87-116% of the true value.

Suspended sediments were separated from water samples by vacuum filtration through preweighed membrane or glass fibre filters (0.45-1.2 μm nominal pore diameter) on the ship. These filters were desiccated, frozen for storage, and in the laboratory they were dried to 100°C and weighed. Chlorophyll methods are reported in Brunskill, Schindler et al. (1979).

As discussed in Brunskill, Elliott et al. (1979) and Brunskill and Graham (1979) stations were located by radar, gyrocompass, and sonar on Canadian Hydrographic Service Charts 6241 and 6240. Station locations referred to in the following tables are shown in Fig. 3.

RESULTS

Table 1 gives the results of our measurements of Secchi disc visibility and water color against the white disc at half the depth of Secchi visibility. Minimal values of 0.1-0.2 m were recorded in the plume of the Red River throughout the open water season, and maximal values of 2-3.5 m were observed in the central part of the north basin. The variation of Secchi disc visibility over the open-water season of 1969 is shown in Figs. 4-9. Although the secchi rope was marked in 0.1 m segments, the measurement of secchi visibility in the very turbid waters of the southern part of the south basin, and along the north and west lake shores of the north basin, will have greater

relative error than in less turbid parts of the lake. Secchi visibility in the Red River was often <0.1 m, whereas in the Winnipeg and Saskatchewan Rivers, Secchi visibility was in the range 1-3 m. Visual color at half secchi depth was usually yellow-brown in turbid water areas of the south basin. Orange-brown colored water occurred near tributaries draining Precambrian Shield to the east of the lake. With the exception of the nearshore or near-island regions, north basin water was green or blue-green.

Table 2 shows results of our beam transmittance measurements, given as percent transmission per path length ($\%T_\lambda$) and a horizontal extinction coefficient (K_α). High K_α 's (ranging from 12-46 m^{-1}) were found in the Red River plume, near shores and islands. Lower values of K_α occurred in the central part of the north basin, and occasionally in the central and northern part of the south basin. These lower values ranged from 0.6 to 3 m^{-1} . The variation of K_α is also shown in Figs. 10-13. Replicate measurements, varying light path length and water depth at Station 35, gave similar results (Table 2). From these measurements, we concluded that light transmittance should be relatively uniform with depth at stations away from shores and river plumes.

Table 3 gives estimates of vertical extinction coefficients (K_ω) obtained from measurements of the percentage of surface radiation reaching several water depths. Some of these measurements were done *in situ* from the unshaded side of the ship (K_ω (IS)), and other measurements were made in lake water-flushed deck tanks (K_ω (DT)) (see Fig. 1) open to sunlight. K_ω (DT) values were always greater than K_ω (IS), probably because of the exclusion of laterally scattered radiation by the black-painted walls of the deck tank. Throughout this paper, K_ω (IS) will be used as our best estimate of the vertical extinction coefficients. Higher values of K_ω (IS) occurred in the Red River plume and near shores and islands, ranging from 1.5-4.5 m^{-1} . Lower values occurred in the center of the north basin, where values of K_ω (IS) were from 0.5 to 1 m^{-1} . K_ω (DT) varied in a similar manner but was usually 2 to 3 fold greater than the value of K_ω (IS).

Table 4 gives results of the measurement of percentage transmission of fluorescent light (Fig. 1, 2) through 10 cm of Lake Winnipeg waters at selected wavelength ($\%T_\lambda$), and an extinction coefficient for each wave band ($K_{SR\lambda}$) over $\lambda_i = 405$ to 1,000 μm . It is difficult to generally discuss this data, as considerable variation occurs in spectra for similar water masses. Highest $K_{SR\lambda}$ were usually in the region $\lambda = 405$ -600 μm , and lower values were often in the region $\lambda = 640$ -900 μm . During the season when this data was collected (September-October), the discharge of the Red River was greatly reduced (Brunskill, Elliott et al. 1979), and the difference between the turbidity of the south and north basin is not as great as in June, July and August (see Figs. 4-6, and compare Figs. 7-9). Nevertheless, $K_{SR\lambda}$ for most λ were much lower in open water north basin stations compared to south basin stations (Figs. 14, 15). Humic acid colored water from Precambrian Shield drainage exhibited high $K_{SR\lambda}$ at $\lambda = 430$ -440 μm (Fig. 16) and decreased to lower values of $K_{SR\lambda}$

above 700 μm . Data for K_λ (IS) (Table 3) and $K_{SR\lambda}$ data were not collected simultaneously, unfortunately.

As we shall see in the discussion, most of our light transmission data are closely correlated with suspended sediment concentrations. We therefore give suspended sediment data in Table 5 and Figs. 17-21. Suspended sediment concentrations were highest (5-80 g m^{-3}) near the mouth of the Red River and near the west and southwest shores of the south basin. Lower values (1-20 g m^{-3}) are found throughout most of the narrows and north basin. The Red River (Station 0 in Table 5) is obviously a major source of sediments for the south basin of Lake Winnipeg, whereas the Saskatchewan (Station 29) and Winnipeg Rivers (Station 7) contribute much less, despite their larger annual discharges (Brunskill, Elliott et al. 1979). The Red River drains a sedimentary watershed with agricultural activities and relatively large populations of humans and livestock, whereas the Winnipeg River drains a Precambrian Shield watershed which has little agriculture and smaller human populations. The Saskatchewan River drains a sedimentary watershed, but most of its sediment load is deposited in its delta (Cedar and Moose Lake area) and recently in reservoirs along its course (Brunskill, Elliott et al. 1979).

Hourly, daily, and monthly solar radiation data is given in Tables 6 and 7, and Fig. 22a-h, for the Winnipeg airport and The Pas, the closest weather stations.

DISCUSSION

For purposes of estimating the attenuation of photosynthetically useful radiation in Lake Winnipeg water, the vertical extinction coefficients measured *in situ* with the Whitney submersible photometer (K_ω (IS)) will be considered the best value. The other measurements of light attenuation will now be related to these data (Table 3).

Secchi disc visibility (Table 1) was closely related to K_ω (IS), as shown in Fig. 23. This is a commonly-found relationship in diverse waters (Jones and Wills 1956; Tyler 1968; Holmes 1970; Schindler 1971). Measurements of vertical extinction coefficients with the Whitney submersible photometer in the deck tank (K_ω (DT), Table 3) was also highly correlated with K_ω (IS) (Fig. 24), but with more variation about the regression line. As previously mentioned, this larger variation and larger absolute value of K_ω (DT) is likely due to the interaction of the black-painted walls of the deck tank and the varying angle of the sun, limiting the amount of lateral and downward scattered radiation. There was an excellent relationship between horizontal extinction (beam transmittance from the transmissometer, K_α , Table 2), and K_ω (IS) (Fig. 25). A different slope and intercept was found for the correlation between K_ω (IS) and K_ω (A) measured on the attenuated circuit of the transmissometer (Fig. 26) in extremely turbid waters. Statistically significant correlations were found between K_ω (IS) and chlorophyll a , pheophytin, and algal biomass (Table 8), but the confidence limits of the regression lines are too great for practical use. The algal biomass, chlorophyll a , and the

pheophytin data used here are taken from Brunskill, Schindler et al. (1970). K_w (IS) was highly correlated with suspended sediment concentration (Fig. 27).

The equations (Table 8) from the above correlations (Figs. 23-27) were used to estimate a vertical extinction coefficient (\bar{K}_w) where K_w (IS) was not measured. This allowed the estimation of K_w for stations done at night and during rough weather. These data are given in Table 9, along with a code to indicate the origin of the estimate. The equations in Table 8 are listed in their order of reliability, such that K_w predicted from algal biomass, chlorophyll a , and pheophytin are to be least trusted. K_w is shown also in Figs. 28-32.

Ideally K_w (IS) and \bar{K}_w should estimate the attenuation of radiation due to absorption and backscattering by water, dissolved organic matter, and particulate matter. K_x and K_x (A) should estimate the attenuation of radiation due to absorption and total scattering by water, dissolved organic matter, and particulate matter. The transmissometer data (K_α) should then be more sensitive to particulate matter (scattering) in the water column. This appears to be true for the values of K_x and K_w in our data, as K_x varied from 2 to 35 m^{-1} in the south basin of Lake Winnipeg in mid July (Fig. 10), whereas \bar{K}_w or K_w (IS) was in the range 1 to 4.6 m^{-1} (Fig. 29). In both cases, however, the variation of K_w and K_α was 4-16 fold across this gradient of suspended sediments (Fig. 17). K_α responded to a largely inorganic suspended sediment gradient in the northern part of the north basin (Fig. 10 and Table 2, Stations 23C to 33) by a 25 fold change (0.8 to 25 m^{-1}), whereas K_w responded with a 15 fold change (0.4 to 6.7 m^{-1} , see Fig. 29 and Table 9). We attempted to partition the components contributing to the measured extinction by approaches similar to those of Tyler (1975), but the heterogeneous suspended sediment parameter completely overwhelmed any derived relationships. A crude and relative estimate of the proportion of K_α that is due to scattering by suspended sediments is obtained by subtracting K_w (IS) from K_α . That ($K_\alpha - K_w$ (IS)) is well correlated with suspended sediment concentration is no surprise (Fig. 33), but this computation indicates that a large fraction of K_α (=50-80%) is related to light scattering by the water column.

Another way to illustrate the role of suspended sediments (SS) in controlling extinction coefficients of Lake Winnipeg waters is given in Fig. 34. This relationship is derived from equation D in Table 8 where:

$$K_w(\text{IS}) = 0.5809 + 0.1227 \text{SS},$$

and:

$$\% \text{ attenuation due to SS} = \frac{0.1227}{K_w(\text{IS})} \times 100$$

This manipulation indicates that scattering and absorbance explained by SS accounts for 50-90% of K_w (IS) over the upper range of SS common in Lake Winnipeg (Table 5). The composition of SS varied considerably. Most samples of SS appeared under the microscope to be largely inorganic matter, whereas some samples contained dense phytoplankton blooms. A similar manipulation on

equation G in Table 8 allowed the construction of Fig. 35, which indicates that 10-60% of K_w (IS) can be controlled by the range of chlorophyll a concentrations observed (1 to 20 mg Chl a m^{-3} , see Brunskill, Schindler et al. 1979). Figures 34 and 35 indicate that at the concentrations of chlorophyll a , phytoplankton biomass, and suspended sediments considered here, phytoplankton are more effective in attenuating light than are suspended sediments, on a per unit weight basis. Ganf (1974) used a variation of this approach in his studies on Lake George (Uganda).

The data on attenuation of light by wavelength in Table 4 and Figs. 14-16 indicate the extreme effect of scattering of light by high suspended sediment concentrations. Attenuation maxima usually occurred in the violet, blue and green wavebands (405-546 μm), and lower extinction coefficients were usually found in the red and infrared (640-900 μm). These spectra are greatly different from the "standard distribution" curves of Vollenweider (1961). According to Hutchinson (1957) and Strickland (1958), particulate matter is very important in scattering and attenuating light below 500 μm , and less effective for wavelengths $>600 \mu\text{m}$. Since there is less energy in solar radiation at the lake surface above 700 μm (see Fig. 2), and most of the violet, blue and green light is strongly attenuated (Figs. 15 and 16), the yellow, orange, and red wavelengths (546-700 μm) will have maximum penetration in the turbid areas of Lake Winnipeg. As indicated in Table 1, lake water color in most of the south basin, narrows, and near shore north basin was observed as yellow-brown (or orange-brown in humic acid-rich waters). Clearer waters of the central north basin had relatively flat and lower extinction spectra over 405-1,000 μm (Fig. 14), and were observed as being green or blue-green in visual color over the secchi disc. Even the lower values for vertical extinction coefficients given here are quite high compared to other lake data (Hutchinson 1957; James and Birge 1938; Whitney 1938; Thomson and Jerome 1975; Thomson et al. 1974). In some cases in Lake Winnipeg (extreme turbidity in the Red River plume or near shore, or in the midst of dense algal blooms), water color is more likely due to the color of suspended matter ("seston color" of Hutchinson (1957, p. 417)). We observed large ($\approx 200 \times 400 \text{ m}$) patches of dark water in both the north and south basin of the lake, which proved to be near surface (0-50 cm) accumulations of insect exuviae. During dense algal blooms in calm weather in July and August in the north and south basins, the surface of the water appeared to be covered with opaque, light green or yellow-green paint. The algal bloom was in the upper 30 cm of the water column, and often large clumps of cells and filaments floated on the surface.

The spectroradiometer data for waters of the Red River plume (Stations 05, 59, 59-60, 61, 62 in Table 4) appear to corroborate the Cree name for the Red River (Mikwakumew sepe = Bloody River) according to La Verendrye in 1735 (see Crouse 1972, p. 42; Hamilton, no date, p. 235), who was told the river had the color of vermillion. Wavelengths of maximum transmission for these waters are in the red. Seasonal variation in winds, river discharge, shore erosion, and algal biomass are likely factors controlling the seasonal variation in extinction coefficients in Lake Winnipeg.

Future work on the optical characteristics of this turbid lake should investigate many of the curious extinction spectra in Table 4. It seems possible that more precise relationships between suspended inorganic matter, detrital organic matter, dissolved organic matter, and algal cells could be obtained if these parameters could be partitioned and sized (Haffner and Evans 1974; Burt 1955), including determinations of particle surface areas, volume, and indices of refraction.

ACKNOWLEDGMENTS

Captain Kris Thorsteinsen and the crew of the Bradbury are gratefully thanked for their patience with our demands to hold the ship in one location without stirring up bottom mud. F. A. J. Armstrong, F. C. Elder, and D. Povoledo gave advice and criticism. B. W. Graham, J. Shearer, and R. V. Schmidt got very cold and wet taking light measurements when they could have been having tea in the galley. We thank Bob Chapel of Canada Centre for Inland Waters for the loan of the spectroradiometer. C. Jones, W. Burton, and B. van der Veen constructed the deck tank and light bank. They also repaired it after each storm on the lake. S. Zettler, L. Davies, and L. Taite did the drafting. C. Anderson, A. Prefontaine, M. Langtry and D. Laroque typed the manuscript. J. R. W. Vallentyne provided spiritual guidance. R. Hecky and E. Fee are thanked for critically reviewing the manuscript.

REFERENCES

- BACK, G. 1836. Narrative of the Arctic Land Expedition to the mouth of the Great Fish River...in 1833, 34, and 35. John Murray, London. 663 p.
- BAJKOV, A. 1930. Biological conditions of Manitoban lakes. *Contrib. Can. Biol. Fish.* 5(12): 383-422.
- . 1934. The plankton of Lake Winnipeg drainage system. *Int. Rev. Gesamten Hydrobiol. Hydrogr.* 31: 239-272.
- BRUNSKILL, G. J. 1973. Rates of supply of nitrogen and phosphorus to Lake Winnipeg, Manitoba, Canada. *Int. Ver. theor. angew. Limnol. Verh.* 18: 1755-1759.
- BRUNSKILL, G. J., S. E. M. ELLIOTT, and P. CAMPBELL. 1979. Morphometry, hydrology, and watershed data pertinent to the limnology of Lake Winnipeg. *Can. Fish. Mar. Serv. MS Rep. (in prep.)*.
- BRUNSKILL, G. J., and B. W. GRAHAM. 1979. The offshore sediments of Lake Winnipeg. *Can. Fish. Mar. Serv. MS Rep. (in prep.)*.
- BRUNSKILL, G. J., D. W. SCHINDLER, S. K. HOLMGREN, H. KLING, P. CAMPBELL, M. P. STAINTON, and F. A. J. ARMSTRONG. 1979. Nutrients, chlorophyll, phytoplankton, and primary production in Lake Winnipeg. *Can. Fish. Mar. Serv. MS Rep. (in prep.)*.
- BURT, W. V. 1955. Interpretation of spectro-photometer readings on Chesapeake Bay water. *J. Mar. Res.* 14: 33-46.
- CROUSE, N. M. 1972. La Verendrye, fur trader and explorer. Kennikat Press, Port Washington, N.Y. 247 p.
- EINARSSON, E., and A. B. LOWE. 1968. Seiches and set-up on Lake Winnipeg. *Limnol. Oceanogr.* 13(2): 257-271.
- FRANKLIN, J. 1823 (1969). Narrative of a journey to the shores of the Polar Sea in the years 1819, 20, 21, and 22. John Murray, London (Reprint by M. G. Hurtig). 768 p.
- GANF, G. G. 1974. Incident solar irradiance and underwater light penetration as factors controlling the chlorophyll *a* content of a shallow equatorial lake (Lake George, Uganda). *J. Ecol.* 62(2): 593-609.
- HAFFNER, G. D., and J. H. EVANS. 1974. Relation of light penetration to particle distribution in vertically mixed lacustrine environments. *Br. Phycol. J.* 9: 261-267.
- HAMILTON, G. No date. In the beginning. Derksen Printers Ltd., Steinbach, Manitoba. 362 p.
- HOLMES, R. W. 1970. The secchi disc in turbid coastal waters. *Limnol. Oceanogr.* 15(5): 688-694.
- HUTCHINSON, G. E. 1957. A treatise on limnology. Vol. 1. John Wiley & Sons, Inc., London. 1015 p.
- JAMES, H. R., and E. A. BIRGE. 1938. A laboratory study of the absorption of light by lake waters. *Trans. Wis. Acad. Sci. Arts Lett.* 31: 1-154.
- JONES, D., and M. S. WILLS. 1956. The attenuation of light in sea and estuarine waters in relation to the concentration of suspended solid matter. *J. Mar. Biol. Assoc. U. K.* 35: 431-444.
- SCHINDLER, D. W. 1971. Light, temperature, and oxygen regimes of selected lakes in the Experimental Lakes Area, northwestern Ontario. *J. Fish. Res. Board Can.* 28(2): 157-169.
- STRICKLAND, J. D. H. 1958. Solar radiation penetrating the ocean. A review of requirements, data, and methods of measurement, with particular reference to photosynthetic productivity. *J. Fish. Res. Board Can.* 15(3): 453-493.
- THOMSON, K. P. B., and J. JEROME. 1975. Transmissometer measurements of the Great Lakes. *Can. Dep. Environ. Inland Waters Div., Sci. Ser.* 53: 10 p.
- THOMSON, K. P. B., J. JEROME, and W. R. MCNEIL. 1974. Optical properties of the Great Lakes (IFYGL). *Proc. Conf. Great Lakes Res.* 17: 811-822.
- TYLER, J. E. 1968. The Secchi Disc. *Limnol. Oceanogr.* 13(1): 1-6.
- . 1975. The *in situ* quantum efficiency of natural phytoplankton populations. *Limnol. Oceanogr.* 29(6): 976-980.
- VOLLENWEIDER, R. A. 1961. Photometric studies in inland waters: I. Relations existing in the spectral extinction of light in water. *Mem. Ist Ital. Idrobiol.* 13: 87-113.
- . (ed.). 1969. A manual on methods for measuring primary production in aquatic environments. IBP Handb. Blackwell Scientific Publ., Edinburgh. 213 p.
- WHITNEY, L. V. 1938. Transmission of solar energy and the scattering produced by suspensoids in lake waters. *Trans. Wis. Acad. Sci. Arts Lett.* 31: 201-221.

p.
n

Table 1. Secchi disc visibility and visual estimates of color at half the depth of Secchi visibility, for stations on Lake Winnipeg in 1969. See Fig. 3 for station locations. TU = Turbid.

STATION NO. DATE SECCHI DISC
VISIBILITY (METERS)

COLOUR AT HALF SECCHI DEPTH
(BA=BLACK;BL=BLUE;BR=BRWN;G,GR=GREEN;
GA=GRAY;HU=HUMIC;NITE=NIGHT;OR=ORANGE;
RU=RUST;Y,YF=YELLOW)

0	27OCT68	0.70	YF TUP
0	3JUN69	0.28	GYRRTU
0A	5NOV68	*****	YHRGTU
0B	5NOV68	*****	YORTU
1	4JUN69	0.30	YFRRTU
1	9JUL69	0.35	YERR
1	3OCT69	0.50	YFRR
2	28OCT68	0.05	
2	23APR69	*****	RRTU
2	4JUN69	0.25	YFARTU
2	9JUL69	0.30	
2	24JUL69	0.30	
2	2SEP69	0.50	YFRR
2	3OCT69	0.20	YFRR
2	17MAR70	*****	TU
3	9JUL69	0.25	
3	24JUL69	0.35	YFRR
3	10SEP69	0.20	BR
3	3OCT69	0.10	YFRR
3C	9JUL69	0.25	YFRR
3C	24JUL69	0.20	
3C	10SEP69	0.70	GRYFRR
3C	3OCT69	0.40	YFRRGA
4	9JUL69	0.30	YFRR
4	10SEP69	0.25	BR
4	3OCT69	0.45	YFRR
4	31OCT69	0.30	YFRR
5	4JUN69	0.20	YFRTU
5	9JUL69	0.60	
5	25JUL69	0.60	YFRR
5	9SEP69	0.50	YFRR
5	4OCT69	0.65	GRYERR
6	4JIN69	0.32	YFRTU
6	10JUL69	0.60	BR
6	26JUL69	0.80	YFRR
6	9SEP69	0.80	BROR
6	4OCT69	0.45	YFRR
7	28OCT68	0.80	YFLBRC
7	5JUN69	1.00	
7	10JUL69	0.80	YFRR
7	26JUL69	0.75	GRRR
7	9SEP69	0.50	BR
7	4OCT69	0.80	
7	31OCT69	1.00	BR
8	28OCT68	0.15	
8	5JUN69	0.50	YFARTU
8	10JUL69	0.80	GRRRHII
8	26JUL69	0.85	YFRR
8	9SEP69	0.75	YFORRR
8	4OCT69	0.40	YFRR
8	31OCT69	0.90	BR
9	10JUL69	0.70	YFRR
9	26JUL69	0.75	YFRR
9	4OCT69	0.20	YFRR
10	5JUN69	0.60	YFRTU
10	10JUL69	0.90	YFRR
10	26JUL69	1.30	YFRR
10	8SEP69	0.50	ORRR
10A	28OCT68	0.30	BROTUR
11	5JUN69	0.90	YFRTII
11	10JUL69	1.30	
11	27OCT69	0.50	YFRTU
12	5JUN69	1.50	YF GR
12	10JUL69	1.10	
12	26JUL69	1.00	GRYFRR
12	8SEP69	0.50	ORRR

STATION NO. DATE SECCHI DISC
VISIBILITY (METERS)

COLOUR AT HALF SECCHI DEPTH
(BA=BLACK;BL=BLUE;BR=BRWN;G,GR=GREEN;
GA=GRAY;HU=HUMIC;NITE=NIGHT;OR=ORANGE;
RU=RUST;Y,YE=YELLOW)

12	30OCT69	0.40	
13B	10JUL69	1.15	YFRRTU
13C	29OCT69	1.00	YFRPHU
14	5JUN69	1.50	YFRPTU
14	10JUL69	1.00	YF GR
14	26JUL69	0.70	YFRR
14	8SEP69	0.50	YFRR
14	5OCT69	0.35	BR
14	30OCT69	0.30	RRTU
16	6JUN69	1.20	GRYERR
16	11JUL69	0.60	YFRRHII
16	27JUL69	*****	YFRR
16	3SEP69	0.05	BR
16	5OCT69	0.40	YFRRTU
16	28OCT69	0.35	YFRRTU
17	6JUN69	0.70	GYPRTU
17	12JUL69	0.75	
17	27JUL69	1.30	GR
18	6JUN69	0.70	YFRRTU
18	11JUL69	0.60	YFRR
18	27JUL69	1.10	GRYE
18	3SEP69	0.80	GR
19	7JUN69	0.85	
19	13JUL69	1.00	GRYE
19	28JUL69	1.20	GR
19	3SEP69	1.00	
20	7JUN69	0.90	YERRGR
20	13JUL69	1.10	GR
20	28JUL69	1.35	GRYE
21	7JUN69	0.95	GYRRTU
21	13JUL69	0.90	GR
21	28JUL69	1.80	GR
21	4SEP69	0.60	GP
21	11OCT69	1.00	GP
22	7JUN69	1.20	GR TU
22	13JUL69	1.30	GR
22	28JUL69	1.40	GRFFN
22	4SEP69	1.50	
22	11OCT69	0.90	GR
23B	7JUN69	*****	GR
23C	7JUN69	*****	GR
23C	13JUL69	2.50	GR
23C	28JUL69	1.80	GRTU
23C	6SEP69	2.00	GRRR
23C	11OCT69	1.40	GP
23D	7JUN69	1.10	
23E	7JUN69	1.30	GR
23E	15JUL69	2.00	GP
23E	30JUL69	1.80	GRFFN
23E	6SEP69	1.60	GRRR
23E	11OCT69	1.00	GR
24	7JUN69	1.90	
25	7JUN69	1.90	GRFFN
25	13JUL69	2.50	GR
25	28JUL69	3.00	GRTU
25	4SEP69	1.50	BLGR
25	29OCT69	1.00	GR
26	14JUL69	2.50	GR
26	29JUL69	3.00	BL YEGY
26	4SEP69	1.50	BLGR
26	9OCT69	0.40	YFRRTU
27	14JUL69	1.10	GR
27	29JUL69	1.20	YFGRTU
27	9OCT69	0.25	YFRRTU
28	13JUL69	2.50	BLGR
28	28JUL69	2.20	BLGR

STATION NO.	DATE	SECCHI DISC VISIBILITY (METERS)	COLOUR AT HALF SECCHI DEPTH (BA=BLACK;BL=BLUF;BR=BRWNIG;GR=GREFN; GA=GRAY;HU=HUMIC;NITE=NIGHT;OR=ORANGE; RU=RUST;Y,YELLLOW)
28	4SFP69	1.00	GR
28	29OCT69	1.30	
29	20MAR69	*****	BL
29	7OCT69	1.50	BLGR
30	18MAR69	*****	YFGRRR
31	9JUN69	1.50	
31	14JUL69	1.50	GR
31	29JUL69	1.90	GRTU
31	6SFP69	0.90	GRRR
31	8OCT69	*****	TU
33	9JUN69	0.50	GRTU
33	14JUL69	0.30	YFAR
33	29JUL69	0.60	GRTU
35	18MAR69	*****	GR
35	9JUN69	2.20	GRTU
35	14JUL69	2.00	GR
35	29JUL69	2.10	GR
35	6SEP69	1.80	GP
35	8OCT69	1.30	GRTU
39	1APR69	*****	GRTU
39	15JUL69	2.40	GR
39	30JUL69	2.60	GREFN
39	6OCT69	1.00	
39	29OCT69	0.85	GR
41	10JUN69	0.90	GRYFTU
41	15JUL69	2.00	GR
41	30JUL69	2.40	GREFN
41	7SFP69	1.90	GRYE
41	6OCT69	0.85	GRTU
43	10JUN69	0.40	GRYFTU
43	15JUL69	1.80	GR
43	30JUL69	1.40	GRTU
43	6OCT69	0.45	GR
45	15JUL69	2.30	GP
45	30JUL69	2.50	GRYE
45	7SFP69	1.20	GR
45	6OCT69	1.10	GRTU
48	2APR69	*****	GRTU
48	15JUL69	0.90	GRYF
48	30JUL69	1.00	YFGRTU
48	7SEP69	1.00	YFAR
48	12OCT69	0.85	GRTU
50B	11JUL69	0.50	YFAR
50B	27JUL69	0.50	YFAR
50C	11JUL69	0.50	YFAR
50C	27JUL69	0.60	YFAR
50C	3SFP69	0.50	GRYFBR
51	30OCT68	1.00	BR ORA
51	11JUN69	0.60	RED-PR
51	11JUL69	0.90	HURR
51	27JUL69	0.85	BRHU
51	3SFP69	0.25	BRBA
51	5OCT69	0.80	BROR
52	11JUN69	0.60	RURRTU
52	11JUL69	0.60	HUYFAR
52	3SFP69	0.25	
52	5OCT69	0.30	YFARTU
54	11JUN69	0.80	YFRRGR
54	11JUL69	0.70	YFAR
54	8SFP69	0.45	
54	5OCT69	0.30	YFARTU
54	30OCT69	0.30	YFARTU
54B	29OCT68	0.30	YFARTU
54B	28OCT69	0.25	YFARTU
55	11JUN69	0.50	
56	29OCT68	0.60	YFARTU

STATION NO.	DATE	SECCHI DISC VISIBILITY (METERS)	COLOUR AT HALF SECCHI DEPTH (BA=BLACK; BL=PLUS; BR=BRWN; G=GREEN; GA=GRAY; HU=HUMIC; NITE=NIGHT; OR=ORANGE; RU=RUSTY; YF=YELLOW)
57	31OCT68	0.70	
57	12JUN69	0.60	GYRRTU
57	10JUL69	0.90	YERR
57	25JUL69	0.90	YFGRRR
57	8SEP69	0.50	GRRR
58	12JUN69	0.30	GYBRTU
58	10JUL69	0.40	YFPR
58	25JUL69	0.40	YEGRPP
58	9SEP69	0.15	YFRR
59	31OCT68	0.50	
59	12JUN69	0.50	
59	9JUL69	0.70	
59	25JUL69	0.95	GRYFRR
59	9SEP69	1.00	GRYFRR
59	4OCT69	1.00	GRYE
59	31OCT69	0.45	YFRR
60	23APR69	3.50	GR TUR
60	12JUN69	0.30	YFRTTU
60	9JUL69	0.40	
60	24JUL69	0.90	YFGRRR
60	2SEP69	1.00	
60	9SEP69	0.70	GRRR
60	31OCT69	0.55	YFRR
60B	9JUL69	0.25	
60B	24JUL69	0.35	YEGRHP
60B	9SEP69	0.45	GRRR
60B	4OCT69	0.40	YFRR
60C	28OCT68	0.15	
60C	24JUL69	0.25	YFRR
60C	9SEP69	0.35	RR
60C	18MAR70	*****	TU
61	12JUN69	0.20	YFRTTU
61	9JUL69	0.40	
61	24JUL69	0.75	GRYFRR
61	2SEP69	0.90	YFRR
61	10SEP69	0.90	YFRR
61	3OCT69	0.45	YFRR
61	31OCT69	0.45	GR
62	31OCT68	0.12	
62	12JUN69	0.20	
62	9JUL69	0.25	
62	24JUL69	0.50	GRYFRR
62	10SEP69	0.50	YFRR
62	3OCT69	0.20	YFRR
63	12JUN69	0.10	
63	17JUL69	0.40	YFRR
63	1AUG69	0.20	YFRTTU
63	10SEP69	0.20	RR
63	13OCT69	0.20	YFRTTU
63	31OCT69	0.35	YFAR
64	30OCT68	0.50	YFRTTU
64	15JUL69	0.90	GR
64	31JUL69	0.50	YFRTTU
64	7SEP69	0.50	YFRR
64	12OCT69	0.30	YFARTU
64B	12OCT69	*****	RPTU
65	16JUL69	0.50	GHYETU
65	31JUL69	0.55	YFRTTU
66B	31JUL69	0.70	GRTU
67	2APR69	*****	GR
68	16JUL69	0.50	GRYFTU
68	31JUL69	0.40	YGTURR
68	8SEP69	0.40	YFRR
68	12OCT69	0.35	YFRTTU
68	30OCT69	0.30	YFRTTU
68B	12OCT69	*****	YFRTTU

STATION NO.	DATE	SECCHI DISC VISIBILITY (METERS)	COLOUR AT HALF SECCHI DEPTH (BA=BLACK;BL=BLUF;BR=BRWN;G=GREEN; GA=GRAY;HU=HUMIC;NITF=NIGHT;OR=ORANGE; RU=RUST;Y,YF=YELLOW)
69	16JUL69	0.40	GRYF
69	31JUL69	0.50	YBRTU
69	8SEP69	0.50	YERR
69	12OCT69	0.25	YFARTU
69B	28OCT69	0.20	YFARTU
70	20MAR69	*****	GR

Table 2. Measurements of the percentage of light ($\% T_a$) transmitted from a battery-powered light source to a photosensitive cell over a variable, horizontal path length in the lake water, and computed horizontal extinction coefficients (K_a) based on this $\% T_a$ data. All measurements were made *in situ* at 1 meter water depth, unless otherwise indicated. The notation (A) indicates the use of an alternate, amplified circuit of the transmissometer, which was specified for use in extremely turbid waters.

STATION	DATE	PATH LENGTH (meters)	% TRANSMITTANCE ($\% T_a$, path length $^{-1}$)	K_a (m^{-1})
1	9 Jul. 69	0.25	2	15.65
	30 Oct. 69	0.10	22	15.14
2	24 Jul. 69	0.10	24 (A)	14.27
	2 Sept. 69	0.10	32	11.39
	2 Sept. 69	0.10	30 (A)	12.04
	3 Oct. 69	0.10	2.5	36.89
3	9 Jul. 69	0.10	5	29.96
	24 Jul. 69	0.10	9.5 (A)	23.54
	10 Sept. 69	0.10	14	19.66
	3 Oct. 69	0.10	1	46.05
3 C	24 Jul. 69	0.10	16 (A)	18.33
	10 Sept. 69	0.10	44	8.21
	3 Oct. 69	0.10	34	10.79
4	9 Jul. 69	0.10	22 (A)	15.14
	24 Jul. 69	0.10	26 (A)	13.47
	2 Sept. 69	0.10	48	7.34
	10 Sept. 69	0.10	25	13.86
	3 Oct. 69	0.10	25	13.86
	31 Oct. 69	0.10	23	14.70
5	9 Jul. 69	0.10	42 (A)	8.68
	25 Jul. 69	0.10	48	7.34
	25 Jul. 69	0.10	46 (A)	7.77
	9 Sept. 69	0.10	48	7.34
	4 Oct. 69	0.10	49	7.13
6	10 Jul. 69	0.10	48.5 (A)	7.24
	26 Jul. 69	0.10	61	4.94
	26 Jul. 69	0.10	58 (A)	5.45
	9 Sept. 69	0.10	52	6.54
	4 Oct. 69	0.10	35	10.50
	30 Oct. 69	0.10	33	11.09
7	10 Jul. 69	0.10	42 (A)	8.68
	26 Jul. 69	0.10	55	5.98
	26 Jul. 69	0.10	50 (A)	6.93
	9 Sept. 69	0.10	47	7.55
	4 Oct. 69	0.10	61	4.94
	31 Oct. 69	0.10	73	3.15
8	10 Jul. 69	0.10	60 (A)	5.11
	9 Sept. 69	0.10	61	4.94
	4 Oct. 69	0.10	35	10.50
	31 Oct. 69	0.10	64	4.46
9	10 Jul. 69	0.10	55	5.98
	10 Jul. 69	0.10	67 (A)	4.00
	26 Jul. 69	0.10	48	7.34
	26 Jul. 69	0.10	46 (A)	7.77
	4 Oct. 69	0.10	3	35.07

Table 2. (cont'd)

STATION	DATE	PATH LENGTH (meters)	% TRANSMITTANCE (%T _a /path length ⁻¹)	K _a (m ⁻¹)
10	10 Jul. 69	0.10	62	4.78
	10 Jul. 69	0.10	72 (A)	3.29
	26 Jul. 69	0.10	81	2.11
	26 Jul. 69	0.10	81 (A)	2.11
	8 Sept. 69	0.10	36	10.22
	4 Oct. 69	0.10	27	13.09
	30 Oct. 69	0.10	21	15.61
11				
	10 Jul. 69	0.10	55	5.98
	10 Jul. 69	0.10	73 (A)	3.15
12	27 Oct. 69	0.10	33	11.09
	10 Jul. 69	0.10	64	4.46
	10 Jul. 69	0.10	76 (A)	2.74
	26 Jul. 69	0.10	76	2.74
	26 Jul. 69	0.10	77 (A)	2.61
	8 Sept. 69	0.10	31	11.71
	4 Oct. 69	0.10	15	18.97
13 B	30 Oct. 69	0.10	14	19.66
	10 Jul. 69	0.10	66	4.16
14	10 Jul. 69	0.10	76 (A)	2.74
	10 Jul. 69	0.10	54	6.16
	10 Jul. 69	0.10	85 (A)	1.63
	26 Jul. 69	0.10	58	5.45
	26 Jul. 69	0.10	58 (A)	5.45
	8 Sept. 69	0.10	38	9.68
	5 Oct. 69	0.10	24	14.30
	30 Oct. 69	0.10	10	23.03
16	11 Jul. 69	0.10	35	10.50
	11 Jul. 69	0.10	42 (A)	8.68
	27 Jul. 69	0.10	32	11.39
	27 Jul. 69	0.10	28 (A)	12.73
	27 Jul. 69	0.10	3	35.07
	3 Sept. 69	0.10	21	15.61
	5 Oct. 69	0.10	13	24.40
17	28 Oct. 69			
	27 Jul. 69	0.10	73	3.15
	27 Jul. 69	0.10	72 (A)	3.29
	27 Jul. 69	0.50	20	3.22
	27 Jul. 69	0.50	18 (A)	3.43
18	5 Oct. 69	0.10	62	4.78
	11 Jul. 69	0.10	50	6.93
	11 Jul. 69	0.10	50 (A)	6.93
	27 Jul. 69	0.10	73	3.15
	27 Jul. 69	0.10	73 (A)	3.15
	3 Sept. 69	0.10	72	3.29
19	5 Oct. 69	0.10	32	11.39
	13 Jul. 69	0.10	61	4.94
	13 Jul. 69	0.10	58 (A)	5.45
	28 Jul. 69	0.50	19	3.32
	28 Jul. 69	0.50	17 (A)	3.54
	3 Sept. 69	0.10	40	9.16
20	11 Oct. 69	0.10	67	4.00
	13 Jul. 69	0.10	72	3.29
	13 Jul. 69	0.10	72 (A)	3.29

Table 2. (cont'd)

STATION	DATE	PATH LENGTH (meters)	% TRANSMITTANCE (%T _a , path length ⁻¹)	K _a (m ⁻¹)
21	28 Jul. 69	0.50	25	2.77
	28 Jul. 69	0.50	23 (A)	2.94
	3 Sept. 69	0.10	78	2.49
	11 Oct. 69	0.10	50	6.93
	28 Oct. 69	0.10	55	5.98
22	13 Jul. 69	0.10	70	3.57
	13 Jul. 69	0.10	72 (A)	3.29
	13 Jul. 69	0.10	68	3.86
	13 Jul. 69	0.10	70.5 (A)	3.50
	13 Jul. 69	0.10	70	3.57
	13 Jul. 69	0.10	72 (A)	3.29
	28 Jul. 69	0.50	34	2.16
	28 Jul. 69	0.50	31 (A)	2.34
	4 Sept. 69	0.10	68	3.86
	11 Oct. 69	0.10	71	3.42
23 C	28 Oct. 69	0.10	72	3.29
	13 Jul. 69	0.10	82	1.98
	13 Jul. 69	0.10	82 (A)	1.98
	28 Jul. 69	0.50	32	2.28
	28 Jul. 69	0.50	30 (A)	2.41
	4 Sept. 69	0.10	31	11.71
23 E	11 Oct. 69	0.10	78	2.48
	29 Oct. 69	0.10	72	3.29
	13 Jul. 69	0.10	93	0.73
	13 Jul. 69	0.50	60	1.02
	28 Jul. 69	0.50	50	1.39
25	28 Jul. 69	0.50	51	1.35
	6 Sept. 69	0.50	36	2.04
	11 Oct. 69	0.10	83	1.86
	29 Oct. 69	0.10	72	3.29
	15 Jul. 69	0.50	32	2.28
	30 Jul. 69	0.50	40	1.83
26	30 Jul. 69	0.50	38 (A)	1.94
	6 Sept. 69	0.50	33	2.22
	11 Oct. 69	0.10	79	2.36
	13 Jul. 69	0.50	48	1.47
	28 Jul. 69	0.50	66	0.83
27	28 Jul. 69	0.50	66 (A)	0.83
	4 Sept. 69	0.10	50	6.93
	6 Oct. 69	0.50	1	9.21
	29 Oct. 69	0.10	72	3.29
	14 Jul. 69	0.50	53	1.27
28	29 Jul. 69	0.50	62	0.96
	29 Jul. 69	0.50	62 (A)	0.96
	4 Sept. 69	0.10	77	2.61
	9 Oct. 69	0.10	14	19.66
	14 Jul. 69	0.50	9	4.82
29	29 Jul. 69	0.50	30	2.41
	29 Jul. 69	0.50	28 (A)	2.55
	9 Oct. 69	0.10	13	20.40
	28 Jul. 69	0.50	66	0.83
30	28 Jul. 69	0.50	66 (A)	0.83
	4 Sept. 69	0.10	55	5.98
	6 Oct. 69	0.10	54	6.16
	29 Oct. 69	0.10	79	2.36

Table 2 . (cont'd)

STATION	DATE	PATH LENGTH (meters)	% TRANSMITTANCE (%T _a , path length ⁻¹)	K _a (m ⁻¹)
29	7 Oct. 69	0.10	82	1.98
31	14 Jul. 69	0.10	39	9.42
	14 Jul. 69	0.50	3	7.01
	29 Jul. 69	0.50	58	1.09
	29 Jul. 69	0.50	56 (A)	1.16
	6 Sept. 69	0.50	13	4.08
	8 Oct. 69	0.10	22	15.14
33	14 Jul. 69	0.10	8	25.26
	29 Jul. 69	0.50	8	5.05
	29 Jul. 69	0.50	6 (A)	5.63
	8 Oct. 69	0.10	6	28.13
35	14 Jul. 69	0.50	34	2.16
	29 Jul. 69	0.50	46	1.55
	29 Jul. 69	0.50	46 (A)	1.55
	6 Sept. 69	0.10	95	0.51
	6 Sept. 69	0.50	36	2.04
	8 Oct. 69 (1m)	0.10	73	3.15
	8 Oct. 69 (1m)	0.10	80	2.23
	8 Oct. 69 (1m)	0.10	73	3.15
	8 Oct. 69 (1m)	0.10	83	1.86
	8 Oct. 69 (1m)	0.10	82	1.98
	8 Oct. 69 (1m)	0.50	22	3.03
	8 Oct. 69 (1m)	0.50	22	3.03
	8 Oct. 69 (2m)	0.10	78	2.49
	8 Oct. 69 (2m)	0.10	73	3.15
	8 Oct. 69 (3m)	0.10	78	2.48
	8 Oct. 69 (3m)	0.10	73	3.15
	8 Oct. 69 (3m)	0.50	19	3.32
	8 Oct. 69 (3m)	0.50	20	3.22
	8 Oct. 69 (3m)	0.50	74	3.01
	8 Oct. 69 (5m)	0.10	72	3.29
	8 Oct. 69 (5m)	0.10	78	2.48
	8 Oct. 69 (5m)	0.10	80	2.23
	8 Oct. 69 (5m)	0.10	18	3.43
	8 Oct. 69 (5m)	0.50	19	3.32
	8 Oct. 69 (5m)	0.50	73	3.15
	8 Oct. 69 (7m)	0.10	71	3.42
	8 Oct. 69 (7m)	0.10	18	3.43
	8 Oct. 69 (7m)	0.50	18	3.43
	8 Oct. 69 (10m)	0.10	70	3.57
	8 Oct. 69 (10m)	0.10	73	3.15
	8 Oct. 69 (10m)	0.10	78	2.48
	8 Oct. 69 (10m)	0.10	77	2.61
	8 Oct. 69 (10m)	0.50	18	3.43
	8 Oct. 69 (10m)	0.50	18	3.43
	8 Oct. 69 (10m)	0.50	55	5.98
	8 Oct. 69 (15m)	0.10	76	2.74
	8 Oct. 69 (15m)	0.10	17	3.54
39	15 Jul. 69	0.50	53	1.27
	30 Jul. 69	0.50	64	0.89
	30 Jul. 69	0.50	63 (A)	0.92
	6 Sept. 69	0.50	15	3.79
	6 Oct. 69	0.50	9	4.82
	29 Oct. 69	0.10	60	5.11
41	15 Jul. 69	0.50	48	1.47
	30 Jul. 69	0.50	64	0.89
	30 Jul. 69	0.50	64 (A)	0.89

Table 2. (cont'd)

STATION	DATE	PATH LENGTH (meters)	% TRANSMITTANCE (%T ₀ , path length ⁻¹)	K _a (m ⁻¹)
	7 Sept. 69	0.50	39	1.88
	6 Oct. 69	0.50	8	5.05
43	15 Jul. 69	0.50	33	2.22
	30 Jul. 69	0.50	16	3.67
	30 Jul. 69	0.50	14 (A)	3.93
	6 Oct. 69	0.50	1	9.21
45	15 Jul. 69	0.50	52	1.31
	30 Jul. 69	0.50	72	0.66
	30 Jul. 69	0.50	72 (A)	0.66
	7 Sept. 69	0.50	18	3.43
	6 Oct. 69	0.10	80	2.23
	6 Oct. 69	0.50	14	3.93
48	15 Jul. 69	0.50	8	5.05
	30 Jul. 69	0.50	27	2.62
	30 Jul. 69	0.50	27 (A)	2.62
	7 Sept. 69	0.50	12	4.24
	12 Oct. 69	0.10	67	4.00
	29 Oct. 69	0.10	18	17.15
50 B	11 Jul. 69	0.10	27 (A)	13.09
	27 Jul. 69	0.10	42	8.68
	27 Jul. 69	0.10	40 (A)	9.16
50 C	11 Jul. 69	0.10	32	11.39
	11 Jul. 69	0.10	38 (A)	9.68
	27 Jul. 69	0.10	50	6.93
	27 Jul. 69	0.10	49 (A)	7.13
	3 Sept. 69	0.10	55	5.98
	5 Oct. 69	0.10	18	17.15
51	11 Jul. 69	0.10	77	2.61
	11 Jul. 69	0.10	72 (A)	3.29
	27 Jul. 69	0.10	64	4.46
	27 Jul. 69	0.10	65 (A)	4.31
	3 Sept. 69	0.10	49	7.13
	5 Oct. 69	0.10	65	4.31
52	11 Jul. 69	0.10	42	8.68
	11 Jul. 69	0.10	49 (A)	7.13
	27 Jul. 69	0.10	42	8.68
	27 Jul. 69	0.10	40 (A)	9.16
	3 Sept. 69	0.10	19	16.61
	5 Oct. 69	0.10	18	17.15
54	11 Jul. 69	0.10	30	12.04
	11 Jul. 69	0.10	62 (A)	4.78
	27 Jul. 69	0.10	61	4.94
	27 Jul. 69	0.10	58 (A)	5.45
	8 Sept. 69	0.10	30	12.04
	5 Oct. 69	0.10	22	15.14
	30 Oct. 69	0.10	12	21.20
S4 B	28 Oct. 69	0.10	10	23.03
57	10 Jul. 69	0.10	58	5.45
	10 Jul. 69	0.10	80 (A)	2.23
	25 Jul. 69	0.10	69 (A)	3.71

Table 2. (cont'd)

STATION	DATE	PATH LENGTH (meters)	% TRANSMITTANCE (%T _o , path length ⁻¹)	K _a (m ⁻¹)
58	8 Sept. 69	0.10	42	8.68
	4 Oct. 69	0.10	34	10.79
58	10 Jul. 69	0.10	22	15.14
	10 Jul. 69	0.10	22 (A)	15.14
	25 Jul. 69	0.10	25 (A)	13.86
	9 Sept. 69	0.10	8	25.26
	4 Oct. 69	0.10	1	46.05
59	9 Jul. 69	0.10	55	5.98
	25 Jul. 69	0.10	80	2.23
	25 Jul. 69	0.10	73 (A)	3.15
	9 Sept. 69	0.10	73	3.15
	4 Oct. 69	0.10	70	3.57
60	31 Oct. 69	0.10	29	12.38
	9 Jul. 69	0.10	30.5	11.87
	24 Jul. 69	0.10	69 (A)	3.71
	2 Sept. 69	0.10	72	3.29
	9 Sept. 69	0.10	69	3.71
60 B	3 Oct. 69	0.10	38	9.68
	31 Oct. 69	0.10	50	6.93
	9 Jul. 69	0.10	3 (A)	35.07
	24 Jul. 69	0.10	28 (A)	12.73
	9 Sept. 69	0.10	45	7.89
60 C	4 Oct. 69	0.10	20	16.09
	24 Jul. 69	0.10	6.5 (A)	27.97
	9 Sept. 69	0.10	28	12.73
	3 Oct. 69	0.10	44	8.21
	9 Jul. 69	0.10	15.5 (A)	18.64
61	24 Jul. 69	0.10	64 (A)	4.46
	2 Sept. 69	0.10	63	4.62
	10 Sept. 69	0.10	58	5.47
	3 Oct. 69	0.10	38	9.68
	31 Oct. 69	0.10	32	11.39
62	9 Jul. 69	0.10	4 (A)	32.19
	24 Jul. 69	0.10	25 (A)	13.86
	10 Sept. 69	0.10	32	11.39
	3 Oct. 69	0.10	3	35.07
	17 Jul. 69	0.10	41	8.92
63	1 Aug. 69	0.10	1	46.05
	1 Aug. 69	0.10	1 (A)	46.05
	10 Sept. 69	0.10	16	18.33
	13 Oct. 69	0.10	2	39.12
	31 Oct. 69	0.10	31	11.71
64	15 Jul. 69	0.10	50	6.93
	15 Jul. 69	0.50	7	5.32
	31 Jul. 69	0.10	45	7.99
	31 Jul. 69	0.10	44 (A)	8.21
	31 Jul. 69	0.50	2	7.82
	31 Jul. 69	0.50	2 (A)	7.82
	7 Sept. 69	0.10	44	8.21
	12 Oct. 69	0.10	18	17.15
	29 Oct. 69	0.10	24	14.27

Table 2. (cont'd)

STATION	DATE	PATH LENGTH (meters)	% TRANSMITTANCE (%T _a , path length ⁻¹)	K _a (m ⁻¹)
65	16 Jul. 69	0.10	50	6.93
	31 Jul. 69	0.10	50	6.93
	31 Jul. 69	0.10	49 (A)	7.13
66 B	31 Jul. 69	0.10	70	3.57
	31 Jul. 69	0.10	68 (A)	3.86
68	16 Jul. 69	0.10	62	4.78
	31 Jul. 69	0.10	40	9.16
	31 Jul. 69	0.10	39 (A)	9.42
	8 Sept. 69	0.10	30	12.04
	12 Oct. 69	0.10	14	19.66
	30 Oct. 69	0.10	12	21.20
69	16 Jul. 69	0.10	49	7.13
	31 Jul. 69	0.10	40	9.16
	31 Jul. 69	0.10	38 (A)	9.68
	8 Sept. 69	0.10	34	10.79
	12 Oct. 69	0.10	10	23.03
69 B	28 Oct. 69	0.10	8	25.26

Table 3. Computed values for vertical extinction coefficients, based on measurements of the percentage of surface light reaching several depths in the water column. These measurements were made with a Whitney submersible photometer (Montedoro Corp. model LMT-8A) with no filters over the sensor. Vertical extinction coefficients were obtained as the slope of regressions between (ln % surface light) and water depth, ignoring the depth interval 0-10 cm. These estimates of extinction coefficients were computed from depth profiles of light in lake water-flushed deck tanks [$K_w(DT)$] on the ship, and from depth profiles of light in the lake [$K_w(IS)$]. r^2 is the correlation coefficient of the regression.

STATION	DATE	TIME	$K_w(DT)$ m ⁻¹	r^2	$K_w(IS)$ m ⁻¹	r^2
00	03Jun69	-	4.06	0.966		
00	17Jul69	1335	6.69	0.998		
01	09Jul69	-	7.24	0.996	4.55	0.965
	17Jul69	1145	6.14	0.999		
02	04Jun69	1730	3.42	0.897		
	04Jun69	1730	3.14	0.827		
	09Jul69	1510	5.99	0.992		
04	09Jul69	1710	5.78	0.989		
06	26Jul69	0600	-	-		
07	05Jun69	0830	1.83	0.971		
	10Jul69	1002	3.36	0.989		
	26Jul69	0800	4.08	0.992	2.00	0.956
08	26Jul69	0910	4.69	0.997	2.06	0.973
10	05Jun69	1145	2.44	0.873		
	10Jul69	1235	3.53	0.976	1.88	0.996
	26Jul69	1220	3.25	0.996	1.40	0.956
13	05Jun69	1545	2.06	0.938		
13B	05Oct69	1015	6.56	0.997		
14	05Jun69	1930	1.57	0.917		
	26Jul69	1653	3.52	0.997		
15	06Jun69	1005	2.26	0.965		
16	06Jun69	1115	1.66	0.971		
	11Jul69	1015	3.40	0.988		
	27Jul69	1100	3.80	0.996	2.98	0.994
16B	05Oct69	1515	6.14	0.992		
17	12Jul69	0930	9.33	0.883	1.71	0.985
	12Jul69	1317	5.62	0.998		
	12Jul69	1400			1.44	0.946
	12Jul69	1500	5.88	0.999		
18	06Jun69	1420	2.26	0.988		
	27Jul69	1900			1.47	0.996
20	07Jun69	0815	1.16	0.697	1.64	0.899
	07Jun69	0815			1.79	0.921
	13Jul69	0830			1.46	0.982
	13Jul69	1010	2.76	0.971		
	28Jul69	0900			1.23	0.998

Table 3 . (cont'd)

STATION	DATE	TIME	K_w (DT) m ⁻¹	r^2	K_w (IS) m ⁻¹	r^2
21	07Jun69	1015	1.50	0.953	1.31 0.822	0.999 0.998
	13Jul69	1100				
	28Jul69	1127				
	28Jul69	1140	1.48	0.931		
22	13Jul69	1130	2.72	0.959		
23A	13Jul69	1415	1.43	0.977		
23C	28Jul69	1630			0.835	0.999
	28Jul69	1655	2.36	0.992		
23D	07Jun69	1430	2.16	0.971		
23E	07Jun69	1530	2.39	0.980	0.827 0.631	0.996 0.898
	30Jul69	0825				
25	07Jun69	2000	0.595	0.890	0.656 0.591	0.991 0.996
	13Jul69	1825	1.79	0.960		
	28Jul69	1836				
28	08Jun69	1045	1.35	0.982		
29	08Jun69	1500	1.92	0.901		
	07Oct69	1045	2.37	0.997		
	07Oct69	1600	2.98	0.983		
33	09Jun69	1305	1.69	0.987		
35	14Jul69	0905			0.935 0.959 0.924	0.995 0.993 0.994
	14Jul69	0905				
	14Jul69	0905				
	14Jul69	0930	2.84	0.779	0.834 0.709 0.802	0.999 0.943 0.998
	14Jul69	1245	2.50	0.971		
	14Jul69	1420	2.60	0.933		
	26Jul69	0930	2.72	0.919		
	29Jul69	1540				
	29Jul69	1550	2.17	0.992	0.938	0.960
	08Oct69	1305	4.35	0.999		
	08Oct69	1305	1.92	0.997		
	08Oct69	1405	1.17	0.996		
	08Oct69	1600			0.938	0.960
	08Oct69	1815	3.44	0.991		
39	15Jul69	0950	3.94	0.952	0.509	0.937
	30Jul69	1039				
41	15Jul69	1140	2.68	0.918	0.640 0.534	0.998 0.986
	30Jul69	1255				
	30Jul69	1259	1.72	0.998		
	06Oct69	1700	3.37	0.993		
43	15Jul69	1420	2.20	0.992		
45	15Jul69	1545	1.90	0.997		
	06Oct69	1245	2.92	0.970		

Table 4. Percentage transmission ($\%T_\lambda$) of light (from a bank of fluorescent tubes) at various wavelengths (λ_i) through a 0.1 meter water column, vertical extinction coefficients ($K_{SR\lambda}$) for each wave length over the 0.1 meter water column, and computed vertical extinction coefficients (\bar{K}_{SR}) and percent transmission ($\bar{\%T}$) for $\lambda = 400$ to $1000 \text{ m}\mu$ through a 1.0 meter water column. These data were determined with a 16 channel submersible spectroradiometer, with the sensor held just above, and 10 cm below the water surface in a large deck tank flushed with lake water. A relatively constant light source was provided by fluorescent tubes affixed to a lid over the deck tank. See Fig. 1 , and methods.

Station	05 02Sep69 1300	12 02Sep69 1740	13B 05Oct69 1030	16 03Sep69 1145		
Wave Length (λ_i) (m μ)	$\%T_\lambda$ (0.1 m $^{-1}$)	$K_{SR\lambda}$ (m $^{-1}$)	$\%T_\lambda$ (0.1 m $^{-1}$)	$K_{SR\lambda}$ (m $^{-1}$)	$\%T_\lambda$ (0.1 m $^{-1}$)	$K_{SR\lambda}$ (m $^{-1}$)
405	24.7	14.0	54.2	6.12	63.4	4.56
436	18.0	17.1	6.59	27.2	36.1	10.2
460	39.8	9.20	38.2	9.62	51.2	6.70
480	32.4	11.3	30.6	11.9	50.6	6.82
500	31.7	11.5	33.3	11.0	50.4	6.86
520	47.2	7.51	46.4	7.68	53.1	6.33
546	60.3	5.06	61.0	4.95	53.9	6.17
577	52.5	6.44	52.0	6.54	57.1	5.60
600	49.8	6.97	46.3	7.70	55.6	5.88
640	50.0	6.93	50.0	6.93	59.7	5.16
660	53.3	6.29	53.3	6.29	60.5	5.02
691	56.8	5.65	56.8	5.65	67.4	3.95
730	55.0	5.98	58.0	5.45	66.4	4.09
800	50.0	6.93	50.0	6.93	75.0	2.88
900	50.0	6.93	-	-	82.7	1.90
1000	0.0	-	0.0	-	82.0	1.99
$\bar{\%T}$	0.043 m $^{-1}$		0.034 m $^{-1}$		0.188 m $^{-1}$	0.008 m $^{-1}$
\bar{K}_{SR}	7.74		7.99		6.27	9.43

Station	48 07Sep69 1530	50C-51 03Sep69 1700	51 05Oct69 1845	58 09Sep69 0845		
Wave Length (λ_i) (m μ)	$\%T_\lambda$ (0.1 m $^{-1}$)	$K_{SR\lambda}$ (m $^{-1}$)	$\%T_\lambda$ (0.1 m $^{-1}$)	$K_{SR\lambda}$ (m $^{-1}$)	$\%T_\lambda$ (0.1 m $^{-1}$)	$K_{SR\lambda}$ (m $^{-1}$)
405	38.6	9.53	34.9	10.5	59.2	5.24
436	42.4	8.57	4.39	31.3	30.7	11.8
460	89.4	1.12	23.6	14.4	46.9	7.58
480	85.2	1.60	41.7	8.76	46.4	7.68
500	55.8	5.83	25.8	13.5	46.3	7.70
520	61.6	4.85	32.8	11.1	51.5	6.63
546	74.0	3.02	46.0	7.76	52.9	6.36
577	68.8	3.75	43.8	8.27	56.3	5.75
600	48.1	7.33	36.8	8.29	64.8	4.34
640	60.0	5.11	43.6	6.60	67.7	3.90
660	68.3	3.81	51.7	7.40	75.0	2.88
691	86.4	1.47	47.7	7.55	71.5	3.35
730	0.0	-	47.0	81.0	1.75	84.8
800	-	-	39.3	9.34	90.0	1.05
900	-	-	-	-	88.0	1.28
1000	28.6	12.5	0.0	-	-	-
$\bar{\%T}$	1.64 m $^{-1}$		0.004 m $^{-1}$		0.182 m $^{-1}$	0.032 m $^{-1}$
\bar{K}_{SR}	4.11		10.1		6.31	5.55

Table 4. (cont'd)

Station	35		35		35		45	
Date	06Sep69	1200	08Oct69	1300	08Oct69	1825	06Oct69	1315
Time								
Wave Length (λ_i) (m)	%T _λ (0.1 m ⁻¹)	K _{SRλ} (m ⁻¹)	%T _λ (0.1 m ⁻¹)	K _{SRλ} (m ⁻¹)	%T _λ (0.1 m ⁻¹)	K _{SRλ} (m ⁻¹)	%T _λ (0.1 m ⁻¹)	K _{SRλ} (m ⁻¹)
405	26.5	13.3	78.3	2.45	84.9	1.65	73.5	3.08
436	65.9	4.18	69.5	3.63	71.2	3.40	61.9	4.79
460	66.7	4.06	80.4	2.19	80.6	2.15	74.1	3.00
480	78.7	2.40	79.1	2.34	82.0	1.99	73.5	3.08
500	54.2	6.13	77.3	2.57	80.5	2.17	73.0	3.15
520	62.4	4.72	77.1	2.60	83.6	1.80	75.9	2.75
546	57.1	5.60	81.3	2.08	82.4	1.94	75.7	2.79
577	57.0	5.62	78.9	2.36	83.3	1.82	76.8	2.63
600	39.6	9.25	78.1	2.48	80.0	2.23	74.4	2.96
640	45.5	7.89	82.0	1.99	84.1	1.74	76.9	2.62
660	48.3	7.27	80.2	2.21	83.5	1.81	75.7	2.79
691	59.1	5.26	83.0	1.87	86.0	1.51	78.3	2.45
730	27.0	13.1	75.5	2.81	83.5	1.80	75.0	2.88
800	-	-	86.6	1.44	86.4	1.47	81.5	2.05
900	75.0	2.88	87.9	1.29	85.0	1.63	80.9	2.12
1000	7.62	25.7	86.9	1.41	87.2	1.37	88.6	1.21
%T̄	0.278 m ⁻¹		8.75 m ⁻¹		12.1 m ⁻¹		4.76 m ⁻¹	
K̄ _{SR}	5.89		2.44		2.11		3.04	

Station	16		17		23C		25	
Date	05Oct69	1530	03Sep69	1230	06Sep69	1750	04Sep69	1600
Time								
Wave Length (λ_i) (m)	%T _λ (0.1 m ⁻¹)	K _{SRλ} (m ⁻¹)	%T _λ (0.1 m ⁻¹)	K _{SRλ} (m ⁻¹)	%T _λ (0.1 m ⁻¹)	K _{SRλ} (m ⁻¹)	%T _λ (0.1 m ⁻¹)	K _{SRλ} (m ⁻¹)
405	63.8	4.50	38.6	9.53	37.3	9.85	54.2	6.12
436	40.0	9.16	4.39	31.3	54.6	6.05	34.1	10.7
460	52.9	6.37	27.6	12.9	69.9	3.58	52.0	6.53
480	53.3	6.29	41.7	8.76	76.9	2.63	63.0	4.63
500	50.0	6.93	28.3	12.6	51.7	6.60	43.3	8.36
520	54.3	6.11	36.0	10.2	62.4	4.72	45.6	7.85
546	54.2	6.13	46.0	7.76	73.7	3.06	63.5	4.54
577	55.2	5.95	43.8	8.27	66.3	4.12	56.3	5.75
600	55.7	5.85	38.6	9.52	43.2	8.40	50.9	6.76
640	58.5	5.37	46.4	7.69	56.4	5.73	54.3	6.06
660	61.2	4.91	55.8	5.83	61.7	4.83	60.0	5.11
691	66.5	4.07	51.4	6.66	72.7	3.19	59.1	5.20
730	71.4	3.37	52.0	6.54	17.0	17.7	55.0	5.98
800	80.9	2.13	45.7	7.83	53.6	6.24	57.1	5.66
900	87.5	1.34	-	-	83.3	1.82	-	-
1000	87.1	1.38	0.0	-	14.3	19.5	0.0	-
%T̄	0.217 m ⁻¹		0.006 m ⁻¹		0.608 m ⁻¹		0.179 m ⁻¹	
K̄ _{SR}	6.14		9.76		5.10		6.51	

Table 4. (cont'd)

Station	59	59-60		59	59	
Date	09Sep69	02Sep69		04Oct69	04Oct69	
Time	0945	1340		0940	1850	
Wave Length (λ_i) (μm)	%T _{λ} (0.1 m^{-1})	K _{SRλ} (m^{-1})	%T _{λ} (0.1 m^{-1})	K _{SRλ} (m^{-1})	%T _{λ} (0.1 m^{-1})	K _{SRλ} (m^{-1})
405	37.7	9.75	34.9	10.5	55.4	5.91
436	22.7	14.8	17.6	17.4	37.3	9.85
460	55.3	5.92	42.3	8.61	49.7	7.00
480	54.5	6.06	35.2	10.4	48.6	7.22
500	53.9	6.18	34.2	10.7	49.2	7.10
520	52.7	6.40	50.8	6.77	51.4	6.65
546	63.3	4.57	63.7	4.52	52.2	6.50
577	63.0	4.65	56.3	5.75	54.8	6.02
600	53.9	6.18	51.9	6.55	53.3	6.29
640	65.0	4.31	53.6	6.23	58.5	5.36
660	67.9	3.88	55.0	5.98	59.4	5.21
691	75.8	2.77	58.2	5.42	65.8	4.18
730	-	-	56.0	5.80	64.0	4.46
800	71.7	3.32	48.2	7.30	79.1	2.55
900	40.0	9.16	55.0	5.98	83.2	1.84
1000	60.0	5.11	0.0	-	82.8	1.89
%T	0.292 m^{-1}		0.078 m^{-1}		0.146 m^{-1}	0.254 m^{-1}
K _{SR}	5.84		7.16		6.55	5.98

Station	61	62		68	68	
Date	10Sep69	04Sep69		08Sep69	12Oct69	
Time	0920	1007		0700	1440	
Wave Length (λ_i) (μm)	%T _{λ} (0.1 m^{-1})	K _{SRλ} (m^{-1})	%T _{λ} (0.1 m^{-1})	K _{SRλ} (m^{-1})	%T _{λ} (0.1 m^{-1})	K _{SRλ} (m^{-1})
405	40.6	9.01	60.2	5.07	80.7	2.14
436	0.0	-	75.2	3.12	34.2	11.6
460	61.8	4.81	81.3	2.07	87.0	1.39
480	60.2	5.08	-	-	80.6	2.16
500	60.6	5.01	75.0	2.88	56.7	5.68
520	63.1	4.60	80.0	2.23	51.2	6.69
546	67.7	3.90	88.9	1.18	62.5	4.69
577	70.3	3.53	87.5	1.34	59.5	5.19
600	64.5	4.38	84.2	1.72	44.2	8.16
640	74.5	2.94	81.8	2.01	61.8	4.81
660	75.9	2.76	90.0	1.05	70.0	3.57
691	81.6	2.03	81.8	2.01	82.7	1.90
730	-	-	80.0	2.23	0.0	-
800	70.0	3.57	64.3	4.42	60.7	4.99
900	76.7	2.66	-	-	83.3	1.82
1000	55.5	5.88	0.0	-	19.0	16.6
%T	0.894 m^{-1}		17.1 m^{-1}		0.344 m^{-1}	0.452 m^{-1}
K _{SR}	4.72		1.77		5.67	5.47

Table 5. Suspended sediment (SS) total dissolved solids (TDS), and total solids (TS) for Lake Winnipeg and Red River stations. Station locations can be found in Fig. 3.

STATION NO.	DATE	SAMPLING DEPTH (METERS)	(GRAMS TDS)	PER SS	CURIC MFTER) TS
0	27OCT68	1.0	145.00	*****	*****
0	3MAR69	1.0	440.00	110.00	550.00
0	18MAR69	1.0	425.00	90.00	515.00
0	31MAR69	1.0	400.00	11.00	411.00
0	15APR69	1.0	240.00	1900.00	2140.00
0	28APR69	1.0	250.00	285.00	535.00
0	12MAY69	1.0	320.00	159.00	479.00
0	26MAY69	1.0	530.00	200.00	730.00
0	9JUN69	1.0	490.00	201.00	691.00
0	23JUN69	1.0	530.00	50.00	580.00
0	7JUL69	1.0	615.00	75.00	690.00
0	21JUL69	1.0	605.00	54.00	659.00
0	5AUG69	1.0	470.00	28.00	508.00
0	18AUG69	1.0	450.00	43.00	493.00
0	2SEP69	1.0	530.00	25.00	555.00
0	15SEP69	1.0	455.00	15.00	470.00
0	29SEP69	1.0	450.00	42.00	492.00
0	14OCT69	1.0	470.00	23.00	493.00
0	27OCT69	1.0	480.00	9.00	489.00
0	10NOV69	1.0	470.00	20.00	490.00
0	24NOV69	1.0	540.00	20.00	560.00
0	8DEC69	1.0	620.00	10.00	630.00
0	29DEC69	1.0	560.00	10.00	570.00
0	19JAN70	1.0	550.00	< 10.00	*****
0	3FEB70	1.0	560.00	10.00	570.00
0	16FEB70	1.0	540.00	*****	*****
0	2MAR70	1.0	480.00	< 10.00	*****
0	30MAR70	1.0	540.00	10.00	550.00
0	13APR70	1.0	480.00	44.00	524.00
0	21APR70	1.0	280.00	780.00	1060.00
0	27APR70	1.0	280.00	251.00	531.00
0	11MAY70	1.0	400.00	23.00	423.00
0	25MAY70	1.0	420.00	174.00	594.00
0	9JUN70	1.0	340.00	218.00	558.00
0	23JUN70	1.0	420.00	132.00	552.00
0	6JUL70	1.0	600.00	195.00	795.00
0	20JUL70	1.0	500.00	134.00	634.00
0	5AUG70	1.0	600.00	33.00	633.00
0	17AUG70	1.0	630.00	30.00	660.00
0	3SEP70	1.0	670.00	20.00	690.00
0	14SEP70	1.0	570.00	10.00	580.00
0	28SEP70	1.0	492.00	16.00	508.00
0	13OCT70	1.0	550.00	27.00	577.00
0	26OCT70	1.0	507.00	11.00	518.00
0	9NOV70	1.0	528.00	86.00	614.00
0	24NOV70	1.0	580.00	23.00	603.00
0	7DEC70	1.0	708.00	17.00	725.00
0	1FEB71	1.0	1524.00	5.00	1529.00
0	15FEB71	1.0	573.00	8.00	581.00
0	1MAR71	1.0	943.00	21.00	964.00
0	15MAR71	1.0	564.00	31.00	595.00
0A	3MAR69	1.0	355.00	205.00	560.00
0A	18MAR69	1.0	360.00	105.00	465.00
0A	31MAR69	1.0	375.00	13.00	388.00
0A	15APR69	1.0	235.00	130.00	1615.00
0A	28APR69	1.0	345.00	150.00	545.00
0A	12MAY69	1.0	425.00	110.00	535.00
0A	26MAY69	1.0	490.00	195.00	685.00
0A	9JUN69	1.0	450.00	174.00	584.00
0A	23JUN69	1.0	450.00	110.00	560.00
0A	7JUL69	1.0	575.00	195.00	770.00
0A	21JUL69	1.0	490.00	78.00	568.00
0A	5AUG69	1.0	350.00	225.00	575.00
0A	18AUG69	1.0	380.00	48.00	428.00
0A	2SEP69	1.0	415.00	18.00	433.00
0A	15SEP69	1.0	335.00	27.00	362.00
0A	29SEP69	1.0	395.00	28.00	423.00
0A	14OCT69	1.0	415.00	25.00	440.00
0A	27OCT69	1.0	570.00	16.00	586.00
0A	10NOV69	1.0	400.00	20.00	420.00

STATION NO.	DATE	SAMPLING DEPTH (METERS)	(GRAMS TDS)	PER SS	CUBIC METER)	MTFR) TS
0A	24NOV69	1.0	420.00	20.00	440.00	
0A	8DEC69	1.0	470.00	10.00	480.00	
0A	29DEC69	1.0	440.00	< 10.00	*****	
0A	19JAN70	1.0	400.00	10.00	410.00	
0A	3FEB70	1.0	380.00	10.00	390.00	
0A	16FEB70	1.0	350.00	*****	*****	
0A	2MAR70	1.0	340.00	20.00	370.00	
0A	30MAR70	1.0	390.00	10.00	400.00	
0A	13APR70	1.0	400.00	47.00	467.00	
0A	27APR70	1.0	280.00	298.00	578.00	
0A	11MAY70	1.0	390.00	12.00	402.00	
0A	25MAY70	1.0	410.00	135.00	545.00	
0A	9JUN70	1.0	290.00	196.00	486.00	
0A	23JUN70	1.0	330.00	129.00	459.00	
0A	6JUL70	1.0	480.00	148.00	668.00	
0A	20JUL70	1.0	380.00	243.00	663.00	
0A	5AUG70	1.0	470.00	50.00	520.00	
0A	17AUG70	1.0	470.00	30.00	500.00	
0A	3SEP70	1.0	470.00	20.00	490.00	
0A	14SEP70	1.0	400.00	20.00	420.00	
0A	28SEP70	1.0	469.00	29.00	498.00	
0A	13OCT70	1.0	440.00	29.00	469.00	
0A	26OCT70	1.0	389.00	16.00	405.00	
0A	9NOV70	1.0	433.00	32.00	465.00	
0A	24NOV70	1.0	500.00	26.00	526.00	
0A	7DEC70	1.0	492.00	15.00	507.00	
0A	1FEB71	1.0	444.00	6.00	450.00	
0A	15FEB71	1.0	355.00	9.00	364.00	
0A	1MAR71	1.0	583.00	7.00	590.00	
0A	15MAR71	1.0	460.00	97.00	557.00	
0B	3MAR69	1.0	820.00	40.00	860.00	
0B	18MAR69	1.0	765.00	70.00	835.00	
0B	31MAR69	1.0	730.00	2.00	732.00	
0B	15APR69	1.0	320.00	450.00	770.00	
0B	28APR69	1.0	260.00	240.00	510.00	
0B	12MAY69	1.0	310.00	226.00	536.00	
0B	26MAY69	1.0	580.00	205.00	785.00	
0B	9JUN69	1.0	570.00	145.00	715.00	
0B	23JUN69	1.0	615.00	130.00	745.00	
0B	7JUL69	1.0	730.00	170.00	900.00	
0B	21JUL69	1.0	705.00	223.00	928.00	
0B	5AUG69	1.0	710.00	340.00	1070.00	
0B	18AUG69	1.0	565.00	360.00	925.00	
0B	2SEP69	1.0	595.00	145.00	740.00	
0B	15SEP69	1.0	630.00	58.00	688.00	
0B	29SEP69	1.0	630.00	58.00	688.00	
0B	14OCT69	1.0	710.00	25.00	735.00	
0B	27OCT69	1.0	740.00	14.00	754.00	
0B	10NOV69	1.0	790.00	20.00	810.00	
0B	24NOV69	1.0	440.00	20.00	460.00	
0B	8DEC69	1.0	920.00	20.00	940.00	
0B	29DEC69	1.0	930.00	10.00	940.00	
0B	19JAN70	1.0	880.00	10.00	890.00	
0B	3FEB70	1.0	890.00	10.00	900.00	
0B	16FEB70	1.0	900.00	*****	*****	
0B	2MAR70	1.0	830.00	10.00	840.00	
0B	30MAR70	1.0	3650.00	20.00	3670.00	
0B	13APR70	1.0	920.00	33.00	953.00	
0B	27APR70	1.0	360.00	343.00	703.00	
0B	11MAY70	1.0	450.00	61.00	511.00	
0B	25MAY70	1.0	440.00	199.00	629.00	
0B	9JUN70	1.0	510.00	113.00	623.00	
0B	23JUN70	1.0	630.00	96.00	726.00	
0B	6JUL70	1.0	740.00	163.00	903.00	
0B	20JUL70	1.0	670.00	305.00	975.00	
0B	5AUG70	1.0	710.00	249.00	979.00	
0B	17AUG70	1.0	790.00	200.00	990.00	
0B	3SEP70	1.0	820.00	220.00	1040.00	
0B	14SEP70	1.0	790.00	70.00	860.00	
0B	28SEP70	1.0	750.00	59.00	809.00	

STATION NO.	DATE	SAMPLING DEPTH (METERS)	(GRAMS TDS)	PER	CUBIC	METER)
				SS	TS	
08	13OCT70	1.0	650.00	79.00	689.00	
08	26OCT70	1.0	553.00	79.00	632.00	
08	9NOV70	1.0	766.00	56.00	822.00	
08	24NOV70	1.0	865.00	49.00	914.00	
08	7DEC70	1.0	940.00	41.00	1021.00	
08	1FEB71	1.0	935.00	12.00	947.00	
08	15FEB71	1.0	913.00	24.00	937.00	
08	1MAR71	1.0	402.00	7.00	409.00	
08	15MAR71	1.0	214.00	30.00	244.00	
1	27OCT68	1.0	95.00	*****	*****	
1	9JUL69	1.0	422.00	30.64	452.64	
1	2SEP69	1.0	*****	27.90	*****	
1	27OCT69	1.0	*****	74.20	*****	
2	28OCT68	1.0	205.00	*****	*****	
2	23APR69	3.0	235.00	153.00	388.00	
2	9JUL69	1.0	250.00	27.41	277.41	
2	24JUL69	1.0	*****	21.50	*****	
2	2SF69	1.0	*****	22.30	*****	
2	3OCT69	1.0	260.00	44.04	304.04	
2	27OCT69	1.0	*****	30.50	*****	
2	17MART0	1.0	296.00	2.29	298.29	
2	17MART0	5.0	266.00	1.68	257.68	
3	9JUL69	1.0	394.00	91.44	485.44	
3	3OCT69	1.0	280.00	138.73	418.73	
3C	9JUL69	1.0	*****	29.87	*****	
3C	3OCT69	1.0	140.00	19.88	159.88	
3C	17MART0	1.0	299.00	1.97	300.97	
3C	17MART0	4.0	285.00	0.87	285.87	
4	9JUL69	1.0	281.00	16.72	297.72	
4	24JUL69	1.0	*****	13.98	*****	
4	2SF69	1.0	*****	21.19	*****	
4	3OCT69	1.0	180.00	25.83	205.83	
4	31OCT69	1.0	*****	22.07	*****	
4	18MART0	1.0	235.00	1.38	236.38	
4	18MART0	6.0	265.00	1.35	266.35	
5	9JUL69	1.0	*****	10.40	*****	
6	10JUL69	1.0	95.00	8.00	103.00	
6	10JUL69	10.0	*****	12.64	*****	
6	3OCT69	1.0	*****	16.30	88.22	
7	10JUL69	1.0	77.00	11.22		
7	9SF69	1.0	*****	14.42	*****	
7	4OCT69	1.0	65.00	11.85	76.85	
7	31OCT69	1.0	*****	6.67	*****	
8	10JUL69	1.0	83.00	7.42	90.42	
8	26JUL69	1.0	*****	9.33	*****	
8	9SEP69	1.0	*****	8.36	*****	
8	4OCT69	1.0	45.00	17.38	62.38	
8	31OCT69	1.0	*****	6.68	*****	
9	10JUL69	1.0	*****	9.98	*****	
10	10JUL69	1.0	106.00	6.39	112.39	
10	10JUL69	11.0	*****	4.23	*****	
10	8SF69	1.0	*****	10.14	*****	
10	4OCT69	1.0	140.00	20.13	160.13	
10	3OCT69	1.0	*****	23.62	*****	
11	10JUL69	1.0	*****	5.41	*****	
11	27OCT69	1.0	*****	15.23	*****	
12	10JUL69	1.0	*****	5.39	*****	
12	8SEP69	1.0	*****	11.32	*****	
12	4OCT69	1.0	*****	25.56	*****	
12	3OCT69	1.0	*****	28.40	*****	
13C	29OCT68	1.0	135.00	*****	*****	
14	10JUL69	1.0	85.00	7.66	92.66	
14	26JUL69	1.0	*****	11.62	*****	
14	8SEP69	1.0	*****	14.90	*****	
14	5OCT69	1.0	135.00	21.08	156.08	
14	3OCT69	1.0	*****	26.73	*****	
15	31JUL69	0.0	*****	42.13	*****	
15	31JUL69	1.0	*****	6.32	*****	
16	11JUL69	1.0	71.00	8.76	79.76	
16	27JUL69	1.0	*****	12.58	*****	

STATION NO.	DATE	SAMPLING DEPTH (METERS)	(GRAMS TDS)	PER SS	CURTC METFR) TS
16	3SFP69	1.0	*****	49.15	*****
16	5OCT69	1.0	80.00	22.63	102.63
16	28OCT69	1.0	*****	18.60	*****
17	5OCT69	1.0	135.00	*****	*****
18	11JUL69	1.0	107.00	9.17	116.17
18	27JUL69	1.0	*****	4.70	*****
18	27JUL69	15.0	*****	4.64	*****
18	3SFP69	1.0	*****	7.54	*****
18	5OCT69	1.0	85.00	13.54	98.54
18	28OCT69	1.0	*****	32.07	*****
19	28JUL69	1.0	*****	4.07	*****
19	28JUL69	15.0	*****	11.98	*****
19	3SFP69	1.0	*****	8.64	*****
19	3SFP69	16.0	*****	4.94	*****
19	11OCT69	1.0	155.00	8.92	163.92
19	28OCT69	1.0	*****	18.60	*****
20	13JUL69	1.0	139.00	6.33	145.33
20	3SFP69	1.0	*****	6.93	*****
20	11OCT69	1.0	180.00	13.02	193.02
20	28OCT69	1.0	*****	10.15	*****
20	28OCT69	1.0	*****	11.25	*****
20	28OCT69	14.0	*****	14.21	*****
21	28JUL69	1.0	*****	3.11	*****
21	4SEP69	1.0	*****	5.97	*****
21	11OCT69	1.0	165.00	7.10	172.10
21	28OCT69	1.0	*****	6.78	*****
22	4SFP69	1.0	*****	4.30	*****
22	11OCT69	1.0	175.00	4.63	179.63
22	29OCT69	1.0	*****	7.75	*****
23A	28JUL69	0.0	*****	4.70	*****
23C	13JUL69	1.0	188.00	2.17	190.17
23C	28JUL69	1.0	*****	3.22	*****
23C	6SFP69	1.0	*****	2.87	*****
23C	11OCT69	1.0	170.00	10.52	190.52
23C	29OCT69	1.0	*****	7.60	*****
23C	29OCT69	16.0	*****	9.63	*****
23E	15JUL69	1.0	148.00	3.85	151.85
23E	30JUL69	1.0	*****	2.18	*****
23E	11OCT69	1.0	170.00	10.20	180.20
25	28JUL69	1.0	*****	2.40	*****
25	4SFP69	1.0	*****	6.64	*****
25	6OCT69	1.0	100.00	10.20	200.20
25	29OCT69	1.0	*****	6.40	*****
28	13JUL69	1.0	199.00	4.15	203.15
28	6OCT69	1.0	230.00	*****	*****
28	29OCT69	1.0	*****	5.95	*****
29	20MAR69	0.0	305.00	2.00	307.00
29	28JUL69	1.0	*****	3.55	*****
29	4SFP69	1.0	*****	4.06	*****
29	7OCT69	1.0	230.00	7.67	237.67
29	29OCT69	1.0	*****	3.11	*****
30	18MAR69	2.0	200.00	5.00	205.00
30	18MAR69	5.0	200.00	4.00	204.00
30	18MAR69	11.0	200.00	4.00	204.00
35	18MAR69	2.0	200.00	3.00	203.00
35	18MAR69	15.0	100.00	3.00	193.00
35	14JUL69	1.0	148.00	2.75	150.75
35	29JUL69	1.0	*****	4.75	*****
35	6SEP69	1.0	*****	3.50	*****
35	6SFP69	15.0	*****	3.07	*****
35	8OCT69	1.0	185.00	6.93	191.93
35	8OCT69	16.0	140.00	15.54	205.54
39	1APR69	2.0	195.00	2.00	197.00
39	1APR69	8.0	195.00	2.00	197.00
39	1APR69	17.0	195.00	2.00	197.00
39	30JUL69	1.0	*****	6.05	*****
39	6SFP69	1.0	*****	7.90	*****
39	6OCT69	1.0	170.00	9.00	179.00
39	6OCT69	15.0	175.00	10.85	185.85
39	29OCT69	1.0	*****	8.98	*****

STATION NO.	DATE	SAMPLING DEPTH (METERS)	(GRAMS TDS)	PER SS	CURIC TS	METER) TS
39	29OCT69	15.0	*****	21.70	*****	
41	15JUL69	1.0	148.00	2.61	150.61	
41	30JUL69	1.0	*****	1.24	*****	
41	30JUL69	14.0	*****	4.46	*****	
41	6OCT69	1.0	190.00	10.57	200.57	
45	15JUL69	1.0	167.00	2.58	169.58	
45	6OCT69	1.0	195.00	8.69	203.69	
48	2APR69	2.0	210.00	7.00	217.00	
48	2APR69	10.0	220.00	6.00	226.00	
48	2APR69	16.0	210.00	7.00	217.00	
48	15JUL69	1.0	120.00	5.83	125.83	
48	30JUL69	1.0	*****	2.48	*****	
48	7SEP69	1.0	*****	5.33	*****	
48	7SEP69	15.0	*****	3.90	*****	
48	12OCT69	1.0	155.00	9.16	164.16	
48	12OCT69	16.0	135.00	9.30	144.30	
48	29OCT69	1.0	*****	22.33	*****	
48	29OCT69	15.0	*****	22.33	*****	
50C	5OCT69	1.0	60.00	*****	*****	
51	11JUL69	1.0	39.00	4.76	43.76	
51	3SEP69	1.0	*****	4.38	*****	
51	5OCT69	1.0	70.00	5.87	75.87	
52	5OCT69	1.0	60.00	20.23	80.23	
54	11JUL69	1.0	56.00	8.10	64.10	
54	27JUL69	1.0	*****	8.70	*****	
54	8SEP69	1.0	*****	11.64	*****	
54	5OCT69	1.0	105.00	23.25	128.25	
54	30OCT69	1.0	*****	26.28	*****	
54B	28OCT69	1.0	*****	28.10	*****	
55	30OCT68	0.0	135.00	*****	*****	
55B	31JUL69	1.0	*****	6.44	*****	
56	29OCT68	1.0	130.00	*****	*****	
57	31OCT68	1.0	165.00	*****	*****	
57	23APR69	2.0	90.00	1.00	91.00	
57	23APR69	6.0	150.00	2.00	152.00	
57	23APR69	10.0	330.00	1.00	331.00	
57	10JUL69	1.0	*****	7.41	*****	
57	25JUL69	1.0	*****	6.26	*****	
57	4OCT69	1.0	355.00	17.30	372.30	
57	27OCT69	1.0	*****	13.00	*****	
58	10JUL69	1.0	*****	19.10	*****	
59	31OCT68	1.0	135.00	*****	*****	
59	9JUL69	1.0	145.00	7.84	152.84	
59	9JUL69	10.0	*****	8.32	*****	
59	25JUL69	1.0	*****	4.17	*****	
59	2SF69	1.0	*****	34.33	*****	
59	9SEP69	1.0	*****	5.74	*****	
59	9SEP69	10.0	*****	5.24	*****	
59	4OCT69	1.0	170.00	7.60	177.60	
59	4OCT69	10.0	160.00	9.00	169.00	
59	31OCT69	1.0	*****	13.24	*****	
59	31OCT69	10.0	*****	17.40	*****	
60	23APR69	1.0	145.00	32.00	177.00	
60	23APR69	4.0	155.00	4.00	159.00	
60	23APR69	9.0	160.00	7.00	167.00	
60	9JUL69	1.0	224.00	13.56	237.56	
60	24JUL69	1.0	*****	6.20	*****	
60	9SEP69	1.0	*****	5.54	*****	
60	3OCT69	1.0	260.00	19.25	279.25	
60	3OCT69	9.0	280.00	19.38	299.38	
60	31OCT69	1.0	*****	10.10	*****	
60B	9JUL69	1.0	*****	33.64	*****	
60B	4OCT69	1.0	280.00	29.85	309.85	
60C	18M/R70	1.0	286.00	3.46	289.46	
61	9JUL69	1.0	209.00	21.60	230.60	
61	9JUL69	8.0	*****	19.65	*****	
61	24JUL69	1.0	*****	6.34	*****	
61	10SEP69	1.0	*****	6.88	*****	
61	10SEP69	8.0	*****	6.75	*****	
61	3OCT69	1.0	235.00	20.42	255.42	

STATION NO.	DATE	SAMPLING DEPTH (METERS)	(GRAMS PER CUBIC METER)		
			TDS	SS	TS
61	31OCT69	1.0	*****	16.40	*****
61	18MAR70	1.0	253.00	4.72	257.72
61	18MAR70	7.0	387.00	16.06	403.06
62	31OCT68	1.0	150.00	*****	*****
62	9JUL69	1.0	*****	27.70	*****
62	30CT69	1.0	310.00	85.90	395.90
63	17JUL69	1.0	257.00	17.76	274.76
63	1AUG69	1.0	*****	84.47	*****
63	10SFP69	1.0	*****	46.60	*****
63	13OCT69	1.0	*****	49.67	*****
63	31OCT69	1.0	*****	23.17	*****
64	30OCT68	0.0	115.00	*****	*****
64	15JUL69	1.0	153.00	*****	*****
64	31JUL69	1.0	*****	11.86	*****
64	7SFP69	1.0	*****	9.82	*****
64	12OCT69	1.0	135.00	14.74	149.74
64	29OCT69	1.0	*****	18.34	*****
65	16JUL69	1.0	207.00	9.83	216.83
65	31JUL69	1.0	*****	8.59	*****
65	31JUL69	7.0	*****	10.05	*****
66B	31JUL69	1.0	*****	7.78	*****
66C	31JUL69	1.0	*****	15.05	*****
67	2APR69	1.0	1440.00	6.00	1446.00
68	31JUL69	1.0	*****	10.92	*****
68	12OCT69	1.0	*****	27.88	*****
68	30OCT69	1.0	*****	26.85	*****
69	16JUL69	1.0	70.00	6.04	76.04
69	31JUL69	1.0	*****	12.00	*****
69	12OCT69	1.0	*****	38.20	*****
69B	28OCT69	1.0	*****	30.55	*****
70	20MAR69	1.0	210.00	2.00	212.00

Table 6. Daily solar radiation at Winnipeg and The Pas for days of our research on Lake Winnipeg. The weather station at The Pas began recording radiation data in 1973, and we include this data, compared to 1973 Winnipeg data, to illustrate differences in solar radiation over the north and south basin of Lake Winnipeg. This data is derived from Canada Dept. of Transport, Met. Branch, Monthly Radiation Summary, 1969, 1970, 1973, and 1974.

Date	Winnipeg g cal cm ⁻² day ⁻¹	The Pas g cal cm ⁻² day ⁻¹ (1973-74)	Winnipeg g cal cm ⁻² day ⁻¹ (1973-74)
18 Mar 69	361	155	191
19 Mar 69	325	212	175
20 Mar 69	457	346	263
1 Apr 69	452	260	466
2 Apr 69	408	436	501
23 Apr 69	596	299	509
24 Apr 69	596	-	378
4 Jun 69	630	358	-
5 Jun 69	622	462	492
6 Jun 69	287	338	653
7 Jun 69	693	212	485
8 Jun 69	742	-	465
9 Jun 69	-	110	528
10 Jun 69	-	399	527
11 Jun 69	580	615	705
12 Jun 69	514	603	745
9 Jul 69	608	20	724
10 Jul 69	471	663	639
11 Jul 69	671	590	443
12 Jul 69	660	309	676
13 Jul 69	597	384	520
14 Jul 69	326	623	642
15 Jul 69	383	439	673
16 Jul 69	664	590	626
17 Jul 69	645	403	421

Table 6. (cont')

Date	Winnipeg g cal cm ⁻² day ⁻¹	The Pas g cal cm ⁻² day ⁻¹ (1973-74)	Winnipeg g cal cm ⁻² day ⁻¹ (1973-74)
24 Jul 69	645	211	382
25 Jul 69	644	287	350
26 Jul 69	107	285	290
27 Jul 69	483	573	437
28 Jul 69	637	355	560
29 Jul 69	607	355	498
30 Jul 69	268	620	317
31 Jul 69	661	581	606
1 Aug 69	628	476	626
2 Sep 69	302	468	163
3 Sep 69	369	164	81
4 Sep 69	282	305	258
5 Sep 69	346	209	357
6 Sep 69	465	472	405
7 Sep 69	291	461	482
8 Sep 69	212	311	407
9 Sep 69	445	225	321
10 Sep 69	365	441	499
3 Oct 69	41	198	110
4 Oct 69	172	226	337
5 Oct 69	111	209	323
6 Oct 69	113	273	336
7 Oct 69	158	120	108
8 Oct 69	333	237	111
9 Oct 69	155	274	63
10 Oct 69	-	195	224
11 Oct 69	-	266	165
12 Oct 69	-	144	99
13 Oct 69	-	113	293

Table 6. (cont')

Date	Winnipeg g cal cm ⁻² day ⁻¹	The Pas g cal cm ⁻² day ⁻¹ (1973-74)	Winnipeg g cal cm ⁻² day ⁻¹ (1973-74)
26 Oct 69	152	66	33
27 Oct 69	177	60	95
28 Oct 69	232	155	211
29 Oct 69	66	44	105
30 Oct 69	79	86	110
31 Oct 69	81	32	185
17 Mar 70	428	174	282
18 Mar 70	430	-	444

Table 7. Total monthly solar radiation expressed as g cal cm⁻² at Winnipeg and The Pas for 1969 to 1975.

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
WINNIPEG												
1969	3565	5656	11377	14010	16399	14760	15314	14136	9060	4619	3030	2635
1970	3689	5992	11873	11070	13113	15990	16213	15624	9150	5952	2790	2976
1971	4278	5600	9176	12000	16771	14820	-	14477	9570	5084	3540	2728
1972	4030	6583	9672	12390	15996	18870	16461	14322	8850	6851	3300	3286
1973	4061	5684	8680	12570	16399	15810	16895	14477	9870	5890	3210	2697
1974	3999	6328	10757	13080	12462	19080	18166	13330	9150	6355	2970	2635
1975	3348	6132	10137	10500	16213	14850	17701	13020	9450	5952	3840	2790
THE PAS												
1973	2511	4676	8866	12090	16895	12510	14973	13609	9090	4309	2040	2449
1974	2759	4816	9331	14340	13020	15750	15283	12090	7620	5146	2400	1581
1975	2170	4788	10261	12630	13733	13920	15748	11470	7410	5084	2610	2821

Table 8. Regression equations relating turbidity related parameters to in situ vertical extinction coefficients [K_ω (IS)] for solar radiation in Lake Winnipeg waters. K_α is a horizontal extinction coefficient derived from transmissometer measurements, and $K_\alpha(A)$ is from the same transmissometer on the attenuated scale, in units of m^{-1} . SS is concentration of suspended sediments, in $g\ m^{-3}$. $K_\omega(DT)$ is a measurement of vertical extinction coefficient in a deck tank of Lake Winnipeg water, utilizing the Whitney submersible photometer, in units of m^{-1} . Chl-a is a fluorometric measurement of chlorophyll a from extracts of seston on a filter paper. F, degrees of freedom, and correlation coefficients are given for each relationship. The letter codes refer to estimated vertical extinction coefficients given in Table 9.

<u>Code</u>	<u>Predicted \hat{K}_ω</u>	<u>Predictors</u>	<u>Significance</u>
(A)	$K_\omega(\text{IS}) = 0.3970 + 0.9632 \ln K_\alpha(A)$		$F_{1,30} = 303 \ r = 0.95$
(B)	$K_\omega(\text{IS}) = 0.4576 + 0.2472 K_\alpha$		$F_{1,45} = 435 \ r = 0.95$
(C)	$\ln K_\omega(\text{IS}) = 0.3412 - 0.8549 \ln \text{Secchi (m)}$		$F_{1,38} = 307 \ r = 0.94$
(D)	$K_\omega(\text{IS}) = 0.5809 + 0.1227 \text{SS (g/m}^3)$		$F_{1,28} = 108 \ r = 0.89$
(E)	$K_\omega(\text{IS}) = -0.2228 + 0.5158 K_\omega(\text{DT})$		$F_{1,29} = 58.5 \ r = 0.82$
(F)	$K_\omega(\text{IS}) = 0.9820 + 2.2497 \times 10^{-4} \text{ Algal Biomass (mg/m}^3)$	$F_{1,19} = 12.2 \ r = 0.62$	
(G)	$K_\omega(\text{IS}) = 1.2239 + 0.5950 \ln \text{Chl-a (mg/m}^3)$	$F_{1,24} = 8.05 \ r = 0.50$	
(H)	$K_\omega(\text{IS}) = 1.3631 + 0.5608 \ln \text{Phaeophyton (mg/m}^3)$	$F_{1,30} = 5.45 \ r = 0.39$	

Table 9. Measured and estimated values for vertical extinction coefficients (\hat{K}_w) of the transmission of incident light into Lake Winnipeg waters at selected stations. The measured values (without code notation) are derived from in situ % of incident light measurements in lake water, using a Whitney submersible photometer [K_w (IS) in previous tables]. The estimated values (with code notation suffix) are derived from regression equations given in Table , utilizing the parameters Secchi visibility, suspended sediments, algal biomass, chlorophyll a, phaeophytin, horizontal extinction coefficients (K_o), and vertical extinction coefficients measured in deck tanks of lake water [K_w (DT)].

Station	CRUISE				
	4-12 June 1969	9-17 July	24 July - 1 Aug.	2-10 Sept.	3-13 Oct.
1	-	4.55	3.45(C)	4.00(D)	2.54(C)
2	4.60(C)	3.94(C)	2.96(A)	2.79(A)	9.58(B)
3	-	7.86(B)	3.44(A)	5.32(B)	11.84(B)
3B	-	1.61(G)	-	-	-
5C	-	4.60(C)	3.20(A)	2.49(B)	3.12(B)
4	-	3.01(A)	2.90(A)	3.08(B)	3.88(B)
5	5.57(C)	2.48(A)	2.37(A)	2.27(B)	2.21(B)
5B	-	2.47(G)	-	-	-
6	3.73(C)	2.30(A)	2.03(A)	2.07(B)	3.05 (B)
7	1.41(C)	2.48(A)	2.00	2.32(B)	1.62 (B)
8	2.54(C)	1.97(A)	2.06	1.68(B)	3.05(B)
9	-	1.73(A)	2.37(A)	-	9.13(B)
10	2.18(C)	1.88	1.40	2.98(B)	3.69(B)
11	1.54(C)	1.50(A)	-	-	-
12	0.995(C)	1.37(A)	1.32(A)	3.35(B)	5.15 (B)
13	0.840(E)	-	-	1.59(H)	1.69(H)
13B	-	1.37(A)	1.28(G)	1.81(G)	3.16(E)
14	0.995(C)	0.868(A)	2.03(A)	2.85(B)	3.99(B)
15	0.943(E)	1.42(G)	5.75(D)	1.38(G)	-
15B	-	-	-	-	1.50(G)
16	1.20(C)	2.48(A)	2.98	9.13(B)	4.52(B)
17	1.91(C)	1.58	1.56(A)	1.42(G)	-
18	1.91(C)	2.26(A)	1.47	1.27(B)	3.27(B)
19	1.62(C)	1.68(B)	1.28(B)	2.72(B)	1.45(B)
20	1.72	1.46	1.23	1.07(B)	2.17(B)

Table 9 . (cont'd)

Station	CRUISE				
	4-12 June 1969	9-17 July	24 July-1 Aug.	2-10 Sept.	3-15 Oct.
20B	-	-	-	-	1.50(G)
21	1.47(C)	1.31	0.822	1.41(B)	1.30(B)
22	1.20(C)	1.05(A)	1.24(A)	3.35(B)	1.07(B)
23A	-	0.515(E)	1.16(D)	-	-
23B	-	-	-	2.12(G)	-
23C	-	0.416(A)	0.835	0.962(B)	0.917(B)
23D	1.30(C)	-	1.54(G)	-	-
23E	0.827	1.02(B)	0.631	1.01(B)	1.04(B)
24	0.813(C)	-	-	-	-
25	0.813(C)	0.656	0.591	2.17(B)	2.73(B)
26	-	0.643(C)	0.550(C)	0.995(C)	3.08(C)
27	-	1.65(B)	1.30(A)	-	5.50(B)
28	0.474(E)	0.643(C)	0.218(A)	1.94(B)	1.98(B)
29	0.768(E)	-	1.02(D)	1.08(D)	0.947(B)
31	0.995(C)	2.49(B)	0.540(A)	1.47(B)	4.20(B)
33	2.54(C)	6.70(B)	2.06(A)	-	7.41(B)
34	-	-	-	-	2.05(G)
35	0.717(C)	0.913	0.736	0.774(B)	0.938
39	1.65(F)	0.772(B)	0.509	1.39(B)	1.65(B)
41	1.54(C)	0.640	0.534	0.922(B)	1.71(B)
43	3.08(C)	1.01(B)	1.36(B)	-	2.73(B)
45	-	0.781(B)	-	1.31(B)	1.22(B)
47	-	-	-	2.61(G)	2.25(G)
48	1.45(F)	1.71(B)	1.32(A)	1.51(B)	1.45(B)
50	-	1.93(E)	-	2.33(H)	-
50B	-	2.99	2.53(A)	-	-
50C	-	2.58(A)	2.29(A)	1.94(B)	4.70(B)
50D	-	-	-	2.30(H)	-
51	-	1.54(A)	1.80(A)	2.22(B)	1.52(B)
52	2.18(C)	2.29(A)	2.53(A)	4.56(B)	4.70(B)

Table 9. (cont'd)

Station	CRUISE				
	4-12 June 1969	9-17 July	24 July-1 Aug.	2-10 Sept.	3-13 Oct.
53	-	1.42(G)	-	-	-
54	1.70(C)	1.90(A)	1.72	3.43(B)	4.20(B)
54B	-	-	-	-	1.38(G)
55	2.54(C)	1.38(G)	1.64(G)	1.28(G)	-
55B	-	-	1.37(D)	-	-
56	-	1.69(G)	1.54(G)	1.54(G)	-
57	2.18(C)	1.17(A)	1.66(A)	2.60(B)	3.12(B)
57B	-	-	-	1.67(G)	-
58	3.94(C)	3.01(A)	2.93(A)	6.70(B)	11.8(B)
59	2.54(C)	1.94(B)	1.82	1.24(B)	1.55
60	3.94(C)	3.39(B)	1.66(A)	1.32(B)	2.85(B)
60B	-	3.82(A)	2.85(A)	2.41(B)	4.44(B)
60C	-	-	3.61(A)	3.60(B)	2.49(B)
61	5.57(C)	5.49	1.84(A)	1.71(B)	2.85(B)
62	5.57(C)	3.74(A)	2.93(A)	3.27(B)	9.13(B)
62B	-	1.72(G)	-	-	-
63	10.1(C)	2.27	4.09(A)	4.99(B)	10.1(B)
64	-	1.97(B)	2.36	2.49(B)	4.70(B)
64B	-	1.52(E)	1.47(G)	-	1.54(G)
65	-	1.73	2.15	-	-
65B	-	-	1.36(H)	-	-
65C	-	-	1.63(H)	-	-
66A	-	-	1.42(G)	-	-
66B	-	-	1.40	-	-
66C	-	-	2.43(D)	-	-
68	-	1.64(B)	2.81	3.43(B)	5.32(B)
68B	-	-	1.54(E)	1.61(G)	1.81(H)
69	-	2.22(B)	2.96	3.12(B)	6.15(B)
69B	-	-	1.61(G)	-	-





Fig. 1. The deck tank used on the wheel-house deck of the Bradbury for light measurements and primary production work.



X = FLUORESCENT LIGHT
 □ = SOLAR RADIATION

Fig. 2. Comparison of the distribution of light energy of the sun (on a cloudy day) with the light bank of fluorescent tubes shown in Fig. 1.




Fig. 3. Location of stations where radiation measurements were made on Lake Winnipeg. See also Brunskill and Graham (1979) for longitude and latitude of these stations.




Fig. 4. Isopleths of the depths (meters) of secchi disc visibility in Lake Winnipeg during 4-12 June 1969.



Fig. 5. Isopleths of the depths (meters) of secchi disc visibility in Lake Winnipeg during 9-17 July 1969.




Fig. 6. Isopleths of the depths (meters) of secchi disc visibility in Lake Winnipeg during 24 July - 1 August 1969.




Fig. 7. Isopleths of the depths (meters) of secchi disc visibility in Lake Winnipeg during 2-10 September 1969.




Fig. 8. Isopleths of the depths (meters) of secchi disc visibility in Lake Winnipeg during 3-13 October 1969.




Fig. 9. Isopleths of the depths (meters) of secchi disc visibility in Lake Winnipeg during 26-31 October 1969.

CITY OF WINNIPEG



Fig. 10. Isopleths of horizontal (beam transmittance) extinction coefficients (K_a , in m^{-1}) for Lake Winnipeg during 9-17 July 1969.






Fig. 12. Isopleths of horizontal (beam transmittance) extinction coefficients (K_α , in m^{-1}) for Lake Winnipeg during 2-10 September 1969.



Fig. 13. Isopleths of horizontal (beam transmittance) extinction coefficients (K_α , in m^{-1}) for Lake Winnipeg during 3-13 October 1969.





Fig. 14. Spectral extinction of radiation at 10 cm water depth in the fluorescent-lighted deck tank (Fig. 1) at Station 35 in the center of the north basin of Lake Winnipeg (Fig. 3) on 8 October 1969.



STATION 58, 9 SEPT 69
0845 HRS.

Fig. 15. Spectral extinction of radiation at 10 cm water depth in the fluorescent-lighted deck tank (Fig. 1) at Station 58 in the northern part of the south basin (Fig. 3) on 9 September 1969.




Fig. 16. Spectral extinction of radiation at 10 cm water depth in the fluorescent-lighted deck tank (Fig. 1) at Station 51 in the mouth of Berens River (Fig. 3) on October 1969.




Fig. 17. Isopleths of concentrations of suspended sediments (g m^{-3}) in Lake Winnipeg during 9-17 July 1969.




Fig. 18. Isopleths of concentrations of suspended sediments (g m^{-2}) in Lake Winnipeg during 24 July - 1 August 1969.




Fig. 19. Isopleths of concentrations of suspended sediments (g m^{-3}) in Lake Winnipeg during 2-10 September 1969.




Fig. 20. Isopleths of concentrations of suspended sediments (g m^{-3}) in Lake Winnipeg during 3-13 October 1969.




Fig. 21. Isopleths of concentrations of suspended sediments (g m^{-3}) in Lake Winnipeg during 26-31 October 1969.




Fig. 22a. Distribution of daily radiation during the days of our research on Lake Winnipeg in 1969 and winter 1970.





Fig. 22b. Distribution of daily radiation during the days of our research on Lake Winnipeg in 1969 and winter 1970.



28 JULY 69






Fig. 22d. Distribution of daily radiation during the days of our research on Lake Winnipeg in 1969 and winter 1970.



6 SEPT 69

63

Fig. 22e. Distribution of daily radiation during the days of our research on Lake Winnipeg in 1969 and winter 1970.






Fig. 22g. Distribution of daily radiation during the days of our research on Lake Winnipeg in 1969 and winter 1970.

17 MARCH 70




Fig. 22h. Distribution of daily radiation during the days of our research on Lake Winnipeg in 1969 and winter 1970.




Fig. 23. The relationship between the depth of secchi disc visibility (m) and vertical extinction coefficients ($K_w(15)$) for Lake Winnipeg in 1969.




Fig. 24. The relationship between vertical extinction coefficients measured in the deck tank (Fig. 1) open to solar radiation ($K_W(DT)$) and an *in situ* measurement of vertical extinction coefficients ($K_W(15)$) in Lake Winnipeg waters in 1969.





Fig. 25. The relationship between horizontal (beam transmittance) extinction coefficients (K_α) and K_W (IS) in Lake Winnipeg waters in 1969.






Fig. 27. The relationship between vertical extinction coefficients ($K_w(15)$) and the concentration of suspended sediments (g m^{-3}) in Lake Winnipeg waters in 1969.




Fig. 28. Isopleths of predicted vertical extinction coefficient (K , in m^{-1}) in Lake Winnipeg during 4-12 June 1969.




Fig. 29. Isopleths of predicted vertical extinction coefficient (K_w , in m^{-1}) in Lake Winnipeg during 9-17 July 1969.




Fig. 30. Isopleths of predicted vertical extinction coefficient (K_w , in m^{-1}) in Lake Winnipeg during 24 July - 1 August 1969.




Fig. 31. Isopleths of predicted vertical extinction coefficient (R_w , in m^{-1}) in Lake Winnipeg during 2-10 September 1969.




Fig. 32. Isopleths of predicted vertical extinction coefficient (k_w , in m^{-1}) in Lake Winnipeg during 3-13 October 1969.




Fig. 33. The relationship between $(K_\alpha - K_w(\text{IS}))$ as an estimator of scattering, and the concentration of suspended sediments in g m^{-3} .
 □ = South Basin, X = North Basin.




Fig. 34. The relationships between suspended sediment concentrations, $K_w(\text{IS})$, and the percentage of $K_w(\text{IS})$ due to suspended sediment.

K_W(15) (%)

PERCENTAGE OF K_W(IS) DUE




Fig. 35. The relationships between chlorophyll α , phytoplankton biomass, $K_w(\text{IS})$, and the percentage of $K_w(\text{IS})$ due to chlorophyll α or phytoplankton biomass.