

Identification of Ecologically or Biologically Significant Marine Areas (EBSAs) in Areas Beyond National Jurisdiction (ABNJ): the Northwest Pacific Seamounts

Cherisse Du Preez, Diva J. Amon, Amy R. Baco, Merlin Best, Georgia Clyde, Ana Colaço, Heidi Gartner, Rachel M. Lauer, Beth N. Orcutt, Anna Metaxas, and Verena Tunnicliffe

Fisheries and Oceans Canada
Science Branch, Pacific Region
Institute of Ocean Sciences
Sidney, British Columbia
V8L 5T5

2023

**Canadian Technical Report of
Fisheries and Aquatic Sciences 3571**



Fisheries and Oceans
Canada

Pêches et Océans
Canada

Canada

Canadian Technical Report of Fisheries and Aquatic Sciences

Technical reports contain scientific and technical information that contributes to existing knowledge but which is not normally appropriate for primary literature. Technical reports are directed primarily toward a worldwide audience and have an international distribution. No restriction is placed on subject matter and the series reflects the broad interests and policies of Fisheries and Oceans Canada, namely, fisheries and aquatic sciences.

Technical reports may be cited as full publications. The correct citation appears above the abstract of each report. Each report is abstracted in the data base *Aquatic Sciences and Fisheries Abstracts*.

Technical reports are produced regionally but are numbered nationally. Requests for individual reports will be filled by the issuing establishment listed on the front cover and title page.

Numbers 1-456 in this series were issued as Technical Reports of the Fisheries Research Board of Canada. Numbers 457-714 were issued as Department of the Environment, Fisheries and Marine Service, Research and Development Directorate Technical Reports. Numbers 715-924 were issued as Department of Fisheries and Environment, Fisheries and Marine Service Technical Reports. The current series name was changed with report number 925.

Rapport technique canadien des sciences halieutiques et aquatiques

Les rapports techniques contiennent des renseignements scientifiques et techniques qui constituent une contribution aux connaissances actuelles, mais qui ne sont pas normalement appropriés pour la publication dans un journal scientifique. Les rapports techniques sont destinés essentiellement à un public international et ils sont distribués à cet échelon. Il n'y a aucune restriction quant au sujet; de fait, la série reflète la vaste gamme des intérêts et des politiques de Pêches et Océans Canada, c'est-à-dire les sciences halieutiques et aquatiques.

Les rapports techniques peuvent être cités comme des publications à part entière. Le titre exact figure au-dessus du résumé de chaque rapport. Les rapports techniques sont résumés dans la base de données *Résumés des sciences aquatiques et halieutiques*.

Les rapports techniques sont produits à l'échelon régional, mais numérotés à l'échelon national. Les demandes de rapports seront satisfaites par l'établissement auteur dont le nom figure sur la couverture et la page du titre.

Les numéros 1 à 456 de cette série ont été publiés à titre de Rapports techniques de l'Office des recherches sur les pêcheries du Canada. Les numéros 457 à 714 sont parus à titre de Rapports techniques de la Direction générale de la recherche et du développement, Service des pêches et de la mer, ministère de l'Environnement. Les numéros 715 à 924 ont été publiés à titre de Rapports techniques du Service des pêches et de la mer, ministère des Pêches et de l'Environnement. Le nom actuel de la série a été établi lors de la parution du numéro 925.

Canadian Technical Report of
Fisheries and Aquatic Sciences 3571

2023

IDENTIFICATION OF ECOLOGICALLY OR BIOLOGICALLY SIGNIFICANT MARINE AREAS
(EBSAS) IN AREAS BEYOND NATIONAL JURISDICTION (ABNJ):
THE NORTHWEST PACIFIC SEAMOUNTS

By

Cherisse Du Preez^{1,2}, Diva J. Amon^{3,4}, Amy R. Baco⁵, Merlin Best¹, Georgia Clyde¹, Ana Colaço⁶, Heidi Gartner¹, Rachel M. Lauer⁷, Beth N. Orcutt⁸, Anna Metaxas⁹, and Verena Tunnicliffe²

¹Fisheries and Oceans Canada, Science Branch, Pacific Region, Institute of Ocean Sciences, 9860 West Saanich Road, Sidney, BC, Canada V8L 4B2

²University of Victoria, 3800 Finnerty Road, Victoria, BC, Canada V8P 5C2

³Marine Science Institute, University of California, Santa Barbara, Santa Barbara, CA, USA

⁴SpeSeas, D'Abadie, Trinidad and Tobago

⁵Florida State University, Department of Earth, Ocean and Atmospheric Sciences, 1011 Academic Way, Tallahassee, FL, United States 32306

⁶Institute of Marine Sciences - Okeanos, University of the Azores, Rua Professor Doutor Frederico Machado 4, 9901-862 Horta, Portugal

⁷University of Calgary, 2500 University Drive NW, Calgary, AB, Canada T2N 1N4

⁸Bigelow Laboratory for Ocean Sciences, 60 Bigelow Drive, East Boothbay, ME, United States 04544

⁹Dalhousie University, Department of Oceanography, Halifax, NS, Canada B3H4R2

© His Majesty the King in Right of Canada, as represented by the Minister of the Department of Fisheries and Oceans, 2023

Cat. No. Fs97-6/3571E-PDF ISBN 978-0-660-68326-3 ISSN 1488-5379

Correct citation for this publication:

Du Preez, C., Amon, D. J., Baco, A. R., Best, M., Clyde, G., Colaço, A., Gartner, H., Lauer, R.M., Orcutt, B. N., Metaxas, A. and Tunnicliffe, V. 2023. Identification of Ecologically or Biologically Significant Marine Areas (EBSAs) in Areas Beyond National Jurisdiction (ABNJ): the Northwest Pacific Seamounts. Can. Tech. Rep. Fish. Aquat. Sci. 3571: vi + 21 p.

Table of Contents

Table of Contents	iii
List of Tables	iv
List of Figures	iv
Abstract	v
Résumé.....	vi
Introduction	1
Location.....	2
Feature description of the area.....	3
Feature condition and future outlook of the area.....	4
Assessment of the area against Convention on Biological Diversity Ecological or Biological Significant Area criteria	5
Conclusions	14
Acknowledgements	15
Literature cited.....	15

List of Tables

Table 1. The Convention on Biological Diversity Ecological or Biological Significant Area scientific criteria and descriptions, and the rankings and supporting information for the Northwest Pacific seamount complex in Areas Beyond National Jurisdiction.....	5
Table 2. Summary of Ecological or Biological Significant Area assessment of the Northwest Pacific seamounts in Areas Beyond National Jurisdiction against the seven criteria	14

List of Figures

Figure 1. The Northwest Pacific Ocean Seamounts Ecologically or Biologically Significant Area	3
Figure 2. The Northwest Pacific Ocean Seamounts support abundance and diverse communities.....	13

Abstract

Du Preez, C., Amon, D. J., Baco, A. R., Best, M., Clyde, G., Colaço, A., Gartner, H., Lauer, R.M., Orcutt, B. N., Metaxas, A. and Tunnicliffe, V. 2023. Identification of Ecologically or Biologically Significant Marine Areas (EBSAs) in Areas Beyond National Jurisdiction (ABNJ): the Northwest Pacific Seamounts. Can. Tech. Rep. Fish. Aquat. Sci. 3571: vi + 21 p.

Marine Areas Beyond National Jurisdiction (ABNJ) are the common heritage of humankind and require coordinated protection, conservation, restoration, and sustainable use by the international community. A common first step in marine area-based management is the identification of Ecologically or Biologically Significant Marine Areas (EBSAs) using the United Nations Convention of Biological Diversity (CBD) scientific criteria. While possessing many EBSA qualities, a published EBSA assessment for the Northwest (NW) Pacific seamounts is lacking. Herein, the CBD EBSA criteria are applied to a seamount complex within ABNJ between the United States (Mariana Islands, Guam, Wake Island Marine National Monument), Federated States of Micronesia, Marshall Islands, and Japan (Ogasawara and Minamitori Island). The EBSA assessment concluded that these Magellan, Marcus-Wake, and Marshall seamounts rank *High* for all EBSA criteria except one (*Medium* ranking) and strongly identify these NW Pacific seamounts as an EBSA, including their surrounding water. Existing or anticipated threats to the seamounts include but are not limited to seabed mining and climate change. This NW Pacific Ocean Seamounts EBSA identification synthesizes the best available science and provides a robust assessment and thorough review to support marine area-based management and future CBD workshops on the region.

Résumé

Du Preez, C., Amon, D. J., Baco, A. R., Best, M., Clyde, G., Colaço, A., Gartner, H., Lauer, R.M., Orcutt, B. N., Metaxas, A. and Tunnicliffe, V. 2023. Identification of Ecologically or Biologically Significant Marine Areas (EBSAs) in Areas Beyond National Jurisdiction (ABNJ): the Northwest Pacific Seamounts. Can. Tech. Rep. Fish. Aquat. Sci. 3571: vi + 21 p.

Les zones situées au-delà de la zone de compétence nationale constituent le patrimoine commun de l'humanité et nécessitent une protection, une conservation, une restauration et une utilisation durable coordonnées par la communauté internationale. Une première étape commune de la gestion marine par zone est la désignation des zones marines d'importance écologique ou biologique (ZIEB) à l'aide des critères scientifiques de la Convention des Nations Unies sur la diversité biologique (CDB). Bien que les monts sous-marins du Pacifique Nord-Ouest présentent de nombreuses qualités propres aux ZIEB, leur évaluation par rapport aux critères de désignation des ZIEB reste à publier. Dans le présent document, les critères de désignation des ZIEB établis dans la CDB sont appliqués à un complexe de monts sous-marins dans la ZIEB située entre les États-Unis (îles Mariannes, Guam, monument national marin de l'île), les États fédérés de Micronésie, les îles Marshall et le Japon (archipel d'Ogasawara et île Minamitori). L'évaluation par rapport aux critères de désignation des ZIEB a permis de conclure que les monts sous-marins Magellan, Marcus-Wake et Marshall se classent à un niveau *élevé* pour tous les critères, sauf un (*moyen*) et qu'il est pleinement justifié qu'ils forment une ZIEB avec leurs eaux environnantes. Les menaces actuelles ou prévues pour les monts sous-marins sont notamment l'exploitation minière des fonds marins et les changements climatiques. Le processus de désignation des monts sous-marins du Pacifique Nord-Ouest en tant que ZIEB synthétise les meilleures données scientifiques disponibles et constitue une évaluation solide et un examen approfondi à l'appui de la gestion marine par zone et des futurs ateliers de la CDB dans la région.

Introduction

Marine Areas Beyond National Jurisdiction (ABNJ) are the common heritage of humankind and require coordinated protection, conservation, restoration, and sustainable use by the international community. There are multiple international agreements supporting this objective (e.g., the Sustainable Development Goals and the Biodiversity Beyond National Jurisdiction ‘treaty’; Christiansen et al. 2022), the success of which relies on all signatories to uphold the principles, rights, duties, and obligations, such as applying precautionary and ecosystem-based approaches and using science-based, transparent management.

A common first step in marine area-based management is the identification of Ecologically or Biologically Significant Marine Areas (EBSAs) using the United Nations Convention of Biological Diversity (CBD) scientific criteria as defined in Annex I of Decision IX/20 of its 9th Conference of Parties (UNEP/CBD 2008). The seven CBD EBSA criteria are (1) uniqueness or rarity, (2) special importance for life-history stages of species, (3) importance for threatened, endangered or declining species and/or habitats, (4) vulnerability, fragility, sensitivity, or slow recovery, (5) biological productivity, (6) biological diversity, and (7) naturalness. A marine feature is identified as an EBSA if it scores *High* on at least one criterion or above average across all (i.e., *Medium* or *High* for cumulative importance; DFO 2004, 2011). The possible rankings are: *No information*, *Low*, *Medium*, and *High*.

Seamounts are underwater mountains—usually volcanic in origin—that provide many essential ecosystem functions and services (summarized in DOSI 2023). Seamounts are most abundant in the North Pacific Ocean and are notably concentrated in three regions (Kitchingman et al. 2007; FAO 2023). The CBD has evaluated two of these three North Pacific seamount regions and designated large complexes of seamount chains and clusters as EBSAs in ABNJ: the ‘Emperor Seamount Chain and Northern Hawaiian Ridge’ (central North Pacific) and the ‘Northeast Pacific Ocean Seamounts’. These EBSAs were identified because the seamount chains have unique geographic features, provide habitats for a variety of benthic species (including vulnerable and long-lived cold-water corals and sponges), are fish spawning grounds, their biological productivity is higher than that of the neighbouring ocean floor, and their chains and configurations—spanning thousands of kilometres—facilitates migration (CBD 2016a,b). Despite sharing many of these EBSA qualities—as well as holding global distinction among seamounts—a published EBSA assessment for the Northwest (NW) Pacific seamounts is lacking.

Herein, the CBD EBSA criteria are applied to the region of NW Pacific seamounts within ABNJ, i.e. the complex of Magellan, Marcus-Wake, and Marshall seamounts. This EBSA assessment was completed by an international working group of scientists led by Fisheries and Oceans Canada. Members of the group participated in two international science workshops on the topic: the Workshop on the Development of a Regional Environmental Management Plan (REMP) for the Area of the Northwest Pacific (hosted by the International Seabed Authority (ISA), online, October 26th to November 6th 2020) and the Global Ocean Biodiversity Initiative (GOBI) EBSAs in ABNJ workshop (hosted by GOBI, in Santa Cruz, California, November 6th to 9th 2022).

At REMP workshop, invited experts reviewed data and information on seamounts in the area, much of which were pre-compiled in documents to support the discussions. The ISA Workshop Report (ISA 2022) captures the approach adopted. The workshop produced a predictive model of the areas and abundances of the region’s seamounts (n = 205) (using the Benthic Terrain Modeler; Walbridge et al. 2018). EBSA-related assessments were conducted for three surveyed seamounts and the seamount complex as a whole, all scoring *Medium* to *High* rankings. At the EBSA in ABNJ workshop, invited experts reviewed and assessed EBSAs in ABNJ using the CBD EBSA workshop approach (GOBI 2023). An extended draft EBSA assessment for the NW Pacific seamount complex was discussed and refined using new publications and additional relevant biological and environmental information and datasets. The EBSA assessment for the ABNJ NW Pacific seamounts is now presented here in its completed form.

Ecological research on these ABNJ NW Pacific seamounts is still in its early stages, as indicated by the frequent discovery of “species new to science” during each expedition (e.g., Wang et al. 2016; Dong et al. 2017; Xu et al. 2017; Dong et al. 2019; Hestetun et al. 2019; Zhang et al. 2019, 2020; Kou et al. 2020; Gan et al. 2020; Shen et al. 2020; Chen et al. 2021a, b; Grischenko 2022). Very few of the seamounts have been visually surveyed, and fewer still have openly accessible data (e.g., Ocean Exploration 2016 expedition dive on Alba Seamount; <https://www.ncei.noaa.gov/waf/oceanos-rov-cruises/ex1606/>). Public access to video surveys of the benthos by seabed mining companies and contractors (referenced in ISA 2022) would help fill existing knowledge gaps. However, the growing body of published literature, knowledge from international subject matter experts (workshop participants), and openly available large-scale datasets (e.g., the GEBCO maps and Undersea Feature Names database (IHO-IOC GEBCO Gazetteer of Undersea Feature Names, www.gebco.net), seamount classification systems (e.g., Yesson et al. 2011), and global distributions of cold-water corals (e.g., Yesson et al. 2012, 2017)) provide more than sufficient information to complete a robust EBSA assessment of the seven criteria (e.g., CBD 2016a,b).

Location

EBSA region: NW Pacific Ocean (Figure 1).

Description of location: The Magellan, Marcus-Wake, and Marshall seamounts within ABNJ (labelled on Figure 1), in an area approximately 1,500 km in diameter, between the Northern Mariana Islands and Guam of the USA, the Federated States of Micronesia, the Marshall Islands, Wake Island Unit of the Pacific Remote Islands National Marine Monument of the USA, and Ogasawara and Minamitori Islands of Japan (labelled on Figure 1).

In alignment with the other North Pacific seamount EBSA assessments (CBD 2016a,b), the NW Pacific seamount complex is evaluated here as a single EBSA. The NW Pacific seamounts function as a unit or network, having similar geological origins and forming a dense configuration of mountainous massifs (details provided in the assessment below). The buffering and grouping of seamounts by the EBSA boundaries is in alignment with the established practices of the Government of Canada and the Convention on Biological Diversity (Ban et al. 2016; CBD 2016a,b). The EBSA includes all the seamounts (regardless of summit depth) in the complex, their footprint and the water column above (Ban et al. 2016; CBD 2016a,b).

Approximate extent: 26°0'00"N 145°0'00"E (NW corner), 26°0'00"N 165°0'00"E (NE corner), 10°0'00"N 165°0'00"E (SE corner), 10°0'00"N 145°0'00"E (SW corner).




Figure 1. The Northwest (NW) Pacific Ocean Seamounts Ecologically or Biologically Significant Area (EBSA). The EBSA contains Magellan, Marcus-Wake, and Marshall seamounts in Areas Beyond National Jurisdiction (ABNJ) between the Exclusive Economic Zones (EEZs) of the United States (Northern Mariana Islands, Guam, the Wake Island Unit of the Pacific Remote Islands National Marine Monument), the Federated States of Micronesia, the Marshall Islands, and Japan (Minamitori Island and Ogasawara). Seamounts adapted from ISA 2022.

Feature description of the area

The assessed seamount complex includes the oldest seamounts and oceanic crust in the world (Clouard and Bonneville 2005; Müller et al. 2016) and some of the tallest seamounts, rising from some of the deepest basins in the world (e.g., east Marianas Basin; Stewart and Jamieson 2019) to summits as shallow as 550 m depth (e.g., Alba Guyot; Mel’nikov et al. 2016). This area also contains the highest density of seamounts in the world (Kitchingman et al. 2007; FAO 2023). Here, the Magellan, Marshall, and Marcus-Wake seamounts intersect and form a complex of volcanotectonic massifs containing approximately 205 seamounts within ABNJ (ISA 2022; only about half are named features, www.gebco.net). While abundant in number, the cumulative area of the seamounts is small and highly fragmented. Within this area, the bare rock seamounts account for only ~4.3% of the ocean floor (ISA 2022) and represent the only shallow bathyal zone habitat (between 3000 and 800 m depth; UNESCO 2009; Watling et al. 2013; ISA 2022). In contrast, the other 95.7% of the ocean floor is predominantly muddy abyssal plains and nodule fields between 6500 and 3500 m depth (UNESCO 2009; ISA 2022).

The flow of currents around and over the seamount complex is not yet resolved or well understood (ISA 2022), but it is known that these seamounts can alter local hydrodynamics, and most are within range of directly interacting with and/or impacting at least their immediate neighbours (Jiang et al. 2021; Xie et al.

2022). Prevailing currents that limit or prohibit the movement of organisms between seamounts may promote geographically isolated metapopulations or potentially endemic species. In contrast, currents that facilitate movement may result in seamount stepping stones that form a pathway for gene flow among metapopulations and migration of species (Rowden et al. 2010; Shank 2010; Miller and Gunasekera 2017).

The heights and the natural variability of the seamounts within the complex (e.g., seamount size, geomorphology, latitude, etc., summarized in ISA 2022) will drive high beta diversity within and among seamounts, and these species turnover patterns have been observed on the surveyed seamounts within the complex (e.g., Chen et al. 2021a; Shen et al. 2021). The seamounts are home to diverse fields of habitat-forming cold-water corals and sponges, dense brittle star mats, fish, sea stars, crinoids, and sea cucumbers (e.g., as summarized in ISA 2022). The seamounts are used as stopover points during migrations (e.g., as feeding grounds; Rogers 2018), and there are many transient species known to pass over the seamount complex, including commercial species (e.g., tuna and billfish; WCPFC 2019) and species of conservation concern (e.g., endangered whales, sea birds, and turtles; Block et al. 2011; IWC 2020; IUCN 2023).

Feature condition and future outlook of the area

The future outlook of the area is highly uncertain. There are cobalt-rich ferromanganese crusts (CFC) and polymetallic nodule (PMN) deep-sea mining exploration contracts from the ISA and reserve areas on dozens of the seamounts with a bias for the tallest, shallowest, flattest, and largest of the features in the complex (ISA 2022) (i.e., those most likely to have highest EBSA rankings; e.g., EBSA indicator characteristics listed in Taranto et al. 2012). These contracts are held by multiple mining entities (five companies: Beijing Pioneer Hi-Tech Development Corporation (BPHDC; China), the Government of the Republic of Korea, the China Ocean Mineral Research and Development Association (COMRA), Japan Oil, Gas, and Metals Corporation (JOGME), and the Government of the Russian Federation (<https://www.isa.org.jm/maps/clarion-clipperton-fracture-zone-2/>)). If mineral exploitation begins, the mining-related stressors (e.g., extreme physical disturbance, habitat removal and crushing, noise and light pollution, sediment plumes, increased ship traffic) are expected to be significant and far-reaching (i.e., on the benthos, mid-water, surface water, at and surrounding the mining sites) (e.g., Levin et al. 2016; Gollner et al. 2017; Drazen et al. 2019; and references cited therein). The concurrence of this, the first seamount mining contract and reserve areas in the world, and the EBSA qualities of these seamounts are not a coincidence. Many of the environmental conditions that make the area attractive for mining exploitation are unique, rare, and unusual biophysical and ecological seamount characteristics (e.g., geological age, environmental stability, shallow depths, accelerated currents, dense assemblage of seamounts, expansive exposed hard substrate with low sedimentation, etc.).

Additional existing and anticipated stressors in the NW Pacific include (but are not limited to) overfishing (e.g., close proximity to fishing nations), shipping traffic, pollution (including marine debris (Amon et al., 2020), microplastics, chemicals, radionuclides (e.g., Fukushima wastewater dumping; Liu and Hoskin 2023), noise and light, and climate change (e.g., Murray et al. 2015). There is evidence of overfishing of large predator species (e.g., tuna, marlin, sharks in the region; WCPFC 2019) and illegal, unreported, and unregulated fishing over these NW Pacific seamounts (Welch et al. 2022). Climate change is rapidly impacting the global ocean. It is altering temperature, acidity, oxygen concentration, currents, species distributions, marine diseases, food webs, surface conditions, etc. (IPCC 2019). Seamounts in the NW Pacific are already experiencing rapid climate change effects. For example, one of the strongest global changes in the ocean heat content in the upper 2000 m depth range occurred in this area (Cheng et al. 2021). Seamount fauna face heightened vulnerability to changing conditions due to multiple life history traits, including their limited ability to relocate (e.g., anchored in place or have a finite area of suitable habitat; Ross et al. 2020). It is unknown what cumulative and synergistic effects these local, regional, and

global stressors will have on seamount-associated species, but many deep-sea studies and models forecast local and even global extinctions (e.g., Levin et al. 2020; Ross et al. 2020; Cheung et al. 2022).

Assessment of the area against the Convention on Biological Diversity (CBD) Ecological or Biological Significant Area (EBSA) criteria

The NW Pacific seamounts in ABNJ were assessed against the seven CBD EBSA criteria (Table 1).

Table 1. The Convention on Biological Diversity (CBD) Ecological or Biological Significant Area (EBSA) scientific criteria and descriptions, and the rankings and supporting information (justifications) for the Northwest (NW) Pacific seamount complex in Areas Beyond National Jurisdiction (ABNJ). Possible rankings: *No information*, *Low*, *Medium*, and *High*.

CBD EBSA Criteria	Description	Ranking of criterion relevance			
		No info.	Low	Medium	High
Uniqueness or rarity	Area contains either (i) unique (“the only one of its kind”), rare (occurs only in few locations) or endemic species, populations or communities, and/or (ii) unique, rare or distinct, habitats or ecosystems; and/or (iii) unique or unusual geomorphological or oceanographic features.				X
<p>Justification</p> <p>The ABNJ NW Pacific Ocean seamounts contain (i) unique, rare, and potentially endemic species (e.g., many species new to science), (ii) regionally and globally rare and distinct habitats and ecosystems (e.g., shallow, rocky, and current-swept habitats), and (iii) globally unique and unusual geomorphological and oceanographic features (e.g., some of the oldest, deepest, densest, tallest, most encrusted seamounts in the world). Therefore, the uniqueness and rarity of these seamounts is <i>High</i> (details are provided below).</p> <p>The interaction of the geographic isolation of these NW Pacific seamounts and the deep-water currents within the area may be contributing to intraspecific differentiation (Kou et al. 2020) and, ultimately, endemic species. Future research is required to resolve whether new species discoveries are endemic or representative of rare species in the under-sampled region (e.g., Rowden et al. 2010). That said, the NW Pacific seamounts are proving to be a hot spot for new species discoveries and specimen collections of species that are rare or endemic species on and around the seamounts include: bryozoa, sea stars, brittle stars, sponges, corals, shrimp, barnacles, and other crustaceans (Wang et al. 2016; Dong et al. 2017; Xu et al. 2017; Dong et al. 2019; Hestetun et al. 2019; Zhang et al. 2019, 2020; Kou et al. 2020; Gan et al. 2020; Shen et al. 2020; Chen et al. 2021a,b; Grischenko 2022). This discussion on rare and endemic species also applies to the next section (i.e., ‘Special importance for life-history stages of species’).</p> <p>In general, seamounts provide a distinctly high diversity of regionally rare habitats in close proximity, supporting a high diversity of seamount-associated species. They provide deep to shallow ocean floor, hard substrate, complex topography, altered oceanographic conditions, enhanced currents, sites of biogenic habitats (e.g., cold-water corals and sponges; Figure 2.a,b), etc. (e.g., Pitcher and Bulman 2007; Mel’nikov et al., 2016; ISA 2022; DOSI 2023). These NW Pacific seamounts, in particular, are regionally rare habitats compared to the surrounding abyssal plains. Based on analysis from the ISA 2020 workshop, only 4.26% of the ocean floor within this region of ABNJ rises over 3,000 m depth and all of it is on seamounts (ISA 2022):</p> <ul style="list-style-type: none"> ● 1,000 to 0 m depth (i.e., on seamounts): 0.12% 					

- 2,000 to 1,000 m depth (i.e., on seamounts): 1.84%
- 3,000 to 2,000 m depth (i.e., on seamounts): 2.30%
- 3,000 to 6,000 m depth: 95.74%

The seamount habitats are also relatively rare on a global scale since the biogeographic province to which the seamounts belong is a relatively small one (i.e., the West Pacific lower bathyal province; UNESCO 2009; Watling et al. 2013), resulting in a rare corresponding faunal assemblage. Furthermore, within the biogeographic province itself, the seamounts represent unusually small and isolated patches (UNESCO 2009; Watling et al. 2013). These seamounts represent different abyssal and lower bathyal biogeographical provinces (UNESCO 2009; Watling et al. 2013) than the existing North Pacific seamount EBSAs (i.e., ‘Emperor Seamount Chain and Northern Hawaiian Ridge’ and ‘North-east Pacific Ocean Seamounts’ EBSAs; www.cbd.int/ebsa/).

The flanks and summits of NW Pacific seamounts are predicted to hold the thickest and highest concentration of ferromanganese (Fe-Mn) crust in the global ocean (Hein et al. 2013). This accumulation is caused by a unique combination of oceanographic processes (e.g., seamount-altered currents, increased deep circulation, upwelling, high nutrient concentration, high primary productivity, sediment-free lava basalt, shallow ocean floor) interacting over 75 million years (Glasby et al. 2007; Ding et al. 2009; Hein et al. 2013). Growth rates of 1-5 mm per million years (Ding et al. 2009; Hein et al. 2013) form crusts up to 200 mm thick on some of the Magellan seamounts (Glasby et al. 2007).

The unusual density and height of these NW Pacific seamounts are discussed in a section below (i.e., ‘Biological productivity’ and ‘Biological diversity’).

Special importance for life-history stages of species	Areas that are required for a population to survive and thrive.	No info.	Low	Medium	High
					X

Justification

These NW Pacific seamounts connect populations and metapopulations of bathyal and benthopelagic species, allowing them to survive and thrive within the ABNJ and likely adjacent Exclusive Economic Zones (EEZs); and the shallower seamounts provide rare habitats for migratory species (oceanic migration and ontogenetic vertical migration). More research is required to understand the specific ecosystem functions provided by the seamounts to associated species, therefore, the special importance for life-history stages of species of these seamounts scored *Medium* without this information but may be determined *High* in the future (details are provided below).

These seamounts are a rare source of hard substrate habitat (exposed lava) and represent the only instances of the shallower bathyal ocean floor in the region (<3000 m depth) (ISA 2022). Therefore, the seamounts are essential habitats for all bathyal benthic and benthopelagic species and communities that are unable to live in the surrounding deep and muddy abyssal plains, such as cold-water corals and sponges, benthic and benthopelagic fish and squid, and potentially hundreds to thousands of other species (many unknown to science). Later discussions on these seamount-associated species also apply to this section.

It is well known that hydrodynamics play an important role in the biological connectivity among deep-sea populations (Shank 2010) and that seamounts influence local and regional hydrodynamics, creating areas of upwelling, enhanced flow, turbulence, tidal rectification, internal waves, eddies, lee waves, etc. (Pitcher and Bulman 2007). Depending on regional oceanographic conditions, groups of seamounts can either function as a collection of isolated submarine islands or as an interconnected network, largely depending

on whether or not larval transport and colonization are facilitated by the prevailing currents (Rowden et al. 2010; Shank 2010; Miller and Gunasekera 2017).

The high density of NW Pacific seamounts generates complex flow patterns around and between seamounts, including trapped and mobile seamount-created mesoscale eddies (measured and modeled data; Jiang et al. 2021; Xie et al. 2022). Such mesoscale eddies can be important for transporting larvae around individual NW Pacific seamounts (trapped eddies) and between neighbouring seamounts (mobile eddies advected past other seamounts) (Jiang et al. 2021). Population connectivity has been documented between the individual NW Pacific seamounts and the different clusters and chains (e.g., Kou et al. 2020; Zhang et al. 2020; Chen et al. 2021a; Na et al. 2021).

Given their central location, configuration, and the uniqueness of the shallower habitats they provide, it is highly likely these ABNJ seamounts and their oceanographic processes also have larger effective oceanographic footprints, potentially influencing and connecting surrounding seamounts and other adjacent shallower bathyal habitats within the EEZs of the United States (Marianas Trench and Volcanic Arc Marine National Monument, Guam, Wake Island Marine National Monument), Federated States of Micronesia, Marshall Islands, Japan (Izu-Bonin Volcanic Arc), and possibly further. In this region, near-surface drifters tracked from the Wake Island longitude all moved westward over the seamount complex and ABNJ and into the Mariana Volcanic Arc (Kendall and Poti 2014). It is currently unknown which habitat patches support source or sink populations, what the connective pathways are, what the environmental limitations and barriers are, and how such connectivity and/or recruitment regimes change over time (e.g., steady supply of recruits, episodic recruitment events, self-recruitment; as described in Parker and Tunnicliffe 1994 and Shank 2010). The dynamics discussed above are currently unresolved but key to structuring the seamount-associated populations, contributing to the justification for the *Medium* ranking.

In addition, globally, seamounts are known to be important habitat and stopover grounds for migrating species, including commercial species, species at risk, and culturally important species (Rowden et al. 2010). Seamounts also facilitate ontogenetic vertical migration into the deep or the benthopelagic zone, with their summits acting as rare shallower nursery grounds for young to settle (e.g., Koslow 1996). As such, the migration discussion in the next section also applies here (i.e., ‘Importance for threatened, endangered or declining species and/or habitats’).

The importance of these offshore habitats for species to survive may increase as climate variables continue to change and impact nearshore populations first, with seamounts becoming potential refugia for some species (e.g., cold-water stony corals; Tittensor et al. 2010; summarized in Rowden et al. 2010).

Importance for threatened, endangered or declining species and/or habitats	Area containing habitat for the survival and recovery of endangered, threatened, declining species or areas with significant assemblages of such species.	No info.	Low	Medium	High
					X

Justification

The ABNJ NW Pacific seamounts contain habitat for the survival and recovery of endangered, threatened, declining species and areas with significant assemblages of such species, including sea turtles, sea birds, whales, sharks, pelagic fish, vulnerable cold-water corals and sponges, etc. Therefore, the importance for threatened, endangered, and declining species and habitat of these seamounts is *High* (details are provided below).

Seamounts are known hotspots of pelagic biodiversity in the open ocean and transient megafauna are

known to gather around or benefit from seamount services (e.g., Rogers et al. 2018), and these NW Pacific seamounts are no exception (e.g. the presence of pelagic predators; Morato et al. 2010). Some species travel specifically to this area (e.g., polar sea birds, potentially as feeding grounds; Harrison et al. 2021), while others use converging pathways through the area to access nearby resources (e.g., Leatherback sea turtle (*Dermochelys coriacea*) nesting grounds; Benson et al. 2011; Figure 2.c). Many of these pelagic species are classified as threatened and endangered species on the International Union for Conservation of Nature (IUCN) Red List (IUCN 2023).

The International Whaling Commission currently lists data for 21 species of interest (IWC 2020), 16 of which have known primary or secondary ranges that overlap the NW Pacific seamount area, including the endangered Blue Whale (*Balaenoptera musculus*), Sei Whale (*Balaenoptera borealis*), and Sperm Whale (*Physeter macrocephalus*; Figure 2.d). Approximately 65% of seabird species in this NW Pacific seamount area are under some degree of threat, with 44% of species classified as near threatened, vulnerable, or endangered (IUCN 2023; e.g., the Short-tailed Albatross, *Phoebastria albatrus* (Figure 2.e), the Phoenix Petrel, *Pterodroma alba*, and the Buller’s Shearwaters, *Ardenna bulleri*). Sea turtle migration pathways have also been documented crossing the seamount area, including that of the vulnerable Leatherback (*D. coriacea*) and endangered Loggerhead (*Caretta caretta*) turtles (Block et al. 2011).

The area supports large pelagic fish and is a known spawning ground for many of these species (Buenafe et al. 2022). The area is surrounded by fishing nations and, as such, it experiences high pelagic fishing. The seamounts overlap the Western and Central Pacific Fisheries Commission (WCPFC) and the North Pacific Fisheries Commission (NPFC) convention areas and are located close to the southwest corner of the North Pacific Anadromous Fish Commission (NPAFC) convention area (Løbach et al. 2020). There are commercially targeted species, including declining or overfished species, in the area. For example, in 2017, the WCPFC authorized between 3,000 and 7,000 vessels to fish up to ten stocks, including Pacific Bluefin Tuna (*Thunnus orientalis*), North Pacific Striped Marlin (*Tetrapturus audax*, also *Kajikia audax*), and Oceanic Whitetip Shark (*Carcharhinus longimanus*), all three of which were assessed as ‘overfished’ and ‘overfishing’ by the WCPFC (2019) (Figure 2.f-h). Some shark, billfish, tuna, and other species had higher probability of capture and higher numbers of fish caught in association with seamounts over the southern Magellan and Marshall seamounts (Morato et al. 2010).

Most deep-sea taxa are not included in the IUCN dataset, but, based on the best available science, many of them would likely also classify as threatened or endangered. This has been recognized by the Convention on the International Trade in Endangered Species (CITES), which lists all species of black corals (Antipatharia) and many genera of stony corals (Scleractinia), coral taxa that are commonly found on seamounts of the NW Pacific, under CITES Appendix II, meaning that they may become at risk of extinction in the near future (<https://cites.org/eng>). In addition, many of the deep-sea species are designated by the Food and Agriculture Organization (FAO) of the United Nations and the Regional Fisheries Management Organizations as Vulnerable Marine Ecosystem (VME) indicator taxa, hence the VME discussion in the next section is also applicable here.

Vulnerability, fragility, sensitivity, or slow recovery	Areas that contain a relatively high proportion of sensitive habitats, biotopes or species that are functionally fragile (highly susceptible to degradation or depletion by human activity or by natural events) or with slow recovery.	No info.	Low	Medium	High
					X

Justification

The ABNJ NW Pacific seamounts contain a relatively high proportion of sensitive habitats, biotopes and

species that are functionally fragile and with slow recovery, such as all cold-water coral and sponges Vulnerable Marine Ecosystems (VMEs) that live above 3000 m in the area, all seamounts supporting these species groups, and regionally specific rare and endemic deep-sea animals. Therefore, the vulnerability, fragility, sensitivity, and slow recovery of these seamounts is *High* (details are provided below).

Cold-water coral and sponges, as well as other emergent and structure-forming invertebrates (e.g., xenophyophores), have life history characteristics such as being long-lived, slow-growing, sessile, and rigid, that make them fragile, vulnerable to impacts, and slow to recover. As such, assemblages of these taxa are considered to be VMEs (FAO 2009b; based on the designation criteria: uniqueness or rarity, functional significance of the habitat, fragility, life-history traits of component species that make recovery difficult, structural complexity). Seamounts that support these species groups are also considered VMEs (FAO 2009b).

NW Pacific seamounts are home to VME taxa, with some surveyed sites described as covered in dense and diverse cold-water coral and sponge fields (Shen et al. 2020; Shen et al. 2021; COMRA and Russian Federation expeditions reported in ISA 2022). Open data sources also confirm VME taxa occur on the surveyed seamounts (e.g., the Deepwater Exploration of the Marianas summarized in Bell et al. 2017; NOAA Deep-Sea Coral Data Portal; <https://deepseacoraldata.noaa.gov/>). The probability of the unexplored NW Pacific seamounts in the area harboring VME taxa is almost certain. Of the hundreds of seamounts surveyed to date in the North and Central Pacific, all have cold-water corals, sponges, and/or other VME taxa (e.g., Baco 2007; Parrish et al. 2015; Parke 2018; Baco et al. 2019; Kennedy et al. 2019; Kelley et al. 2022). These results are consistent with studies in other oceans (Watling and Auster 2017; Rogers 2018). Supporting this prediction, habitat suitability models show extremely high habitat suitability for octocorals and antipatharians on all of the NW Pacific seamounts (Yesson et al. 2012, 2017). Adjacent seamounts in the Wake Island Unit of the Pacific Remote Islands National Marine Monument host a high diversity of such corals (Kelley et al. 2022). Therefore, it is highly likely all the seamounts in the area contain vulnerable, fragile, and sensitive taxa with low resilience and slow recoveries to impacts.

Other discussions in the above sections are also relevant here. For example, rare or endemic species and geographically isolated populations, such as those on the NW Pacific seamounts, are more vulnerable to extirpation and extinction events than cosmopolitan species or populations on continental slopes (e.g., Clark et al. 2010). The species and communities on these NW Pacific seamounts are likely adapted to the extreme environmental and physical stability of the features. The CFC on the seamount slopes and the nodules on the seamount bases require millions of years of the right conditions to form (Hein and Koschinsky 2014) and, again, these are the oldest seamounts in the world. In deep-sea environments, where conditions have typically changed over geological timescales, the associated animals, adapted to these stable conditions, are expected to be highly vulnerable to any change or direct human impact, in particularly long-lived, habitat-forming deep-sea species (as summarized in Ross et al. 2020).

Biological productivity	Area containing species, populations or communities with comparatively higher natural biological productivity.	No info.	Low	Medium	High
					X

Justification

The ABNJ NW Pacific seamounts contain species, populations and communities with comparatively higher natural biological productivity relative to the surrounding open ocean and abyssal plains, and other seamounts in the global ocean. Therefore, the biological productivity of these seamounts is *High* (details are provided below).

Seamounts throughout the world’s oceans are associated with high biological productivity, particularly

compared to the surrounding open ocean and abyssal plains (Pitcher et al. 2007). While it is generally accepted that enhanced primary production occurs over shallow and intermediate depth seamounts (e.g., Dai et al. 2020), it has also been observed in situ over a deep NW Pacific seamount (Dai et al. 2022), in contrast to the otherwise surrounding oligotrophic body of water (Yamaoka et al. 2020). In addition, visual observations from these NW Pacific seamounts of dense and high biomass cold-water coral and sponge fields and mats of ophiuroids (Chen et al. 2021a; Shen et al. 2021; ISA 2022) are consistent with seamount-associated high biological productivity, as are the enhanced sedimentation rate at the summits of seamounts (relative to the surrounding abyssal plains), organic carbon flux, and bioturbation on and around the Marcus-Wake and Magellan seamounts (Yang et al. 2020; Ota et al. 2022).

While research on these NW Pacific seamounts is still in its infancy, drivers of increased biological productivity on and around the seamounts have already been documented and are consistent with the biological observations mentioned above (i.e., enhanced primary production, sedimentation, organic carbon flux, and biomass of suspension feeders, filter feeders, and deposit feeders on and around these NW Pacific seamounts). For example,

- topographic upward movement of nutrients results in enhanced primary productivity over shallow and deep NW Pacific seamount summits (Dai et al. 2020, 2022);
- the abundance of NW Pacific seamounts creates eddies, which have critical roles in transporting, concentrating, and trapping phytoplankton and zooplankton on and around the seamounts (Jiang et al. 2021);
- current jets around the seamounts can increase the sediment and detrital suspension, enhancing food supply for suspension feeders like deep-sea corals and sponges (Jiang et al. 2021);
- enhanced near-bottom circulation and mixing by surface eddies over the seamounts (as reported by Xie et al. 2022) is expected to also increase sediment and detrital suspension;
- the raised topographic features of hard substrate support dense cold-water corals and sponges benefiting from local current enhancement and food enrichment, which in turn create local biological hotspots (Shen et al. 2021);
- seamount productivity and nutrient-enriched water is transported downstream, potentially to adjacent seamounts, by eddies or other currents (Dai et al. 2022).

Other processes likely contributing to the enhanced productivity on and around seamounts include seamount-induced current accelerations, upwelling, turbulence, and mixing (Leitner et al., 2020 and references therein) and topographic trapping of daily vertical migration (particularly for the shallower seamounts; Genin et al., 1994; Taranto et al. 2012). The spatial and temporal variation of biological productivity on and around seamounts is a complex pattern driven by these processes and others (e.g., Dower et al., 1992; Comeau et al., 1995; Dower and Mackas, 1996; Martin and Christiansen, 2009).

The high concentration of migrating species and predator species attracted to these NW Pacific seamounts discussed in the sections above are also strong indicators of high biological productivity.

The unusual density and height of the NW Pacific seamount complex discussed throughout this assessment are also pertinent in the context of biological productivity. These physical attributes (i.e., seamount size) enhance the available seafloor area for seamount-associated fauna to occupy, setting this complex apart from most other seamount areas in the world, almost all of which are less dense and contain shorter seamounts.

Biological diversity	Area contains comparatively higher diversity of ecosystems, habitats, communities, or species, or has higher genetic diversity.	No info.	Low	Medium	High
					X

Justification

The ABNJ NW Pacific seamounts contain a comparatively higher diversity of ecosystems, habitats, communities, and species, relative to the surrounding open ocean and abyssal plains, and other seamounts in the global ocean. The high alpha and beta diversity are likely owing to, in large part, the enhanced productivity, currents (discussed in the above sections), habitat complexity, and habitat heterogeneity. Therefore, the biological diversity of these seamounts is *High* (details are provided below).

Despite the limited in situ research, seamounts with the ABNJ NW Pacific are known homes to diverse benthic communities (Chen et al. 2021a; Shen et al. 2021; ISA 2022). For example, more than 90 morpho-species were recorded on a single transect between 1100 and 1300 m depth (ISA 2022) and 29 species of ophiuroids in 11 families were recorded on a single seamount (Chen et al. 2021a). The assemblages of cold-water coral and sponge fields (including large gorgonian corals and erect sponges) were described by researchers as having high biodiversity and biomass, and were recorded to completely cover the complex topography of the seamount slope and ridges (ISA 2022). In general, cold-water corals and sponges support a high diversity of invertebrate fauna and fish (e.g., Husebø et al. 2002; Buhl-Mortenson et al. 2010), providing nursery grounds, shelter, sources of food etc. (e.g., Etnoyer and Warrenchuk 2007; Baillon et al. 2012; Cau et al. 2017). Fauna associated with these NW Pacific seamount cold-water coral and sponge habitats include commensal crinoids and ophiuroids, as well as fish, crustaceans, sea stars, sea cucumbers, and sea anemones (Xu et al. 2017; ISA 2022).

These NW Pacific seamounts provide heterogeneous habitats across a range of depths as they rise from the abyssal plains across ~5.5 vertical kilometers (ISA 2022). Based on the wide depth range, in combination with the different well-stratified oceanographic layers (e.g., the western North Pacific Subtropical gyre, North Pacific Intermediate Water, and the North Pacific Deep Water; Korff et al. 2016; Yamaoka et al. 2020), and the generally accepted high depth fidelity of seamount species, we can infer a high spatial turnover (beta diversity) of benthic species on each seamount (similarly to Long and Baco 2014; Schlacher et al. 2014; Du Preez et al. 2016; Victorero et al. 2018; Morgan et al. 2019; Mejia-Mercado et al. 2019). In addition, the variability of habitats within depth bands (e.g., complex and rocky in comparison to flat and muddy) would further enhance the diversity of each seamount (e.g., Morgan et al. 2019). Lastly, with the high natural variability of the 205 seamounts (e.g., seamount size, height, sedimentation, geomorphology, latitude, etc.; ISA 2022) bathed in different oceanographic conditions (ISA 2022), high biological beta diversity between seamounts is reasonable to infer. Examples of these species turnover patterns have already been recorded within these NW Pacific seamounts. Weijia Seamount supports a dense, diverse, distinct, and highly patchy pattern of megafaunal community (inclusive of cold-water corals and sponges) as a function of habitat heterogeneity variables, such as latitude, depth, slope, curvature, rugosity, bathymetric position index, topography type, and substrate type (e.g., Chen et al. 2021a; Shen et al. 2021).

Naturalness	Area with a comparatively higher degree of naturalness as a result of the lack of or low level of human-induced disturbance or degradation.	No info.	Low	Medium	High
					X

Justification

The ABNJ NW Pacific seamounts have a comparatively higher degree of naturalness as a result of the low level of human-induced disturbance and degradation relative to other seamounts (e.g., no formal bottom-contact fishing industry), but disturbance, in the form of seabed mining is under consideration in this location. Therefore, the naturalness of these seamounts is currently *High*, but that may change in the near future (details are provided below).

As summarized in ISA report (2022), these seamounts are relatively far removed from human population centers and, as a result, are isolated from many anthropogenic stressors that impact nearshore areas, such as runoff and chemical pollution. That said, the area is surrounded by fishing nations and there are pelagic fisheries over the seamounts (e.g., drifting longlines), as well as suspected illegal, unreported, and unregulated fishing (Welch et al. 2022). The removal and/or overfishing of large predator species (e.g., the tuna, billfish, and sharks fished over the seamounts; Morato et al. 2010; WCPFC 2019) can have cascading effects on ecosystem trophic structure and the environment itself (Scheffer et al. 2005). However, a survey of the Global Fish Watch database indicates very little bottom trawling effort is present (Automatic Identification System data; Kroodsma et al. 2018) —which has caused irreparable damage to the natural seafloor of many seamounts world-wide, including those in the other North Pacific seamounts EBSAs (CBD 2016a,b). No visual evidence of human activities (including mining exploration and exploitation, bottom trawl fishery, or submarine cable) was found on six surveyed seamounts in 2019 and 2020 during expeditions by COMRA (China Ocean Mineral Resources Research & Development Association) (reported in ISA 2022). No marine debris was observed on the one seamount surveyed by NOAA’s Office of Ocean Exploration and Research in 2016 (Amon et al. 2020). However, whether from future mining, increased fishing, or other stressors, biological communities on seamounts are highly vulnerable to human activities (Clark et al. 2010), and climate change is rapidly impacting the global ocean, including the deep (IPCC 2019).




Figure 2. Northwest (NW) Pacific Ocean Seamounts support abundant and diverse communities. (a-b) Cold-water corals and sponges on ferromanganese-encrusted lava. Threatened and Endangered species include the (c) Leatherback Sea Turtle (*Caretta caretta*), (d) Sperm Whale (*Physeter macrocephalus*), and (e) Short-tailed Albatross (*Phoebastria albatrus*). Overfished species such as the (f) Bluefin Tuna (*Thunnus orientalis*), (g) North Pacific Striped Marlin (*Tetrapturus audax*, also *Kajikia audax*), and (h) Oceanic Whitetip Shark (*Carcharhinus longimanus*). Photos courtesy of (a-b) NOAA Ocean Exploration (www.ncei.noaa.gov/waf/oceanos-rov-cruises/ex1606/), (c-g) iNaturalist (Azure 27014, Wayne and Pam Osborn, 范姜士豪, Tom Mills, Royle Safaris; www.inaturalist.org), (h) Shelton Du Preez and SeaLegacy.

Conclusion

The CBD (as well as the many States that use EBSA assessments, including Canada) identifies a marine feature as an EBSA if it scores High on at least one criterion or above average across all (DFO 2004, 2011; UNEP/CBD 2008). The NW Pacific seamount complex in ABNJ scores *High* for all EBSA criteria except ‘Special importance for life-history stages of species’ (ranked *Medium*) and can, therefore, be classified as an EBSA (Table 2). Notably, these NW Pacific seamounts hold exceptional distinction on a regional scale, in comparison to the surrounding abyssal plains and open ocean, and on a global scale, in comparison to other seamounts and seamount complexes in the world. They are also notable for their diverse habitats, which provide high habitat heterogeneity for a wide range of species, including rare shallow-water and rocky current-swept habitats for cold-water corals and sponges, creating dense and diverse biogenic habitats. The unusual oceanographic conditions, geomorphology, and remarkable natural stability have produced rare physical and biological seamount features, such as thick cobalt-rich ferromanganese crusts, geographically isolated deep-sea populations, and rare or endemic species. The seamounts are stepping stones that form pathways for the migration of species. Pelagic species—many of which are threatened and endangered—migrate through the area and are known to benefit from the ecosystem functions provided by the seamounts. These seamounts support rich, fragile, and vulnerable species and ecosystems—from their base to their summit, the water column, and above.

This NW Pacific Ocean Seamounts EBSA identification synthesizes the best available science and provides a robust assessment and thorough review to support future CBD workshops on the region and critical regional area-based management related to fishing, climate change, seabed mining, the new Biodiversity Beyond National Jurisdiction treaty, and other spatial management processes.

Table 2. Summary of Ecological or Biological Significant Area (EBSA) assessment of the Northwest Pacific seamounts in Areas Beyond National Jurisdiction (ABNJ) against the seven criteria. Possible rankings: *No information*, *Low*, *Medium*, and *High*.

CBD EBSA Criteria	Seamount Rating
Uniqueness or rarity	<i>High</i>
Special importance for life-history stages of species	<i>Medium</i>
Importance for threatened, endangered or declining species and/or habitats	<i>High</i>
Vulnerability, fragility, sensitivity, or slow recovery	<i>High</i>
Biological productivity	<i>High</i>
Biological diversity	<i>High</i>
Naturalness	<i>High</i>
EBSA Identification	Meets Criteria

Acknowledgements

The authors would like to express their gratitude to the following colleagues for their contributions. We value the oceanographic context provided by John Dower to improve the draft manuscript. We are appreciative to Jessica Nephin, Shannon Schmunk, and Tammy Norgard for their thoughtful review of the draft. This work greatly benefited from organizers and invited experts at the Workshop on the Development of a Regional Environmental Management Plan (REMP) for the Area of the Northwest Pacific (hosted by the International Seabed Authority, online, October 26th to November 6th 2020) and the Global Ocean Biodiversity Initiative EBSAs in ABNJ workshop (hosted by GOBI, in Santa Cruz, California, November 6th to 9th 2022).

Literature cited

- Amon, D.J., Kennedy, B.R.C., Cantwell, K., Suhre, K., Glickson, D., Shank, T.M., and R.D. Rotjan. 2020. [Deep-sea debris in the Central and Western Pacific Ocean](#). *Front. Mar. Sci.* 7: 369.
- Baco, A.R. 2007. [Exploration for deep-sea corals on the north Pacific seamounts and islands](#). *Oceanography* 20(4): 108-117.
- Baco, A.R., Roark, E.B., and Morgan, N.B. 2019. [Amid fields of rubble, scars, and lost gear, signs of recovery observed on seamounts on 30-to 40-year time scales](#). *Sci. Adv.* 5: eaaw4513.
- Baillon, S., Hamel, J.F., Wareham, V.E., and Mercier, A. 2012. [Deep cold-water corals as nurseries for fish larvae](#). *Front. Ecol. Environ.* 10: 351-356.
- Ban, S., Curtis, J.M.R., St. Germain, C., Perry, R.I., and Therriault, T.W. 2016. [Identification of Ecologically and Biologically Significant Areas \(EBSAs\) in Canada's Offshore Pacific Bioregion](#). DFO Can. Sci. Advis. Sec. Res. Doc. 2016/034. x + 152 p.
- Bell, K.L.C., J. Flanders, A. Bowman, and N.A. Raineault, eds. 2017. [New frontiers in ocean exploration: The E/V Nautilus, NOAA Ship Okeanos Explorer, and R/V Falkor 2016 field season](#). *Oceanography* 30(1), supplement, 94 pp.
- Benson, S.R., Eguchi, T., Foley, D.G., Forney, K.A., Bailey, H., Hitipeuw, C., Samber, B.P., Tapilatu, R.F., Rei, V., Ramohia, P., Pita, J., Dutton, P.H. 2011. [Large-scale movements and high-use areas of western Pacific leatherback turtles, *Dermochelys coriacea*](#). *Ecosphere* 2(7): 1-27.
- Block, B.A., Jonsen, I.D., Jorgensen, S.J., Winship, A.J., Shaffer, S.A., Bograd, S.J., Hazen, E.L., Foley, D.G., Breed, G.A., Harrison, A.L., and Ganong, J. E. [Tracking apex marine predator movements in a dynamic ocean](#). *Nature* 475: 86–90.
- Buenafe, K.C.V., Everett, J.D., Dunn, D.C., Mercer, J., Suthers, I.M., Schilling, H.T., Hinchliffe, C., Dabalà, A., and Richardson, A.J. 2022. [A global, historical database of tuna, billfish, and saury larval distributions](#). *Sci. Data* 9(1): 1-9.
- Buhl-Mortensen, L., Vanreusel, A., Gooday, A.J., Levin, L.A., Priede, I.G., Buhl-Mortensen, P., Gheerardyn, H., King, N.J., and Raea, M. 2010. [Biological structures as a source of habitat heterogeneity and biodiversity on the deep ocean margins](#). *Mar. Ecol.* 31: 21-50.
- Cau, A., Follasa, M.C., Moccia, D., Bellodi, A., Mulas, A., Bo, M., Canese, S., Angiolillo, M., and Cannas, R. 2017. [Leiopathes glaberrima millennial forest from SW Sardinia as nursery ground for the small spotted catshark *Scyliorhinus canicular*](#). *Aquatic Conserv: Mar. Freshw. Ecosyst.* 27: 731 – 735.
- CBD (Convention on Biological Diversity). 2016a. [Ecologically or Biologically Significant Areas \(EBSAs\): North Pacific Transition Zone](#). Accessed 2023-10-03.
- CBD (Convention on Biological Diversity). 2016b. [Ecologically or Biologically Significant Areas \(EBSAs\): Emperor Seamount Chain and the Northern Hawaiian Ridge](#). Accessed 2023-10-03.

- Cheung, W. W., Wei, C. L., and Levin, L. A. 2022. [Vulnerability of exploited deep-sea demersal species to ocean warming, deoxygenation, and acidification](#). *Environ. Biol. Fishes.* 105(10): 1301-1315.
- Chen, W., Na, J., Shen, C., Zhang, R., Lu, B., Cheng, H., Wang, C., and Zhang, D. 2021a. [Ophiuroid fauna of cobalt-rich crust seamounts in the Northwest Pacific Ocean](#). *Acta Oceanol. Sin.* 40:55-78.
- Chen, W., Na, J., and Zhang, D. 2021b. [Description of three species of ophioplithacids, including a new species, from a deep seamount in the Northwest Pacific Ocean](#). *PeerJ* 9: e11566.
- Cheng, L., Abraham, J., Trenberth, K.E., Fasullo, J., Boyer, T., Locarnini, R., Zhang, B., Yu, F., Wan, L., Chen, X., Song, X., Liu, Y., Mann, M.E., Reseghetti, F., Simoncelli, S., Gouretski, V., Chen, G., Mishonov, A., Reagan, J., and Zhu, J. 2021. [Upper Ocean Temperatures Hit Record High in 2020](#). *Adv. Atmos. Sci.* 38: 523–530 (2021).
- Christiansen, S., Durussel, C., Guilhon, M., Singh, P., and Unger, S. 2022. [Towards an ecosystem approach to management in areas beyond national jurisdiction: REMPs for deep seabed mining and the proposed BBNJ instrument](#). *Front. Mar. Sci.* 9: 720146.
- Clark, M.R., Rowden, A.A., Schlacher, T., Williams, A., Consalvey, M., Stocks, K.I., Rogers, A.D., O’Hara, T.D., White, M., Shank, T.M., and Hall-Spencer, J.M. 2010. [The ecology of seamounts: structure, function, and human impacts](#). *Ann Rev* 2: 253-278.
- Clouard, V., and Bonneville. A. 2005. [Ages of seamounts, islands and plateaus on the Pacific Plate](#). *Geol. Soc. Am. Bull.* 388: 71-90.
- Comeau, L.A., Vézina, A.F., Bourgeois, M., and Juniper, S.K. 1995. [Relationship between phytoplankton production and the physical structure of the water column near Cobb Seamount, northeast Pacific](#). *Deep Sea Res. Part I Oceanogr. Res. Pap.* 42: 993-1005.
- Dai, S., Zhao, Y., Li, X., Wang, Z., Zhu, M., Liang, J., Lui, H., Tian, Z., and Sun, X. 2020. [The seamount effect on phytoplankton in the tropical western Pacific](#). *Mar. Environ. Res.* 162: 105094.
- Dai, S., Zhao, Y., Li, X., Wang, Z., Zhu, M., Liang, J., Liu, H., and Sun, X. 2022. [Seamount effect on phytoplankton biomass and community above a deep seamount in the tropical western Pacific](#). *Mar. Pollut. Bull.* 175: 113354.
- DFO. 2004. [Identification of Ecologically and Biologically Significant Areas](#). DFO Can.Sci. Advis. Sec. Ecosystem Status Rep. 2004/006.
- DFO. 2011. [Ecologically and Biologically Significant Areas – Lessons Learned](#). DFO Can. Sci. Advis. Sec. Sci. Advis. Rep. 2011/049.
- Ding, X., Gao, L., Fang, N., Qu, W., Liu, J., and Li, J. 2009. [The relationship between the growth process of the ferromanganese crusts in the pacific seamount and Cenozoic ocean evolvement](#). *Sci. China Ser. D-Earth Sci.* 52: 1091–1103.
- Dong, D., Li, X. Z., Lu, B., and Wang, C.S. 2017. [Three squat lobsters \(Crustacea: Decapoda: Anomura\) from tropical West Pacific seamounts, with description of a new species of Uroptychus Henderson, 1888](#). *Zootaxa* 4311(3): 389-398.
- Dong, D., Xu, P., Li, X., Wang, C., 2019. [Munidopsis species \(Crustacea: Decapoda: Munidopsidae\) from carcass falls in Weijia Guyot, West Pacific, with recognition of a new species based on integrative taxonomy](#). *PeerJ* 7: e8089.
- DOSI (Deep-Ocean Stewardship Initiative). 2023. [“The Seamount Ecosystem” - Information Sheet](#). Accessed 2023-10-03.

- Dower, J.F., Freeland, H., and Juniper, S.K. 1992. [A strong biological response to oceanic flow past Cobb Seamount](#). Deep Sea Res. Part A Oceanogr. Res. Pap. 39: 1139-1145.
- Dower, J.F., and Mackas, D.L. 1996. [“Seamount effects” in the zooplankton community near Cobb Seamount](#). Deep Sea Res. Part I Oceanogr. Res. Pap. 43: 837-858.
- Drazen, J.C., Smith, C.R., Gjerde, K., Au, W., Black, J., Carter, G., Clark, M.R., Durden, J., Dutrieux, P., Goetze, E., Haddock, S., Hatta, M., Hauton, C., Hill, P., Koslow, J., Leitner, A., Measures, C., Pacini, A., Parrish, F., Peacock, T., Perelman, J., Sutton, T., Taymans, C., Tunnicliffe, V., Watling, L., Yamamoto, H., Young, E., Ziegler, A. 2019. [Report of the workshop Evaluating the nature of midwater mining plumes and their potential effects on midwater ecosystems](#). Res. Ideas Outcomes 5: e33527.
- Du Preez, C., Curtis, J.M., and Clarke, M.E. 2016. [The structure and distribution of benthic communities on a shallow seamount \(Cobb Seamount, Northeast Pacific Ocean\)](#). PLOS ONE 11(10): e0165513
- Etnoyer, P.J., and Warrenchuk, J. 2007. A catshark nursery in a deep gorgonian field in the Mississippi Canyon, Gulf of Mexico. Bull. Mar. Sci. 81(3): 553 – 559.
- FAO (Food and Agriculture Organization). 2023. [PACIFIC, WESTERN CENTRAL \(Major Fishing Area 71\)](#). Fisheries and Aquaculture Division [online]. Rome. Accessed 2023-10-03
- FAO. 2009b. [International Guidelines for the Management of Deep-sea Fisheries in the High Seas](#). Rome. 73p. Accessed 2023-10-03.
- Gan, Z., Xu, P., Li, X. and Wang, C. 2020. [Integrative taxonomy reveals two new species of stalked barnacle \(Cirripedia, Thoracica\) from seamounts of the Western Pacific with a review of barnacles distributed in seamounts world-wide](#). Front. Mar. Sci. 7: 582225.
- Genin, A., Greene, C., Haury, L., Wiebe, P., Gal, G., Kaartvedt, S., Meir, E., Fey, C., and Dawson, J. 1994. [Zooplankton patch dynamics: daily gap formation over abrupt topography](#). Deep Sea Res. Part I Oceanogr. Res. Pap. 41: 941-951.
- Glasby, G.P., Ren, X., Shi, X., and Pulyaeva. 2007. [Co-rich Mn crusts from the Magellan Seamount cluster: the long journey through time](#). Geo-Mar. Lett. 27: 315–323.
- GOBI (Global Ocean Biodiversity Initiative) Secretariat. 2023. [Report of the Global Ocean Biodiversity Initiative \(GOBI\) international workshop on Ecologically or Biologically Significant Marine Areas \(EBSAs\) in Areas Beyond National Jurisdiction \(ABNJ\): 6-9 November 2022, Santa Cruz, California](#). 15 pages. Accessed 2023-10-03.
- Gollner, S., Kaiser, S., Menzel, L., Jones, D.O., Brown, A., Mestre, N.C., Van Oevelen, D., Menot, L., Colaço, A., Canals, M., *et al.* 2017. [Resilience of benthic deep-sea fauna to mining activities](#). Mar. Environ. Res. 129: 76-101.
- Grischenko, A. V. 2022. [New deep-water Bryozoa from the Magellan Seamounts, tropical western Pacific, with a global review of bryozoans associated with ferromanganese crusts](#). Prog. Oceanogr. 205: 102827.
- Harrison, A.L., Woodard, P.F., Mallory, M.L., and Rausch, J. 2022. [Sympatrically breeding congeneric seabirds \(*Stercorarius* spp.\) from Arctic Canada migrate to four oceans](#). Ecol. Evol. 12(1): e8451.
- Hein, J.R., and Koschinsky, A. 2014. [Deep-ocean ferromanganese crusts and nodules](#). In Treatise on Geochemistry, edited by H.D. Holland and K.K. Turekian 2nd Edition. Elsevier: 273-291.
- Hein, J.R., Mizell, K., Koschinsky, A., and Conrad, T.A. 2013. [Deep-ocean mineral deposits as a source of critical metals for high- and green-technology applications: Comparison with land-based resources](#). Ore Geol. Rev. 51: 1-14.

- Hestetun, J.T., Rapp, H.T. and Pomponi, S. 2019. [Deep-sea carnivorous sponges from the Mariana Islands](#). *Fron. Mar. Sci.* 6: 371.
- Husebø, Å., Nøttestad, L., Fosså, J. H., Furevik, D. M., and Jørgensen, S. B. 2002. [Distribution and abundance of fish in deep-sea coral habitats](#). In *biology of cold water corals*, edited by L. Watling and M. Risk. *Hydrobiologia* 471: 91-99.
- IPCC (Intergovernmental Panel on Climate Change). 2019. [Special report of the ocean and cryosphere in a changing climate](#). Accessed 2023-10-03.
- ISA (International Seabed Authority). 2022. [Workshop on the development of a regional environmental management plan for the area of the Northwest Pacific](#). Accessed 2023-10-03.
- IUCN (International Union for Conservation of Nature). 2023. [The IUCN Red List of Threatened Species](#). Version 2022-2. Accessed 2023-10-03
- IWC (International Whaling Commission). 2020. [Whale species](#). Accessed 2023-10-03.
- Jiang, X., Dong, C., Ji, Y., Wang, C., Shu, Y., Liu, L., and Ji, J. 2021. [Influences of deep-water seamounts on the hydrodynamic environment in the Northwestern Pacific Ocean](#). *J. Geophys. Res. Oceans* 126: e2021JC017396.
- Kelley, C.D., Bingo, S., Putts, M., Moriwake, V., Hourigan, T.F., and Cairns, S.D. 2022. [Preliminary List of Deep-Sea Coral Taxa in the Wake Island Unit of the Pacific Remote Islands Marine National Monument \(v. 2021\). Online Supplement to The state of deep-sea coral and sponge ecosystems of the United States](#). Accessed 2023-10-03.
- Kendall, M.S., and Poti, M. 2014. [Potential larval sources, destinations, and self-seeding in the Mariana Archipelago documented using ocean drifters](#). *J. Oceanogr.* 70: 549-557.
- Kennedy, B.R., Cantwell, K., Malik, M., Kelley, C., Potter, J., Elliott, K., Lobecker, E., McKenna Gray, L., Sowers, D., White, M.P., France, S.C., Auscavitch, S., Mah, C., Moriwake, V., Bingo, S.R.D., Putts, M., and Rotjan, R.D. 2019. [The unknown and the unexplored: insights into the pacific deep-sea following NOAA CAPSTONE expeditions](#). *Front. Mar. Sci.* 6: 480.
- Kitchingman, A., Lai, S., Morato, T., and Pauly, D. 2007. [How many seamounts are there and where are they located?](#) In *Seamounts: Ecology, Fisheries & Conservation*, edited by T.J. Pitcher, T. Morato, P.J.B. Hart, M.R. Clark, N. Haggan and R.S. Santos. Oxford: Blackwell Publishing: 26-40.
- Korff, L., von Dobeneck, T., Frederichs, T., Kasten, S., Kuhn, G., Gersonde, R., and Diekmann, B. 2016. [Cyclic magnetite dissolution in Pleistocene sediments of the abyssal northwest Pacific Ocean: Evidence for glacial oxygen depletion and carbon trapping](#). *Paleoceanography* 31: 600–624.
- Koslow, J. A. 1996. [Energetic and life-history patterns of deep-sea benthic, benthopelagic and seamount-associated fish](#). *J. Fish Biol.* 49: 54-74.
- Kou, Q., Xu, P., Poore, G.C.B., Li, X., and Wang, C. 2020. [A new species of the deep-sea sponge-associated Genus *Eiconaxius* \(Crustacea: Decapoda: Axiidae\), with new insights into the distribution, speciation, and mitogenomic phylogeny of axiidean shrimps](#). *Front. Mar. Sci.* 7: 469.
- Kroodsma, D.A., Mayorga, J., Hochberg, T., Miller, N.A., Boerder, K., Ferretti, F., Wilson, A., Bergman, B., White, T.D., and Block, B. A. 2018. [Tracking the global footprint of fisheries](#). *Science* 359: 904-908.
- Leitner, A.B., Neuheimer, A.B., and Drazen, J.C. 2020. [Evidence for long-term seamount-induced chlorophyll enhancements](#). *Sci. Rep.* 10: 1-10.
- Levin, L.A., Mengerink, K., Gjerde, K.M., Rowden, A.A., Van Dover, C.L., Clark, M.R., Ramirez-Llodra, E., Currie, B., Smith, C.R., Sato, K.N., Gallo, N., Sweetman, A.K., Lily, H., Armstrong, C.W.,

- and Brider, J. 2016. [Defining “serious harm” to the marine environment in the context of deep-seabed mining](#). Mar. Policy, 74: 245-259.
- Levin, L.A., Wei, C.L., Dunn, D.C., Amon, D.J., Ashford, O.S., Cheung, W.W., Colaço, A., Dominguez-Carrió, C., Escobar, E.G., Harden-Davies, H.R., and Drazen, J.C. 2020. [Climate change considerations are fundamental to management of deep-sea resource extraction](#). Glob. Change Biol. 26(9): 4664-4678.
- Liu, D., and Hoskin, M. 2023. [Contemporary international Law: Regulating the upcoming fukushima radioactive wastewater discharge](#). Ocean Coast. Manag. 234: 106452.
- Løbach, T., Petersson, M., Haberkon, E., and Mannini, P. 2020. [Regional fisheries management organizations and advisory bodies: Activities and developments, 2000–2017](#). FAO Fish. Aqua. Tech. Pap. 651: 1-112.
- Long, D.L., and Baco, A.R. 2014. [Rapid change with depth in megabenthic structure-forming communities of the Makapu’u deep-sea coral bed](#). Deep Sea Res. Part II Top. Stud.Oceanogr. 99: 158 – 168.
- Martin, B., and Christiansen, B. 2009. [Distribution of zooplankton biomass at three seamounts in the NE Atlantic](#). Deep Sea Res. Part II Top. Stud.Oceanogr. 56: 2671-2682.
- Mejia-Mercado, B.E., Mundy, B., and Baco, A.R. 2019. [Variation in the structure of deep-sea fish assemblages in Necker Island, Northwestern Hawaiian Islands](#). Deep Sea Res. Part I Oceanogr. Res. Pap. 152: 103086.
- Mel’nikov, M., Pletnev, S., Anokhin, V., Sedysheva, T., and Ivanov, V. 2016. [Volcanic edifices on guyots of the Magellan Seamounts \(Pacific Ocean\)](#). Russ. J. of Pac. Geol. 10: 435-442.
- Miller, K.J., and Gunasekera, R.M. 2017. [A comparison of genetic connectivity in two deep sea corals to examine whether seamounts are isolated islands or stepping stones for dispersal](#). Sci. Rep. 7: 1-14.
- Morato, T., Hoyle, S.D., Allain, V., and Nicol, S.J. 2010. [Seamounts are hotspots of pelagic biodiversity in the open ocean](#). PNAS 107(21): 9707-9711.
- Morgan, N.B., Goode, S., Roard, E.B., and Baco, A.R. 2019. [Fine scale assemblage structure of benthic invertebrate megafauna on the North Pacific seamount Mokumanamana](#). Front. Mar. Sci. 6: 715.
- Müller, R. D., Seton, M., Zahirovic, S., Williams, S. E., Matthews, K. J., Wright, N. M., Shephard, G., Maloney, K. T., Barnett-Moore, N., Hosseinpour, M., Bower, D. J., and Cannon, J. 2016. [Ocean basin evolution and global-scale plate reorganization events since Pangea breakup](#). Annu. Rev. Earth Planet. Sci. 44(1): 107-138.
- Murray, C.C., Agbayani, S., Alidina, H.M., and Ban, N.C. 2015. [Advancing marine cumulative effects mapping: An update in Canada’s Pacific waters](#). Mar. Policy 58: 71-77.
- Na, J., Chen, W., Zhang, D., Zhang, R., Lu, B., Shen, C., Zhou, Y., and Wang, C. 2021. [Morphological description and population structure of an ophiuroid species from cobalt-rich crust seamounts in the Northwest Pacific: implications for marine protection under deep-sea mining](#). Acta Oceanol. Sin. 40: 79-89.
- Ota, Y., Suzumura, M., Tsukasaki, A., Suzuki, A., Seike, K., and Minatoya, J. 2022. [Sediment accumulation rates and particle mixing at northwestern Pacific seamounts](#). J. Mar. Syst. 229: 103719.
- Parke, M. 2018. [Deep-sea coral research and technology program: Pacific Islands deep-sea coral and sponge 3-year research wrap-up workshop May 23–24, 2018](#). NOAA Tech Memo. NMFS-PIFSC-78, 30 p.

- Parker, T., and Tunnicliffe, V. 1994. [Dispersal strategies of the biota on an oceanic seamount: implications for ecology and biogeography](#). Biol. Bull. 187(3): 336-345.
- Parrish, F.A., Baco, A.R., Kelley, C., and Reiswig, H. 2015. State of Deep-Sea Coral and Sponge Ecosystems of the U.S. Pacific Islands Region. In The State of Deep-Sea Coral and Sponge Ecosystems of the United States, edited by T.F. Hourigan, P.J. Etnoyer and S. D. Cairns. Silver Spring, MD: NOAA Technical Memorandum.
- Pitcher, T.J., and Bulman, C. 2007. [Raiding the larder: a quantitative evaluation framework and trophic signature for seamount food webs](#). In Seamounts: Ecology, Fisheries & Conservation, edited by T. J. Pitcher, T. Morato, P.J.B. Hart, M.R. Clark, N. Haggan and R.S. Santos. Oxford: Blackwell Publishing: 282-295.
- Pitcher, T.J., Morato, T., Hart, P.J., Clark, M.R., Haggan, N., and Santos, R.S. 2007. [The depths of ignorance: an ecosystem evaluation framework for seamount ecology, fisheries and conservation](#). In Seamounts: ecology, fisheries, and conservation. Blackwell Fisheries and Aquatic Resources Series. Edited by T.J. Pitcher, T. Morato, P.J.B. Hart, M.R. Clark, N. Haggan and R.S. Santos. Blackwell Publishing, Oxford. pp. 476-488.
- Rowden, A.A., Dower, J.F., Schlacher, T.A., Consalvey, M., and Clark, M.R. 2010. [Paradigms in seamount ecology: fact, fiction and future](#). Mar. Ecol. 31: 226-241.
- Rogers, A.D. 2018. [The biology of seamounts: 25 years on](#). Adv. Mar.Biol. 79: 137-224.
- Ross, T., Du Preez, C., and Ianson, D. 2020. [Rapid deep ocean deoxygenation and acidification threaten life on Northeast Pacific seamounts](#). Global Change Biology 26(11): 6424-6444.
- Scheffer, M., Carpenter, S., and de Young, B. 2005. [Cascading effects of overfishing marine systems](#). Trends Ecol. Evol. 20(11): 579-581.
- Schlacher, T.A., Baco, A.R., Rowden, A.A., O'Hara, T.D., Clark, M.R., Kelley, C., and Dower, J.F. 2014. [Seamount benthos in a cobalt-rich crust region of the central Pacific: Conservation challenges for future seabed mining](#). Diversity Distrib. 20: 491-502.
- Shank, T.M. 2010. [Seamounts: deep-ocean laboratories of faunal connectivity, evolution, and endemism](#). Oceanography 23(1): 108-122.
- Shen, C., Zhang, D., Lu, B., and Wang, C. 2020. [A new species of glass sponge \(Hexactinellida: sceptrulophora: uncinateridae\) from the Weijia seamount in the northwestern Pacific Ocean](#). Zootaxa 4878 (2): 322–334.
- Shen, C., Lu, B., Li, Z., Zhang, R., Chen, W., Xu, P., Yao, H., Chen, Z., Pang, J., Wang, C., and Zhang, D. 2021. [Community structure of benthic megafauna on a seamount with cobalt-rich ferromanganese crusts in the northwestern Pacific Ocean](#). Deep Sea Res. Part I Oceanogr. Res. Pap. 178: 103661.
- Stewart, H.A., and Jamieson, A.J. 2018. [Habitat heterogeneity of hadal trenches: Considerations and implications for future studies](#). Prog. Oceanogr. 161: 47– 65.
- Taranto, G. H., Kvile, K. Ø., Pitcher, T. J., and Morato, T. 2012. [An ecosystem evaluation framework for global seamount conservation and management](#). PLoS ONE, 7(8): e42950.
- Tittensor, D.P., Baco, A.R., Hall-Spencer, J.M., Orr, J.C., and Rogers, A.D. 2010. [Seamounts as refugia from ocean acidification for cold-water stony corals](#). Mar. Ecol. 31: 212-225.
- UNEP (United Nations Environment Program)/CBD. 2008. Decision adopted by the conference of the parties to the convention on biological diversity at its ninth meeting. [IX/20 Marine and coastal biodiversity: Annex II](#). Accessed 20223-10-03.

- UNESCO (United Nations Educational, Scientific and Cultural Organization). 2009. [Global Open Oceans and Deep Seabed \(GOODS\): biogeographic classification](#). IOC Technical Series 84: 96 p.
- Victorero, L., Robert, K., Robinson, L.F., Taylor, M.L., and Huvenne, V.A. 2018. [Species replacement dominates megabenthos beta diversity in a remote seamount setting](#). Scientific Reports 8: 4152.
- Walbridge, S., Slocum, N., Pobuda, M., and Wright, D.J. 2018. [Unified geomorphological analysis workflows with benthic terrain modeler](#). Geosci. 8(3): 94.
- Wang, D., Wang, C., Zhang, Y., and Liu, C. 2016. [Three new species of glass sponges Phoronematidae \(Porifera: Hexactinellida\) from the deep-sea of the northwestern Pacific Ocean](#). Zootaxa 4171 (3): 562–574.
- Watling, L., and Auster, P.J. 2017. [Seamounts on the high seas should be managed as vulnerable marine ecosystems](#). Front. Mar. Sci. 4: 14.
- Watling, L., Guinotte, J., Clark, M.R., and Smith, C.R. 2013. [A proposed biogeography of the deep ocean floor](#). Prog. Oceanogr. 111: 91-112.
- WCPFC (Western and Central Pacific Fisheries Commission). 2019. [Overview of Stock Status of Interest to the WCPFC](#). Accessed 2023-10-03.
- Welch, H., Clavelle, T., White, T.D., Cimino, M.A., Van Osdel, J., Hochberg, T., Kroodsmas, D., and Hazen, E.L. 2022. [Hot spots of unseen fishing vessels](#). Sci. Adv. 8: eabb12109.
- Xie, X., Wang, Y., Liu, X., Wang, J., Xu, D., Liu, T., Ji, J., Zhang, D., Wang, C., and Chen, D. 2022. [Enhanced near-bottom circulation and mixing driven by the surface eddies over abyssal seamounts](#). Prog. Oceanogr. 208: 102896.
- Xu, P., Zhou, Y., and Wang, C. 2017. [A new species of deep-sea sponge-associated shrimp from the North-West Pacific \(Decapoda, Stenopodidea, Spongicolidae\)](#). ZooKeys 685: 1-14.
- Yamaoka, K., Suzuki, A., Tanaka, Y., Suzumura, M., Tsukasaki, A., Shimamoto, A., Fukuhara, T., Matsui, T., Kato, S., Okamoto, N., and Igarashi, Y. 2020. [Late summer peak and scavenging-dominant metal fluxes in particulate export near a seamount in the western North Pacific subtropical gyre](#). Front. Earth Sci. 8: 558823.
- Yang, Z., Qian, Q., Chen, M., Zhang, R., Yang, W., Zheng, M., and Qiu, Y. 2020. [Enhanced but highly variable bioturbation around seamounts in the northwest Pacific](#). Deep Sea Res. Part I Oceanogr. Res. Pap. 156: 103190.
- Yesson, C., Clark, M.R., Taylor, M.L., and Rogers, A. D. 2011. [The global distribution of seamounts based on 30 arc seconds bathymetry data](#). Deep Sea Res. Part I Oceanogr. Res. Pap. 58(4), 442-453.
- Yesson C., Taylor, M.L., Tittensor, D.P., Davies, A.J., Guinotte, J., Baco, A., Black, J., Hall-Spencer, J.M., and Rogers, A.D. 2012. [Global habitat suitability of cold-water octocorals](#). J.Biogeogr. 39(7): 1278–1292.
- Yesson, C., Bedford, F., Rogers, A.D., and Taylor, M.L. 2017. [The global distribution of deep-water Antipatharian habitat](#). Deep Sea Res. Part II Top. Stud. Oceanogr. 145: 79 – 86.
- Zhang, R., Wang, C., Zhou, Y., and Zhang, H. 2019. [Morphology and molecular phylogeny of two new species in genus Freyastera \(Asteroidea: Brisingida: Freyellidae\), with a revised key to close species and ecological remarks](#). Deep Sea Res. Part I Oceanogr. Res. Pap. 154: 103163.
- Zhang, R., Zhou, Y., Xiao, N., and Wang, C. 2020. [A new sponge-associated starfish, *Astrolirus patricki* sp. nov. \(Asteroidea: brisingida: Brisingidae\), from the northwestern Pacific seamounts](#). PeerJ 8: e9071.