

MAP NUMBER

Natural Resources Canada, Geological Survey of Canada
Canadian Geoscience Map 459

Canada-Nunavut Geoscience Office
Open File Map 2023-05

TITLE

Bedrock geology, Armit Lake Area, Kivalliq, Nunavut, NTS 56-B and 56-C east

SCALE

1:150 000

CATALOGUE INFORMATION

Catalogue No. M183-1/459-2023E-PDF
ISBN 978-0-660-48721-2
<https://doi.org/10.4095/331889>

COPYRIGHT

© His Majesty the King in Right of Canada, as represented by the Minister of Natural Resources, 2023

Information contained in this publication or product may be reproduced, in part or in whole, and by any means, for personal or public non-commercial purposes, without charge or further permission, unless otherwise specified.

You are asked to:

- exercise due diligence in ensuring the accuracy of the materials reproduced;
- indicate the complete title of the materials reproduced, and the name of the author organization; and
- indicate that the reproduction is a copy of an official work that is published by Natural Resources Canada (NRCan) and that the reproduction has not been produced in affiliation with, or with the endorsement of, NRCan.

Commercial reproduction and distribution is prohibited except with written permission from NRCan. For more information, contact NRCan at nrcan.copyrightdroitdauteur.nrcan@canada.ca.

RECOMMENDED CITATION

Steenkamp, H.M., Wodicka, N., Lawley, C.J.M., Peterson, T., Weller, O.M., Kendrick, J., and Tschirhart, V., 2023. Bedrock geology, Armit Lake area, Kivalliq, Nunavut, NTS 56-B and 56-C east; Geological Survey of Canada, Canadian Geoscience Map 459; Canada-Nunavut Geoscience Office, Open File Map 2023-05, scale 1:150 000.
<https://doi.org/10.4095/331889>

ABSTRACT

New geological mapping in the Tehery Lake–Wager Bay area of northwestern Hudson Bay, Nunavut, frames the emplacement, depositional, and metamorphic histories of the dominant rock types, major structures, and links to neighbouring areas of the central Rae Craton and Chesterfield Block. The area is divided into six domains (Gordon, Lunan, and Kummel Lake domains presented here, and Ukkusiksalik and Douglas Harbour domains and Daly Bay Complex on adjoining maps) defined by large-scale structures and characterized by differing metamorphic assemblages, Sm-Nd and U-Pb isotopic data, and/or specific lithologies. Meso- to Neoarchean granitoid rocks underlie most of the area and are tectonically intercalated with Archean (volcano)sedimentary packages (Kummel Lake, Lorillard, and Paliak belts). These rocks are locally intruded by ca. 2.62 to 2.58 Ga Snow Island suite granite and cut by younger, thin, east-trending diabase dykes. Paleoproterozoic (volcano)sedimentary rocks are preserved in the Kingmirit belt (Daly Bay Complex) and in basement-cover infolds of Ketyet River group-equivalent strata (Douglas Harbour and Ukkusiksalik domains). In the south, the Daly Bay Complex (comprising mostly mafic granulite-facies rocks) and Kummel Lake Domain (a granulite-grade core complex) share some characteristics with rocks of the Kramanituar and Uvauk complexes, which may delineate the northeastern segment of the ca. 1.90 Ga Snowbird tectonic zone. The Paleoproterozoic Trans-Hudson Orogeny had widespread, penetrative structural and metamorphic effects on the area, and led to the intrusion of the ca. 1.85 to 1.81 Ga Hudson suite monzogranite and mafic ultrapotassic rocks, and ca. 1.83 Ga monzodiorite in the Ukkusiksalik and Douglas Harbour domains. The area is cut by large, southeast-trending gabbro dykes of the 1.267 Ga Mackenzie igneous event.

RÉSUMÉ

La nouvelle cartographie géologique dans la région de lac Tehery-baie Wager, au nord-ouest de la baie d'Hudson, au Nunavut, encadre l'histoire de la mise en place, du dépôt et du métamorphisme des types de roches dominants, les principales structures et les liens avec les régions avoisinantes de la partie centrale du craton de Rae et du bloc de Chesterfield. La région est divisée en six domaines (domaines de Gordon, de Lunan et de Kummel Lake présentés ici, ainsi que domaines d'Ukkusiksalik et de Douglas Harbour et complexe de Daly Bay sur les cartes adjacentes) définis par des structures à grande échelle et caractérisés par des assemblages métamorphiques, des données isotopiques Sm-Nd et U-Pb et/ou des lithologies spécifiques différents. Des roches granitoïdes du Mésoarchéen au Néoarchéen couvrent la majeure partie de la région et sont tectoniquement intercalées avec des assemblages (volcano)sédimentaires de l'Archéen (ceintures de Kummel Lake, de Lorillard et de Paliak). Ces roches sont localement injectées par des intrusions de granite de la suite de Snow Island datant d'environ 2,62 à 2,58 Ga et recoupées par des dykes de diabase plus jeunes, minces et de direction est. Des roches (volcano)sédimentaires du Paléoprotérozoïque sont conservées dans la ceinture de Kingmirit (complexe de Daly Bay) et dans des replis de socle-couverture de strates équivalentes au groupe de Ketyet River (domaines de Douglas Harbour et d'Ukkusiksalik). Au sud, le complexe de Daly Bay (composé principalement de roches mafiques du faciès des granulites) et le domaine de Kummel Lake (un complexe à noyau métamorphique du faciès des granulites) partagent certaines caractéristiques avec les roches des complexes de Kramanituar et d'Uvauk, qui pourraient délimiter le segment nord-est de la zone tectonique de Snowbird d'environ 1,90 Ga. L'orogénèse trans-

hudsonienne du Paléoprotérozoïque a eu des effets structuraux et métamorphiques étendus et pénétrants sur la région et a conduit à l'intrusion de monzogranite de la suite d'Hudson d'environ 1,85 à 1,81 Ga et de roches ultrapotassiques mafiques, ainsi que de monzodiorite d'environ 1,83 Ga dans les domaines d'Ukkusiksalik et de Douglas Harbour. La région est recoupée par de grands dykes de gabbro de direction sud-est, issus de l'événement igné de Mackenzie (1,267 Ga).

LICENCE AGREEMENT

View the licence agreement at

<https://open.canada.ca/en/open-government-licence-canada>

ACCORD DE LICENCE

Voir l'accord de licence à

<https://ouvert.canada.ca/fr/licence-du-gouvernement-ouvert-canada>

SHEET 1 OF 1, BEDROCK GEOLOGY

GENERAL INFORMATION

Authors: H.M. Steenkamp, N. Wodicka, C.J.M. Lawley, T. Peterson, O.M. Weller, J. Kendrick, and V. Tschirhart

Geology by H.M. Steenkamp (2015 to 2017), L. Lebeau (2016), A. Lion (2016) and I. Therriault (2017), Canada-Nunavut Geoscience Office; N. Wodicka (2012, 2015 to 2016), D.A. Kellett (2012), J.B. Whalen (2012), C.J.M. Lawley (2015), T. Peterson (2015), J. Beales (2015), W. Garrison (2015 to 2016), J. Kendrick (2016), O.M. Weller (2016), V. Tschirhart (2016), Geological Survey of Canada; K. Hatogina (2015) and R. Bayne (2016), Government of Nunavut; C. Guilmette (2015 and 2017), Université Laval; K.P. Larson (2017), University of British Columbia Okanagan

Geological compilation by H.M. Steenkamp, 2022

Geological data conforms to Bedrock Data Model v. 2.10 (Brouillette et al., 2019).

Geomatics by A. Morin

Cartography by N. Côté

Scientific editing by A. Weatherston

Joint initiative of the Geological Survey of Canada and Canada-Nunavut Geoscience Office, conducted under the auspices of the Rae Province project, Tehery-Wager Bay activity as part of Natural Resources Canada's Geo-mapping for Energy and Minerals (GEM) program and CanNor's Strategic Investments in Economic Development program

Logistical support provided by the Polar Continental Shelf Program (PCSP) as part of its mandate to promote scientific research in the Canadian north, PCSP 053-15, 342-15, 059-16, and 063-17

Map projection Universal Transverse Mercator, zone 15
North American Datum 1983

Base map at the scale of 1:250 000 from Natural Resources Canada, with modifications
Elevations in metres above mean sea level

Shaded-relief image derived from the digital elevation model supplied by Natural Resources Canada

Illumination: azimuth 315°, altitude 45°, vertical factor 5x

Mean magnetic declination 2023, 7°21'W, decreasing 15.4' annually

Readings vary from 9°38'W in the NE corner to 5°07'W in the SW corner of the map.

This map is not to be used for navigational purposes.

Title photograph: Aerial view of a mafic granulite body (unit Agab) near Lunan Lake, Nunavut (photo location 7). Photograph by O.M. Weller. NRCan photo 2022-259

The Geological Survey of Canada welcomes corrections or additional information from users (gscpublications-cgcpublishations@nrcan-rncan.gc.ca).

Data may include additional observations not portrayed on this map. See map info document accompanying the downloaded data for more information about this publication.

This publication is available for free download through GEOSCAN (<https://geoscan.nrcan.gc.ca/>) and Canada-Nunavut Geoscience Office (<https://cngo.ca/>).

MAP VIEWING FILES

The published map is distributed as a Portable Document File (PDF), and may contain a subset of the overall geological data for legibility reasons at the publication scale.

CARTOGRAPHIC REPRESENTATIONS USED ON MAP

This map utilizes ESRI Cartographic Representations in order to customize the display of standard GSC symbols for visual clarity on the PDF of the map only. The digital data still contains the original symbol from the standard GSC symbol set. The following legend features have Cartographic Representations applied:

- Geolines thin lithology
- Depositional, intrusive, or sheared contacts
- Thrust fault, inferred (teeth indicate hanging-wall side)
- Generic fault
- Detachment fault, extensional, approximate
- Shear, generic, steep dip, approximate

- Fold, synform, approximate, upright
- Fold, synform, inferred, upright, first generation
- Fold, antiform, approximate, upright:
- Fold, recumbent, approximate, second generation:
- Structural form line
- Geochronology location
- Photo location

DEFINITION QUERIES USED ON MAP

This map utilizes definition queries in order to customize the display for visualization on the PDF of the map only and does not affect the digital data. The following features have a definition query applied:

- Stations
- Geopoints - planar
- Geopoints - linear
- Geolines - shear
- Geolines - thin lithology
- Geolines - fault
- Geolines - overprint
- Geolines - unit construct line
- Geolines - contact
- Geolines - fold
- Geolines - trace
- Map Units - Overprints

DESCRIPTIVE NOTES

The Tehery-Wager mapping activity was a collaborative, multidisciplinary project led by the Canada-Nunavut Geoscience Office and the Geological Survey of Canada (GSC) through Natural Resources Canada's Geo-mapping for Energy and Minerals (GEM-2) program, and was developed following a two-week reconnaissance survey led by the GSC in the area in 2012. Bedrock geology, surficial geology, lake- and stream-sediment sampling, and ground-gravity transects were conducted with thematic scientific contributions from researchers at the University of British Columbia Okanagan, Dalhousie University, Université du Québec à Montréal, Université Laval, and University of New Brunswick. The aim of this activity was to increase the level of geological knowledge across the area from Tehery Lake to Wager Bay, Nunavut (NTS 56-A, 56-B, and parts of 46-D, 46-E, 56-C, 56-F, 56-G, and 56-H), in addition to evaluating the potential for a variety of commodities including base and precious metals, industrial minerals, diamonds and other gemstones, carving stone, and aggregate deposits (Steenkamp et al., 2015, 2016; Wodicka et al., 2015, 2016, 2017a). The project also aimed to assist northerners by providing geoscience training to college students in the field and temporary employment to locals as camp attendants and wildlife monitors, producing topographic maps that feature traditional Inuktitut place names (Peplinski et al., 2017), and ensuring that new geoscience information is accessible for future land-use decisions.

The 1:150 000 scale bedrock mapping component of the Tehery-Wager activity included three field seasons (2015–2017) and used a helicopter-supported, targeted approach for traverse set-outs based on airborne geophysical data (Geological Survey of Canada, 1978, 1998; Keating et al., 2003; Coyle and Kiss, 2012a–h), satellite imagery,

and historical information from previous mapping campaigns. The first reconnaissance-level geological surveys of the study area were Operation Baker (Wright, 1955; Wright, 1967; Lord and Wright, 1967) and Operation Wager (Heywood, 1967a, b). More detailed geological assessments were later conducted in the Daly Bay area (Gordon and Heywood, 1987; Gordon, 1988; Hanmer and Williams, 2001) and along the coastline of Wager Bay (Henderson et al., 1986; Henderson and Broome, 1990; Henderson and Roddick, 1990; Henderson et al., 1991). Data from those campaigns were digitized and used to inform the geological interpretations across the Tehery Lake–Wager Bay area that are presented here. The geological interpretations also take into consideration the data and observations documented in previous geochronological (Henderson and Roddick, 1990; van Breemen et al., 2007; Therriault, 2019; Therriault et al., 2018, 2022), geophysical (Spratt et al., 2014; Tschirhart et al., 2016; Wodicka et al., 2017a), and geochemical (Peterson et al., 2002; Peterson et al., 2015; Steenkamp et al., 2015) studies that relate to the rock units found within this area.

GEOLOGICAL BACKGROUND

The Tehery Lake–Wager Bay area is situated in the central part of the Rae Craton, which is a major constituent piece of the western Churchill Province of Laurentia. The Rae Craton comprises largely Meso- to Neoproterozoic and Paleoproterozoic metaplutonic and metasedimentary rocks cut by Mesoproterozoic dykes. Greenschist- to amphibolite-facies meta(volcano)sedimentary rocks in the central part of the craton include the 2.74 to 2.68 Ga Woodburn Lake group (Zaleski et al., 2000; Pehrsson et al., 2013), 2.73 to 2.69 Ga Committee Bay belt (Skulski et al., 2003a, b; Sanborn-Barrie et al., 2014), and 2.97 Ga Prince Albert and ≥ 2.765 to 2.75 Ga Roche Bay belts (Wodicka et al., 2011; Corrigan et al., 2013; Richan et al., 2015). The Chesterfield Block to the south of the Rae Craton also comprises Archean greenstone belts, namely the 2.74 to 2.64 Ga Yathkyed, Angikuni, and MacQuoid belt segments, and the northern Rankin Inlet belt (Davis et al., 2006; Hanmer et al., 2006; Sandeman et al., 2006; Lawley et al., 2016; Acosta-Góngora et al., 2018a).

Much of the western Churchill Province is intruded by ca. 2.62 to 2.58 Ga feldspar-porphyrific granitoid rocks of the Snow Island suite, which also has lesser mafic and extrusive components (Peterson et al., 2015, in press). K-feldspar-porphyrific monzogranite with crystallization ages of 2.615 to 2.585 Ga have been identified in the Tehery Lake–Wager Bay area (van Breemen et al., 2007; Wodicka et al., 2017a), confirming the presence of Snow Island suite plutonism. Peterson et al. (in press) recently proposed that the Snow Island suite represents a unified, short-lived, northwest-verging continental arc, and suggested that the Tehery Lake–Wager Bay area south of the Chesterfield shear zone (CSZ), together with the Chesterfield block and Snowbird Domain, may have formed part of a continental terrane that initially formed outboard of the Rae Craton and collided with an oceanic arc (Axis Lake gabbro of the Tantato Domain; Acosta-Góngora et al., 2018b) at ca. 2.62 Ga. At the peak time of Snow Island suite granite emplacement, i.e. at 2.605 Ga, the combined Tantato-Snowbird-Chesterfield-Tehery terrane collided with the proto-Rae block (Peterson et al., in press).

The Snow Island igneous event is interpreted to have had widespread thermotectonic effects that grew metamorphic zircon and monazite at ca. 2.62 to 2.60 Ga in the Uvauk Complex (Mills et al., 2007) just south of the Tehery Lake–Wager Bay area, and monazite at ca. 2.58 Ga in the Committee Bay belt (Berman et al., 2010). The Uvauk Complex also records subsequent tectonometamorphism at ca. 2.56 to 2.50 Ga (Mills et al., 2007)

linked with the MacQuoid Orogeny (Berman, 2010), while the Committee Bay belt was locally affected by the ca. 2.35 Ga Arrowsmith Orogeny (Berman et al., 2010, 2013a). Furthermore, evidence for both the MacQuoid and Arrowsmith orogenies are recorded in supracrustal rocks on Southampton Island, east of the Tehery Lake–Wager Bay area (Berman et al., 2013b).

In central Nunavut, Paleoproterozoic metasedimentary rocks are distributed to the west of the Tehery Lake–Wager Bay area in the <1.96 Ga Amer and <2.00 Ga Ketyet River groups, to the northwest in the <2.19 Ga Montresor and <2.46 Ga Chantrey groups, and to the northeast in the <1.90 Ga Penrhyn group on Melville Peninsula (Hinchey et al., 2007; Rainbird et al., 2010; McEwan, 2012; Partin et al., 2014; Calhoun, 2017; Percival et al., 2017). Rocks preserved in basement-cover infolds in the northern part of the Tehery Lake–Wager Bay area are interpreted as correlative with the lower formations of the Ketyet River group (Panagapko et al., 2003; Ferderber et al., 2013). The base of the lower succession of the Ketyet River group comprises abundant quartz arenite, impure arenite, and greywacke and local shale interbedded with two massive to pillowed mafic volcanic flow sequences (Rainbird et al., 2010; McEwan, 2012). The structures and sedimentology of the siliciclastic rocks are interpreted to represent a lacustrine to fluvial depositional setting. The volcanic rocks near the base of the Ketyet River group display a geochemical signature typical of tholeiitic continental flood basalt and may be related to the ca. 2.19 Ga Tulemalu/MacQuoid dykes and ca. 2.15 Ga Schultz swarm further to the west and southwest (Tella et al., 1997; Kerswill et al., 1999; Pehrsson et al., 2002; Rainbird et al., 2010; McEwan, 2012).

The Paleoproterozoic cover sequence records the intense weathering of a stable yet subsiding supercontinent, *Kenorland*, partial rifting, and extension of the Rae Craton with supercontinent breakup, flooding and deepening of basins, and subsequent foreland basin development during the early amalgamation of Laurentia in the supercontinent *Nuna* at ca. 1.9 Ga (Rainbird et al., 2010). Amalgamation began with the collision of the composite Rae-Chesterfield and Hearne domains, and resulted in the Snowbird Orogeny (Berman et al., 2007; Martel et al., 2008; Thiessen et al., 2018). The Snowbird tectonic zone (Hoffman, 1988), a crustal-scale geophysical lineament characterized by shear zones, mylonitic fabrics, and a string of uplifted mafic granulite domains along the southeastern Rae Craton margin (e.g. Kramanituar (Sanborn-Barrie et al., 2001, 2019), Uvauk (Mills et al., 2007), and Daly Bay (Gordon, 1988; Hanmer and Williams, 2001) complexes, and Hanbury Island shear zone (Tella and Annesley, 1988), formed or was reactivated during the Snowbird Orogeny (e.g. Baldwin et al., 2006; Berman et al., 2007). As an example, Mills et al. (2007) constrained the ca. 2.71 Ga anorthosite–gabbro of the Uvauk Complex as being structurally emplaced over ca. 2.68 Ga tonalite gneiss to the south at ca. 2.56 to 2.50 Ga during the MacQuoid Orogeny, and subsequent high-pressure granulite-facies deformation and magmatism at ca. 1.9 Ga (Snowbird Orogeny) ended with rapid exhumation of the complex to shallower crustal levels.

Syn- to post-orogenic plutonism is ubiquitous across the Rae Craton owing to crustal thickening that occurred during the Paleoproterozoic Trans-Hudson Orogen (THO; Hoffman, 1988; Corrigan et al., 2009). Granitic plutons, sills, and dykes of the Hudson igneous suite cut the dominant regional fabrics and were emplaced at ca. 1.85 to 1.81 Ga at mid-crustal levels across the THO hinterland (van Breemen et al., 2005). These rocks are present across much of the western Churchill Province, but are highly concentrated west of Hudson Bay. Peterson et al. (2002) interpreted the Hudson igneous suite as syn-orogenic to early postorogenic intrusions derived from in situ melting of lower- to middle-

crust wall rocks that contained garnet (giving low Y and HREE characteristics to the melts), and that migrated very short distances in the crust given their low thermal inertia and absence of equivalent extrusive rock types. Ultrapotassic lamprophyre and mafic syenite (Martell syenite), and the Dubawnt minette dykes and related volcanic rocks of the Dubawnt Supergroup (Gall et al., 1992) were produced synchronously with the Hudson igneous suite and have similar isotopic and trace element compositions (Peterson et al., 2002). In the Tehery Lake–Wager Bay area, several large plutons of Hudson monzogranite have been identified and confirmed with crystallization ages of ca. 1.843 to 1.808 Ga (Henderson and Roddick, 1990; Wodicka et al., 2017a; Therriault et al., 2018; Therriault, 2019). Similarly, the occurrence of ultrapotassic and minette rock types has been extended northeastward from the Baker Lake basin into the Tehery Lake–Wager Bay area (Steenkamp et al., 2015, 2016; Wodicka et al., 2015, 2016). Conversely, postorogenic rapakivi granite and its extrusive equivalents (Pitz Formation rhyolite) of the ca. 1.75 Ga Nueltin igneous suite are concentrated where post-THO extensional faulting and basin development is preserved (Peterson et al., 2002), and have therefore not been identified in the Tehery Lake–Wager Bay area.

The final addition of new rock material in the Tehery Lake–Wager Bay area was the emplacement of diabase (gabbro) dykes, including relatively small (1–5 m thick), undated, roughly east-trending porphyritic dykes and much larger (10–50 m wide), southeast-trending dykes that trace northwestward back to the source region for the 1.276 Ga Mackenzie dyke swarm (LeCheminant and Heaman, 1989).

GENERAL GEOLOGY, STRUCTURE, AND METAMORPHISM IN THE TEHERY LAKE–WAGER BAY AREA

The Tehery Lake–Wager Bay area can be broadly separated into six domains (Fig. 1 and 2; Gordon, Lunan, and Kummel Lake domains presented here, and Ukkusiksalik and Douglas Harbour domains, and Daly Bay Complex on adjoining maps) that are delineated by large-scale structures and characterized by differing metamorphic assemblages, Sm-Nd isotopic and U-Pb data, and/or the presence/absence of specific lithologies. Sm-Nd isotopic data from 12 felsic metaplutonic rocks of van Breemen et al. (2007) and 34 new samples collected during the Tehery-Wager activity indicate that the central part of the study area south of the CSZ can be divided into two crustal domains based on Nd model ages (Wodicka et al., 2017a): the Gordon Domain to the east with 3.29 to 2.76 Ga Nd model ages, and Lunan Domain to the west characterized by generally younger 2.97 to 2.72 Ga Nd model ages. The Gordon Domain is also characterized by distinctly older plutonic rocks, including a granodiorite gneiss with a U-Pb Mesoarchean crystallization age of ca. 2860 Ma, in addition to Neoproterozoic plutonic rocks of ca. 2714 to 2676 and 2615 to 2602 Ma age, while Archean plutonic rocks within the Lunan Domain are only of Neoproterozoic age (ca. 2711–2699 and 2584 Ma) (Table 1¹). A transition zone between the two domains (Fig. 1 and 2) includes both evolved and juvenile rocks with ca. 3.09 to 2.75 Ga Nd model ages, and the complexly deformed, upper amphibolite-facies ca. 2700 Ma Archean Lorillard supracrustal belt (Table 1). This zone also coincides with a north-south-trending lineament visible in the magnetic field and regional Bouguer gravity data (Wodicka et al., 2017a), and is marked, at least in part, by numerous 1 to 5 m wide,

¹ Table 1 contains a compilation of U-Pb data and Nd model ages (where applicable) from samples within the Tehery Lake–Wager Bay area and immediate vicinity. For more details on the age data cited in the text (e.g. dating method, age uncertainties, data sources, etc.), the reader is referred to this table.

steeply dipping high-strain zones with local reverse-sense indicators and widespread, strongly developed foliation and mineral lineation fabrics in the field. The Gordon and Lunan domains have been broadly correlated to the north with the Mesoarchean Repulse Bay and Neoproterozoic Committee Bay blocks, respectively (Pehrsson et al., 2013; Spratt et al., 2014; Wodicka et al., 2017a, b), but the extrapolation of the transition zone north of the CSZ and precise connection to those blocks remain unknown owing to a lack of adequate isotopic data in that area and added complexity from structural offsets across the Chesterfield and Wager shear zones. Alternatively, it is possible that the Lunan and Gordon domains initially formed outboard of the proto-Rae Craton (Peterson et al., in press) and therefore have no counterparts in the Rae Craton north of the CSZ. In general, the composite Lunan-Lorillard-Gordon Domain preserves upper amphibolite- to granulite-facies metamorphic assemblages largely attributed to the THO, as well as common evidence of partial retrogression to moderate amphibolite-facies conditions.

The Chesterfield shear zone, which separates the upper amphibolite- to granulite-facies rocks of the Lunan-Lorillard-Gordon Domain in the hanging wall to the south from lower amphibolite-facies rocks of the Douglas Harbour Domain in the footwall to the north (Fig. 1 and 2), is characterized by strongly deformed to mylonitic rocks that dip either steeply to the north-northwest or moderately to steeply to the south-southeast at surface (Steenkamp et al., 2016; Wodicka et al., 2016). By contrast, magnetotelluric, gravity, and magnetic data consistently indicate that the CSZ forms a south-dipping structure at depth (Spratt et al., 2014; Tschirhart et al., 2016). The trace of the CSZ can be loosely followed in magnetic anomaly patterns from the north side of Baker Lake to the southern shore of Wager Bay. Where porphyritic rocks of the Snow Island suite are present along the CSZ, they develop a characteristic augen texture.

The Douglas Harbour Domain (Fig. 1 and 2) preserves the lowest grade metamorphic assemblages (lower to moderate amphibolite-facies) and is therefore interpreted as preserving the structurally highest crustal level of the THO in the Tehery Lake–Wager Bay area. Thick-skinned thrust sheets preserve stacked repetitions of basement gneiss and supracrustal rocks, which are interpreted as correlative with the lower stratigraphy of the Ketyet River group (Panagapko et al., 2003; Ferderber et al., 2013). U-Pb crystallization ages of Archean basement gneiss and plutonic rocks within the Douglas Harbour Domain range in age from ca. 2904 to 2608 Ma (Table 1).

The Douglas Harbour Domain is bounded to the north by the Wager shear zone (WSZ; Fig. 1 and 2), a major dextral strike-slip shear zone with mylonitic fabrics and sub-horizontal mineral lineations that run east to west (Henderson et al., 1986; Henderson and Broome, 1990; Henderson and Roddick, 1990; Henderson et al., 1991; Therriault et al., 2017; Therriault, 2019). Based on magnetic survey data, the western end of the WSZ bends to the northwest and merges with the Amer mylonite zone through a right-lateral extensional step-over (Therriault, 2019). A secondary splay also splits off the WSZ to the southwest and becomes the Quoich River fault (Panagapko et al., 2003). Recent field and geochronological work has led to the identification of a number of plutonic intrusions with ca. 1840 to 1822 Ma crystallization ages that locally cut pre-existing fabrics along the WSZ, and provide evidence of high-temperature ductile deformation in the WSZ that was active as late as ca. 1754 to 1738 Ma (Table 1). The youngest movement along the WSZ is coeval with dextral strike-slip movement along the Amer mylonite zone to the west and widespread post-orogenic plutonism (Kivalliq igneous suite; Therriault, 2019; Therriault et al., 2022).

Rocks of the Ukkusiksalik Domain north of the WSZ (Fig. 1 and 2) contain lithologies similar to those of the Douglas Harbour Domain, including supracrustal rocks of probable Archean age (Paliak belt; Henderson et al., 1986; Jefferson et al., 1991) and of Paleoproterozoic age (lower Ketyet River group-equivalent). Archean basement rocks have been dated at only two localities, at ca. 2707 and 2606 Ma (Table 1). Unlike elsewhere in the Tehery Lake–Wager Bay area, pelitic rocks in the Ukkusiksalik Domain preserve rare, relict high-pressure metamorphic mineral assemblages (kyanite-bearing) that are overprinted by lower-pressure, high-temperature assemblages (sillimanite- and cordierite-bearing), which are common in the Gordon and Lunan domains (H. Steenkamp, work in progress).

The southern part of the Tehery Lake–Wager Bay area preserves two large granulite domains: the Kummel Lake Domain and the Daly Bay Complex (Fig. 1 and 2). The Kummel Lake Domain comprises mostly Archean granodiorite to monzogranite basement gneiss with a greasy green colour, abundant in situ leucosome and magnetite (creating a widespread strongly magnetic anomaly), and local orthopyroxene, including in leucosome (Steenkamp et al., 2016; Wodicka et al., 2016). Preliminary unconstrained 3D inversions of the regional gravity data across this area show the Kummel Lake Domain as a dense body that dips shallowly to the north, below the Lunan-Lorillard-Gordon Domain (Wodicka et al., 2017a). A strong magnetic anomaly defines the surface expression of the northern margin of the Kummel Lake Domain (Fig. 2), which, where mapped, corresponds to an outward-dipping, greenschist-facies fault. The fault geometry, combined with the subsurface geometry of the domain and metamorphic contrast across the domain boundary, suggest that the Kummel Lake Domain has been exhumed as a metamorphic core complex. It is possible that the Kummel Lake Domain shares aspects of the tectonometamorphic histories of the mafic granulite complexes along the Snowbird tectonic zone, but more data are required to test this.

The Daly Bay Complex is a Paleoproterozoic, dominantly mafic granulite domain that is juxtaposed alongside Archean basement orthogneiss via mylonitic shear zones and later, brittle normal faults. It is dominated by pyroxene-plagioclase tonalitic igneous phases with lesser anorthosite to gabbro intrusions, and garnet leucogranite derived from a largely pelitic sedimentary protolith (Gordon, 1988; Hanmer and Williams, 2001). In contrast to the Kummel Lake Domain, the Daly Bay Complex has been modeled as having inward-dipping sheared margins and is interpreted as having been tectonically emplaced over amphibolite-facies basement orthogneiss of the Gordon Domain by northward-directed thrusting (Gordon and Lawton, 1995).

While evidence of early (ca. 2.7, 2.6, 2.55, and 1.95–1.90 Ga) tectonometamorphic events are becoming increasingly recognized with ongoing research (H. Steenkamp, work in progress; N. Wodicka, work in progress), the entire Tehery Lake–Wager Bay area has been penetratively overprinted by the intensely pervasive ca. 1.88 to 1.81 Ga THO, which involved the closure of the Manikewan ocean and culminated with the subduction of the Superior Craton from the south below the crustal collage that makes up the western Churchill Province (e.g. Hoffman, 1988; Corrigan et al., 2009). The THO has forced the transposition of many pre-existing fabrics and the development of new fabrics and structures. Early folds (F_1 ; timing unknown) are refolded by the dominant northeast- or southwest-trending F_2 folds thought to have developed through progressive northwestward compression of the THO. Most F_2 folds plunge shallowly to moderately to the northeast. The pervasive regional deformation at this time also produced the predominant L_2 mineral lineations, and S_2 foliation fabrics and gneissosity. Both the WSZ

and CSZ may have initially formed during the Archean (e.g. Peterson et al., in press), but were reactivated during (and after for the WSZ) the THO based on geochronological and metamorphic data (Henderson and Roddick, 1990; Therriault, 2019; Therriault et al., 2018, 2022; H. Steenkamp, work in progress). A later compressional event created broad, northwest- or southeast-trending open folds (F_3) that, with interference over F_2 folds, created Type 1 refolds (Grasemann et al., 2004) resulting in canoe- and donut-shaped map patterns.

GEOLOGICAL UNITS

Archean orthogneiss (units $\bar{A}t$, $\bar{A}gd$, $\bar{A}ggd$, $\bar{A}mg$)

The basement gneiss across the Gordon, Lunan, Douglas Harbour, and Ukkusiksalik domains is predominantly medium-grained biotite±hornblende±magnetite±muscovite granodiorite (unit $\bar{A}gd$). Lesser biotite±hornblende±epidote tonalite (unit $\bar{A}t$) typically occurs as magnetic lows at the cores of antiformal domes and locally contains thin sheets of supracrustal rocks separating it from other basement rocks. The gneissic to migmatitic $\bar{A}gd$ (Fig. 3) and $\bar{A}t$ units contain medium- to coarse-grained, injected or locally derived monzogranite veins that have been transposed parallel to the regional foliation fabric (S_2). Pods, bands, and lenses of intermediate to mafic rock (e.g. units $\bar{A}gab$, $\bar{A}amph$) are locally present and generally similarly deformed. These rocks may represent older plutonic or volcanic phases, transposed and boudinaged dykes, or intrusions that postdate the emplacement of the basement granodiorite and tonalite; however, in many places, strong overprinting renders identification of the intermediate to mafic protoliths difficult. In the Gordon Domain, an area along the southern coast of Wager Bay (NTS sheets 46-E and 56-H) comprises biotite+magnetite±hornblende granodiorite to monzogranite with local occurrences of orthopyroxene (unit $\bar{A}ggd$), indicating the area underwent granulite-facies metamorphism (e.g. Derome, 1988). Biotite±hornblende±magnetite monzogranite plutons and sheets (unit $\bar{A}mg$) are intercalated with the unit $\bar{A}gd$ granodiorite across much of the area and generate moderate magnetic anomalies. Although this unit is predominantly monzogranite, it locally varies compositionally from syenite to syenogranite, and texturally from homogeneous and medium-grained, to more gneissic and coarse-grained (Fig. 4), to K-feldspar megacrystic. In a few localities south of the CSZ, preserved contact relationships suggest that unit $\bar{A}mg$ monzogranite intrudes supracrustal rocks of the Lorillard supracrustal belt (unit $\bar{A}Lsp$). Unit $\bar{A}t$ tonalite and unit $\bar{A}gd$ granodiorite samples across the area yield U-Pb ages of ca. 2904 to 2903, 2860, and 2714 to 2676 Ma (Table 1), indicating variability in basement orthogneiss-protolith crystallization histories (van Breemen et al., 2007; Wodicka et al., 2017a, b). An orthopyroxene-bearing monzogranite sample from the $\bar{A}ggd$ unit yielded a poorly constrained crystallization age of ca. <2720 to >2703 Ma, while a few dated unit $\bar{A}mg$ monzogranite plutons give uniform ages of ca. 2699 to 2689 Ma (Table 1).

Archean Kummel Lake Domain (units $\bar{A}KLggd$, $\bar{A}KLgmg$, $\bar{A}KLsp$, $\bar{A}KLan$)

The Kummel Lake Domain is underlain predominantly by granulite-grade granodiorite (unit $\bar{A}KLggd$; Fig. 5) and monzogranite (unit $\bar{A}KLgmg$) that are characterized as having a greasy green overall colour, abundant magnetite and in situ melt leucosome, migmatitic textures, and the local presence of orthopyroxene (including in leucosome). It also preserves discontinuous, folded panels of supracrustal rocks, termed the Kummel Lake belt (unit $\bar{A}KLsp$), that correspond to magnetic-low anomalies (Fig. 2) in the otherwise strongly magnetic, granulite-facies basement rocks (units $\bar{A}KLggd$, $\bar{A}KLgmg$). The

Kummel Lake belt (unit $\bar{A}KLsp$) comprises garnet±clinopyroxene amphibolite, garnet-biotite semipelite–pelite with abundant leucosome (Fig. 6), and lesser silicate-facies iron-formation. A garnet-biotite semipelite sampled from the Kummel Lake belt yielded a maximum depositional age of ca. 2725 to 2670 Ma, and contains evidence of metamorphism at about 2612 to 2570 Ma, 2554 to 2505 Ma, and 1840 Ma (Table 1). Gabbroic anorthosite (unit $\bar{A}KLan$) is locally associated with the supracrustal rocks in the Kummel Lake Domain. Unit $\bar{A}KLgd$ granodiorite and unit $\bar{A}KLan$ leucogabbro yield preliminary crystallization ages of ca. 2711 and 2696 Ma, respectively (Table 1). Supracrustal rocks and ca. 2.71 Ga gabbroic anorthosite in and around the Uvauk Complex to the south of NTS sheet 56-C (Tella, 1993; Tella and Schau, 1994; Mills et al., 2007), as well as the granulite suite of Tella et al. (1993; unit $\bar{A}grn$) present to the south of NTS sheet 56-B (described as quartzofeldspathic granulite interlayered with minor mafic granulite, paragneiss, granitic gneiss, anorthosite, and anorthositic gabbro), appear to continue into the Kummel Lake Domain, although extrapolation of structural features and lithological boundaries require more focused work to confirm this.

Archean Lorillard belt (units $\bar{A}Lsp$, $\bar{A}Lif$)

Supracrustal rocks belonging to the Lorillard belt (unit $\bar{A}Lsp$) lie primarily within the transition zone that separates the Gordon and Lunan domains south of the CSZ (Fig. 1 and 2; Wodicka et al., 2017a, b). These rocks occur as discontinuous panels and layers surrounded by monzogranite (unit $\bar{A}mg$) sheets/injections and granodiorite (unit $\bar{A}gd$) basement gneiss, and comprise varying combinations and abundances of sillimanite quartzite, garnet-biotite±cordierite semipelite–pelite, mafic (amphibolite) to intermediate rocks of probable volcanic origin, biotite psammite, garnetite, iron-formation (including unit $\bar{A}Lif$), and lesser calc-silicate and marble (Fig. 7). Metamorphosed gabbro and peridotite are common within the Lorillard belt acting as relatively rigid and commonly boudinaged layers without pervasive internal fabrics. Complete lithological descriptions are presented in Steenkamp et al. (2015, 2016). A U-Pb crystallization age of ca. 2700 Ma has been obtained for an intermediate rock with basaltic andesite geochemistry, while quartzite to psammite samples record U-Pb zircon maximum-depositional ages of ca. 2705 to 2650 Ma (Table 1). Some of the dated supracrustal rocks contain metamorphic zircon or monazite overgrowths and grains at ca. <2700, >2580, 1943, and 1879 to 1857 Ma (Table 1; H. Steenkamp, work in progress). It is unclear whether the Lorillard belt rocks are genetically related to supracrustal rocks in the Kummel Lake Domain (unit $\bar{A}KLsp$) or Paliak supracrustal belt (unit $\bar{A}Psp$); however, based on their geographic positions and interpreted peak metamorphic mineral assemblages, these three packages of rocks are interpreted to have different tectonometamorphic histories.

Archean Paliak belt (unit $\bar{A}Psp$)

Supracrustal rocks intercalated with basement orthogneiss (units $\bar{A}gd$ and $\bar{A}mg$) and exposed north of the CSZ within both the Douglas Harbour and Ukkusiksalik domains are termed the Paliak belt (unit $\bar{A}Psp$; Henderson et al., 1986; Jefferson et al., 1991). These rocks include garnet-biotite±sillimanite±cordierite semipelite to pelite, garnet±clinopyroxene amphibolite, white quartzite, and lesser peridotite and iron-formation. Semipelite is locally gossanous, and both the semipelite and amphibolite have garnet porphyroblasts up to 3 to 4 cm wide. The iron-formation is silicate facies with layers of quartz, garnet, magnetite, and grunerite amphibole. The quartzite commonly contains sillimanite and is only a few metres thick. Distinction between the Paliak belt (unit $\bar{A}Psp$)

and the Paleoproterozoic supracrustal rocks (unit pPsp) is not always straightforward, particularly at the scale of mapping, and it is possible that some of the unit APsp panels identified on the map also contain unit pPsp rocks, or vice versa. However, a key characteristic that helps to distinguish these two units is how highly deformed and intercalated the Paliak belt rocks are with the basement orthogneiss (units Agd and Amg). The Paliak belt also contains similar lithologies compared to the Lorillard belt (unit ALsp), but has slightly lower grade metamorphic-mineral assemblages and a considerably lower abundance of mafic lithologies (Panagapko et al., 2003). A sample of garnet-biotite-sillimanite pelite from an exposure of Paliak rocks (unit APsp) that is surrounded by Paleoproterozoic supracrustal rocks (unit pPsp) yielded a maximum depositional age of ca. 2695 to 2660 Ma and contains metamorphic zircon overgrowths dated at ca. 1872 Ma (Table 1).

Archean igneous rocks (units Aamph, Agab, ASis)

Intermediate and mafic igneous rocks are relatively common across the Tehery Lake–Wager Bay area; most are present as pods, layers, and lenses in basement orthogneiss and supracrustal rocks. However, most occurrences are too small for representation at the current map scale. Unit Aamph encompasses a suite of rock types including intermediate compositions, such as diorite, through more mafic compositions, such as amphibolite. In general, these rocks are medium to coarse grained and commonly contain metamorphic garnet, magnetite, and local biotite. Portable X-ray fluorescence data collected in the field indicate that most intermediate to mafic rocks have variably metasomatized andesitic to basaltic compositions with several subpopulations having chemical characteristics that suggest derivation from either primitive and mantle-like sources or more evolved sources (Lawley et al., 2015; Steenkamp et al., 2015).

Unit Agab represents isolated bodies of coarse-grained gabbro that occur in basement orthogneiss and are only locally deformed at their margins or strongly deformed throughout to produce augen textures, depending on the local strain intensity. Undeformed gabbro comprises cumulate orthopyroxene-clinopyroxene with interstitial plagioclase overprinted by coronitic growth of amphibole, garnet, and local biotite.

Megacrystic K-feldspar monzogranite to monzonite (unit ASis), correlative to the ca. 2.6 Ga Snow Island suite (Peterson et al., 2015; in press), occurs as large plutons and smaller sheets intercalated with basement orthogneiss within the Lunan, Gordon, Douglas Harbour, and Ukkusiksalik domains, inside the Gordon-Lunan transition zone, and along the CSZ. Internally, the plutons contain round K-feldspar grains up to 5 cm wide, locally with white plagioclase rims constituting pseudo-rapakivi texture, in a matrix of plagioclase, quartz, and biotite. Positive magnetic anomalies within the plutons correlate with dioritic and gabbroic phases that contain plagioclase phenocrysts. The compositional variability is believed to be due to magma mixing of mafic and felsic igneous phases (Steenkamp et al., 2015). The plutons are characterized as having strongly sheared to mylonitic outer margins that extend into the underlying granodiorite gneiss, resulting in porphyroclastic K-feldspar and metamorphic amphibole, biotite, and elongate garnet growth parallel to the shear direction. Geophysical modelling of the prominent Borden complex in NTS 56-A, combined with field observations and U-Pb age data, suggest that the southern body represents a thin slice of Snow Island suite porphyry (~800 m of structural thickness) that was displaced from the roof of the northern Borden body and transported south along a flat-lying shear zone (Peterson et al., in press). The smaller unit ASis sheets that are intercalated with basement orthogneiss commonly

display pale pink K-feldspar augen set in a fine-grained, dark grey matrix and local pinhead garnet porphyroblasts. U-Pb zircon age determinations for variably deformed unit ASIs plutonic rocks of felsic to mafic composition across the Tehery Lake–Wager Bay area range between ca. 2615 and 2584 Ma (Table 1).

Paleoproterozoic supracrustal rocks (units pPsp, pPq)

North of the CSZ in the Douglas Harbour and Ukkusiksalik domains are supracrustal rocks (unit pPsp) dominated by white quartzite, biotite psammite, and lesser biotite±garnet±sillimanite±muscovite semipelite and amphibolite that are relatively less deformed than any of the Archean supracrustal packages. These rocks, interpreted to be correlative with the lower stratigraphy of the Paleoproterozoic Ketyet River group, are primarily exposed in imbricated thick-skinned thrust sheets preserved in moderately to shallowly doubly plunging synformal structures (intersecting F₂ and F₃ folds), with the base of each thrust layer comprising Archean granodiorite (unit Ågd) or monzogranite (unit Åmg) basement. Mylonite zones several metres in thickness are present both above and below the thrust planes, which are generally concordant with S₂ mineral foliations and lithologic contact orientations. The white quartzite (unit pPq) is locally up to 170 m thick and contains sillimanite and muscovite. Thick biotite psammite layers are concentrated in the central and eastern exposures of unit pPsp, and locally contain a gneissic texture with thin leucocratic layers. The semipelite is locally gossanous and commonly associated with amphibolite, which contains clinopyroxene and rare garnet. Detrital zircon U-Pb age analysis of unit pPq quartzite and unit pPsp psammite and semipelite illustrates the predominant local sourcing of sediment with prominent ca. 2.72 to 2.67 Ga age peaks (Table 1).

Paleoproterozoic Daly Bay Complex (units ApPDBgd, pPDBpg, pPDBgl, pPDBgb, pPDBpgs, pPDBla, pPDBan, pPDBsp, pPDBq)

Previously restricted to rocks metamorphosed at granulite-facies conditions (Gordon, 1988), the Daly Bay Complex is herein more broadly defined to contain both the inner granulite-facies rocks and the surrounding, concentrically oriented belt of strongly foliated to mylonitic lower-grade rocks (i.e. roughly corresponding to the outer shear zone of Gordon, 1988). This new definition is consistent with the tentative correlation between amphibolite units within the 'outer shear zone' and granulite-facies mafic rocks south of the high-strain zone (W. Garrison, unpub. B.Sc. research project, 2016). The interior of the complex is dominated by orthopyroxene-clinopyroxene tonalite (pyroxene granulite of Gordon, 1988; unit pPDBpg) that, based on field relationships, appears to be the oldest igneous phase in the complex (Gordon, 1988; Gordon and Lawton, 1995). This unit is injected with orthopyroxene(-clinopyroxene) tonalite–diorite–gabbro layers that both cut the predominant fabric of the tonalite and are transposed with that fabric. One of these layers, an orthopyroxene tonalite, yielded a crystallization age of ca. 1918 Ma (Table 1). The interior part of the complex also hosts relatively older, white-weathering garnet±sillimanite±biotite leucogranite (unit pPDBgl) that contains remnant pelitic gneiss enclaves, and younger gabbroic plutons (unit pPDBgb) that cut the fabric of the surrounding orthopyroxene-clinopyroxene tonalite and leucogranite. The core of the complex is mantled by more orthopyroxene-clinopyroxene tonalite and lesser leucogranite that have been more intensely deformed and locally sheared (unit pPDBpgs). Migmatitic granodiorite–granite (unit ApPDBgd) around Bernheimer Bay is thought to represent annealed mylonite units of Archean basement orthogneiss (units Ågd, Åmg)

that may have facilitated some tectonic shuffling inside the complex (Hanmer and Williams, 2001), but this theory has not yet been tested. The inner mantle area also contains plutons of layered gabbroic anorthosite (unit pPDBla) and anorthosite (unit pPDBan) with a U-Pb zircon crystallization age of ca. 1908 Ma and U-Pb titanite cooling age of ca. 1902 Ma (Table 1), as well as supracrustal panels (unit pPDBsp) comprising psammite-semipelite schist and gneiss, orthoquartzite (unit pPDBq where thick enough for representation at this map scale), and local marble. A similar assemblage of supracrustal rocks (Kingmirit belt of Wodicka et al., 2017a) occurs in discontinuous panels in the footwall of the outer extensional fault around the complex but retains only greenschist- to lower amphibolite-facies metamorphic minerals and relatively finer-grained textures (i.e. psammite-semipelite phyllite; W. Garrison, unpub. B.Sc. research project, 2016). A sample of semipelite collected from one of the Kingmirit belt panels gave a maximum depositional age of ca. 1975 Ma (Table 1), and it is currently assumed that the supracrustal rocks inside the complex share a depositional history with those at its margin.

It is unclear how and when the presumed Paleoproterozoic supracrustal rocks (unit pPDBsp) that appear to host the igneous phases of the Daly Bay Complex came to be buried. However, based on the above relationships and ages, heat from the magmatic emplacement of the mantle-derived orthopyroxene-clinopyroxene tonalite (units pPDBpg and pPDBpgs) contributed to localized granulite-facies metamorphism, and consequently melted the supracrustal wall-rocks. Later, orthopyroxene-bearing tonalite–diorite–gabbro injections and anorthositic rocks were emplaced syn-kinematically during the formation of the dominant fabric in the complex (Hanmer and Williams, 2001). This deformation event shortened the complex through north-verging compression (Gordon and Lawton, 1995) and produced internal folding and dislocation of the complex into two components, possibly with the migmatitic rocks around Bernheimer Bay facilitating some of that movement (Hanmer and Williams, 2001). Exactly how the Daly Bay Complex was uplifted and exposed from the middle to lower crust remains to be determined; however, there are likely some similarities to the rapid exhumation processes that affected the Kramanitar and Uvauk complexes (Sanborn-Barrie et al., 2001; Mills et al., 2007) related to the Snowbird Orogeny (e.g. Berman et al., 2007; Martel et al., 2008; Regis et al., 2019; Card et al., 2021).

Paleoproterozoic igneous rocks (units pPHm, pPmd, pPup, pPdb)

Monzogranite to monzonite plutons, sills, and dykes (unit pPHm) are interpreted to be related to the ca. 1.85 to 1.81 Ga Hudson igneous suite (Peterson et al., 2002). The pluton phases are generally homogeneous, medium to coarse grained, and monzogranitic to syenogranitic with minor biotite and magnetite. Xenolith blocks and enclaves comprising basement orthogneiss and supracrustal rocks are locally present near the pluton margins. A weak, outcrop-scale foliation is present in some plutons as preferred exfoliation-weathering planes and biotite alignment. Sills tend to have a slightly more K-feldspar-rich composition, such as syenogranite to monzonite, and locally contain fluorite. Ubiquitous, randomly oriented granitic dykes are the youngest phase of the suite as they are undeformed and cut through all rock units except the Mackenzie dykes. Dated unit pPHm plutons and sills across the Tehery Lake–Wager Bay area yielded U-Pb ages ranging from ca. 1843 to 1808 Ma (Table 1). The Ford Lake batholith, the southern portion of which is exposed in NTS 56-G, gave ages (ca. 1826 to 1823 Ma; Table 1; LeCheminant et al., 1987) similar to other Hudson suite rocks, but is petrographically unique given the

presence of coarse, locally porphyritic textures and higher abundance of mafic minerals (T. Peterson, work in progress).

The Wager pluton (Wodicka et al., 2017a) comprises biotite+magnetite±hornblende monzodiorite (unit pPmd) that intrudes basement gneiss of the Ukkusiksalik Domain and the western segment of the WSZ. It becomes increasingly abundant westward where the WSZ is thought to connect to the Amer mylonite zone via a right-lateral extensional step-over (west of the study area; Therriault, 2019). The monzodiorite is generally undeformed, medium grained, and contains distinctive grey translucent plagioclase crystals with white opaque rims. It is associated with a strong magnetic signal, contains deformed inclusions of granodiorite gneiss and supracrustal rocks, cuts the fabric of the basement orthogneiss units in these areas, and is cut by late, pervasive monzogranitic pegmatite dykes attributed to the Hudson igneous suite. A sample of monzodiorite-granodiorite yielded a U-Pb crystallization age of ca. 1833 Ma (Table 1), within the age range of the Hudson suite.

Medium- to coarse-grained, homogeneous plutons with ultrapotassic compositions (unit pPup) comprise varying proportions of phlogopite clinopyroxenite, biotite-pyroxene syenite (Fig. 8), clinopyroxene melasyenite, leucosyenite, and rare biotite+clinopyroxene+olivine lamprophyre. The more mafic phases occur as rafts that are partly resorbed, and net-veined complexes are common along the margins of large intrusions (Steenkamp et al., 2015). Ultrapotassic plugs of varying sizes are scattered across the Lunan Domain, in the southern portion of the transition zone between the Gordon and Lunan domains, and north of the CSZ. Some plugs are associated with small intrusions of syenogranite (unit pPHm), for example in the cores of doubly plunging fold structures, while all bodies are cut by coarse-grained to pegmatitic monzogranite dykes (unit pPHm). All ultrapotassic intrusions produce strong magnetic signatures. The pPup unit has been dated at two localities, including within the large Armit Lake intrusion in the southern part of NTS 56-B, giving ages of ca. 1827 to 1825 Ma (Table 1). It is believed to be correlative with, and represent the deeper crustal equivalent of (at least south of the CSZ; Wodicka et al., 2015), the Martell syenite suite (Peterson et al., 2002).

Diabase dykes (unit pPdb) that trend roughly east-west are 1 to 5 m wide, weather dark brown, and dominantly contain plagioclase phenocrysts in a dark grey, aphanitic matrix. Gabbroic compositions and homogeneous, fine-grained textures are locally present where the dykes are widest. Although they have not been dated, it is possible that these dykes are related to the ca. 2.19 Ga Tulemalu/MacQuoid dyke swarm south and southwest of Chesterfield Inlet (Tella et al., 1997), the undated Thelon River dyke swarm (Buchan and Ernst, 2004; Wodicka et al., 2016; Jefferson et al., in press) exposed along strike to the west of the Tehery Lake–Wager Bay area, or the ca. 1700 Ma (Bleeker et al., 2011) east- and east-northeast-striking Pelly Bay dyke swarm identified on the mainland south of Boothia Peninsula (Ryan et al., 2009).

Mesoproterozoic dykes (unit mPM)

Diabase to gabbro dykes (unit mPM) of the ca. 1276 Ma Mackenzie igneous event (LeCheminant and Heaman, 1989) cut through all older rock units and trend to the southeast. They are generally medium to coarse grained, homogeneous, 10 to 50 m wide, and weather dark brown.

ACKNOWLEDGMENTS

The Tehery-Wager activity was made possible by logistical contributions from the Polar Continental Shelf Program, Discovery Mining, Ookpik Aviation, and Prairie Helicopters (pilot/engineers E. Polzin and J. Lagimodiere). The 2015 to 2017 field seasons included expert support and assistance from É. Girard, C. Gilbert, R. Buenviaje, C. Guilmette, K. Larson, J. Beales, W. Garrison, K. Hatogina, A. Lion, R. Bayne, L. Lebeau, I. McMartin, S. Day, M. Roy, I. Randour, and J. Byatt. Camp staff are graciously thanked for their cooking, wildlife monitoring, and companionship: L. Levesque, D. Guilfoyle, R. Issaluk, D. Aggark, A. Alooq, and T. Evviuk. The hamlet offices and Hunters and Trappers organizations of Chesterfield Inlet and Baker Lake, Victor Sammurtok School, G. Brocklebank, Baker Lake Lodge, Baker Lake Contracting and Supplies, Aulajuk Ltd., SK Construction, and Peregrine Diamonds Ltd. are thanked for their support and organizational assistance with various aspects of the project, including community engagement meetings. The authors are grateful to D. Regis for providing a helpful and constructive review of the bedrock geology maps and descriptive notes.

REFERENCES

- Acosta-Góngora, P., Pehrsson, S.J., Sandeman, H.A., Martel, E., and Peterson, T.D., 2018a. The Ferguson Lake deposit: an example of Ni-Cu-Co-PGE mineralization emplaced in a back-arc basin setting?; *Canadian Journal of Earth Sciences*, v. 55, p. 958–979. <https://doi.org/10.1139/cjes-2017-0185>
- Acosta-Góngora, P., Pehrsson, S.P., Knox, B., Regis, D., Hulbert, L., Creaser, R.A., and Ashton, K., 2018b. Neoproterozoic convergent margin Ni-Cu mineralization? Axis Lake and Nickel King Ni-Cu deposits in the south Rae Province craton of the Canadian Shield; *Precambrian Research*, v. 316, p. 305–323. <https://doi.org/10.1016/j.precamres.2018.07.016>
- Baldwin, J.A., Bowring, S.A., Williams, M.L., and Mahan, K.H., 2006. Geochronological constraints on the evolution of high-pressure felsic granulites from an integrated electron microprobe and ID-TIMS geochemical study; *Lithos*, v. 88, p. 173–200. <https://doi.org/10.1016/j.lithos.2005.08.009>
- Berman, R.G., 2010. Metamorphic map of the western Churchill Province, Canada; Geological Survey of Canada, Open File 5279, 1 .zip file. <https://doi.org/10.4095/287320>
- Berman, R.G., Davis, W.J., and Pehrsson, S., 2007. Collisional Snowbird tectonic zone resurrected: growth of Laurentia during the 1.9 accretionary phase of the Hudsonian orogeny; *Geology*, v. 35, no. 10, p. 911–914. <https://doi.org/10.1130/G23771A.1>
- Berman, R.G., Sanborn-Barrie, M., Rayner, N., Carson, C., Sandeman, H.A.I., and Skulski, T., 2010. Petrological and in situ SHRIMP geochronological constraints on the tectonometamorphic evolution of the Committee Bay belt, Rae Province, Nunavut; *Precambrian Research*, v. 181, p. 1–20. <https://doi.org/10.1016/j.precamres.2010.05.009>

- Berman, R.G., Pehrsson, S., Davis, W.J., Ryan, J.J., Qui, H., and Ashton, K.E., 2013a. The Arrowsmith orogeny: geochronological and thermobarometric constraints on its extent and tectonic setting in the Rae craton, with implications for pre-Nuna supercontinent reconstruction; *Precambrian Research*, v. 232, p. 44–69. <https://doi.org/10.1016/j.precamres.2012.10.015>
- Berman, R.G., Sanborn-Barrie, M., Rayner, N., and Whalen, J., 2013b. The tectonometamorphic evolution of Southampton Island, Nunavut: insight from petrologic modeling and in situ SHRIMP geochronology of multiple episodes of monazite growth; *Precambrian Research*, v. 232, p. 140–166. <http://dx.doi.org/10.1016/j.precamres.2012.08.011>
- Bleeker, W. and Ernst, R.E., 2011. Does the Slave craton continue into southern Siberia: comparison of their large igneous province (LIP) records; in 39th Annual Yellowknife Geoscience Forum Abstracts, (ed.) B.J. Fischer and D.M. Watson; Northwest Territories Geoscience Office, Yellowknife, NT, YKGSF Abstracts, v. 2011, p. 22–23.
- Brouillette, P., Girard, É., and Huot-Vézina, G., 2019. Geological Survey of Canada Bedrock Data Model and tools: design and user guide documentation including ArcGIS(TM) add-ins; Geological Survey of Canada, Open File 8247, 129 p, 1 .zip file. <https://doi.org/10.4095/314673>
- Buchan, K.L. and Ernst, R.E., 2004. Diabase dyke swarms and related units in Canada and adjacent regions; Geological Survey of Canada, Map 2022A, scale 1:5 000 000. <https://doi.org/10.4095/214883>
- Calhoun, L.J., 2017. Lithostratigraphy and structure of the eastern Amer group, Nu, using combined surface mapping and enhanced sub-surface geophysical data; M.Sc. thesis, University of New Brunswick, Fredericton, New Brunswick, 166 p.
- Card, C.D., Bethune, K.M., Rayner, N., and Ashton, K.E., 2021. Tectonic significance of the Virgin River shear zone of the Canadian Shield and implications for the origin of the Snowbird tectonic zone of Laurentia; *Precambrian Research*, v. 361, 22 p. <https://doi.org/10.1016/j.precamres.2021.106241>
- Corrigan, D., Pehrsson, S., Wodicka, N., and de Kemp, E., 2009. The Paleoproterozoic Trans-Hudson orogen: a prototype of modern accretionary processes; *in* *Ancient Orogens and Modern Analogues*, (ed.) J.B. Murphy, J.D. Keppie, and A.J. Hynes; Geological Society, London, Special Publication 327, p. 457–479. <https://doi.org/10.1144/SP327.19>
- Corrigan, D., Nadeau, L., Brouillette, P., Wodicka, N., Houlié, M.G., Tremblay, T., Machado, G., and Keating, P., 2013. Overview of the GEM Multiple Metals - Melville Peninsula project, central Melville Peninsula, Nunavut; Geological Survey of Canada, Current Research 2013-19, 21 p. <https://doi.org/10.4095/292862>

- Coyle, M. and Kiss, F., 2012a. Residual total magnetic field, aeromagnetic survey of the Tehery Lake area, NTS 56-A/SW, 56-B/SE and parts of 56-A/NW, Nunavut; Geological Survey of Canada, Open File 7203, scale 1:100 000.
<https://doi.org/10.4095/291780>
- Coyle, M. and Kiss, F., 2012b. Residual total magnetic field, aeromagnetic survey of the Tehery Lake area, NTS 56-B/SW, 56-C/SE, and parts of 56-B/NW and 56-C/NE, Nunavut; Geological Survey of Canada, Open File 7204, scale 1:100 000.
<https://doi.org/10.4095/291781>
- Coyle, M. and Kiss, F., 2012c. Residual total magnetic field, aeromagnetic survey of the Tehery Lake area, parts of 56-B/NW, 56-C/NE, 56-F/SE and 56-G/SW, Nunavut; Geological Survey of Canada, Open File 7205, scale 1:100 000.
<https://doi.org/10.4095/291782>
- Coyle, M. and Kiss, F., 2012d. Residual total magnetic field, aeromagnetic survey of the Tehery Lake area, parts of 56-A/NW, 56-B/NE, 56-G/SE and 56-H/SW, Nunavut; Geological Survey of Canada, Open File 7206, scale 1:100 000.
<https://doi.org/10.4095/291783>
- Coyle, M. and Kiss, F., 2012e. First vertical derivative of the magnetic field, aeromagnetic survey of the Tehery Lake area, NTS 56-A/SW, 56-B/SE, and parts of 56-A/NW and 56-B/NE, Nunavut; Geological Survey of Canada, Open File 7207, scale 1:100 000. <https://doi.org/10.4095/291784>
- Coyle, M. and Kiss, F., 2012f. First vertical derivative of the magnetic field, aeromagnetic survey of the Tehery Lake area, NTS 56-B/SW, 56-C/SE, and parts of 56-B/NW and 56-C/NE, Nunavut; Geological Survey of Canada, Open File 7208, scale 1:100 000. <https://doi.org/10.4095/291785>
- Coyle, M. and Kiss, F., 2012g. First vertical derivative of the magnetic field, aeromagnetic survey of the Tehery Lake area, NTS 56-B/NW, 56-C/NE, 56-F/SE and 56-G/SW, Nunavut; Geological Survey of Canada, Open File 7209, scale 1:100 000.
<https://doi.org/10.4095/291786>
- Coyle, M. and Kiss, F., 2012h. First vertical derivative of the magnetic field, aeromagnetic survey of the Tehery Lake area, parts of NTS 56-A/NW, 56-B/NE, 56-G/SE and 56-H/SW, Nunavut; Geological Survey of Canada, Open File 7210, scale 1:100 000. <https://doi.org/10.4095/291787>
- Davis, W.J., Hanmer, S., Tella, S., Sandeman, H.A., and Ryan, J.J., 2006. U-Pb geochronology of the MacQuoid supracrustal belt and Cross Bay plutonic complex: key components of the northwestern Hearne subdomain, western Churchill Province, Nunavut, Canada; *Precambrian Research*, v. 145, p. 53–80.
<https://doi.org/10.1016/j.precamres.2005.11.016>
- DePaolo, D.J., 1981. Neodymium isotopes in the Colorado front range and crust-mantle evolution in the Proterozoic; *Nature*, v. 291, p. 193–196.

- Derome, I., 1988. Evolution of the Wager Bay shear zone, District of Keewatin, N.W.T., Canada; M.Phil. thesis, The University of Exeter, Devon, United Kingdom, 190 p.
- Ferderber, J.L., 2013. Exploring the Tehery region: correlating supracrustal sequences using detrital zircon geochronology, Rae Craton, Nunavut; B.Sc. thesis, University of Ottawa, Ottawa, Ontario, 83 p.
- Ferderber, J., Kellett, D.A., and Wodicka, N., 2013. Exploring the Tehery region: correlating supracrustal sequences using detrital zircon geochronology, Rae Craton, Nunavut; Geological Survey of Canada, Open File 7424. <https://doi.org/10.4095/292709>
- Gall, Q., Peterson, T.D., and Donaldson, J.A., 1992. A proposed revision of early Proterozoic stratigraphy of the Thelon and Baker Lake basins, Northwest Territories; *in* Current Research, Part C; Geological Survey of Canada, Paper 92-1C, p. 129–137. <https://doi.org/10.4095/132856>
- Geological Survey of Canada, 1978. Airborne gamma ray spectrometric map, Armit Lake, District of Keewatin, Northwest Territories; Geological Survey of Canada, Geophysical Series, Map 35256G, scale 1:250 000. <https://doi.org/10.4095/118609>
- Geological Survey of Canada, 1998. Western Churchill NATMAP Project region, decorrugated residual total field magnetics, colour shaded relief; Geological Survey of Canada, scale 1:1 000 000 scale.
- Gordon, T.M., 1988. Precambrian geology of the Daly Bay area, District of Keewatin; Geological Survey of Canada, Memoir 422, 21 p. <https://doi.org/10.4095/126316>
- Gordon, T.M. and Heywood, W.W., 1987. Geology, Daly Bay area, District of Keewatin, Northwest Territories; Geological Survey of Canada, Map 1652A, scale 1:250 000. <https://doi.org/10.4095/126317>
- Gordon, T.M. and Lawton, D.C., 1995. Geometry of the allochthonous Daly Bay Complex, Northwest Territories: a model constrained by geology and a three-dimensional gravity interpretation; Canadian Journal of Earth Sciences, v. 32, p. 1292–1302. <https://doi.org/10.1139/e95-105>
- Grasemann, B., Wiesmayr, G., Draganits, E., and Füsseis, F., 2004. Classification of refold structures; Journal of Geology, v. 112, p. 119–125. <https://doi.org/10.1086/379696>
- Hanmer, S. and Williams, M.L., 2001. Targeted field work in the Daly Bay Complex, Hudson Bay, Nunavut; Geological Survey of Canada, Current Research 2001-C15, 10 p. <https://doi.org/10.4095/212095>

- Hanmer, S., Tella, S., Ryan, J., Sandeman, H., and Berman, R.G., 2006. Late Neoproterozoic thick-skinned thrusting and Paleoproterozoic reworking in the MacQuoid supracrustal belt and Cross Bay plutonic complex, western Churchill Province, Nunavut, Canada; *Precambrian Research*, v. 144, p. 126–139. <https://doi.org/10.1016/j.precamres.2005.10.005>
- Henderson, J.R., LeCheminant, A.N., Jefferson, C.W., Coe, K., and Henderson, M.N., 1986. Preliminary account of the geology around Wager Bay, District of Keewatin; *in* *Current Research, Part A*; Geological Survey of Canada, Paper 86-1A, p. 159–176. <https://doi.org/10.4095/127683>
- Henderson, J.R. and Broome, J., 1990. Geometry and kinematics of Wager shear zone interpreted from structural fabrics and magnetic data; *Canadian Journal of Earth Sciences*, v. 27, p. 590–604. <https://doi.org/10.1139/e90-055>
- Henderson, J.R. and Roddick, J.C., 1990. U-Pb age constraint on the Wager shear zone, District of Keewatin, N.W.T.; *in* *Radiogenic Age and Isotopic Studies: Report 3*; Geological Survey of Canada, Paper 89-2, p. 149–152. <https://doi.org/10.4095/129080>
- Henderson, J.R., Jefferson, C.W., Henderson, M.N., Coe, K., and Derome, I., 1991. Geology of the region around Wager Bay, District of Keewatin (parts of 46E and 56H); Geological Survey of Canada, Open File 2383, scale 1:100 000, 1 .zip file. <https://doi.org/10.4095/131852>
- Heywood, W.W., 1967a. Geology of northeastern District of Keewatin and southern Melville Peninsula, Districts of Franklin and Keewatin; Geological Survey of Canada, Open File 1, scale 1:506 880. <https://doi.org/10.4095/100310>
- Heywood, W.W., 1967b. Geological notes northeastern District of Keewatin and southern Melville Peninsula, District of Franklin, Northwest Territories (parts of 46, 47, 56, 57); Geological Survey of Canada, Paper 66-40, 20 p. <https://doi.org/10.4095/100923>
- Hinchey, A.M., Ryan, J.J., Davis, W.J., Nadeau, L., and James, D.T., 2007. Detrital zircon geochronology of the Archean volcano-sedimentary sequence of the Barclay belt and the Paleoproterozoic marble-quartzite sequence of the Northern Chantrey group, Boothia mainland area, Kitikmeot region, Nunavut; Geological Survey of Canada, *Current Research 2007-C1*, 19 p. <https://doi.org/10.4095/223860>
- Hoffman, P.F., 1988. United plates of America, the birth of a craton: Early Proterozoic assembly and growth of Laurentia; *Annual Reviews of Earth and Planetary Science*, v. 16, p. 543–603. <https://doi.org/10.1146/annurev.ea.16.050188.002551>
- Jefferson, C.W., Smith, J.E.M., and Hamilton, S.M., 1991. Preliminary account of the resource assessment study of proposed national park, Wager Bay–Southampton Island areas, District of Keewatin; Geological Survey of Canada, Open File 2351, 47 p. <https://doi.org/10.4095/131509>

- Jefferson, C.W., Pehrsson, S., Tschirhart, V., Peterson, T., Chorlton, L., Bethune, K., White, J.C., Davis, W., McNicoll, V., Paulen, R.C., and Rayner, N., in press. Geology and metallogeny of the northeast Thelon Basin region and comparison with the Athabasca Basin; *in* Canada's northern shield: new perspectives from the Geomapping for Energy and Minerals Program, (ed.) S. Pehrsson, N. Wodicka, N. Rogers, and J.A. Percival; Geological Survey of Canada, Bulletin 612.
- Keating, P.B., Pilkington, M., and Miles, W.F., 2003. Colour shaded relief of the residual total magnetic field with Keating correlation coefficients, NTS 56 BCFG, Nunavut, Canada; Geological Survey of Canada, Open File 4489, scale 1:500 000. <https://doi.org/10.4095/214605>
- Kerswill, J.A., Kjarsgaard, B.A., Bretzlaff, R., Jenner, G.A., and Samaras, C., 1999. Metallogeny and geology of the Half Way Hills area, central Churchill Province, Northwest Territories (Nunavut); *in* Current Research 1999-C; Geological Survey of Canada, p. 29–41. <https://doi.org/10.4095/210169>
- Lawley, C.J.M., Steenkamp, H.M., and Wodicka, N., 2015. Portable X-ray fluorescence geochemical results from the Tehery Lake–Wager Bay area, western Hudson Bay, Nunavut; Canada-Nunavut Geoscience Office, Geoscience Data Series 2015-011, Microsoft® Excel® file.
- Lawley, C.J.M., McNicoll, V., Sandeman, H., Pehrsson, S., Simard, M., Castonguay, S., Mercier-Langevin, P., and Dubé, B., 2016. Age and geological setting of the Rankin Inlet greenstone belt and its relationship to the gold endowment of the Meliadine gold district, Nunavut, Canada; *Precambrian Research*, v. 275, p. 471–495. <https://doi.org/10.1016/j.precamres.2016.01.008>
- LeCheminant, A.N. and Heaman, L.M., 1989. Mackenzie igneous events, Canada: Middle Proterozoic hotspot magmatism associated with ocean opening; *Earth and Planetary Science Letters*, v. 96, no. 1, p. 38–48. [https://doi.org/10.1016/0012-821X\(89\)90122-2](https://doi.org/10.1016/0012-821X(89)90122-2)
- LeCheminant, A.N., Roddick, J.C., Tessier, A.C., and Bethune, K.M., 1987. Geology and U–Pb ages of early Proterozoic calc-alkaline plutons northwest of Wager Bay, District of Keewatin; *in* Current Research, Part A; Geological Survey of Canada, Paper 87-1A, p. 773–782. <https://doi.org/10.4095/122551>
- Lord, C.S. and Wright, G.M., 1967. Geology, southeastern Barren Grounds, District of Keewatin–District of Mackenzie; Geological Survey of Canada, Map 1216A, scale 1:1 000 000. <https://doi.org/10.4095/108854>
- Martel, E., van Breemen, O., Berman, R.G., and Pehrsson, S., 2008. Geochronology and tectonometamorphic history of the Snowbird Lake area, Northwest Territories, Canada: new insights into the architecture and significance of the Snowbird tectonic zone; *Precambrian Research*, v. 161, Issues 3–4, p. 201–230. <https://doi.org/10.1016/j.precamres.2007.07.007>

- McEwan, B.J., 2012. Structural style and regional comparison of the Paleoproterozoic Ketyet River group in the region north-northwest of Baker Lake, Nunavut; M.Sc. thesis, University of Regina, Regina, Saskatchewan, 161 p.
- Mills, A.J., Berman, R.G., Davis, W.J., Tella, S., Carr, S., Roddick, C., and Hanmer, S., 2007. Thermobarometry and geochronology of the Uvauk complex, a polymetamorphic Neoproterozoic and Paleoproterozoic segment of the Snowbird tectonic zone, Nunavut, Canada; *Canadian Journal of Earth Sciences*, v. 44, p. 245–266. <https://doi.org/10.1139/e06-080>
- Panagapko, D.A., Pehrsson, S., Pilkington, M., and Currie, M., 2003. Geoscience data compilation: Tehery Lake–Wager Bay area, Nunavut (NTS 56 B, C, F, and G), part 1 – base data themes; Geological Survey of Canada, Open File 1809, 1 .zip file. <https://doi.org/10.4095/214767>
- Partin, C.A., Bekker, A., Corrigan, D., Modeland, S., Francis, D., and Davis, D.W., 2014. Sedimentological and geochemical basin analysis of the Paleoproterozoic Penrhyn and Piling groups of Arctic Canada; *Precambrian Research*, v. 251, p. 80–101. <https://doi.org/10.1016/j.precamres.2014.06.010>
- Pehrsson, S.J., Jenner, G., and Kjarsgaard, B., 2002. The Ketyet River group: correlation with Paleoproterozoic sequences of northeastern Rae and implications for Proterozoic orogenesis in the western Churchill Province; Geological Association of Canada, conference abstracts 2002.
- Pehrsson, S., Berman, R.G., and Davis, W.J., 2013. Paleoproterozoic orogenesis during Nuna aggregation: a case study of reworking of the Rae craton, Woodburn Lake, Nunavut; *Precambrian Research*, v. 232, p. 167–188. <https://doi.org/10.1016/j.precamres.2013.02.010>
- Peplinski, L., Ungalaq, Z., and Gilbert, C., 2017. Inuktitut place names, Aksarniq, Nunavut; Canada-Nunavut Geoscience Office, Open File Map 2017-01, scale 1:250 000.
- Percival, J.A., Davis, W.J., and Hamilton, M.A., 2017. U-Pb zircon geochronology and depositional history of the Montresor Group, Rae Province, Nunavut, Canada; *Canadian Journal of Earth Sciences*, v. 54, p. 512–528. <https://doi.org/10.1139/cjes-2016-0170>
- Peterson, T.D., van Breemen, O., Sandeman, H., and Cousens, B., 2002. Proterozoic (1.85–1.75 Ga) igneous suites of the western Churchill Province: granitoid and ultrapotassic magmatism in a reworked Archean hinterland; *Precambrian Research*, v. 119, Issues 1–4, p. 73–100. [https://doi.org/10.1016/S0301-9268\(02\)00118-3](https://doi.org/10.1016/S0301-9268(02)00118-3)
- Peterson, T.D., Jefferson, C.W., and Anand, A., 2015. Geological setting and geochemistry of the 2.6 Ga Snow Island Suite in the central Rae Domain of the Western Churchill Province, Nunavut; Geological Survey of Canada, Open File 7841, 29 p. <https://doi.org/10.4095/296599>

- Peterson, T.D., Wodicka, N., Pehrsson, S.J., Acosta-Gongora, P., Tschirhart, V., Jefferson, C.W., Steenkamp, H., Martel, E., Percival, J., and Corrigan, D., in press. The Rae Province at 2.6 Ga: a Sanukitoid Storm on the Canadian Shield; *in* Canada's northern shield: new perspectives from the Geomapping for Energy and Minerals Program, (ed.) S. Pehrsson, N. Wodicka, N. Rogers, and J.A. Percival; Geological Survey of Canada, Bulletin 612.
- Rainbird, R.H., Davis, W.J., Pehrsson, S.J., Wodicka, N., Rayner, N., and Skulski, T., 2010. Early Paleoproterozoic supracrustal assemblages of the Rae domain, Nunavut, Canada: Intracratonic basin development during supercontinent break-up and assembly; *Precambrian Research*, v. 181, Issues 1–4, p. 167–186. <https://doi.org/10.1016/j.precamres.2010.06.005>
- Regis, D., Davis, W.J., Ryan, J.J., Berman, R.G., Pehrsson, S., Joyce, N.L., and Sandeman, H.A., 2019. Multiple burial–exhumation episodes revealed by accessory phase in high-pressure granulite-facies rocks (Rae Craton, Nunavut, Canada); *Contributions to Mineralogy and Petrology*, v. 174, 25 p. <https://doi.org/10.1007/s00410-019-1572-8>
- Richan, L., Gibson, H.L., Houlié, M.G., and Leshner, C.M., 2015. Mode of emplacement of Archean komatiitic tuffs and flows in the Selkirk Bay area, Melville Peninsula, Nunavut, Canada; *Precambrian Research*, v. 263, p. 174–196. <http://dx.doi.org/10.1016/j.precamres.2015.03.004>
- Ryan, J.J., Nadeau, L., Hinchey, A.M., James, D.T., Sandeman, H.A., Schetselaar, E.M., Davis, W.J., and Berman, R.G., 2009. Bedrock geology of the southern Boothia mainland area (Pelly Bay-Rae Strait-Spence Bay map area), Kitikmeot region, Nunavut; Geological Survey of Canada, Current Research 2009-1, 21 p. <https://doi.org/10.4095/226519>
- Sanborn-Barrie, M., Carr, S.D., and Theriault, R., 2001. Geochronological constraints on metamorphism, magmatism and exhumation of deep-crustal rocks of the Kramanituar Complex, with implications for the Paleoproterozoic evolution of the Archean western Churchill Province, Canada; *Contributions to Mineralogy and Petrology*, v. 141, p. 592–612. <https://doi.org/10.1007/s004100100262>
- Sanborn-Barrie, M., Camacho, A., and Berman, R.G., 2019. High-pressure, ultrahigh-temperature 1.9 Ga metamorphism of the Kramanituar Complex, Snowbird Tectonic Zone, Rae Craton, Canada; *Contributions to Mineralogy and Petrology*, v. 174, 26 p. <https://doi.org/10.1007/s00410-019-1547-9>
- Sanborn-Barrie, M., Davis, W.J., Berman, R.G., Rayner, N., Skulski, T., and Sandeman, H., 2014. Neoproterozoic continental crust formation and Paleoproterozoic deformation of the central Rae craton, Committee Bay belt, Nunavut; *Canadian Journal of Earth Sciences*, v. 51, p. 635–667. <https://dx.doi.org/10.1139/cjes-2014-0010>

- Sandeman, H.A., Hanmer, S., Armitage, A.A., Davis, W.J., and Ryan, J.J., 2006. Petrogenesis of Neoarchean volcanic rocks of the MacQuoid supracrustal belt: A back-arc setting for the northwestern Hearne subdomain, western Churchill Province, Canada; *Precambrian Research*, v. 144, p. 140–165. <https://doi.org/10.1016/j.precamres.2005.11.001>
- Siivola, J. and Schmidt, R., 2007. Recommendations by the IUGS Subcommittee on the Systematics of Metamorphic Rocks: Web version 01.02.07. <<https://www.semanticscholar.org/paper/Recommendations-by-the-IUGS-Subcommission-on-the-of-Siivola-Schmid/db0016b6295591bf9cb2bd567a8964c607379ac3>> [accessed March 30, 2023]
- Skulski, T., Sanborn-Barrie, M., and Sandeman, H., 2003a. Geology, Walker Lake and Arrowsmith River area, Nunavut; Geological Survey of Canada, Open File 3777, scale 1:100 000, 1 .zip file. <https://doi.org/10.4095/214769>
- Skulski, T., Sandeman, H.A., Sanborn-Barrie, M., MacHattie, T., Young, M., Carson, C., Berman, R.G., Brown, J., Rayner, N., Panagapko, D., Byrne, D., and Deyell, C., 2003b. Bedrock geology of the Ellice Hills map area and new constraints on the regional geology of the Committee Bay area, Nunavut; Geological Survey of Canada, Current Research 2003-C22, 11 p. <https://doi.org/10.4095/214204>
- Spratt, J.E., Skulski, T., Craven, J.A., Jones, A.G., Snyder, D.B., and Kiyani, D., 2014. Magnetotelluric investigations of the lithosphere beneath the central Rae Craton, mainland Nunavut, Canada; *Journal of Geophysical Research, Solid Earth*, v. 119, p. 2415–2439. <https://doi.org/10.1002/2013JB010221>
- Steenkamp, H.M., Wodicka, N., Lawley, C.J.M., Peterson, T.D., and Guilmette, C., 2015. Overview of bedrock mapping and results from portable X-ray fluorescence spectrometry in the eastern part of the Tehery Lake–Wager Bay area, western Hudson Bay, Nunavut; *in* Summary of Activities 2015; Canada-Nunavut Geoscience Office, p. 121–134.
- Steenkamp, H.M., Wodicka, N., Weller, O.M., and Kendrick, J., 2016. Overview of bedrock mapping in the Tehery Lake–Wager Bay area, western Hudson Bay, Nunavut; *in* Summary of Activities 2016; Canada-Nunavut Geoscience Office, p. 27–40.
- Tella, S., 1993. Geology, Chesterfield Inlet, District of Keewatin, Northwest Territories; Geological Survey of Canada, Open File 2756, scale 1:250 000. <https://doi.org/10.4095/183967>
- Tella, S. and Annesley, I.R., 1988. Hanbury Island Shear Zone, a deformed remnant of a ductile thrust, District of Keewatin, N.W.T.; *in* Current Research, Part C; Geological Survey of Canada, Paper 1988-1C, p. 283–289. <https://doi.org/10.4095/122643>

- Tella, S. and Schau, M., 1994. Geology, Gibson Lake east-half, District of Keewatin, Northwest Territories; Geological Survey of Canada, Open File 2737, scale 1:250 000, 1 .zip file. <https://doi.org/10.4095/193350>
- Tella, S., Schau, M., Armitage, A.E., and Loney, B.C., 1993. Precambrian geology and economic potential of the northeastern parts of Gibson Lake map area, District of Keewatin, Northwest Territories; *in* Current research, Part C; Geological Survey of Canada, Paper 93-C1, p. 196–208. <https://doi.org/10.4095/134246>
- Tella, S., LeCheminant, A.N., Sanborn-Barrie, M., and Venance, K.E., 1997. Geology and structure of parts of MacQuoid Lake map area, District of Keewatin, Northwest Territories; *in* Current Research 1997-C; Geological Survey of Canada, p. 123–132. <https://doi.org/10.4095/208638>
- Thiessen, E.J., Gibson, H.D., Regis, D., and Pehrsson, S.J., 2018. Deformation and extensional exhumation of 1.9 Ga high-pressure granulites along the Wholdaia Lake shear zone, south Rae Craton, Northwest Territories, Canada; *Lithosphere*, v. 10, no. 5, p. 641–661. <https://doi.org/10.1130/L704.1>
- Therriault, I., 2019. Structural characterisation and geochronological constraints on the Wager shear zone, northwestern Hudson Bay, Nunavut; M.Sc. thesis, University of British Columbia Okanagan, Kelowna, British Columbia, 228 p.
- Therriault, I., Steenkamp, H.M., and Larson, K.P., 2017. New mapping and initial structural characterization of the Wager shear zone, northwestern Hudson Bay, Nunavut; *in* Summary of Activities 2017; Canada-Nunavut Geoscience Office, p. 1–12.
- Therriault, I., Steenkamp, H.M., Larson, K.P., and Cottle, J.M., 2018. Geochronological constraints on deformation in the Wager shear zone, northwestern Hudson Bay, Nunavut. *in* Summary of Activities 2018; Canada-Nunavut Geoscience Office, p. 1–14.
- Therriault, I., Larson, K.P., Steenkamp, H.M., Apen, F.E., Graziani, R., Soret, M., Guilmette, C., and Cottle, J.M., 2022. Characterization of the Wager shear zone, Nunavut, Canada: Insights from microstructures and geochronology; *Canadian Journal of Earth Sciences*, v.60, p 78–96. <https://doi.org/10.1139/cjes-2022-0031>
- Tschirhart, V.L., Wodicka, N., and Steenkamp, H.M., 2016. Shallow crustal structure of the Tehery Lake–Wager Bay area, western Hudson Bay, Nunavut, from potential-field datasets; *in* Summary of Activities 2016; Canada-Nunavut Geoscience Office, p. 41–50.
- van Breemen, O., Peterson, T.D., and Sandeman, H.A., 2005. U–Pb zircon geochronology and Nd isotope geochemistry of Proterozoic granitoids in the western Churchill Province: intrusive age pattern and Archean source domains; *Canadian Journal of Earth Sciences*, v. 42, p. 339–377. <https://doi.org/10.1139/e05-007>

- van Breemen, O., Pehrsson, S., and Peterson, T.D., 2007. Reconnaissance U-Pb SHRIMP geochronology and Sm-Nd isotope analyses from the Tehery-Wager Bay gneiss domain, western Churchill Province, Nunavut; Geological Survey of Canada, Current Research 2007-F2, 15 p. <https://doi.org/10.4095/224019>
- Wodicka, N., Corrigan, D., Nadeau, L., and Erdmann, S., 2011. New U-Pb geochronological results from Melville Peninsula: unravelling the Archean and early Paleoproterozoic magmatic history of the north-central Rae Craton; *in* Ottawa 2011 GAC-MAC-SEG-SGA Joint Annual Meeting, Ottawa, Abstracts Volume 34, p. 236.
- Wodicka, N., Steenkamp, H.M., Lawley, C.J.M., Peterson, T.D., Guilmette, C., Girard, É., and Buenviaje, R., 2015. Report of activities for the bedrock geology and economic potential of the Tehery-Wager area: GEM-2 Rae project; Geological Survey of Canada, Open File 7970, 14 p. <https://doi.org/10.4095/297294>
- Wodicka, N., Steenkamp, H.M., Weller, O.M., Kendrick, J., Tschirhart, V.L., Peterson, T.D., and Girard, É., 2016. Report of 2016 Activities for the Bedrock Geology and Economic Potential of the Tehery-Wager Area: GEM-2 Rae Project; Geological Survey of Canada, Open File 8149, 21 p. <https://doi.org/10.4095/299392>
- Wodicka, N., Steenkamp, H.M., Peterson, T.D., McMartin, I., Day, S.J.A., and Tschirhart, V.L., 2017a. Report of 2017 activities for the geology and economic potential of the Tehery-Wager area, Nunavut: GEM-2 Rae Project; Geological Survey of Canada, Open File 8318, 20 p. <https://doi.org/10.4095/305979>
- Wodicka, N., Steenkamp, H.M., Peterson, T., Whalen J., and Lawley, C.J.M., 2017b. Neoproterozoic to Paleoproterozoic evolution of the south-central Rae margin, Tehery-Wager area, Nunavut: Insights from field mapping, U-Pb geochronology, and Sm-Nd isotope data; Kingston 2017: Geological Association of Canada-Mineralogical Association of Canada, Kingston, Ontario, Program with abstracts, v. 40, p. 423.
- Wright, G.M., 1955. Geological notes on central District of Keewatin, Northwest Territories; Geological Survey of Canada, Paper 55-17, 21 p. <https://doi.org/10.4095/101294>
- Wright, G.M., 1967. Geology of the southeastern Barren Grounds, parts of Mackenzie and Keewatin, Operations Keewatin, Baker, Thelon; Geological Survey of Canada, Memoir 350, 91 p. <https://doi.org/10.4095/101544>
- Zaleski, E., Pehrsson, S., Duke, N., Davis, W.J., L'Heureux, R., Greiner, E., and Kerswill, J.A., 2000. Quartzite sequences and their relationships, Woodburn Lake group, western Churchill Province, Nunavut; Geological Survey of Canada, Current Research 2000-C7, 12 p. <https://doi.org/10.4095/211098>

ADDITIONAL INFORMATION

The Additional Information folder of this product's digital download contains figures and tables that appear in the map surround as well as additional geological information not depicted on the map, nor this document, nor the geodatabase.

-PDF(s) of each figure that appears in the CGM surround.

-PDF of Table 1 that do not appear in the CGM surround

AUTHOR CONTACT

Questions, suggestions, and comments regarding the geological information contained in the data sets should be addressed to:

N. Wodicka

601 Booth Street

Ottawa ON

K1A 0E8

natasha.wodicka@NRCan-RNCan.gc.ca

COORDINATE SYSTEM

Projection: Universal Transverse Mercator

Units: metres

Zone: 15

Horizontal Datum: NAD83

Vertical Datum: mean sea level

BOUNDING COORDINATES

Western longitude: 93°00'00"W

Eastern longitude: 90°00'00"W

Northern latitude: 65°00'00"N

Southern latitude: 64°00'00"N

SOFTWARE VERSION

Data has been originally compiled and formatted for use with ArcGIS™ desktop version 10.8.2 developed by ESRI®.

DATA MODEL INFORMATION

Bedrock

Based on a data-centric approach, the GSC Bedrock Model was designed using the ESRI ArcGIS® environment. The model architecture is almost entirely tailored to the proprietary functionalities of the ESRI® File Geodatabase such as *SubTypes*, *Domain Values* and *Relationship Classes*.

Consult PDFs in Data folder for complete description of the model with its feature classes, tables, attributes, and domain values.

Note: the PDF document is not intended to describe the entire GSC Bedrock Model, but it provides a complete and detailed description of a subset of the model representing the published dataset.

For a more in depth description of the data model please refer to the official publication:

Brouillette, P., Girard, É., and Huot-Vézina, G., 2019. Geological Survey of Canada Bedrock Data Model and tools: design and user guide documentation including ArcGIS(TM) add-ins; Geological Survey of Canada, Open File 8247, 129 p, 1 .zip file. <https://doi.org/10.4095/314673>