

Proposed Maximum Residue Limit

PMRL2024-02

Pyroxsulam

(publié aussi en français)

2 February 2024

This document is published by the Health Canada Pest Management Regulatory Agency. For further information, please contact:

Publications
Pest Management Regulatory Agency
Health Canada
2 Constellation Drive
8th floor, A.L. 2608 A
Ottawa, Ontario K1A 0K9

Internet: canada.ca/pesticides pmra.publications-arla@hc-sc.gc.ca

Information Service: 1-800-267-6315 pmra.info-arla@hc-sc.gc.ca



ISSN: 1925-0835 (print) 1925-0843 (online)

Catalogue number: H113-24/2024-2E (print version)

H113-24/2024-2E-PDF (PDF version)

© His Majesty the King in Right of Canada, as represented by the Minister of Health Canada, 2024

All rights reserved. No part of this information (publication or product) may be reproduced or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, or stored in a retrieval system, without prior written permission of Health Canada, Ottawa, Ontario K1A 0K9.

Purpose of consultation

A maximum residue limit (MRL)¹ is being proposed for the pesticide pyroxsulam, as part of the following application for Canadian use, under submission number 2022-0479.

Under the authority of the *Pest Control Products Act*, Health Canada's Pest Management Regulatory Agency (PMRA) is proposing acceptability of the requested application to add the new commodity of rye (belonging to wheat crop subgroup 15-21A) to the product label of Simplicity GoDRI Herbicide containing technical grade pyroxsulam and safener cloquintocet acid, to control or suppress certain weeds. The specific uses approved in Canada are detailed on this product label, Pest Control Products Act Registration Number 31916.

The evaluation of this pyroxsulam application indicated that the end-use product has value, and the human health and environmental risks associated with the new use are acceptable. Dietary risks from the consumption of foods listed in Table 1 were shown to be acceptable when pyroxsulam is used according to the supported label directions. Therefore, foods containing residues resulting from this use are safe to eat, and an MRL is being proposed as a result of this assessment. A summary of the field trial data used to support the proposed MRL can be found in Appendix I.

Dietary health assessment

In assessing the risk of a pesticide, Health Canada combines information on pesticide toxicity with information on the degree and duration of dietary exposure to the pesticide residue from food. The risk assessment process involves four distinct steps:

- 1) Identifying the toxicology hazards posed by the pesticide;
- 2) Determining the "acceptable dietary level" for Canadians (including all vulnerable populations), which is protective of adverse health effects;
- 3) Estimating human dietary exposure to the pesticide from all applicable sources (domestic and imported commodities); and
- 4) Characterizing health risk by comparing the estimated human dietary exposure to the acceptable dietary level.

Before registering a pesticide for food use in Canada, Health Canada must determine the quantity of residues that could remain in or on the food when the pesticide is used according to label directions and that such residues will not be a concern to human health (Steps 3 and 4 above). If estimated human exposure is less than or equal to the acceptable level (developed in Step 2 above), Health Canada concludes that consuming residues resulting from use according to approved label directions is not a health concern. The proposed MRL is then subject to consultation to legally specify it as an MRL. An MRL applies to the identified raw agricultural food commodity as well as to any processed food product that contains it, except for certain

An MRL is the maximum amount of residue that may remain in or on food when a pesticide is used according to label directions.

instances where different MRLs are specified for the raw agricultural commodity and its processed product(s).

Consultation on the proposed MRL for pyroxsulam is being conducted via this document. Consultation on the proposed MRL for cloquintocet-mexyl (to cover residues of the safener cloquintocet acid) will be addressed under a separate action. Health Canada invites the public to submit written comments on the proposed MRL for pyroxsulam in accordance with the process outlined in the Next steps section of this document.

To comply with Canada's international trade obligations, consultation on the proposed MRL is also being conducted internationally by notifying the <u>World Trade Organization</u>, as coordinated by the <u>Canada's Notification Authority and Enquiry Point</u>.

Proposed MRL

The proposed MRL, to replace the MRL already established for pyroxsulam, is summarized in Table 1.

 Table 1
 Proposed Maximum Residue Limit for Pyroxsulam

Common name	Residue definition	MRL (ppm) ¹	Food commodity
Pyroxsulam	<i>N</i> -(5,7-dimethoxy[1,2,4]triazolo[1,5- a]pyrimidin-2-yl)-2-methoxy-4- (trifluoromethyl)-3-pyridinesulfonamide	0.01	Wheat (crop subgroup 15-21A) ²

ppm = parts per million

The commodities included in the listed crop subgroup can be found on the <u>Residue Chemistry Crop Groups</u> webpage in the <u>Pesticides section</u> of Canada.ca.

MRLs established in Canada may be found using the <u>Maximum Residue Limit Database</u> on the <u>Maximum Residue Limits for Pesticides</u> webpage. The database allows users to search for established MRLs, regulated under the *Pest Control Products Act*, both for pesticides or for food commodities.

International situation and trade implications

The MRL proposed for pyroxsulam in Canada is the same as the corresponding American tolerance for wheat grain as listed in the <u>Electronic Code of Federal Regulations</u>, 40 CFR Part 180, by pesticide. Currently, there are no Codex MRLs² listed for pyroxsulam in or on any commodity on the Codex Alimentarius <u>Pesticide Index</u> webpage.

The MRL is proposed to replace the currently established 0.01 ppm MRLs for wheat and triticale with an MRL for all food commodities in the crop subgroup at the same MRL value.

The Codex Alimentarius Commission is an international organization under the auspices of the United

Next steps

Health Canada invites the public to submit written comments on the proposed MRL for pyroxsulam up to 75 days from the date of publication of this document (by 17 April 2024). Please forward your comments to Publications (see the contact information on the cover page of this document). Health Canada will consider all comments received and a science-based approach will be applied in making a final decision on the proposed MRL. Comments received will be addressed in a separate document linked to this PMRL. The established MRL will be legally in effect as of the date that it is entered into the Maximum Residue Limit Database.

Nations that develops international food standards, including MRLs.

Appendix I

Summary of field trial data used to support the proposed maximum residue limit

Previously reviewed residue data from field trials conducted in/on wheat were reassessed in the framework of this petition on rye.

Dietary risk assessment results

Studies in laboratory animals showed no acute health effects relevant to dietary exposure. Consequently, a single dose of pyroxsulam is not likely to cause acute health effects in the general population (including infants and children).

Chronic dietary (food plus drinking water) intake estimates indicated that the general population and all population subgroups are exposed to less than 1% of the acceptable daily intake, and therefore there are no health concerns.

Maximum residue limit

The recommendation for a maximum residue limit (MRL) for pyroxsulam was based upon the submitted field trial data, and the guidance provided in the OECD MRL Calculator. Table A1 summarizes the residue data for pyroxsulam used to calculate the proposed MRL for the wheat crop subgroup (15-21A).

Table A1 Summary of field trial and processing data used to support the MRL

Commodity	Application method/Total application rate (g a.i./ha) ¹	Preharvest interval (days)	Lowest average field trial residues (ppm)	Highest average field trial residues (ppm)	Experimental processing factor
Wheat grain	Foliar broadcast/ 14.3–15.6	50–110	<0.01	<0.01	No quantifiable residues observed when treated at exaggerated rate

g a.i./ha = grams of active ingredient per hectare

Following the review of all available data, the MRL proposed in Table 1 is recommended, in order to cover residues of pyroxsulam. Dietary risks from exposure to residues of pyroxsulam in these crop commodities at the proposed MRL were shown to be acceptable for the general population and all subpopulations, including infants, children, adults and seniors. Thus, the foods that contain residues as listed in Table 1 are considered safe to eat.

References

None.