

# Amended Recovery Strategy for the Spotted Owl *caurina* subspecies (*Strix occidentalis caurina*) in Canada

## Spotted Owl *caurina* subspecies



2025



Government  
of Canada

Gouvernement  
du Canada

Canada

**Recommended citation:**

Environment and Climate Change Canada. 2025. Amended Recovery Strategy for the Spotted Owl *caurina* subspecies (*Strix occidentalis caurina*) in Canada. *Species at Risk Act Recovery Strategy Series*. Environment and Climate Change Canada, Ottawa. x + 53 pp.

**Official version**

The official version of the recovery documents is the one published in PDF. All hyperlinks were valid as of date of publication.

**Non-official version**

The non-official version of the recovery documents is published in HTML format and all hyperlinks were valid as of date of publication.

For copies of the recovery strategy, or for additional information on species at risk, including the Committee on the Status of Endangered Wildlife in Canada (COSEWIC) Status Reports, residence descriptions, action plans, and other related recovery documents, please visit the [Species at Risk \(SAR\) Public Registry](#)<sup>1</sup>.

**Cover illustration:** © Jared Hobbs

Également disponible en français sous le titre  
« Programme de rétablissement modifié de la Chouette tachetée de la sous-espèce *caurina* (*Strix occidentalis caurina*) au Canada »

© His Majesty the King in Right of Canada, represented by the Minister of Environment and Climate Change, 2025. All rights reserved.

ISBN 978-0-660-75641-7

Catalogue no. En3-4/216-2025E-PDF

*Content (excluding the illustrations) may be used without permission, with appropriate credit to the source.*

---

<sup>1</sup> [www.canada.ca/en/environment-climate-change/services/species-risk-public-registry.html](http://www.canada.ca/en/environment-climate-change/services/species-risk-public-registry.html)

## Preface

The federal, provincial, and territorial government signatories under the [Accord for the Protection of Species at Risk \(1996\)](#)<sup>2</sup> agreed to establish complementary legislation and programs that provide for effective protection of species at risk throughout Canada. Under the *Species at Risk Act* (S.C. 2002, c.29) (SARA), the federal competent ministers are responsible for the preparation of recovery strategies for listed Extirpated, Endangered, and Threatened species and are required to report on progress within five years after the publication of the final document on the Species at Risk Public Registry.

The “Recovery Strategy for the Northern Spotted Owl (*Strix occidentalis caurina*) in British Columbia” (Chutter et al. 2004), and two addendums prepared by the Canadian Spotted Owl Recovery Team and Environment Canada, respectively, was originally adopted as the federal recovery strategy for the Spotted Owl *caurina* subspecies, and posted on the Species at Risk Public registry in October 2006 (Environment Canada 2006). An amendment was required because the identification of critical habitat in the 2006 version of the recovery strategy was insufficient to meet the population and distribution objectives, and a schedule of studies required to complete the identification of critical habitat was included.

This Amended Recovery Strategy for the Spotted Owl *caurina* subspecies (*Strix occidentalis caurina*) in Canada includes updated information on the species population, distribution, and threats, and a refined population and distribution objective that addresses the reasons behind the at-risk status of the species, includes short-term recovery statements, and has a clear timeline goal in accordance with the Policy on Survival and Recovery (Government of Canada 2020). The previous objectives from 2006 were developed before these guidelines were in place. The identification of critical habitat has also been updated to include new information and modeling, informed by the updated population and distribution objective.

The Minister of Environment and Climate Change is the competent minister under SARA for the Spotted Owl *caurina* subspecies and has prepared this amended recovery strategy, as per section 37 of SARA. To the extent possible, it has been prepared in cooperation with the province of British Columbia, as per section 39(1) of SARA.

Success in the recovery of this species depends on the commitment and cooperation of many different constituencies that will be involved in implementing the directions set out in this strategy and will not be achieved by Environment and Climate Change Canada or any other jurisdiction alone. All Canadians are invited to join in supporting and implementing this strategy for the benefit of the Spotted Owl *caurina* subspecies and Canadian society as a whole.

This recovery strategy will be followed by one or more action plans that will provide information on recovery measures to be taken by Environment and Climate Change Canada and other jurisdictions and/or organizations involved in the conservation of the species. Implementation of this strategy is subject to appropriations, priorities, and budgetary constraints of the participating jurisdictions and organizations.

The recovery strategy sets the strategic direction to arrest or reverse the decline of the species, including identification of critical habitat to the extent possible. It provides all Canadians with information to help take action on species conservation. When critical habitat is identified, either

---

<sup>2</sup> [www.canada.ca/en/environment-climate-change/services/species-risk-act-accord-funding.html#2](http://www.canada.ca/en/environment-climate-change/services/species-risk-act-accord-funding.html#2)

in a recovery strategy or an action plan, SARA provides a legal framework that enables the protection of that critical habitat.

In the case of critical habitat identified for terrestrial species, including migratory birds, SARA requires that critical habitat identified in a federal protected area<sup>3</sup> be described in the *Canada Gazette* within 90 days after the recovery strategy or action plan that identified the critical habitat is included in the public registry. A prohibition against destruction of critical habitat under ss. 58(1) will apply 90 days after the description of the critical habitat is published in the *Canada Gazette*.

For critical habitat located on federal lands that are not a federal protected area, as in SARA ss. 58(2), the competent minister must make an order applying the ss. 58(1) prohibition against destruction of critical habitat if it is not already legally protected by a provision in, or measure under, SARA or any other Act of Parliament. If the competent minister does not make the order, a statement must be included on the Species at Risk Public Registry setting out how the critical habitat, or portions of it are legally protected on those federal lands.

If there are portions of critical habitat of a migratory bird to which the following applies:

- 1) Habitat to which the Migratory Birds Convention Act, 1994 applies, AND
- 2) Not on federal land, within the exclusive economic zone, or on the continental shelf of Canada, AND
- 3) Not within a migratory bird sanctuary

SARA requires that the Minister recommend that the Governor in Council make an order to prohibit destruction of critical habitat, if the competent minister forms the opinion that there are no provisions in, or measures under, SARA or other Acts of Parliament that legally protect them. If the competent minister does not make the recommendation, a statement must be included on the Public Registry setting out how those portions of critical habitat for the migratory bird are legally protected. Spotted Owl is not a migratory bird under the Migratory Birds Convention Act, 1994.

For any part of critical habitat located on non-federal lands, if the competent minister forms the opinion that any portion of critical habitat is not protected by provisions in or measures under SARA or other Acts of Parliament, or the laws of the province or territory, SARA requires that the Minister recommend that the Governor in Council make an order to prohibit destruction of critical habitat. The discretion to protect critical habitat on non-federal lands that is not otherwise protected rests with the Governor in Council.

---

<sup>3</sup> These federal protected areas are: a national park of Canada named and described in Schedule 1 to the *Canada National Parks Act*, The Rouge National Park established by the *Rouge National Urban Park Act*, a marine protected area under the *Oceans Act*, a migratory bird sanctuary under the *Migratory Birds Convention Act, 1994* or a national wildlife area under the *Canada Wildlife Act* see ss. 58(2) of SARA.

## Acknowledgments

Many people are to be acknowledged for their involvement in the federal recovery planning process for the Spotted Owl *caurina* subspecies. Development of this recovery strategy was coordinated by Megan Harrison and Kella Sadler (Environment and Climate Change Canada, Canadian Wildlife Service - Pacific Region (ECCC-CWS-PAC). Greg Rickbeil, Danielle Yu, and Leon McCarthy (ECCC-CWS-PAC) provided geospatial modelling and mapping support. This amended recovery document borrows significantly from the original Recovery Strategy for the Northern Spotted Owl (*Strix occidentalis caurina*) in British Columbia (Chutter et al. 2004), which was adopted as part of the federal Recovery Strategy for the species in 2006. All those involved in the development of that original document are gratefully acknowledged. In addition, thanks are owed to Jared Hobbs and Glenn Sutherland (independent experts); Joe Buchanan (Washington Department of Fish and Wildlife); Ian Blackburn, Nicola Bickerton, Joel Gillis (B.C. Ministry of Water, Land and Resource Stewardship [WLRS]); and Louise Waterhouse (B.C. Ministry of Forests) who provided expert input into this updated document. Helpful review comments were also provided by numerous individuals from the federal and provincial governments, First Nations, and members of the public, on earlier versions of this document including proposed versions posted on the Species at Risk Public Registry in January 2023 and July 2024.

## Executive Summary

This document builds on the original Recovery Strategy for the Northern Spotted Owl (*Strix occidentalis caurina*) in British Columbia (Chutter et al. 2004), which was adopted as part of the federal Recovery Strategy for the species in 2006. The 2006 document contains more comprehensive information on the species' life history and early recovery measures, so it should be consulted for background. Detailed planning/strategy documents published since the original recovery strategy (e.g., Sutherland et al. 2007; Fenger et al. 2007) should also be consulted for additional background.

The Spotted Owl *caurina* subspecies (henceforth, the Spotted Owl) is a medium-sized owl with dark brown plumage patterned by small pale spots over most of the body. The species was first assessed by the Committee on the Status of Endangered Wildlife in Canada (COSEWIC) as Endangered in 1986. Its status was re-assessed and re-confirmed in 1999, 2000, and 2008. The species was designated as Endangered due to catastrophic population decline (driven by habitat loss and fragmentation and competition from the closely-related Barred Owl [*Strix varia*]), severely depressed population size, and continued vulnerability to ongoing threats.

The Spotted Owl once occurred throughout mixed-coniferous old-growth forests in southwestern British Columbia (B.C.) and may have numbered as many as 500 pairs prior to the impacts of significant human activity. Its historical range spans three ecological sub-regions that differ in their mean annual precipitation and corresponding habitat characteristics: the wet 'Maritime', moist 'Sub-maritime', and dry 'Continental'. Both the population and the distribution of the Spotted Owl have declined precipitously from historical estimates, with only one wild-born individual detected in its natural habitat in 2022. As of March 2024, there are 37 known Spotted Owls in British Columbia: the wild-born individual, two owls released from a breeding facility into the wild in July 2023, and 34 owls in a breeding facility.

Throughout its range, the Spotted Owl is strongly associated with mixed-coniferous forests that are characterized by: an uneven-aged cohort of trees; a multi-layered, relatively closed canopy; numerous large trees with broken tops, deformed limbs, and large cavities; and numerous large snags and accumulations of logs and downed woody debris. The full set of features and attributes needed to support all life functions (nesting, roosting, foraging, and safe movement/dispersal) are most typically associated with old-growth forests. Mature forests more often contain only a subset of these attributes, which may for example support foraging and safe movement/dispersal, but not other life functions such as nesting.

The primary threats to the Spotted Owl in B.C. include: problematic native species (i.e., competition from Barred Owls), logging and wood harvesting, roads and railroads (including logging roads), and fire and fire suppression.

The population and distribution objective is to recover the Spotted Owl in Canada by restoring a stable population of at least 250 mature individuals distributed within a connected network of habitat representative of all three sub-regions within the species' historical Canadian range, and linked to the larger population in the U.S.A.

Recognizing that the population and distribution objective will take >50 years to achieve, the following short-term recovery statements toward meeting the population and distribution objective have been established:

1. Immediately mitigate human-caused threats that would cause loss of habitat needed for recovery (i.e., the critical habitat).
2. By 2030, re-establish a wild Spotted Owl population in British Columbia of four to six resident individuals, utilizing individuals bred from the Northern Spotted Owl Breeding Program.
3. Complete regular Barred Owl surveillance at sites occupied by Spotted Owl and/or where reintroductions are planned and remove all Barred Owls that are detected.

Broad strategies and general approaches toward addressing the primary threats to the survival and recovery of the species, as well as key knowledge gaps, are presented in section 6. Successful implementation of these broad strategies and approaches will be required for the population and distribution objective to be met.

Critical habitat has been identified to the extent possible, based on the best available information for the Spotted Owl. It is recognized that the acoustic critical habitat<sup>4</sup> identified is insufficient to achieve the population and distribution objectives for the species. A schedule of studies (Section 7.2) has been developed to provide the information necessary to complete the identification of acoustic critical habitat that will be sufficient to meet population and distribution objectives.

Three performance indicators were developed to measure progress towards meeting the population and distribution objective. One or more action plans for the Spotted Owl will be posted on the Species at Risk Public Registry within five years of the final posting of the recovery strategy.

---

<sup>4</sup> One of the two variations of critical habitat identified within this document. Acoustic critical habitat is the critical habitat surrounding nesting areas that functions to maintain the acoustic environment within those areas during the breeding season. See Section 7.1.

## Recovery Feasibility Summary

Based on the following three criteria that Environment and Climate Change Canada uses to establish recovery feasibility, it is considered to be biologically and technically feasible to recover the Spotted Owl in Canada (see *Species at Risk Policy on Recovery and Survival* [Environment and Climate Change Canada 2021]), although there are some significant uncertainties associated with this determination. In keeping with the *Guidelines on Characterizing Recovery and Developing Population and Distribution Objectives* under the *Species at Risk Policy on Recovery and Survival* (Environment and Climate Change Canada 2021), where there is a range of uncertainty associated with the full extent of improvements that are biologically and technically feasible, the determination defaults to the upper bound of what is considered to be within the scope of biological and technical feasibility.

- 1. Survival characteristics:** Can survival characteristics be addressed to the extent that the species is no longer at significantly greater risk of extinction or extirpation as a result of human activity?

**YES, with uncertainty:** The Spotted Owl is currently assessed as Endangered on the basis of four key survival characteristics: (i) resilience (D1 COSEWIC quantitative criteria<sup>5</sup>) – its wild population is estimated to be very small (well below the 250-individual threshold for Endangered status) and is in decline; (ii) redundancy and connectivity (linked with COSEWIC B2ab indicators) – its habitat and associated distribution is in decline and fragmented (iii) stability (linked with COSEWIC A2ac, C1+2a, and E indicators) - there is an ongoing decline in the number of mature individuals and area/quality of habitat, and a quantitative analysis showing high probability of extirpation, and (iv) continuing impacts caused by ongoing human-caused threats.

- Resilience: Prior to impacts from human activity (i.e., in its natural condition), the Spotted Owl population in Canada may have been as many as 500 breeding pairs (Blackburn et al. 2002). This small population size would have meant the species' persistence was somewhat precarious even in its natural condition (e.g., would still be assessed as Threatened under the COSEWIC D1 quantitative criteria, which apply to a species with <1000 mature individuals). However, the results of human activity have put the species at a significantly greater risk of extirpation, such that it is now assessed as Endangered on the D1 indicator (i.e., population below 250-individuals). For recovery to be considered feasible, it must be biologically and technically feasible to improve the resilience of the Spotted Owl such that it exceeds the 250-individual D1 threshold associated with Endangered status, and thus returns to a status of Threatened (on the basis of D1 criteria).

As of mid-2022, only one wild-born female owl was thought to remain in its natural habitat (Government of B.C. 2022). The status of that owl was unknown as of October 2024. The Spotted Owl is known to suppress its calling in the presence of the closely related competitor, the Barred Owl (*Strix varia*; Kelly et. al 2003; Crozier et al. 2006; Van Lanen et al. 2011; Yackulic et al. 2019) and Barred Owls have been detected at all 10 sites surveyed in 2020 that were previously occupied by Spotted Owls, so it is possible that some owls are present at surveyed sites, potentially including the wild-

---

<sup>5</sup> For COSEWIC quantitative criteria and guidelines see [www.cosewic.ca/index.php/en-ca/assessment-process/wildlife-species-assessment-process-categories-guidelines/quantitative-criteria](http://www.cosewic.ca/index.php/en-ca/assessment-process/wildlife-species-assessment-process-categories-guidelines/quantitative-criteria).

born female that was last detected in 2022, but are going undetected by standard call-playback survey methods.

Comprehensive, range-wide surveys have also not been undertaken in recent years, so some owls may still exist in un-surveyed areas, although given known rates of juvenile survivorship and recruitment this seems unlikely. However, even accounting for uncertainties about undetected individuals, the wild population is extremely small and would have correspondingly low genetic diversity. The population is apparently incapable of recovering on its own, so resilience cannot be addressed without the reintroduction of owls from a breeding program (Fenger et al. 2007).

A Spotted Owl breeding and reintroduction program has been in operation in B.C. since 2007 (Northern Spotted Owl Breeding Program n.d., Government of B.C. 2021). The first release occurred in August 2022, when three facility-bred male owls were released into two Wildlife Habitat Areas; one was later found injured and returned to the breeding facility, and the other two died of unknown causes (Government of B.C. 2023a). In late June 2023, the previously injured owl and one additional male were released (Government of B.C. 2023b). As of March 2024, there were 27 adults and seven chicks born in 2023 at the breeding facility (McCulligh 2024). Based on minimum targets<sup>6</sup> of releasing ~4 individuals/year from 2023 to 2024, ~9 individuals/year from 2025-2030, and ultimately ~14 individuals/year thereafter, the provincial government's draft release plan projected that it is within the scope of biological and technical feasibility to restore a stable population of  $\geq 250$  mature individuals within 50 years (B.C. MFLNRORD 2021). The release plan is being adaptively managed following review of the results of the first two years (Government of B.C. 2023b).

- Redundancy, connectivity, and stability: Prior to impacts from human activity (i.e., in its natural condition), the Spotted Owl had a relatively restricted Canadian distribution, concentrated in southwestern B.C. Although the precise bounds of its historical range (including extent of occurrence and area of occupancy) are unknown, it would have included three ecologically distinct sub-regions (the wet 'Maritime', moist 'Sub-maritime', and dry 'Continental' sub-regions), with connectivity in habitat within these sub-regions and to the U.S.A, to support a stable and genetically diverse population. With connectivity in habitat, and stability in population and distribution characteristics, it is unlikely that any of the COSEWIC quantitative criteria associated with redundancy, fragmentation, and/or stability for assessment as Endangered would have been met for the species in its natural condition. For recovery to be considered feasible, it must be biologically and technically feasible to reverse the declines in population and distribution characteristics, and to ensure there is a connected network of habitat that will support at least 250 mature individuals. Through breeding and reintroduction (see above) and threat mitigation, it is considered to be within the scope of biological and technical feasibility to restore a stable population. Through habitat protection and threat mitigation, it is considered to be within the scope of biological and technical feasibility to ensure that there are no further habitat declines. Based on the configuration of regenerating and existing Spotted Owl habitat (see section 7 - Critical Habitat), it is considered to be within the scope of biological and technical feasibility to achieve a connected network of habitat sufficient to support  $\geq 250$  mature individuals within 50 years. Connectivity to the U.S.A. will always be

---

<sup>6</sup> This number is derived from projections by the provincial government (B.C. MFLNRORD 2021) but is subject to adjustment following the pilot phase of the reintroduction (2022-2025), based on the actual annual reproductive output of pairs in the breeding facility and the survival outcomes of released individuals.

reduced relative to historical conditions due to permanent habitat loss within developed portions of the Lower Mainland and Fraser Valley; however, some connectivity still exists and/or can be restored through long-term protection of maturing forest. In its recovered condition, the Spotted Owl would not meet the quantitative criteria for assessment as Endangered or Threatened based on A,B,C, or E COSEWIC indicators (see Table 2 in COSEWIC 2021).

- Protection from human-caused threats: There are ongoing human-caused threats that must be addressed (ceased, mitigated, or avoided) in order for the preceding key survival characteristics to be met and for recovery to be feasible. The most significant ongoing human-caused threats include: competition with a problematic native species (i.e., the closely-related Barred Owl); habitat impacts caused by logging and wood harvesting, roads and railroads (including logging roads), and fire and fire suppression. It is biologically and technically feasible to cease or mitigate the human-caused threats of logging, road-building; and fire / increased fuel loads resulting from fire suppression. Barred Owls are now considered to be one of the highest-level threats to the Spotted Owl both in B.C. and range-wide in North America (section 4 of this document; USFWS 2011) and may currently pose a greater biological and technical challenge than habitat loss and availability. Barred Owl control programs (translocation and lethal removal) have been initiated within both the U.S.A. and B.C. (Diller et al. 2016, Gillis and Waterhouse 2020, Wiens et al. 2021, USFWS 2024) as an important component of Spotted Owl conservation efforts. Post-treatment monitoring studies in the U.S.A. have shown increases in local Spotted Owl site occupancy, survivorship and productivity within 4.5 years of (lethal) Barred Owl removal (Diller et al. 2016; Wiens et al. 2021, Hofstadter et al. 2022). In more northern study areas there were longer lag times before reduced Barred Owl colonization rates and increased Spotted Owl responses were measured (Wiens et al. 2021). The longer lag times in northern sites have been attributed to two potential causes: 1) more established Barred Owl populations in these areas, and 2) lower numbers of Spotted Owls available to recolonize empty territories (Diller et al. 2016; Yackulic et al. 2019). Discernable Spotted Owl responses to Barred Owl removals have not yet been reported in Canada (Gillis and Waterhouse 2020); however, the planned re-introduction of Spotted Owls could help improve re-colonization rates. While there is uncertainty, it is still considered to be within the scope of biological and technical feasibility that impacts of Barred Owl can be managed successfully, to the extent that the preceding survival characteristics can be addressed.

- 2. Independence:** Is the species currently able to persist in Canada independent of deliberate human interventions, and/or will it eventually be able to achieve and maintain independence in the state where condition (1) is met, such that it is **not reliant on significant, direct, ongoing human intervention?**

**YES, with uncertainty.** The Spotted Owl is currently nearing extirpation in Canada and requires significant, direct human interventions (i.e., population augmentation through the Northern Spotted Owl Breeding Program, and Barred Owl control) in the short-to-medium term (i.e., within the next 20 years), in order for condition '1' of recovery feasibility to be met. Barred Owl control is the primary intervention that may need to continue for a longer period (i.e., beyond 20 years). Although there is uncertainty, habitat improvement/recovery is expected to improve Spotted Owl persistence in combination with Barred Owl control and may help reduce the necessary level of investment in Barred Owl removals in the future (Yackulic et al. 2019). The release of facility-bred owls into the wild began in 2022, and the

continued success of facility-bred owls after release is not well understood. A period of learning and adaptive management is expected. Although there is a high level of uncertainty, it is considered to be within the scope of biological and technical feasibility that a point will be reached in the longer term (up to 50 years), where the Spotted Owl population has recovered such that it can remain stable in the absence of ongoing human interventions.

**3. Improvement:** Can the species' condition be improved over when it was assessed as at risk?

**YES, with uncertainty.** It is biologically and technically feasible to meaningfully improve the condition of the Spotted Owl in Canada through addressing one or more key survival characteristics (as they pertain to results of human activity) such that the species' risk of extinction or extirpation is reduced. Population stability and resilience may be improved, and population/habitat connectivity and redundancy restored, through a) ensuring a connected network of habitat, so that the habitat needed to support all life functions for a population of >250 mature individuals is available on the landscape when the recovering/recovered population needs it; b) continuing the breeding and reintroduction program, so that protected habitat is repopulated, and c) continuing Barred Owl control efforts, so that Spotted Owls can survive and reproduce successfully within protected habitats.

## Table of Contents

Preface.....	i
Acknowledgments .....	iii
Executive Summary .....	iv
Recovery Feasibility Summary.....	vi
1. COSEWIC Species Assessment Information.....	1
2. Species Status Information .....	1
3. Species Information .....	2
3.1 Species Description .....	2
3.2 Species Population and Distribution .....	2
3.2.1 Population.....	2
3.2.2 Distribution.....	3
3.3 Needs of the Spotted Owl .....	7
4. Threats.....	13
4.1 Description of Threats.....	15
5. Population and Distribution Objectives.....	21
6. Broad Strategies and General Approaches to Meet Objectives .....	22
6.1 Actions Already Completed or Currently Underway .....	22
6.2 Strategic Direction for Recovery .....	25
7. Critical Habitat.....	27
7.1 Identification of the Species' Critical Habitat .....	27
7.1.1 Information and methods used to identify critical habitat .....	30
7.1.2 Geospatial Location of Areas Containing Critical Habitat.....	32
7.2 Schedule of Studies to Identify Critical Habitat .....	39
7.3 Activities Likely to Result in the Destruction of Critical Habitat.....	40
8. Measuring Progress .....	41
9. Statement on Action Plans .....	41
10. References .....	42
11. Personal Communications.....	52
Appendix A: Effects on the Environment and Other Species .....	53

## 1. COSEWIC\* Species Assessment Information

**Date of Assessment:** April 2008

**Common Name (population):** Spotted Owl *caurina* subspecies

**Scientific Name:** *Strix occidentalis caurina*

**COSEWIC Status:** Endangered

**Reason for Designation:** This owl requires old-growth forests for its survival and has suffered a catastrophic population decline over the past 50 years as habitat is lost and fragmented. With the severely depressed population, an additional threat is the recent arrival of the closely related Barred Owl as a breeding bird in B.C.; this species competes with and hybridizes with the present species. Its historical population of about 500 adult owls in Canada has been reduced to 19, and only 10 of these are in breeding pairs. All adults are old and near the end of their breeding age and there is no recruitment of young owls into the population. If current trends are not reversed, extirpation will likely occur within the next decade.

**Canadian Occurrence:** British Columbia

**COSEWIC Status History:** Designated Endangered in April 1986. Status re-examined and confirmed in April 1999, May 2000, and April 2008.

\* COSEWIC (Committee on the Status of Endangered Wildlife in Canada)

The above summary reflects population status information as of the 2008 COSEWIC assessment. Since 2008 there has been new information about historical and current population levels (summarized in section 3.2 – species population and distribution).

## 2. Species Status Information

The legal designation for the Spotted Owl on SARA Schedule 1 is Endangered (2003). Approximately 8% of the global (historical) range of the Spotted Owl is located in Canada (COSEWIC 2008). The species' status ranks, globally and in the different parts of its range, are summarized in Table 1.

**Table 1.** List and description of various conservation status ranks for the Spotted Owl (*caurina* subspecies) (NatureServe 2021).

Global (G) Rank	National (N) Rank	Sub-national (S) Rank	COSEWIC Status
Rounded global rank (of G3G4T3) = T3 (vulnerable)	Canada (N1 – critically imperiled) United States (N3 - vulnerable)	BC (S1) California (S2) Oregon (S1S2) Washington (S1)	EN (Endangered)

S1: Critically Imperiled; S2: Imperiled; S3: Vulnerable; S4: Apparently Secure; S5: Secure; SNR: Unranked; SNA: Not Applicable; B: Breeding.

### 3. Species Information

Spotted Owl, or Northern Spotted Owl, are English names given to the species. First Nations have other names for the species that reflect the diversity of Indigenous languages throughout its historic range.

Environment and Climate Change Canada recognizes that this recovery strategy does not currently reflect the deep cultural and spiritual significance of the species to some Indigenous peoples. We welcome future collaborations that respectfully explore other languages, and the connections between these owls, their relatives in human form, and the broader environment in the context of describing additional perspectives on the species, and to inform ongoing planning and implementation of recovery actions.

#### 3.1 Species Description

The Spotted Owl is a medium-sized owl averaging 45 cm in length and 90 cm in wingspan. Plumage is dark overall with brown feathers patterned by small pale spots over most of the body. The tail has narrow white horizontal bars and there are no “ear” tufts. Eyes are large, dark brown and are set within lighter brown facial disks (Forsman 1981; Gutiérrez et al. 1995). Age cohorts can be identified by differences in plumage characteristics. Juveniles <5 months old are identified by visible down feathers. Sub-adults (1-2 years old) and adults (>2 years) may be differentiated based on tail feathers; sub-adults have pointed tail feathers with white tips whereas adult tail feathers are rounded and usually mottled in colour (Forsman 1981). Males and females have similar plumage but females are ~15% larger (Blakesley et al. 1990; Gutiérrez et al. 1995).

#### 3.2 Species Population and Distribution

##### 3.2.1 Population


The global population of the Spotted Owl was estimated at roughly 6000 breeding pairs in the late 1980s (Thompson et al. 1990), with the bulk of the population (>90%) occurring in the U.S.A. (COSEWIC 2008). Local population declines were observed at demographic study areas within Washington, Oregon, and California between 1985 and 2013, with an overall mean annual rate of decline of 3.8% (Dugger et al. 2015). Although no formal global population estimates have been published in recent decades, an approximate current population estimate can be deduced using the annual rates of decline observed in long-term study areas; assuming a 6000-pair starting population and a 3.8% mean annual decline from 1985-2021, the global population would now be <1500 pairs. An updated analysis of the long-term study area data including data from 1995 to 2017 (Franklin et al. 2021) suggests that declines may have become even sharper than the 3.8% mean decline reported in Dugger et al. (2015), with some long-term study sites exhibiting mean annual declines as high as 9%. Declines have been most pronounced in Washington, Oregon and B.C., and less pronounced in California (Blackburn and Godwin 2003; Dugger et al. 2015; Franklin et al. 2021).

Before European settlement, the Canadian Spotted Owl population likely did not exceed 500 breeding pairs, or ~10% of the global population (Blackburn et al. 2002). In 1991, it was estimated at fewer than 100 potential breeding pairs (Dunbar et al. 1991; Dunbar and Blackburn 1994) and by 2002 it had declined further to fewer than 33 (Blackburn and Godwin 2003). A

survey of 10 previously-occupied sites in 2020 found one pair and one single owl at two sites (J. Gillis pers. comm. 2020). Continued occupancy of those two sites was reconfirmed in 2021 (J. Gillis pers. comm. 2021). As of mid-2022, only one female owl was thought to remain in the wild (Government of B.C. 2022). This suggests a decline of over 99% from estimated historical levels in Canada, with Canada now supporting <0.01% of the global (combined Canada and U.S.A.) population.

As outlined in the Recovery Feasibility Summary, a Spotted Owl breeding and reintroduction program has been in operation in B.C. since 2007. The first release occurred in August 2022, when three facility-bred male owls were released into two Wildlife Habitat Areas; one was later found injured and returned to the breeding facility, and the other two died of unknown causes (Government of B.C. 2023a). In late June 2023, the previously injured owl and one additional male were released (Government of B.C. 2023b). As of September 2023, there were 37 known Spotted Owls in B.C.: one wild-born owl last confirmed in the wild during surveys in 2022, the two facility-bred owls released in 2023, and 34 owls in the breeding facility (27 adults and seven chicks born in 2023) (Government of B.C. representatives pers. comm. 2023)


The combined wild and facility-bred population was relatively static between 2004 and 2020, where declines in the wild population in part resulted from individuals periodically being taken into the Northern Spotted Owl Breeding Program (Figure 1).



**Figure 1.** Known Spotted Owl population in Canada from 2004 to 2020 (Government of B.C. 2020). Note that annual inventory effort has varied for wild population counts.

### 3.2.2 Distribution

The Spotted Owl *caurina* subspecies is one of three subspecies of Spotted Owls found within North America (Figure 2). The *caurina* subspecies is distributed from the southwest mainland of B.C. through western Washington, western Oregon and the west coast of California, south to San Francisco Bay.



**Figure 2.** Approximate historical year-round range of Spotted Owls (three subspecies) in North America. (BirdLife International 2018). The *caurina* subspecies is often also referred to as the Northern Spotted Owl.

Historically, the Spotted Owl's range in B.C. extended from the U.S.A. border north ~200 km to Carpenter Lake, and ~160 km from Howe Sound in the west to the Cascade Range in the east (Figure 3; Chutter et al. 2004). Within this range, there are three ecological sub-regions that differ in their mean annual precipitation and corresponding habitat characteristics: the wet 'Maritime', moist 'Sub-maritime', and dry 'Continental'. Permanent range contraction occurred within the Lower Mainland and Lower Fraser Valley where once suitable habitat has been irreversibly lost to human development (Chutter et al. 2004; Figure 3); however, habitat remains within the rest of the historical range and could potentially be re-occupied by Spotted Owls. The remaining known wild individual (as of 2022) was found within the Sub-maritime sub-region.




Figure 3. Approximate historical distribution of the Spotted Owl *caurina* subspecies in B.C.

### 3.3 Needs of the Spotted Owl

Historically, Spotted Owls occurred primarily within the Coastal Western Hemlock (CWH) and Interior Douglas Fir (IDF) biogeoclimatic zones in B.C. (SOMIT 1997; Sutherland et al. 2007). Their historical range would also have included parts of the Coastal Douglas Fir (CDF) zone, although that habitat has been permanently lost to human development (Chutter et al. 2004; Sutherland et al. 2007; Figure 3). The species is associated with mixed-coniferous forests characterized by: an uneven-aged cohort of trees; a multi-layered, relatively closed canopy; numerous large trees with broken tops, deformed limbs, and large cavities; and, numerous large snags and accumulations of logs and downed woody debris (Thomas et al. 1990; USDI 1992). These old-growth forests have the potential to buffer local warming, and thus function as local refugia for species reliant on cooler conditions (Spies et al. 2018; Dinerstein et al. 2019).

#### Habitat configuration

##### *Landscape-level configuration*

In order for a stable Spotted Owl population to exist within a landscape, habitat must be configured such that it can support all critical life functions (breeding, roosting, foraging, and safe movement/dispersal) for the entire population. This requires patches of forested habitat capable of supporting the year-round needs of breeding pairs and resident individuals, as well as an overall configuration of both the year-round habitat patches and seasonally-used dispersal habitat that maximizes survival/success of dispersing individuals. Juvenile Spotted Owls disperse from their natal site in the late summer / early fall of their first year, and then may disperse several more times and persist in the background as “floaters” for up to five years before settling and beginning to breed (Forsman et al. 2002). Breeding-age owls also occasionally disperse to new locations, particularly when their original location has been disturbed (Forsman et al. 2002; Jenkins et al. 2019; Jenkins et al. 2021).

Simulated landscape population modelling results for the species suggest that larger year-round habitat clusters are most likely to support stable long-term (100-year) occupancy by Spotted Owls due to the ability for individuals to disperse within their natal/original cluster versus having to leave their natal cluster and disperse across a less hospitable matrix (Lamberson et al. 1994; Marcot et al. 2013). Marcot et al. (2013) also found that clusters were more likely to exhibit stable long-term occupancy by Spotted Owls when they were spaced more closely together (<15 km). Being a highly-mobile species, Spotted Owls are capable of dispersing long distances. In an analysis of 1534 dispersal events in Oregon and Washington, Hollenbeck et al. (2018) reported a maximum dispersal distance of 177 km and a mean of 23.8 km ( $\pm$  19.2 km standard deviation). Dispersal has also been observed across a range of habitat types. Large non-forested valleys, high-elevation subalpine forest, alpine tundra and large water bodies are the only features suspected to act as complete barriers to dispersal (Forsman et al. 2002; Chutter et al. 2004; I. Blackburn pers. comm. 2021). However, Spotted Owls must feed and escape predation to survive dispersal, and in moving through areas that lack foraging resources and security features, dispersing individuals are expected to incur an increased energetic/survival cost (Lamberson et al. 1994; Buchanan 2004; Sutherland et al. 2007; Marcot et al. 2013; Conlisk et al. 2020). In their simulated landscape population modelling, Marcot et al. (2013) found that as more of the landscape becomes suitable (i.e., overall habitat more contiguous), all cluster size/spacing configuration options become sufficient to achieve low dispersal mortality and high long-term stability. There have been few empirical studies of dispersal habitat use and demographic associations; however, in a study in western Oregon,

Miller (1997) showed that juveniles that used more clearcut areas during dispersal had higher mortality rates than those using more intact forest habitat. Similarly, in their analysis of 1534 successful juvenile dispersal events in the U.S.A., Hollenbeck et al. (2018) found that dispersal pathways tended to coincide with the distribution of forested areas along mountain ranges, as opposed to non-forested areas. Overall population stability is therefore most likely when a landscape includes not just large, closely configured habitat patches to support year-round occupancy, but also habitat occurring in between year-round patches that provides foraging and security opportunities for dispersing birds.

### *Home range-level configuration*

Within suitable landscapes, areas that adult/resident Spotted Owls occupy year-round are represented as home ranges. Home ranges can be occupied by unpaired resident birds, or by a breeding pair. A certain amount of habitat must be present in these areas to support nesting, roosting, and foraging life history functions (as described below). Further, this habitat must not be too fragmented, so that it can be accessed without excess energy expenditure and/or exposure to predation (Carey et al. 1992; Courtney et al. 2004; Sutherland et al. 2007). The mean area of habitat estimated to support a resident Spotted Owl home range varies between sub-regions: Maritime – 3010 ha, Sub-maritime – 2224 ha, Continental – 1907 ha (Chutter et al. 2004; Sutherland et al. 2007). In locations with contiguous mature or old-growth forested habitat, these numbers also represent minimum home range sizes. Home ranges become larger as habitat is more fragmented (Carey et al. 1992). The maximum energetically-viable home range sizes in fragmented landscapes could be more than three times larger than in contiguous habitats, and are estimated at 11,047, 7258, and 6305 ha in the Maritime, Sub-maritime, and Continental sub-regions, respectively (Sutherland et al. 2007). In continuous habitat, adjacent home ranges may overlap up to 25% (Sutherland et al. 2007).

During the breeding season, pairs concentrate their activities within a smaller area of their home range, in close vicinity to the nest grove. In Canada, most breeding season activities are estimated to occur within ~500 m of the nest tree (Blackburn et al. 2009).

### *Patch-level configuration*

Due to a combination of both natural and anthropogenic disturbances, remnant Spotted Owl habitat in Canada exists in a range of patch<sup>7</sup> sizes, from large contiguous expanses to patches <1 ha in size. A patch's size may impact whether it can provide functional habitat for a Spotted Owl. Ten hectares has been estimated by experts within Canada as the minimum habitat patch size within which preferred prey can persist and thus Spotted Owls can successfully forage (reviewed in Sutherland et al. 2007). In addition to absolute size, the irregularity of a patch may also impact its utility for Spotted Owls. Research from Pacific Northwest forests has shown that microclimate (including humidity and solar exposure) can be impacted up to ~100 m from an edge (Kremsater and Bunnell 1999). These impacts may be particularly pronounced for species of fungi and lichens, which are often adapted to the cooler, moister, darker conditions associated with interior forest (Crockatt 2012; Gauslaa et al. 2018). Spotted Owls in Canada feed disproportionately (>40% of diet) on Northern Flying Squirrels (*Glaucomys sabrinus*; Horoupian et al. 2004), which in turn feed preferentially on fungi and lichens associated with coniferous forested habitats (Carey 1991; Carey et al. 1992; Waters and Zabel 1995). In small or highly irregular forest habitat patches with high edge-to-interior ratios, the conditions necessary to sustain foraging resources for Northern Flying Squirrels may not exist.

---

<sup>7</sup> A discrete area of habitat.

Competitors (of Spotted Owls) that are better adapted to foraging within diverse habitats may also over-exploit preferred prey species in openings and along edges, further reducing prey availability for Spotted Owls in small or irregular habitat patches (Wilson and Forsman 2013; Wiens et al. 2014).

## Habitat attributes

### *Nesting*

Spotted Owls do not build their own nests but depend on naturally occurring or previously constructed (by other raptor species) nest sites (Chutter et al. 2004, Waterhouse et al. 2012; Wilk et al. 2018). Nest sites include broken treetops, tree cavities, abandoned raptor nests, mistletoe brooms, and debris accumulations captured in branches (Forsman et al. 1984, Dawson et al. 1986, Waterhouse et al. 2012; Wilk et al. 2018). In the breeding facility, nesting has occurred in artificial cavities (McCulligh 2019, see also Gutierrez et al. 1995). In general, cavities are more often used in moist climates and platforms are more frequently used in drier climates, particularly where cavities in trees >50 cm in diameter are not available (Chutter et al. 2004). In a survey of 14 known nest trees in B.C., Waterhouse et al. (2012) found that nest cavities were in trees averaging 88 cm in diameter ( $\pm 26.8$  cm standard deviation). A variety of different tree species are used for nesting within the species' range although large Douglas-fir may be selected more frequently in the drier regions (Waterhouse et al. 2012; Wilk et al. 2018). In the wetter regions, Western Hemlock and Western Redcedar have been used in equal proportion to Douglas-fir (Forsman and Giese 1997; Wilk et al. 2018). Nest site fidelity<sup>8</sup> is high and re-use of nest structures is common (Forsman et al. 1984).

Breeding Spotted Owls may experience stress, lower reproductive output, and disrupted nesting behaviours when exposed to acute noise within their nesting areas (Wasser et al. 1997, Hayward et al. 2011, USFWS 2020), therefore, in order to successfully carry out breeding functions they also require nesting areas to be free of significant acoustic disturbance during the breeding season. Acoustic disturbance significant enough to impact nesting functions can result from activities that result in an overall sound level above 90 db (e.g., operation of large machinery, use of chainsaws, blasting, operation of large engines and engine brakes, operation of motorized recreational vehicles) or that increase the sound level above ambient conditions by over 20 db (USFWS 2020).

### *Roosting and escape*

Spotted Owls require roosting sites that provide good protective cover from both inclement weather and predators. The multi-storied nature and high percentage canopy closure of old-growth forests enables thermoregulation and escape from inclement weather, as well as providing protection from predators (Blackburn et al. 2009). The Spotted Owl is easily subjected to heat stress and reduces its exposure by moving between roosting habitats in different parts of the canopy (Barrows 1981). The closed canopies of old-growth habitats also provide refuge from rain and snow (North et al. 2000). Great Horned Owls (*Bubo virginianus*) are the primary predator of the Spotted Owl (Gutierrez et al. 1995), and favour edge habitats and openings where they have greater access to prey (Artuso et al. 2013). Susceptibility to Great Horned Owl predation may thus be minimized in areas with intact, contiguous old-growth/mature forest (Johnson 1993).

---

<sup>8</sup> Tendency to return to the same nesting location and/or re-use the same nest structure in subsequent years.

### *Foraging*

Spotted Owls require habitat with characteristics that promote abundant and accessible prey, which is primarily comprised of arboreal and semi-arboreal small mammals (Chutter et al. 2004; Wiens et al. 2014). Studies in western Washington and B.C. showed that Northern Flying Squirrels, Bushy-tailed Woodrats (*Neotoma cinerea*), and Deer Mice (*Peromyscus* sp.) were the most common prey for Spotted Owls in the northern extent of the Spotted Owl's range (Forsman et al. 2001; Horoupian et al. 2004; Wiens et al. 2014). The abundant large coarse woody debris (CWD), standing snags, and diverse shrub layers present within old-growth forests support prey populations by providing moist microclimates, protective cover for movement, nest/burrow sites, and food in the form of fungi, plants and invertebrates (Carey 1991; Carey et al. 1997; Carey et al. 1999; Wilson and Forsman 2013). The open mid-storey structure of old-growth habitats also enables Spotted Owls to have more efficient access to those prey through providing longer sightlines and unimpeded flight paths (Chutter et al. 2004; D'Anjou et al. 2015).

### *Safe movement / dispersal*

Like resident Spotted Owls, dispersing individuals require available prey and security features, therefore, old-growth and mature forests (i.e., the same forests that support nesting/roosting and foraging) are understood to provide ideal conditions (reviewed in Buchanan 2004). Where no habitat capable of supporting foraging and security exists between two natal patches, dispersal success between those patches is likely to be reduced, ultimately reducing long-term patch occupancy and population stability. Safe movement/dispersal is best-supported by nesting/foraging quality habitat located either within year-round forested habitat patches (enabling within-patch dispersal) or in between those patches. Spotted Owls may traverse forested habitat in other seral stages during dispersal; however, it is not yet clear what other habitat attributes/configurations may contribute to dispersal success (Buchanan 2004). Research will be required to evaluate drivers of dispersal success in Canada and determine whether additional habitats should be identified as important for supporting safe movement.

### **Distribution of competitors**

In addition to habitat amount, quality and configuration, the distribution and abundance of the Spotted Owl's primary competitor, the Barred Owl, has been shown to strongly influence Spotted Owl occupancy across the landscape (Dugger et al. 2011; Dugger et al. 2015; Yackulic et al. 2019; Jenkins et al. 2019). Barred Owls reduce Spotted Owl occupancy of otherwise suitable habitat through both competition for prey (exploitative competition) and territorial displacement (Dugger et al. 2011; Wiens et al. 2014; Jenkins et al. 2021). See Section 4 (Threats) for more details.

## Classification of habitat for the Spotted Owl

The Vegetation Resource Inventory<sup>9</sup> (VRI) geospatial database provides detailed information about the characteristics of forests in B.C. The VRI attributes used to classify forests as potentially suitable for the Spotted Owl in B.C. (not accounting for configuration considerations) are summarized in Table 2. 'Nesting' quality habitat is characterized by old, tall, low elevation stands, and 'foraging' quality habitat is characterized by mature, moderately tall stands that may extend further upslope. Both 'nesting' and 'foraging' quality habitats are considered to have characteristics that also support roosting and safe movement / dispersal. Nesting quality habitat is used disproportionately relative to its availability on the landscape, whereas foraging quality habitat is used in proportion with its availability on the landscape (Forsman et al. 1984; Carey et al. 1990; Carey et al. 1992).

The provincial government is also continuing to develop and refine habitat classification approaches as part of its Stewardship Baseline Objectives Tool (Government of B.C. 2021).

---

<sup>9</sup> A photo-based, two-phased vegetation inventory design consisting of 1) 1:20,000-scale air photo interpretation and 2) ground sampling. See <https://www2.gov.bc.ca/gov/content/industry/forestry/managing-our-forest-resources/forest-inventory>.

**Table 2.** Summary of attributes used to classify forests as potentially suitable for the Spotted Owl in B.C. using the Vegetation Resource Inventory (VRI) geospatial database (from Sutherland et al. 2007). Note that this does not account for habitat configuration, competitor distribution, or the locations where facility-bred Spotted Owls will be released, which ultimately determine the likelihood that habitat will support recovery of the Spotted Owl.

Function / Class	Attribute	VRI polygon-level selection thresholds					
		Maritime sub-region (CWHdm, CWHvm1&2, CWHxm1)*		Sub-maritime sub-region (CWHds1, CWHms1, IDFww)		Continental sub-region (IDFdc, IDFdk1,2&3, IDFww1, IDFxc, IDFxh1&2, PPxh2)	
		Structure present**	Structure absent**	Structure present	Structure absent	Structure present	Structure absent
Nesting, roosting and safe movement	Stand age	≥ 140 years	≥ 200 years	≥ 110 years	≥ 200 years	≥ 110 years	≥ 200 years
	Stand height	≥ 28.5 m		≥ 23 m		≥ 23 m	
	Elevation	≤ 900 m		≤ 1000 m		≤ 1200 m***	
Foraging, roosting and safe movement	Stand age	≥ 120 years	≥ 140 years	≥ 100 years	≥ 120 years	≥ 80 years	≥ 100 years
	Stand height	≥ 19.5 m		≥ 19.5 m		≥ 19.5 m	
	Elevation	No limit, other than BEC		No limit, other than BEC		No limit, other than BEC	

\*Biogeoclimatic Ecosystem Classification (BEC) zones and variants within which selection occurred. Note: re-mapping of BEC variants in the Continental sub-region since 2004 has resulted in some additions/deletions to the selected variants from the Sutherland et al. (2007) version. For descriptions and definitions see: <https://www.for.gov.bc.ca/hre/becweb/resources/classificationreports/index.html>

\*\*This distinction is relevant to future projections only, in determining whether a stand that was previously harvested will have the structural characteristics of nesting and foraging class habitat within the 50-year recovery timeframe. Stands that were of natural disturbance origin or that were harvested prior to the advent of clear cut harvesting are assumed to have remnant old forest structure present, and so are expected to have all the attributes required to support Spotted Owl nesting and/or foraging at a younger age. In comparison, stands harvested since the advent of clearcut harvesting will not have old forest structure remaining, so will take longer to re-acquire these characteristics.

\*\*\*Increased from 1100 m (Sutherland et al. 2007) to 1200 m to accommodate nests found more recently >1100 m in the Continental sub-region (Hobbs 2004).

## 4. Threats

The Spotted Owl threat assessment is based on the IUCN-CMP (World Conservation Union–Conservation Measures Partnership) unified threats classification system. Threats are defined as the proximate activities or processes that have caused, are causing, or may cause in the future the destruction, degradation, and/or impairment of the entity being assessed (population, species, community, or ecosystem) in the area of interest (global, national, or subnational). Limiting factors are not considered during this assessment process. For purposes of threat assessment, only present and future threats are considered. Historical threats, indirect or cumulative effects of the threats, or any other relevant information that would help understand the nature of the threats are presented in the Description of Threats section.

**Table 3.** Threat calculator assessment.

Threat #	Threat description	Impact <sup>a</sup>	Scope <sup>b</sup>	Severity <sup>c</sup>	Timing <sup>d</sup>
1	Residential & commercial development	Low	Small	Extreme	High
1.1	Housing & urban areas	Low	Small	Extreme	High
1.2	Commercial & industrial areas	Low	Small	Extreme	High
1.3	Tourism & recreation areas	Low	Small	Extreme	High
2	Agriculture & aquaculture	Negligible	Negligible	Extreme	High
2.1	Annual & perennial non-timber crops	Negligible	Negligible	Extreme	High
2.2	Wood & pulp plantations	Negligible	Negligible	Extreme	High
2.3	Livestock farming & ranching	Negligible	Negligible	Slight	High
3	Energy production & mining	Low	Small	Extreme	High
3.1	Oil & gas drilling	Negligible	Negligible	Moderate	High
3.2	Mining & quarrying	Low	Small	Extreme	High
3.3	Renewable energy	Negligible	Negligible	Extreme	High
4	Transportation & service corridors	Medium	Restricted	Extreme	High
4.1	Roads & railroads	Medium	Restricted	Extreme	High
4.2	Utility & service lines	Low	Small	Extreme	High
4.4	Flight paths	Negligible	Negligible	Negligible	High
5	Biological resource use	High	Large	Extreme	High
5.1	Hunting & collecting terrestrial animals	Negligible	Negligible	Negligible	High
5.2	Gathering terrestrial plants	Negligible	Negligible	Negligible	High
5.3	Logging & wood harvesting	High	Large	Extreme	High
6	Human intrusions & disturbance	Low	Restricted	Slight	High
6.1	Recreational activities	Low	Restricted	Slight	High
6.2	War, civil unrest & military exercises	Negligible	Negligible	Negligible	High

Threat #	Threat description	Impact <sup>a</sup>	Scope <sup>b</sup>	Severity <sup>c</sup>	Timing <sup>d</sup>
7	Natural system modifications	Medium	Restricted	Extreme	High
7.1	Fire & fire suppression	Medium	Restricted	Extreme	High
7.2	Dams & water management/use	Negligible	Small	Negligible	High
8	Invasive & other problematic species & genes	Very High	Pervasive	Extreme	High
8.1	Invasive non-native/alien species/diseases	Negligible	Negligible	Negligible	High
8.2	Problematic native species/diseases	Very High	Pervasive	Extreme	High
8.3	Introduced genetic material	Negligible	Negligible	Negligible	High
8.4	Problematic species/diseases of unknown origin	Unknown	Unknown	Unknown	Unknown
8.5	Viral/prion-induced diseases	Unknown	Unknown	Unknown	Unknown
8.6	Diseases of unknown cause	Unknown	Unknown	Unknown	Unknown
9	Pollution	Negligible	Negligible	Negligible	High
9.1	Domestic & urban waste water	Negligible	Negligible	Negligible	High
9.2	Industrial & military effluents	Negligible	Negligible	Negligible	High
9.3	Agricultural & forestry effluents	Negligible	Negligible	Negligible	High
9.5	Air-borne pollutants	Negligible	Negligible	Slight	High
9.6	Excess energy	Negligible	Negligible	Negligible	High
10	Geological events	Negligible	Negligible	Moderate	High
10.3	Avalanches/landslides	Negligible	Negligible	Moderate	High
11	Climate change & severe weather	Unknown	Unknown	Unknown	Unknown
11.1	Habitat shifting & alteration	Unknown	Unknown	Unknown	Unknown
11.2	Droughts	Unknown	Unknown	Unknown	Unknown
11.3	Temperature extremes	Unknown	Unknown	Unknown	Unknown
11.4	Storms & flooding	Unknown	Unknown	Unknown	Unknown

<sup>a</sup> **Impact** – The degree to which a species is observed, inferred, or suspected to be directly or indirectly threatened in the area of interest. The impact of each threat is based on Severity and Scope rating and considers only present and future threats. Threat impact reflects a reduction of a species population or decline/degradation of the area of an ecosystem. The median rate of population reduction or area decline for each combination of scope and severity corresponds to the following classes of threat impact: Very High (75% declines), High (40%), Medium (15%), and Low (3%). Unknown: used when impact cannot be determined (e.g., if values for either scope or severity are unknown); Not Calculated: impact not calculated as threat is outside the assessment timeframe (e.g., timing is insignificant/negligible or low as threat is only considered to be in the past); Negligible: when scope or severity is negligible; Not a Threat: when severity is scored as neutral or potential benefit.

<sup>b</sup> **Scope** – Proportion of the species that can reasonably be expected to be affected by the threat within 10 years. Usually measured as a proportion of the species' population in the area of interest. (Pervasive = 71–100%; Large = 31–70%; Restricted = 11–30%; Small = 1–10%; Negligible < 1%).

<sup>c</sup> **Severity** – Within the scope, the level of damage to the species from the threat that can reasonably be expected to be affected by the threat within a 10-year or three-generation timeframe. Usually measured as the degree of reduction of the species' population. (Extreme = 71–100%; Serious = 31–70%; Moderate = 11–30%; Slight = 1–10%; Negligible < 1%; Neutral or Potential Benefit ≥ 0%).

<sup>d</sup> **Timing** – High = continuing; Moderate = only in the future (could happen in the short term [ $< 10$  years or 3 generations]) or now suspended (could come back in the short term); Low = only in the future (could happen in the long term) or now suspended (could come back in the long term); Insignificant/Negligible = only in the past and unlikely to return, or no direct effect but limiting.

## 4.1 Description of Threats

Based on IUCN threat evaluation criteria, the overall range-wide threat impact for the Spotted Owl in Canada is assessed as 'very high'. There is one threat that is assessed as 'very high' impact, one that is assessed as 'high' impact, two threats that are assessed as 'medium' impact, seven threats that are assessed as 'low' impact, and numerous threats that were evaluated as having 'negligible' or 'unknown' impacts, within the 10-year IUCN assessment timeframe (Table 3). Although threats related to habitat loss from sources other than logging & wood harvesting (e.g. energy production and mining, transportation and service corridors, recreational activities) are assessed as less than 'high' impact, all sources of habitat loss have cumulatively greater impacts than when assessed individually.

The Northern Spotted Owl Breeding Program facility near Langley, B.C., represents an ex-situ location (i.e. not part of the wild population), and as such, threats to this location are considered separately, and not included in the development of the threat calculation table. Potential threats to the ex-situ location include:

- noise disturbance (e.g. from construction, and railway, aircraft, and vehicle traffic) during the breeding season, which can increase individual stress levels and reduce reproductive output when it occurs near nesting areas (Wasser et al. 1997, Hayward et al. 2011) as well as potentially altering nesting behaviours (USFWS 2020); and
- disease and other health concerns, and natural disasters, the potential impacts of which are elevated due to all Northern Spotted Owl Breeding Program owls being in a single geographic location. These threats are mitigated at the breeding facility through biosecurity protocols, staff training and experience, and separation of individual owls around the property (McCulligh 2024).

### Very High Impact Threats

#### *IUCN 8.2 – Problematic native species*

The Barred Owl is native to eastern Canada but has expanded its range westward and southward. This is hypothesized as being a consequence of human activities that either directly or indirectly resulted in the introduction of trees across the previously tree-less prairie regions of central North America, e.g., through European settlers excluding fires historically set by First Nations, suppressing wildfires, extirpating American Bison (*Bison bison*), and planting trees. In the 1960s Barred Owls began to overlap the range of the Spotted Owl in B.C. (Campbell et al. 1990; Dunbar et al. 1991). Barred Owls were detected at all 10 of the previously-occupied Spotted Owl survey sites visited in 2019 (J. Gillis pers. comm. 2019). They have also been detected extensively along general owl survey routes throughout the Spotted Owl's historical range. Barred Owls thrive in a variety of forest types and seral stages and have adapted to more varied food sources than have Spotted Owls (Livezey et al. 2009a&b; Wiens et al. 2014; Diller et al. 2016; Dugger et al. 2015). Barred Owls threaten the Spotted Owl primarily through competition for habitat and prey (Dugger et al. 2011). This resource competition and competitive displacement has been found to reduce Spotted Owl fecundity and recruitment, leading to overall population declines (Jenkins et al. 2021). Additionally, a study of a managed redwood forest in an area of California not yet colonized by Barred Owls showed that the number of territories occupied by Spotted Owl pairs remained relatively constant over a 25 year period (Kroll et al 2016). Hybridization and predation have also been observed on rare occasions (Leskiw and Gutiérrez 1998; Kelly and Forsman 2004); however, these are not considered serious threats (USFWS 2011).

In recognition of the severity of this threat, Barred Owl control programs have been initiated within the range of both the American and Canadian Spotted Owl populations (Diller et al. 2016; Dugger et al. 2015; Gillis and Waterhouse 2020; Wiens et al. 2021). American programs have employed lethal removal and the B.C. program has employed a combination of translocation and lethal removal. In 2024, the U.S. Fish and Wildlife Service finalized a comprehensive Barred Owl Management Strategy that involves the targeted lethal removal of Barred Owls in specific areas of Northern and California Spotted Owl range, effectiveness monitoring, and adaptive management (USFWS 2024).

Results from Barred Owl removal studies have varied, with more immediate success at the southern edge of the range and slower results at the northern edge of the range. In one California study, the annual Spotted Owl population growth rate four years after (lethal) removals was 1.029 (increasing) on removal sites versus 0.870 (declining) on control sites (Diller et al. 2016), and in another, Spotted Owls recolonized 56% of formerly occupied territories within one year after Barred Owls were removed (Hofstadter 2022). In Oregon and Washington, increases in Spotted Owl occupancy and fecundity and decreases in local extinction rates were observed 4.5 years following Barred Owl removals (Wiens et al. 2021). However, a longer lag time was observed in the more northern sites in Oregon and in Washington (Wiens et al. 2021) and in B.C. (lethal and non-lethal) removal efforts have not yet been sufficient to offset Barred Owl recolonization rates (Gillis 2016a&b; Gillis and Waterhouse 2020). One model predicted that long term recovery of Spotted Owl populations was dependent on high intensities of Barred Owl removal and was negatively affected in areas where Barred Owl population growth is increasing (Perlman 2017). The dual approach of continued Barred Owl removal combined with management actions that maintain high quality habitat indicate improvements in habitat occupancy by Spotted Owls (Long and Wolfe 2019, Wiens et. al 2020).

Diller et al. (2016) suggested that Spotted Owl populations further north may experience slower recovery following Barred Owl removal because Barred Owl populations are more well-established (so require more intensive and sustained removal efforts to overcome recolonization by floaters/dispersers) and Spotted Owl populations are too small to recover quickly (fewer floaters/dispersers waiting to take up available territories). The supplementation of the B.C. Spotted Owl population through re-introduction may counter this effect. Supplemental feeding of released individuals may also bolster post-release survival. Habitat improvement/recovery is also expected to improve Spotted Owl persistence in combination with Barred Owl control and may help reduce the necessary level of investment in Barred Owl removals in the future (Yackulic et al. 2019). However, it is unknown whether this threat can be mitigated or avoided to the extent that the Spotted Owl can recover without ongoing human intervention (Bodine and Capaldi 2017). Research and adaptive management will be required to determine if and how threats from Barred Owl can be effectively addressed over the long term and thus whether a recovered Spotted Owl population can be sustained in the absence of ongoing Barred Owl control.

## High Impact Threats

### *IUCN 5.3 – Logging and wood harvesting*

Logging has had and continues to have severe impacts on the Spotted Owl, including direct loss of old forest habitat (loss of nesting, roosting, and foraging habitat attributes) and fragmentation (COSEWIC 2008, Chutter et al. 2004). The primary impact of forestry-related habitat fragmentation appears to relate to foraging energetics (reviewed in Courtney et al. 2004). As foraging patches become more dispersed following forest harvest, they may no longer be accessible within an individual's energetic budget, and so the individual may starve or be forced to disperse to a new location (Sovern et al. 2014; Jenkins et al. 2019). Further, as residual patches become smaller and more irregular, they may no longer be able to support adequate numbers of the Spotted Owls' preferred prey species (see section 3.3 – Needs of the Spotted Owl). Additional impacts of logging can include noise disturbance associated with logging operations, when operations take place within 400 m of nesting areas (Wasser et al. 1997, Hayward et al. 2011, USFWS 2020). Fragmentation and the conversion of the landscape from old-growth coniferous forest to other habitat types may also increase the exposure of Spotted Owls to their primary predator, the Great Horned Owl (Johnson 1993). Competitive pressure may also be greater within harvested landscapes as Barred Owls are better able to adapt to the more varied seral stages and food sources present in harvested landscapes than are Spotted Owls (Hamer et al. 2007; Wiens et al. 2014; Yackulic et al. 2019).

Improved forestry practices on Crown Land under the provincial *Forest and Range Practices Act* as well as Spotted Owl-specific habitat protection initiatives under the Spotted Owl Management Plans (1 and 2) have partially reduced forestry impacts on Spotted Owl by requiring or promoting the retention of veteran trees, snags, and riparian areas; reducing cut block size; increasing retention area size; and providing some measure of habitat protection for tracts of old forest through the designation of Wildlife Habitat Areas (WHAs), Old Growth Management Areas (OGMAs) and Ungulate Winter Ranges (UWRs) (Government of B.C. 2009). However, a large amount of nesting and foraging class habitat within the Spotted Owl's range still falls within the unprotected portions of the Timber Harvesting Land Base (THLB), and harvesting continues to both remove and isolate habitat.

## Medium Impact Threats

### *IUCN 4.1 – Roads and railroads*

Spotted Owl nesting habitat is located within low-land forests where there has been increasing concentration of roads for logging and other purposes. Major railway corridors also fall in these areas. Road-building and expansion results in direct and often permanent habitat loss through eliminating old forest habitat within the immediate road surface and managed right-of-way. Roads and railways also expose individuals to risk of collisions (Forsman et al. 2002), and noise disturbance from road and rail traffic can increase individual stress levels and reduce reproductive output when it occurs near nesting areas (Wasser et al. 1997, Hayward et al. 2011) as well as potentially altering nesting behaviours (USFWS 2020). Great Horned Owls may also be more prevalent along linear corridors such as roads and railways, putting Spotted Owls at greater risk of predation when these features transect their habitat (Johnson 1993). Road-building will continue to accompany resource extraction/development activities (e.g., forest harvesting). New rail lines are not being planned within the Spotted Owl range.

*IUCN 7.1 – Fire and fire suppression*

Within the drier Sub-maritime and Continental sub-regions, vigorous fire protection by the B.C. Forest Service between the 1960s and 1990s extended fire return intervals well beyond their historical range, creating an accumulation of woody fuels, which can lead to more intense, stand-replacing wildfires (Wong et al. 2003, ESTR Secretariat 2014). Within the American portion of the range, Davis et al. (2016) estimated that 191,900 ha of nesting and roosting habitat on federal lands had been lost to wildfires between 1994 and 2013, four times the amount of habitat that was harvested. A similar analysis in the Canadian portion of the species' range by the Canadian Wildlife Service using annual fire disturbance mapping from 1985 to 2015 (Hermosilla et al. 2015a&b, 2017), indicated that 47,915 ha of forests within the areas classified as suitable for the Spotted Owl has been detectably<sup>10</sup> impacted by fire across that 30-year period, primarily within the drier Sub-maritime and Continental sub-regions, with annual burn areas as large as 4156 ha. Fire impacts are expected to increase in the Spotted Owl range under climate change (reviewed in Spies et al. 2018). Within the wetter regions (i.e., Maritime sub-region in Canada), overall area impacted by fire is expected to remain relatively low due to the naturally very low fire incidence there, even when multiplied according to climate projections (Littell et al. 2010). However, in the drier sub-regions, where existing fire intervals are shorter and fire extents larger, the increase will translate into more significant habitat impacts (reviewed in Spies et al. 2018). Applying an assumption that future annual burn rates under climate change are likely to approximate the upper end of the annual burn rates observed in the previous 30 years (i.e., up to 4156 ha per year), it is estimated that as much as 207,800 ha of Spotted Owl habitat within Canada will be impacted by fire within the 50-year recovery timeframe. This projection was supported during the 2021 fire season when as much as 7700 ha of Spotted Owl habitat may have been impacted by fire (based on B.C. Fire Perimeters mapping). Although not all of these fires will be stand-destroying and result in long-term habitat loss, projections of increasing incidence and area of catastrophic fire under climate change do indicate that fire will be a significant driver of habitat loss in the future (reviewed in Spies et al. 2018; Price and Daust 2016).

Wildfire risk reduction efforts could counter this risk; however, such efforts also have the potential to impact Spotted Owl habitat directly (through loss of potential nesting trees and the features required to support prey populations) when crews target downed wood (CWD) and snags for removal (Wilson and Forsman 2013). Addressing the risk of fire and climate change-mediated increases in fire impacts to the Spotted Owl will require a number of strategies. These include: increasing the overall area conserved in support of Spotted Owl recovery to account for projected fire impacts, including adequate representation within and connectivity to the wetter western portion of the range (where disturbance rates are expected to be lower), ensuring a high level of connectivity as well as alternate connections to provide refugia and enable recolonization/recovery following disturbance, and employing carefully-managed wildfire risk reductions efforts (e.g., avoiding irreplaceable old forest elements such as snags and CWD) in more fire-prone regions that have surplus fuel loads as a consequence of historical fire suppression.

---

<sup>10</sup> Fire impacts were significant enough to result in changes to the forest canopy that could be detected within satellite imagery.

## Low Impact Threats

### *IUCN 1.1 – Housing and urban areas & IUCN 1.2 – Commercial and industrial areas*

Historically (prior to the 1930s), urbanization (and associated commercial and industrial development) resulted in broad-scale loss of mixed-coniferous forests throughout the Lower Mainland (Boyle et al. 1990) as well as portions of the Lower Fraser Valley where agricultural development did not predate urbanization. However, most old forest habitat within range of these population centers has now been converted to urban areas (Chutter et al. 2004; Sutherland et al. 2007), so this is not expected to represent a significant, broad-scale threat in the next decade.

### *IUCN 1.3 – Tourism and recreation areas*

Several large ski resorts exist within the Maritime sub-region in areas with habitat for the Spotted Owl. Expansion of resort infrastructure within existing ski areas could lead to additional, localized, habitat loss. Planning is also underway for one new ski resort in the Sub-maritime sub-region, although proposed development is largely within the footprint of an existing mine, so additional habitat impacts may be minimal. Use of provincial parks and other accessible Crown lands within all three sub-regions has also increased dramatically in the last decade (B.C. Parks 2018; J. Hirner, pers. comm. 2020), creating pressure to expand trails and park infrastructure into potential Spotted Owl habitat. However, this threat applies to a relatively small percentage of the species' range, so the overall impact is assessed as low.

### *IUCN 3.2 – Mining and quarrying*

Mining and mineral exploration activities are uncommon in the Spotted Owl range; however, because they are exempt from the prohibitions on forest harvest under the General Wildlife Measures in WHAs (Government of B.C. 2019), such activities have the potential to cause habitat loss even in areas under timber harvest constraints. Any Spotted Owls nesting or foraging within the vicinity of mining or quarrying operations could also be disturbed by operational noise, with similar impacts as from noise associated with wood harvesting and road construction or traffic. However, this threat applies to a relatively small percentage of the species' range, so the overall impact is assessed as low.

### *IUCN 4.2 – Utility and service lines*

As with roads, habitat clearing associated with utility and service line construction (which includes pipelines) will result in some direct habitat loss and the linear edge habitats created could impact prey populations and increase predator exposure. Any Spotted Owls nesting or foraging within the vicinity of utility or service lines during construction or maintenance could also be disturbed by machine noise, with similar impacts as from noise associated with wood harvesting and road construction or traffic. However, this threat applies to a relatively small percentage of the species' range (one major utility line project was being planned when the threats calculator was prepared, the construction of which was completed in 2023), so the overall impact is assessed as low.

### *IUCN 6.1 – Recreational activities*

Backcountry recreation use has increased dramatically within Southern B.C. Visitor numbers at B.C. Parks in southern regions increased by 60% between 2007 and 2017 (B.C. Parks 2018).

Recreational use of other accessible Crown lands has also increased dramatically in the last decade (J. Hirner, pers. comm. 2020). As more backcountry users visit parks and recreation areas where Spotted Owls nest, the potential for human disturbance increases. Motorized recreation, in particular, could disturb Spotted Owls nesting in the vicinity of recreational trails/areas. However, this threat applies to a relatively small percentage of the species' range, so the overall impact is assessed as low.

### **Negligible and Unknown Impact Threats**

Eleven individual threats or complete IUCN threat categories were classified as having a negligible impact on the Spotted Owl based on limited spatial overlap with the species' range and habitat and/or no anticipated impacts within the 10-year IUCN-CMP assessment timeframe.

A further five threats were classified as having unknown impacts within the 10-year assessment timeframe, most related to climate change. Climate change impacts could be significant, particularly within the 50-year recovery timeframe, but there remains considerable uncertainty around the direction and magnitude of climate change-mediated shifts in weather, natural disturbance, and forest health within the Spotted Owl range, as well as the likely response of Spotted Owls to those changes (reviewed in Courtney et al. 2004; Spies et al. 2018).

A comprehensive review of climate modelling research has been undertaken for the Northwest Forest Plan (in the U.S.A.), which is focused on management of old-growth forests for Spotted Owl recovery (Spies et al. 2018). Most models assessed within that review predicted warmer, drier summers and potentially warmer and wetter winters. Conditions are projected to exceed the 20th-century range of variability by the 2050s. These predictions are supported by modelling that also covers the Canadian portion of the Spotted Owl's range (Wang et al. 2016). A comprehensive analysis of Spotted Owl survival and recruitment in relation to predictors including climate (Dugger et al. 2015) found an association between climate variables and both juvenile recruitment and adult annual survival. Recruitment was lowest when conditions during the previous winter were cold and wet, and highest when the previous winter was cold and dry. Observed survival rates were higher when winters were relatively warmer and drier. Summer temperature extremes could also impact recruitment rates; the heat dome of 2021 had significant impacts on juveniles in the fledge stage (J. Gillis, pers. comm. 2021). However, given that predicted temperature and precipitation patterns under climate change could lead to both positive and negative changes to different demographic rates, it is difficult to generate an overall prediction of how Spotted Owl populations may be impacted.

When it comes to habitat impacts from climate change, lower elevation, moist vegetation zones (e.g., those within much of the Maritime sub-region in Canada) are expected to experience decreased growth and productivity, especially where tree species are already water limited during the growing season (reviewed in Spies et al. 2018). Within drier forests (e.g., those within the Continental sub-region and some portions of the Sub-maritime) most models predict an increased role of fire, including more area burned and larger patches of high-severity fire (reviewed in Spies et al. 2018; Price and Daust 2016), which will increase the rate of fire-related habitat loss, relative to past decades (e.g., see IUCN-CMP 7.1, above). A preliminary assessment of anticipated climate change vulnerability for a number of species in B.C. was conducted in 2016 (Price and Daust 2016). Although the Spotted Owl was not amongst the species assessed, other old forest-associated species with similar ranges were assessed as having moderate-high climate change vulnerability, primarily due to increased climate change-mediated natural disturbance within their old forest habitats.

## 5. Population and Distribution Objectives

### Population and Distribution Objective:

To recover the Spotted Owl in Canada by restoring a stable population of at least 250 mature individuals distributed within a connected network of habitat representative of all three sub-regions within the species' historical Canadian range, and linked to the larger population in the U.S.A.

### *Rationale:*

Historically, the Spotted Owl's restricted range and small population size would have made it naturally precarious (i.e., naturally falling within COSEWIC's Threatened status); however, the population was believed to be large enough to be stable, with connectivity/representation across its range. In contrast, the species is now assessed as Endangered on the basis of compromised stability, redundancy, connectivity, and resilience.

There has been permanent loss of habitat within the Lower Mainland and Lower Fraser Valley (now a major human population center), which both reduces the overall area available to the species in Canada and restricts the potential for continued gene flow between Canada and the U.S.A. (Chutter et al. 2004). However, portions of habitat within the remainder of the range, across all three historically-occupied sub-regions are either still intact or are close to re-acquiring the attributes of Spotted Owl habitat, such that if they are maintained now they will contribute to habitat patch size, quality and connectivity, and help to restore historical representation. Other limitations to recovery are also believed to be manageable over the long term, given current/anticipated tools (Chutter et al. 2004; Government of B.C. 2020; B.C. MFLNRORD 2021). Assuming that planned actions are undertaken to i) protect and restore sufficient Spotted Owl habitat; ii) control Barred Owls to reduce interspecific competition; iii) continue the Northern Spotted Owl Breeding Program, and iv) release facility-born Spotted Owls to supplement wild populations and achieve successful breeding of reintroduced owls in the wild, the provincial government projects that it is within the scope of biological and technical feasibility to achieve the COSEWIC threshold for Threatened status (i.e.,  $\geq 250$  mature individuals) within 50 years (B.C. MFLNRORD 2021). The amount, configuration and attributes of habitat that is necessary to achieve this population target in context of distribution objectives (connectivity and representation) are described in section 7 of this document. Recovery of Spotted Owls will require significant targeted interventions in the form of population augmentation and competitor control in the short- to medium-term (10-30 years), therefore, short-term recovery statements toward meeting the objective are set out below, to facilitate recovery implementation.

### Short-term Recovery Statements Toward Meeting the Population and Distribution Objective:

1. Immediately mitigate human-caused threats that would cause loss of habitat needed for recovery (i.e., the critical habitat).
2. By 2030, re-establish a wild Spotted Owl population in British Columbia of four to six resident individuals, utilizing individuals bred from the Northern Spotted Owl Breeding Program.
3. Complete regular Barred Owl surveillance at sites occupied by Spotted Owls and/or where reintroductions are planned, and remove all Barred Owls that are detected.

## 6. Broad Strategies and General Approaches to Meet Objectives

### 6.1 Actions Already Completed or Currently Underway

#### Habitat protection, enhancement and stewardship

##### *Management planning*

In 1997, a Spotted Owl Management Plan (SOMP) was developed with a goal of stabilizing (and optimistically improving) the population over the long-term, without exceeding a 4% reduction in the Timber Harvesting Land Base (THLB) (i.e., SOMIT 1997). SOMP established 21 Special Resource Management Zones (SRMZs) that included pre-existing protected areas as well as Crown forest land. Within the SRMZs that fell outside protected areas, 67% of the habitat was to remain suitable for the Spotted Owl, while the remaining 33% was eligible for harvest using specific prescriptions.

In 2009, an updated version of SOMP ('SOMP2') was released, which involved transferring most SRMZs into Long Term Owl Habitat Areas (LTOHAs; managed for Spotted Owl conservation) and Managed Future Habitat Areas (MFHAs; managed for forest harvest with consideration for long-term Spotted Owl habitat development), adjusting some managed area boundaries, and creating updated harvesting guidelines/designations (Blackburn et al. 2009; Government of B.C. 2009). The requirement to limit impacts on the THLB to <4% remained, so there was no increase in the area managed for Spotted Owl recovery under the new plan. In 2012, the LTOHA and MFHA areas under SOMP2 became legally-designated Wildlife Habitat Areas (WHA) with General Wildlife Measures (GWM). Thirty-two WHAs are now in place to provide a measure of protection to areas large enough to support one or more breeding pairs of Spotted Owls (Government of B.C. 2023). Within the LTOHA WHAs forest harvest is largely prohibited, and within the MFHA WHAs harvest is permitted subject to conditions.

##### *Additional regulatory measures*

In addition to provincial WHAs, other protected area designations<sup>11</sup> provide some measure of protection for Spotted Owl habitat. These include: Provincial/Municipal/Regional Parks; Provincial Protected Areas, Recreation Areas, Ecological Reserves and Conservancy Areas; Sea-to-Sky Wildland Areas; Metro Vancouver Watersheds; Ungulate Winter Ranges, Old Growth Management Areas, and Wildlife Habitat Areas for other species; and National Wildlife Areas.

---

<sup>11</sup> These forms of habitat protection do not necessarily qualify as effective protection of critical habitat under SARA. Such a determination can only be made following a Critical Habitat Protection Assessment (Environment and Climate Change Canada 2016).

## Active population management

### *Northern Spotted Owl Breeding Program and Release*

In 2007, the Spotted Owl Population Enhancement Team, an arm's-length independent panel that was established by the provincial government, determined that the wild population was so small and isolated that extirpation was a certainty. It therefore made the recommendation to capture either all or a subset of the remaining wild individuals and establish a facility-bred population whose offspring could be re-introduced into the wild (Fenger et al. 2007). The provincial government elected to capture only a subset of the remaining wild individuals to establish the Northern Spotted Owl Breeding Program and allow a small wild population to persist. The home ranges that owls were removed from to establish the Northern Spotted Owl Breeding Program were all designated as LTOHAs at that time. They were then converted into WHAs in 2012.

The Northern Spotted Owl Breeding Program has been in operation in B.C. since 2007 (Government of B.C. 2021). It had slow initial success rates, but a now younger breeding population and improvements in husbandry techniques have resulted in higher breeding output in recent years (McCulligh 2023, Northern Spotted Owl Breeding Program n.d.). However, given the small size of the breeding population at the facility, even modest increases in adult mortality or declines in the average egg fertility rate could have significant consequences for the recovery program.

As of spring 2023, the breeding program had produced 31 facility-born owls (McCulligh 2024). The first trial release occurred in August 2022, when three facility-bred male owls were released into two Wildlife Habitat Areas; one was later found injured and returned to the breeding facility, and the other two died of unknown causes (Government of B.C. 2023a). Additional details about the 2022 release considerations and protocols are included in McCulligh 2023. In late June 2023, the previously released owl and one additional male were released following updated protocols (Government of B.C. 2023b). Following those releases, as of March 2024, there were 34 Spotted Owls at the breeding facility (McCulligh, 2024).

### *Barred Owl control*

In 2007, the provincial government initiated a Barred Owl removal program, with target sites including active Spotted Owl territories and sites planned for re-establishment through the release of facility-bred owls (Fenger et al. 2007). Between 2007 and fall 2022, a total of 207 Barred Owls were removed from active (i.e., currently occupied) Spotted Owl territories and from proposed Spotted Owl re-establishment sites (Gillis and Waterhouse 2020; Government of B.C. representatives, pers.comm. 2023). Removals were a combination of capture and translocation (at re-establishment sites) and lethal removal via shooting (at active sites). Between 2007-2021, 108 Barred Owls were captured and translocated away from proposed re-establishment sites and 80 were removed from active Spotted Owl sites using lethal methods. The combined removal effort reduced the number of detected Barred Owls overall, but as of 2016 had not been sufficient to overcome local re-colonization rates. Moving forward, adaptations to removal methods could improve the effectiveness of Barred Owl removal efforts in B.C. (Gillis and Waterhouse 2020). The augmentation of the wild population through the release of facility-bred owls may also increase the rate of Spotted Owl recolonization of removal sites. Habitat improvement/recovery is also expected to improve Spotted Owl persistence in

combination with Barred Owl control, reducing the necessary level of investment in Barred Owl removals in the future (Yackulic et al. 2019).

## **Inventory, monitoring, and population evaluation**

### *Owl population inventory and monitoring*

From the 1990s to 2008, inventories were conducted to determine the range, distribution, and abundance of the Spotted Owl in B.C., as well as to assist in resource management decisions (Blackburn et al 2002; Hobbs 2004&2005; J. Gillis pers. comm. 2019). An organized banding program (attaching unique leg bands) was initiated in 1998 to identify individuals and monitor their movements and habitat occupancy. Between 1998 and 1999, transmitters were affixed to several breeding pairs to monitor habitat use and home range sizes (Chutter et al. 2004). Between 2003 and 2014, juvenile owls were affixed with transmitters to ascertain their dispersal movements and overwinter survival (Hobbs 2004&2005; J. Gillis pers. comm. 2019). Beginning in 2015, inventory/monitoring efforts became focused on revisiting previously known Spotted Owl sites to assess re-occupancy, as well as inventory of sites identified for potential re-introduction through release of facility-bred owls (Gillis 2016a&b; 2017; 2018). Starting in 2016, a pilot program was launched to assess the utility of Autonomous Recording Units (ARUs) in Spotted Owl and Barred Owl monitoring (Gillis 2016a&b; 2017; 2018). ARUs continue to be used for monitoring purposes (McCulligh 2024), although Spotted Owls are known to suppress their vocalizations in the presence of Barred Owls (Dugger et al. 2011; Wiens et al. 2014). As such, the absence of Spotted Owl detections from ARU data is not conclusive evidence of absence of Spotted Owls. ARU data informs the implementation and effectiveness monitoring of the Barred Owl control program (McCulligh 2024).

### *Habitat and population evaluation*

In 2007, the Canadian Spotted Owl Recovery Team (CSORT), with the support of Cortex Consultants and Andrew Fall Gowlland Technologies Ltd, developed an integrated modelling framework designed to inform the Spotted Owl recovery program in B.C. and associated habitat management (Sutherland et al. 2007). The framework included models for spatial landscape projection, ecological classification, cross-scale habitat assessment, population dynamics, and reserve selection. This work informed changes / refinements in habitat protection under SOMP 2 (Government of B.C. 2009 & 2020) as well as the approach for the identification of critical habitat in this document.

## 6.2 Strategic Direction for Recovery

**Table 4.** Recovery Planning Table.

Threat or Limitation	Priority <sup>a</sup>	Broad Strategy to Recovery	General Description of Research and Management Approaches
Habitat loss and fragmentation (IUCN #1, 3.2, 4.1, 4.2, 5.3, 7.1)	High	Habitat protection, enhancement and stewardship	Work between governments (federal, Indigenous, provincial) to establish or confirm protection <sup>b</sup> for identified critical habitat.
	Medium		Pursue wildfire risk-reduction efforts that align with Spotted Owl habitat requirements.
	Low		Develop/expand silvicultural guidelines to create, enhance and/or maintain suitable conditions for Spotted Owls within younger forests that fall within or between critical habitat patches.
	Low		Promote habitat stewardship with forest companies that operate within the Spotted Owl range in Canada.
	Low		Promote Spotted Owl population stewardship with stakeholders.
Barred Owls (IUCN #8.2)	High	Active population management	Continue the operational Barred Owl control program with annual adaptations informed by results of the B.C. program and those of similar efforts within the U.S.A.
Lack of natural recruitment	High		Continue the Northern Spotted Owl Breeding Program and release / reintroduction program, including post-release measures such as supplemental feeding and satellite tracking of released individuals, and develop strategic partnerships with First Nations communities to continually evolve the adaptive management approach for Spotted Owl population recovery, supporting collaboration where there is mutual interest, and fostering knowledge exchange.
	Medium	Work with government agencies within the U.S.A. to improve international coordination of Spotted Owl recovery efforts and increase the likelihood of cross-border immigration/gene flow, support collaboration where there is mutual interest, and foster knowledge exchange for the benefit of the Spotted Owl population in B.C. and more broadly.	
Knowledge gaps	High	Research, inventory, monitoring and population evaluation	Continue to pilot new monitoring technologies such as ARUs in order to enable comprehensive inventory of the entire Spotted Owl range and timely detection/monitoring of Barred Owls.
	Medium		Establish a periodic, recurrent, standardized survey (counts by age class; number of active territories, recruitment surveys; DNA samples) to monitor the status and composition of the Spotted Owl population.
	Low		Pursue additional research on impacts of acoustic disturbance on Spotted Owls, including impacts outside of the breeding season.
	Low		Pursue research on the relative contribution of forested habitats of different seral stages to survival of dispersing Spotted Owls, to improve management of dispersal corridors.
	Low		Pursue research on climate change impacts for Spotted Owls and better-integrate climate change resilience/mitigation strategies into recovery planning.

<sup>a</sup> “Priority” reflects the degree to which the broad strategy contributes directly to the recovery of the species or is an essential precursor to an approach that contributes to the recovery of the species. High priority measures are considered those most likely to have an immediate and/or direct influence on attaining the population and distribution objective for species and thus considered to be most urgently needed to ensure the species survival or of highest importance for the species’ recovery. In some cases, a high priority action may need the completion of another stated high priority action before it can be accomplished. Medium priority measures may have a less immediate or less direct influence on reaching the population and distribution objectives, but are still important for recovery of the population. Low priority recovery measures will likely have an indirect or gradual influence on reaching the recovery objectives, but are considered important contributions to the knowledge base and/or public involvement and acceptance of species. This may be reflected in the timeline for completion.

<sup>b</sup> Legal or effective protection under SARA (Environment and Climate Change Canada 2016).

### 6.3 Narrative to support recovery planning table

The success of Spotted Owl recovery is dependent on both the protection and recruitment of suitable habitat and the successful release of facility-born Spotted Owls. Released owls must establish territories and form breeding pairs in order to achieve recovery as a self-sustaining population.

Several management actions are required and must be sustained, likely over several decades, to achieve recovery. Given the timeframe and uncertainty, an incremental management approach should be used to review the effectiveness of Spotted Owl releases and the Barred Owl control program in supporting the establishment, survival and breeding success of facility-born Spotted Owls, as well as the assessment of habitat use.

Recruitment of younger forest (e.g. burned or previously harvested) involves time for these areas to become suitable. Within those previously disturbed areas, targeted, site-specific forest management that increases the amount of coarse woody debris and retains large diameter standing trees may decrease recruitment time relative to standard silvicultural practices (Irwin et al. 2024).

Although it is difficult to predict the full magnitude of climate change-mediated impacts on Spotted Owls and their habitat, it is possible to anticipate and to implement strategies for reducing/mitigating those impacts. Within B.C., there is significant spatial overlap between the low-elevation old-forest-dominated habitat throughout the Spotted Owl range and landscapes with high predicted climate change resilience and macrorefugia potential (Beckers and Carroll 2020). Ensuring protection of highly-connected networks of old-forest Spotted Owl habitat, which could function as refugia in an increasingly disturbance-prone landscape, will be an essential component of mitigating climate change-mediated disturbance and maintaining climate change resilience for Spotted Owls (Gayton 2008; Price and Daust 2016; Spies et al. 2018; Thom et al. 2019; Krawchuk et al. 2020; USGS 2021; Lesmeister et al. 2021).

Addressing the risk of fire and climate change-mediated increases in fire impacts to the Spotted Owl will require a number of strategies, including ensuring the overall availability of sufficient habitats that have characteristics consistent with Spotted Owl recovery to account for projected fire impacts, and mitigating the impacts of fire risk reduction efforts within habitat. Some wildfire risk reduction activities could help mitigate large-scale and high severity wildfires (Pritchard et al. 2021, BC Wildfire Service 2024), but activities will need to be conducted so that they are not likely to destroy critical habitat (Section 7.3). Actions should include ensuring adequate representation within and connectivity to the wetter western portion of the range (where disturbance rates are expected to be lower); a high level of connectivity as well as alternate connections to provide refugia and enable recolonization/recovery following disturbance; and

employing carefully-managed wildfire risk reductions efforts (e.g., avoiding irreplaceable old forest elements such as snags and CWD and focused reduction of ladder fuels) in more fire-prone regions that have surplus fuel loads as a consequence of historical fire suppression.

The effectiveness of Barred Owl lethal removals in established release areas should be assessed using several short- and long-term performance metrics, including a) monitoring changes in Barred Owl density and recolonization via acoustic monitoring, b) surveying changes in Spotted Owl prey populations, and c) assessing the physical health condition, population structure, and diet of removed Barred Owl carcasses. This initial work should occur prior to Spotted Owl releases to ensure sites are suitable for facility-bred Spotted Owls, but also to understand the effects of Barred Owl control in the absence of Spotted Owls. Active monitoring of released Spotted Owls will also help understand the effects of Barred Owl control in the presence of Spotted Owls.

Monitoring will help inform effectiveness of habitat protection and conservation measures, as well as situations where Spotted Owls establish territories outside of protected areas. Results from these monitoring and research efforts will inform revisions to the approach taken for future direct management actions and prioritization and boundaries for spatial habitat protection (e.g., wildlife habitat areas, old growth management areas, etc.) and potential land use objectives.

## 7. Critical Habitat

Section 41 (1)(c) of SARA requires that recovery strategies include an identification of the species' critical habitat, to the extent possible, as well as examples of activities that are likely to result in its destruction. This federal recovery strategy identifies critical habitat to the extent possible, based on the best available information for the Spotted Owl.

It is recognized that the 'acoustic' critical habitat identified below is insufficient to achieve the population and distribution objectives for the species. A schedule of studies (Section 7.2) has been developed to provide the information necessary to complete the identification of acoustic critical habitat (defined below) that will be sufficient to meet population and distribution objectives. For core critical habitat (defined below), boundaries may be revised, or additional core critical habitat may be added in the future, if additional information supports such changes. The identification of critical habitat will be updated when the information becomes available, in a revised recovery strategy.

### 7.1 Identification of the Species' Critical Habitat

The Spotted Owl requires habitat for nesting, roosting, foraging and safe movement (see Section 3.3 – Needs of the Spotted Owl). Some mature and old-growth stands already possess the attributes required to support these functions, and some previously-disturbed habitat has the potential to acquire the necessary attributes within the 50-year timeframe needed to meet the population and distribution objective. A 400-m area surrounding nesting areas must also be protected from acoustic disturbance during the breeding season to ensure that acoustic disturbance does not result in loss of breeding habitat function. The critical habitat required by the Spotted Owl for recovery is therefore comprised of two subtypes:

1. **Core critical habitat:** habitat that either already possesses, or will develop (within a 50-year period), the features required by the owls to successfully nest, roost, forage and move safely.
2. **Acoustic critical habitat:** habitat surrounding nesting areas that functions to maintain the acoustic environment within those areas during the breeding season.

The geospatial areas that may contain critical habitat for the Spotted Owl (totaling 416, 258 ha) are presented in Figures 4-9. Within these geospatial areas, critical habitat is identified wherever the following biophysical attributes occur, or would be expected to occur within the 50-year timeframe needed to meet the population and distribution objective.

### **Biophysical features and attributes of critical habitat**

A description of the known biophysical features and attributes of the species' habitat that are required to support life-cycle processes (functions) are summarized in Section 3.3, Needs of the Spotted Owl, and form the basis of the biophysical attribute description in Table 5 below.

**Table 5.** Functions, biophysical features, and attributes of Spotted Owl critical habitat. Attributes represented within VRI mapping act as criteria for selecting core critical habitat polygons (see Table 2). The presence of these attributes must be assessed at the scale of the component VRI polygon. Some or all of the attributes listed here are expected to either be present, or in the process of developing (within the 50-year recovery timeframe), within the core critical habitat polygons; however, due to the scale of VRI, there can be some uncertainty, so on-the-ground verification of attributes is important. Minimum quantitative thresholds are from the minimum definition of ‘moderate/suitable’ habitat in Appendix 5 of Chutter et al. (2004). This should not be confused with quantitative definitions of ‘superior’ habitat (e.g., in Blackburn et al. 2009; Waterhouse et al. 2012; D’Anjou et al. 2015).

Type	Function	Biophysical features	Attributes	
			Maritime sub-region	Sub-maritime and Continental sub-regions
Core critical habitat	Nesting	Nest trees	Large (>50 cm dbh) snags or trees with deformities (e.g., large cavities, broken tops, dwarf mistletoe infections)	Large (>30 cm dbh) snags or trees with deformities (e.g., large cavities, broken tops, dwarf mistletoe infections)
	Roosting and safe movement	Closed, multi-storey canopy to provide thermoregulation opportunities and protection from inclement weather and predators	>60% canopy closure	>50% canopy closure
			≥2 horizontal canopy layers	
	Foraging and safe movement	An open understory structure (characteristic of stands dominated by tall, large-diameter trees) to enable efficient access to prey	Canopy dominated by overstorey trees >50 cm dbh	Canopy dominated by overstorey trees >30 cm dbh
			≥19.5 m stand height	
			Abundant CWD and a diverse shrub layer	
Acoustic critical habitat	Maintenance of suitable acoustic levels within nesting areas	Anthropogenic noise level that does not interfere with life functions within nesting areas, resulting in loss of habitat availability or function.	Noise level not exceeding 90 db and/or not exceeding ambient conditions by >20 db during the Spotted Owl nesting season (February 1 <sup>st</sup> to July 31 <sup>st</sup> )	

Within the geospatial areas mapped as **core critical habitat** only unsuitable areas that do not possess the features and attributes required by the Spotted Owl at any time - either currently, or within the 50-year recovery timeframe - are excluded from identification as critical habitat. Examples of excluded areas include cultivated and/or landscaped areas, buildings, roads and artificial surfaces, or forested areas that have been recently harvested or subject to stand-destroying disturbance (e.g., catastrophic fire), and so will not acquire the critical features and attributes of critical habitat within the 50-year recovery timeframe (see Table 2 for stand age thresholds specific to each sub-region).

### 7.1.1 Information and methods used to identify critical habitat

The location and spatial configuration of critical habitat is based on three principle assumptions:

- **Quantity:**
  - *Minimum starting polygon size:* to support foraging and nesting, noting that the species' arboreal prey has a home range size of ~10 ha (Sutherland et al. 2007), core critical habitat must be configured around nesting class habitat polygons that are at least 10 ha in size.
  - *Overall core critical habitat area:* to support the long-term population objective, the summed area of core critical habitat identified must be sufficient to:
    - support the home ranges of at least 125 pairs<sup>12</sup> of owls, noting that the mean amount of habitat to support a pair's home range is estimated at 3010, 2224, and 1907 ha for the Maritime, Sub-maritime, and Continental sub-regions, respectively, and that adjacent home ranges may overlap up to 25% (Chutter et al. 2004 & 2007; Sutherland et al. 2007); and
    - account for up to 207,800 ha of habitat being impacted by fire within the 50-year recovery period (see Section 4.1).
- **Support for all critical life functions (i.e., biological value):** To support all critical life functions for a recovering population, habitat must be prioritized for inclusion within the core critical habitat geospatial area based on its:
  - a) contribution to a core patch that can support multiple home ranges and enable successful nesting, roosting, foraging, and safe movement within the patch:
    - i. vicinity/degree of connectivity to recovery origins (existing locations and proposed re-introduction sites) or historical anchors (historical locations of resident birds);
    - ii. proportion of nesting class habitat; and
    - iii. size; or

---

<sup>12</sup> The 250-adult-bird population target has been translated into a critical habitat target of 125 home ranges for simplicity; however, it is acknowledged that 250 adult birds does not necessarily equate to 125 pairs. A normal population includes a certain number of adult, non-breeding, non-territorial 'floaters' whose habitat use may overlap with that of territorial breeders (Franklin 2001). Further, not all resident adults are paired, so some home ranges may be occupied by single birds. However, 125 home ranges is being interpreted as a reasonable benchmark for the amount of habitat needed to support 250 adults, in the absence of more detailed information on eventual population structure and space use by the recovered population.

- b) contribution to a potential connective corridor that can enable safe movement between core patches.
- **Representation across the three ecological sub-regions:** To restore pre-human-impact patterns of representation, critical habitat must be identified across all the three ecological sub-regions within the species' historical range (Maritime, Sub-maritime, Continental).

The geospatial delineation process summarized below aims to create a well connected critical habitat network large enough to support 125 pairs, accounting for home range overlap and anticipated fire impacts, prioritizing habitat with the potential to support all critical life functions and restore pre-human-impact patterns of representation. The geospatial delineation process is founded upon chapter 6 of the integrated population and habitat modelling framework developed under the direction of CSORT (Sutherland et al. 2007). It is summarized below and outlined in greater detail in an accompanying technical document (available upon request).

The information used in the geospatial delineation of critical habitat for the Spotted Owl includes:

- a) provincial VRI mapping (2018 version);
- b) a Spotted Owl *caurina* subspecies habitat suitability classification produced by CSORT to be applied to VRI (Table 2);
- c) a 50-year future projection of VRI (BC MFLNRORD 2019);
- d) a least-cost/resistance landscape created through applying habitat-specific cost categories<sup>13</sup> to VRI polygons;
- e) a set of contiguous habitat clusters created through:
  - i. applying a connectivity analysis to link all  $\geq 10$ -ha nesting class polygons via least-cost pathways (based on the resistance landscape);
  - ii. selecting all habitat (either nesting class or foraging class) from the 50-year future projected VRI that intersects the least cost pathways; and
  - iii. dissolving polygons to create discrete clusters;
- f) origins/anchors of recovery comprised of:
  - i. habitat occurring within the estimated maximum home range area of extant and historical locations of resident Spotted Owls, both within Canada (I. Blackburn, pers. comm. 2021) and in northern Washington; and
  - ii. habitat occurring within the estimated maximum home range area of the currently-proposed re-introduction locations for facility-bred Spotted Owls (I. Blackburn, pers. comm. 2021); and
- g) potential connective corridors created through:
  - i. applying a connectivity analysis to link all origins/anchors of recovery via least-cost pathways (based on the resistance landscape); and
  - ii. selecting all habitat (either nesting class or foraging class) from the 50-year future projected VRI that is located within 500 m of a least-cost pathway.

A summary of the geospatial delineation process is as follows:

---

<sup>13</sup> Based on Table 1 in Sutherland et al. (2007) with adjustments to treatment of roads and high elevation areas following expert review.

1. Merge origins/anchors with any contiguous habitat clusters that intersect them to delineate origin/anchor habitat patches. Retain those that contain enough habitat to support at least one home range, within an area no larger than the maximum home range area estimated for that sub-region (see Section 3 - Species Needs).
2. Assess the biological value of the contiguous habitat clusters occurring outside of origin/anchor habitat patches and potential connective corridors.
3. Build core critical habitat that will accommodate 125 home ranges (accounting for 25% overlap between adjacent ranges), enable safe movement between origin patches, and remain sufficient to support home range and safe movement targets after accounting for anticipated fire impacts (up to 207,800 ha; see Section 4.1 - Threats):
  - a. include all functional<sup>14</sup> origin/anchor habitat: 272,793 ha (125 home ranges + ~65,000 ha towards fire impacts);
  - b. include all functional<sup>16</sup> potential corridor habitat linking origins/anchors: 99,585 ha (safe movement + fire impacts); and
  - c. include additional contiguous habitat clusters occurring outside of origin/anchor and corridor habitat, by biological value score, until the fire impact target is met: 43,387 ha.
4. Build acoustic critical habitat to counter acoustic disturbance within nesting areas:
  - a. delineate nesting areas as all habitat polygons intersecting a 500-m area around nest sites (Blackburn et al. 2009); and
  - b. establish a 400-m (horizontal distance) acoustic influence zone around the delineated nesting area(s).
5. Apply critical habitat sub-type classifications:
  - a. designate all habitat identified through steps 1-3 as core critical habitat; and
  - b. designate the habitat identified through step 4 as acoustic critical habitat.

### 7.1.2 Geospatial Location of Areas Containing Critical Habitat


Critical habitat for Spotted Owl is identified within three sub-regions in B.C. (Figures 4-9):

- Overview map summary (all sub-regions): Figure 4
- Maritime sub-region: Figure 5
- Sub-maritime sub-region: Figures 6-8
- Continental sub-region: Figure 9


The 10 km x 10 km UTM grid overlay shown on these figures is a standardized national grid system that highlights the general geographic area containing critical habitat, for land use planning. The currently identified acoustic critical habitat is based on the location of a nest site that was discovered in 2019, and is the only known breeding location in the wild within the last 5 years.

---


<sup>14</sup> Habitat polygons with insufficient interior habitat could fail to support adequate prey (see section 3.3 – Needs of the Spotted Owl). Within origin/anchor patches and dispersal corridors, an additional 100-m area of surrounding habitat was included around all polygons that had an interior area <10 ha to ensure that all habitat patches identified as critical habitat will maintain sufficient interior habitat to support prey, even if forested habitat is removed along their boundaries. Patches that were still <10 ha even after the additional surrounding habitat was added were not considered functional, so were excluded from potential identification.




**Figure 4.** Overview of the critical habitat for the Spotted Owl within B.C. Critical habitat is represented by the yellow shaded and black outlined polygons where the criteria and methodology set out in this section are met. The area below the hatched line is the U.S.A. land base.




**Figure 5.** Critical habitat for the Spotted Owl within the Maritime sub-region is represented by the yellow shaded polygons where the criteria and methodology set out in this section are met. The 10 km x 10 km UTM grid overlay shown on this figure is a standardized national grid system that indicates the general geographic area containing critical habitat. Federal Protected Areas that overlap with critical habitat are also shown.




**Figure 6.** Critical habitat for the Spotted Owl within the northern portion of the Sub-maritime sub-region is represented by the yellow shaded polygons where the criteria and methodology set out in this section are met. The 10 km x 10 km UTM grid overlay shown on this figure is a standardized national grid system that indicates the general geographic area containing critical habitat.



**Figure 7.** Critical habitat for the Spotted Owl within the central portion of the Sub-maritime sub-region is represented by the yellow shaded and black outlined polygons where the criteria and methodology set out in this section are met. The 10 km x 10 km UTM grid overlay shown on this figure is a standardized national grid system that indicates the general geographic area containing critical habitat.



**Figure 8.** Critical habitat for the Spotted Owl within the southern portion of the Sub-maritime sub-region is represented by the yellow shaded and black outlined polygons where the criteria and methodology set out in this section are met. The 10 km x 10 km UTM grid overlay shown on this figure is a standardized national grid system that indicates the general geographic area containing critical habitat. The hatched line in the southern extent of the map is the border with the Continental U.S.A.



**Figure 9.** Critical habitat for the Spotted Owl within the Continental sub-region is represented by the yellow shaded polygons where the criteria and methodology set out in this section are met. The 10 km x 10 km UTM grid overlay shown on this figure is a standardized national grid system that indicates the general geographic area containing critical habitat.

## 7.2 Schedule of Studies to Identify Critical Habitat

The following schedule of studies (Table 6) is required to complete the identification of acoustic critical habitat for the Spotted Owl.

**Table 6.** Schedule of studies to complete the identification of acoustic critical habitat for the Spotted Owl.

Description of Activity	Rationale	Timeline
Identify <b>acoustic critical habitat</b> surrounding additional nesting areas as they become established.	Currently it is unknown precisely where and when Spotted Owl reintroductions will be successful and where the recovering population will establish nesting areas. As new nesting areas become established, additional <b>acoustic critical habitat</b> must be identified to support those breeding pairs.	2025-2075

### 7.3 Activities Likely to Result in the Destruction of Critical Habitat

**Table 7.** Description of activities likely to result in destruction of the two critical habitat subtypes for the Spotted Owl.

Description of Activity	Description of Effect	Details of Effect
Any activity involving removal or disruption of natural vegetation and ground cover within <b>core critical habitat</b> , e.g., logging and wood harvesting; road-building; residential and commercial development; deliberate setting of stand-replacing fires	Activities resulting in destruction or removal of natural vegetation and ground cover (vegetation, snags, CWD) can result in destruction of core critical habitat through causing direct and permanent loss of the critical features and attributes required for all life functions (nesting, roosting, foraging, and safe movement).	Related IUCN-CMP Threat # 1, 4, 5.3, 7.1  The collective features and attributes of core critical habitat take >100 years to develop and are required annually (i.e., nest trees) or year-round (i.e., roosting and foraging attributes), so cannot be removed without resulting in destruction.
Fire management activities that involve removal of snags and CWD within old forest portions of <b>core critical habitat</b>	Removal of downed wood (CWD) and snags during fire management activities can result in destruction of core critical habitat through causing direct and permanent loss of the critical features and attributes required for nesting (i.e., nest trees) and foraging (i.e., features required to support prey populations).	Related IUCN-CMP Threat # 7.1  In some cases, it may be necessary to safeguard the longer-term integrity of core critical habitat in areas that are at high risk of catastrophic fire as a consequence of long-term fire suppression, through wildfire risk reduction practices. These may be undertaken without resulting in destruction of core critical habitat provided that removal of irreplaceable old forest attributes such as snags and CWD is avoided.
Activities emitting sounds resulting in an overall sound level $\geq 90$ db or in an increase above ambient levels by $>20$ db* within <b>acoustic critical habitat</b> (e.g., operation of large machinery, use of chainsaws, blasting, operation of large engines and engine brakes, operation of motorized recreational vehicles)	Acoustic disturbance can result in destruction of core critical habitat within nesting areas through displacing Spotted Owls from the habitat and/or disrupting their behaviour such that they can no longer successfully carry out nesting functions.	Related IUCN-CMP Threat # 1, 4, 5.3  Applies only during the Spotted Owl breeding season (February 1 <sup>st</sup> to July 31 <sup>st</sup> ).

\*See [https://www.fws.gov/arcata/es/birds/nso/documents/2020\\_MAMU\\_NSQ\\_Disturbance\\_Guide\\_Combined\\_Final\\_signed.pdf](https://www.fws.gov/arcata/es/birds/nso/documents/2020_MAMU_NSQ_Disturbance_Guide_Combined_Final_signed.pdf) for guidance on interpretation.

## 8. Measuring Progress

The performance indicators presented below provide a way to define and measure progress toward achieving the population and distribution objectives.

1. Human-caused threats that would result in the further loss of habitat needed for recovery (i.e., the critical habitat) have been mitigated.
2. A wild Spotted Owl population of four to six resident individuals exists by 2030.
3. Barred Owl surveillance has regularly taken place at all sites occupied by Spotted Owls and/or where reintroductions are planned, and all detected Barred Owls have been removed.

## 9. Statement on Action Plans

One or more action plans for the Spotted Owl will be published on the Species at Risk Public Registry within 5 years of the finalization of this document.

## 10. References

- Artuso, C., C.S. Houston, D.G. Smith, and C. Rohner. 2013. Great Horned Owl (*Bubo virginianus*), version 2.0. In *The Birds of North America* (A. F. Poole, Editor). Cornell Lab of Ornithology, Ithaca, NY, USA. <https://doi.org/10.2173/bna.372>.
- Barrows, C.W. 1981. Roost selection by Spotted Owls: An adaptation to heat stress. *Condor* 83:302–309.
- Bart, J. 1995. Amount of suitable habitat and viability of Northern Spotted Owls. *Conservation Biology* 9:943–946.
- Blackburn, I.R., A.S. Harestad, J.N.M. Smith, S. Godwin, R. Hentze, and C.B. Lenihan. 2002. Population assessment of the Northern Spotted Owl in British Columbia 1992–2001. BC Ministry of Water, Land and Air Protection, Surrey, BC. 22pp.
- Blackburn, I.R., and S. Godwin. 2003. Status of the Northern Spotted Owl in British Columbia. BC Ministry of Water, Land and Air Protection, Surrey, BC.
- Blackburn, I., B. D’Anjou, J. Fisher, C. Galliazzo, J. Jonker, A. Peter, and L. Waterhouse. 2009. Best Management Practices for Managing Spotted Owl Habitat – A component of the Spotted Owl Management Plan 2. Prepared for the B.C. Ministry of Environment and Ministry of Forests and Range, Victoria, British Columbia. 118 pp (the “B.C. Habitat Management Practices”)
- Blakesley, J.A., A.B. Franklin and R.J. Gutierrez. 1990. Sexual dimorphism in Northern Spotted Owls from northwest California. *Journal of Field Ornithology* 61:320-327.
- Bodine, E., and A. Capaldi. 2017. Can culling Barred Owls save a declining Northern Spotted Owl population? *Natural Resource Modelling*. 2017;30:e12131.
- Boyle, C.A., L. Lavkulich, H. Schreier, and E. Kiss. 1990. Changes in Land Cover and Subsequent Effects on Lower Fraser Basin Ecosystems from 1827 to 1990. *Environmental Management* Vol. 21, No. 2, pp. 185–196.
- B.C. Ministry of Forests, Lands, Natural Resource, Operations and Rural Development (MFLNRORD). 2021. Release Strategy for Spotted Owls in British Columbia: Five-year plan for Barred Owl lethal control and Spotted Owl release trials 2021-2025 – DRAFT Version 1.0. 42 pp.
- B.C. MFLNRORD - Forest Analysis and Inventory Branch. 2019. Variable Density Yield Projection 7 [Online]. <https://www2.gov.bc.ca/gov/content/industry/forestry/managing-our-forest-resources/forest-inventory/growth-and-yield-modelling/variable-density-yield-projection-vdyp>.
- B.C. Parks. 2018. BC Parks 2017/18 Statistics Report. [https://bcparks.ca/research/statistic\\_report/statistic-report-2017-2018.pdf?v=1617494400061](https://bcparks.ca/research/statistic_report/statistic-report-2017-2018.pdf?v=1617494400061).
- BC Wildfire Service, 2024: [Wildfire Season Summary - Province of British Columbia](#)
- Buchanan, J.B. 2004. Managing habitat for dispersing Northern Spotted Owls – are the current management strategies adequate? *Wildlife Society Bulletin* 32:1333-1345.

Campbell, R.W., N.K. Dawe, I. McTaggart-Cowan, J.M. Cooper, G.W. Kaiser, and M.C.E. McNall. 1990. The Birds of British Columbia, Vol. 2, Nonpasserines: Diurnal birds of prey through woodpeckers. Royal British Columbia Museum and Canadian Wildlife Service. Mitchell Press, Vancouver, BC. 636pp.

Carey, A.B. 1991. The biology of arboreal rodents in Douglas-fir forests. General Technical Report. PNW-GTR-276. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 46 pp. (Huff, Mark H.; Holthausen, Richard S.; Aubry, Keith B., tech. coords.; Biology and management of old-growth forests).

Carey, A. B., Horton, S. P., and B.L. Biswell. 1992. Northern Spotted Owls: Influence of Prey Base and Landscape Character. *Ecological Monographs* 62(2): 223–250. <https://doi.org/10.2307/2937094>.

Carey, A.B., J.K. Kershner, B.L. Biswell, and L.D. de Toledo. 1999. Ecological scale and forest development: squirrels, dietary fungi, and vascular plants in managed and unmanaged forests. *Wildlife Monographs* 63(1): 223-250.

Carey, A.B., and K.C. Peeler. 1995. Spotted Owls: Resource and Space Use in Mosaic Landscapes. *Journal of Raptor Research* 29(4): 223-239.

Carey, A.B., T.M. Wilson, C.C. Maguire, and B.L. Biswell. 1997. Dens of Northern Flying Squirrels in the Pacific Northwest. *Journal of Wildlife Management* 61(3):684-699

Chutter, M.J., Blackburn, I., Bonin, D., Buchanan, J., Costanzo, B., Cunnington, D., Harestad, A., Hayes, T., Heppner, D., Kiss, L., Surgenor, J., Wall, W., Waterhouse, L., and Williams, L. 2004. Recovery Strategy for the Northern Spotted Owl (*Strix occidentalis caurina*) in British Columbia. Prepared for the BC Ministry of Environment, Victoria, BC. 74 pp.

Chutter, M.J., I. Blackburn, D. Bonin, J.B. Buchanan, D. Cunnington, L. Feldes, A. Harestad, D. Heppner, L. Kiss, S. Leech, J. Smith, J. Surgenor, W. Wall, L. Waterhouse, and L. Williams, 2007. Guidance and some components of action planning for the Northern Spotted Owl (*Strix occidentalis caurina*) in British Columbia. B.C. Ministry of Environment, Victoria, British Columbia. 92 pp.

Committee on the Status of Endangered Wildlife in Canada (COSEWIC). 2008. COSEWIC assessment and update status report on the Spotted Owl (*Strix occidentalis Caurina*) subspecies, in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa. vii + 48 pp. [www.SARRegistry.gc.ca/status/status\\_e.cfm](http://www.SARRegistry.gc.ca/status/status_e.cfm).

Committee on the Status of Endangered Wildlife in Canada (COSEWIC). 2021. COSEWIC Assessment Process, Categories and Guidelines. Unpublished. Available at: <https://www.cosewic.ca/index.php/en/assessment-process/cosewic-assessment-process-categories-and-guidelines/quantitative-criteria.html>

Conlisk, E., Haeuser, E., Flint, A., Lewison, R.L., and M.K. Jennings. 2020. Pairing functional connectivity with population dynamics to prioritize corridors for Southern California spotted owls. *Diversity and Distributions* 00:1–13. DOI: 10.1111/ddi.13235

Courtney, S.P., J.A. Blakesley, R.E. Bigley, M.L. Cody, J.P. Dumbacher, R.C. Fleischer, A.B. Franklin, J.F. Franklin, R.J. Gutierrez, J.M. Marzluff, and L. Sztukowski. 2004. Scientific evaluation of the status of the Northern Spotted Owl. Sustainable Ecosystems Institute, Portland, Oregon. <http://www.sei.org/owl/finalreport/finalreport.htm>.

Crockatt, M.A. 2012. Are there edge effects on forest fungi and if so do they matter? Fungal Biology Reviews 26: 94-101.

Crozier, M.L., M. E. Seamans, R.J. Gutierrez, P.J. Loschl, R.B. Horn, S.G. Sovern, and E.D. Forsman. 2006. Does the presence of Barred Owls suppress the calling behaviour of Spotted Owls? The Condor 108: 760-769.

Dawson, W., J. Ligon, J. Murphy, J. Myers, D. Simberloff, and J. Verner. 1986. Report of the scientific advisory panel on the Spotted Owl. Condor 89: 205–229.

D'Anjou, B., F. L. Waterhouse, M. Todd, and P. Braumberger. 2015. A systematic review of stand-level forest management for enhancing and recruiting Spotted Owl habitat in British Columbia. Prov. B.C., Victoria, B.C. Tech. Rep. 091. [www.for.gov.bc.ca/hfd/pubs/Docs/Tr/Tr091.htm](http://www.for.gov.bc.ca/hfd/pubs/Docs/Tr/Tr091.htm).

Davis, R.J., Hollen, B., Hobson, J., Gower, J.E. and D. Keenum. 2016. Northwest Forest Plan—the first 20 years (1994–2013): status and trends of northern spotted owl habitats. Gen. Tech. Rep. PNW-GTR-929. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 111 pp.

Diller L.V., K.A. Hamm, E.A. Desiree, K.M. Dugger, C.B. Yackulic, C.J. Schwarz, P.C. Carlson, and T.L. McDonald. 2016. Demographic Response of Northern Spotted Owls to Barred Owl Removal. Journal of Wildlife Management 80(4): 1-17. DOI: 10-1002/jwmg.1046.

Dinerstein, E., C. Vynne, E. Sala, A. R. Joshi, S. Fernando, T. E. Lovejoy, J. Mayorga, D. Olson, G. P. Asner, J. E. M. Baillie, N. D. Burgess, K. Burkart, R. F. Noss, Y. P. Zhang, A. Baccini, T. Birch, N. Hahn, L. N. Joppa, E. Wikramanayake. 2019. A Global Deal For Nature: Guiding principles, milestones, and targets. Sci. Adv.5, eaaw2869. DOI: 10.1126/sciadv.aaw286

Dugger, K.M., R.G. Anthony, and L. S. Andrews. 2011. Transient dynamics of invasive competition: Barred Owls, Spotted Owls, habitat, and the demons of competition present. 2011. Ecological Applications 21(7): 2459-2468.

Dugger, K.M., E.D. Forsman, A.B. Franklin, R.J. Davis, G.C. White, C.J. Schwarz, K.P. Burnham, J.D. Nichols, J.E. Hines, C. B. Yackulic, P.F. Doherty Jr., L. Bailey, D.A. Clark, S.H. Ackers, L.S. Andrews, B. Augustine, B.L. Biswell, J. Blakesley, P.C. Carlson, M.J. Clement, L.V. Diller, E.M. Glenn, A. Green, S.A. Gremel, D.R. Herter, J.M. Higley, J. Hobson, R.B. Horn, K.P. Huyvaert, C. McCafferty, T. McDonald, K. McDonnell, G.S. Olson, J.A. Reid, J. Rockweit, V. Ruiz, J. Saenz and S.G. Sovern. 2015. The effects of habitat, climate, and Barred Owls on long-term demography of Northern Spotted Owls. The Condor 118(1): 57-116.

Dunbar, D., Booth, B., Forsman, E., Hetherington, A., and D. Wilson. 1991. Status of the Spotted Owl (*Strix occidentalis*) and Barred Owl (*Strix varia*) in southwestern British Columbia. Canadian Field-Naturalist 105:464–468.

Dunbar, D., and I. Blackburn. 1994. Management options for the Northern Spotted Owl in British Columbia. Report of the Canadian Spotted Owl Recovery Team. BC Minist. Environ. Lands and Parks, Surrey, BC. 180 pp.

Environment and Climate Change Canada. 2020. Policy on recovery and survival. *Species at Risk Act* Policies and Guidelines Series. Environment and Climate Change Canada, Ottawa. 7 pp. <https://species-registry.canada.ca/index-en.html#/documents/2985> .

Environment and Climate Change Canada. 2016. Policy on Critical Habitat Protection on Non-federal Lands [Proposed]. *Species at Risk Act: Policies and Guidelines Series*. Environment and Climate Change Canada, Ottawa. 9 pp. <https://species-registry.canada.ca/index-en.html#/documents/2987>.

ESTR Secretariat. 2014. Western Interior Basin Ecozone evidence for key findings summary. Canadian Biodiversity: Ecosystem Status and Trends 2010, Evidence for Key Findings Summary Report No. 11. Canadian Councils of Resource Ministers. Ottawa, ON. viii + 106 pp. <http://www.biodivcanada.ca/default.asp?lang=En&n=137E1147-1>.

Fenger, M., J.B. Buchanan, T.J. Cole, E.D. Forsman, S.M. Haig, K. Martin, and W.A. Rapley. 2007. B.C. Ministry of Forests, Land, Natural Resource Operations, Surrey, BC. Spotted Owl Population Enhancement Team, Government of British Columbia pp 1-50.

Forsman, E.D. 1981. Molt of the Spotted Owl. *Auk* 98:735-742.

Forsman, E.D., E.C. Meslow, and H.M. Wight. 1984. Distribution and biology of the Spotted Owl in Oregon. *Wildlife Monographs* 87:1–64.

Forsman, E.D., and A.R. Giese. 1997. Nests of the Northern Spotted Owl on the Olympic Peninsula, Washington. *Wilson Bulletin* 109:28–41.

Forsman, E.D., I.A. Otto, S.G. Sovern, M. Taylor, D.W. Hays, H. Allen, S.L. Roberts, and D.E. Seaman. 2001. Spatial and temporal variation in diets of Spotted Owls in Washington. *Journal of Raptor Research* 35:141–150.

Forsman, E.D., R.G. Anthony, J.A. Reid, Loschl, P.J., Sovern, S.G., Taylor, M., Biswell, B.L., Ellingson, A., Meslow, E.C., Miller, G.S., Swindle, K.A., Thraikill, J.A., Wagner, F.F., and D.E. Seaman. 2002. Natal and breeding dispersal of northern spotted owls. *Wildlife Monographs*. 149: 1-35.

Franklin, A.B. 2001. Population regulation in Northern Spotted Owls: theoretical implications for management. In: McCullough D.R., Barrett R.H. (eds) *Wildlife 2001: Populations*. Springer, Dordrecht. [https://doi.org/10.1007/978-94-011-2868-1\\_62](https://doi.org/10.1007/978-94-011-2868-1_62)

Franklin, A.B., K.M. Dugger, D.B. Lesmeister, R.J. Davis, J.D. Wiens, G.C. White, J.D. Nichols, J.E. Hines, C.B. Yackulic, C.J. Schwarz, S.H. Ackers, L.S. Andrews, L.L. Bailey, R. Bown, J. Burgher, K.P. Burnham, P.C. Carlson, T. Chestnut, M.M. Conner, K.E. Dillione, E.D. Forsman, E.M. Glenn, S.A. Gremel, K.A. Hamm, D.R. Herter, J.M. Higley, R.B. Horn, J.M. Jenkins, W.L. Kendall, D.W. Lamphear, C. McCafferty, T.L. McDonald, J.A. Reid, J.T. Rockweit, D.C. Simon, S.G. Sovern, J.K. Swingle, and H. Wise. 2021. Range-wide declines of northern spotted owl populations in the Pacific Northwest: A meta-analysis. *Biological Conservation* 259:109168. <https://doi.org/10.1016/j.biocon.2021.109168>.

Gauslaa, Y., P. Bartemucci, and K. Asbjørn Solhauga. 2018. Forest edge-induced damage of cephalo- and cyanolichens in northern temperate rainforests of British Columbia. *Canadian Journal of Forest Research*. 49: 434-9.

Gillis, J. 2016a. The Enhancement of Spotted Owl Habitats by Removing Competition Effects Associated with Barred Owls. Project HSP7141. Final Report prepared for the Habitat Stewardship Program. Prepared by B.C. Ministry of Forests, Lands and Natural Resource Operations.

Gillis, J. 2016b. Restoration of the Lillooet Sub-Population of Spotted Owls in British Columbia - Final Report Project 16.W.BRG.13. Prepared for the Fish and Wildlife Compensation Program by B.C. Ministry of Forests, Lands and Natural Resource Operations.

Gillis, J. 2017. Restoration of the Lillooet Sub-Population of Spotted Owls in British Columbia - Final Report Project COA-F17-W-1314. Prepared for the Fish and Wildlife Compensation Program by B.C. Ministry of Forests, Lands and Natural Resource Operations.

Gillis, J. 2018. Restoration of the Lillooet Sub-Population of Spotted Owls in British Columbia. Final Report Project COA-F18-2409. Prepared for the Fish and Wildlife Compensation Program by BC Ministry of Forests, Lands and Natural Resource Operations.

Gillis, J. and F.L. Waterhouse. 2020. Barred Owl removal report 2007–2016. Prov. B.C., Victoria, B.C. Tech. Rep. 128. [www.for.gov.bc.ca/hfd/pubs/Docs/Tr/Tr128.htm](http://www.for.gov.bc.ca/hfd/pubs/Docs/Tr/Tr128.htm).

Government of British Columbia. 2024. Wildfire Season Summary. [Wildfire Season Summary - Province of British Columbia \(gov.bc.ca\)](https://www2.gov.bc.ca/gov/content/safety/wildfire/wildfire-season-summary)

Government of British Columbia. 2023a. Joint statement on death of two spotted owls released into the wild in 2022. News release. <https://news.gov.bc.ca/releases/2023WLRS0027-000729>

Government of British Columbia. 2023b. Province continues spotted owl recovery efforts with release of two birds into the wild. News release. <https://news.gov.bc.ca/releases/2023WLRS0053-001493>

Government of British Columbia. 2022. World's first recovery effort sees spotted owls released into the wild for the first time. News release. <https://news.gov.bc.ca/releases/2022LWRS0056-001572>

Government of British Columbia. 2021. Stewardship Baseline Objectives Tool (SBOT) – version 1.1 - DRAFT. <https://governmentofbc.maps.arcgis.com/apps/MapSeries/index.html?appid=8e015f61714c410f93ab8f16ce714ae5>.

Government of British Columbia. 2023. Approved Wildlife Habitat Areas (WHAs). <http://www.env.gov.bc.ca/cgi-bin/apps/faw/wharesult.cgi?search=species&species=spotted+owl&speciesname=english&submit2=Search>.

Government of British Columbia. 2009. British Columbia 2009 Spotted Owl Management Plan (SOMP 2): Rationale for Revisions to the 1997 Spotted Owl Management Plan (SOMP 1). <https://www2.gov.bc.ca/assets/gov/environment/natural-resource-stewardship/resource-stewardship-tools/sbot/somp2.pdf>.

Gutiérrez, R.J., A.B. Franklin, and W.S. Lahaye. 1995. Spotted Owl (*Strix occidentalis*), version 2.0. In the Birds of North America (A. F. Poole and F. B. Gill, Editors). Cornell Lab of Ornithology, Ithaca, NY, USA. <https://doi.org/10.2173/bna.179>.

Hamer, T.E., E.D. Forsman, and E.M. Glenn. 2007. Home range attributes and habitat selection of Barred Owls and Spotted Owls in an area of sympatry. *The Condor* 109: 750–768.

Harper, W.L., and R. Milliken. 1994. Other species associated with late successional forests. Appendix C. in D. Dunbar and I. Blackburn, Management options for the Northern Spotted Owl in British Columbia. Report of the Canadian Spotted Owl Recovery Team. BC Minist. Environ. Lands and Parks, Surrey, BC. 180pp.

Hayward, L.S., A. Bowles, J.C. Ha, and S.K. Wasser. 2011. Impacts of acute and long-term vehicle exposure on physiology and reproductive success of the northern spotted owl. *Ecosphere* 2(6):art65. doi:10.1890/ES10-00199.1

Hermosilla, T., M.A. Wulder, J.C. White, N.C. Coops, and G.W. Hobart. 2015a. An integrated Landsat time series protocol for change detection and generation of annual gap-free surface reflectance composites. *Remote Sensing of Environment* 158: 220-234. Data: [https://opendata.nfis.org/mapserver/nfis-change\\_eng.html](https://opendata.nfis.org/mapserver/nfis-change_eng.html).

Hermosilla, T., M.A. Wulder, J.C. White, N.C. Coops, and G.W. Hobart. 2015b. Regional detection, characterization, and attribution of annual forest change from 1984 to 2012 using Landsat-derived time-series metrics. *Remote Sensing of Environment* 170: 121-132. Data: [https://opendata.nfis.org/mapserver/nfis-change\\_eng.html](https://opendata.nfis.org/mapserver/nfis-change_eng.html).

Hermosilla, T., M.A. Wulder, J.C. White, N.C. Coops, and G.W. Hobart. 2017. Updating Landsat time series of surface-reflectance composites and forest change products with new observations. *International Journal of Applied Earth Observation and Geoinformation* 63:104-111. Data: [https://opendata.nfis.org/mapserver/nfis-change\\_eng.html](https://opendata.nfis.org/mapserver/nfis-change_eng.html).

Hobbs, J. 2004. Spotted Owl Nest Site Descriptions (2002 and 2003) and Telemetry and Monitoring of Juvenile Spotted Owls (2003 and 2004). Internal report Prepared for Biodiversity Branch, Ministry of Water, Land and Air Protection. 38 pp.

Hobbs, J. 2005. Spotted Owl Inventory and Nest Site Descriptions (2004) and Telemetry and Monitoring of Juvenile Spotted Owls (2004 and 2005). Internal report Prepared for Biodiversity Branch, Ministry of Water, Land and Air Protection. 40 pp.

Hofstadter, D.F., N.F. Kryshak, C.M. Wood, B.P. Dotters, K.N. Roberts, K.G. Kelly, J.J. Keane, S.C. Sawyer, P.A. Shaklee, H.A. Kramer, R.J. Gutiérrez, and M.Z. Peery. 2022. Arresting the spread of invasive species in continental systems. *Frontiers in Ecol & Environ* 1–7. <https://doi.org/10.1002/fee.2458>

Hollenbeck, J.P., S.M. Haig, E.D. Forsman, and J.D. Wiens. 2018. Geographic variation in natal dispersal of Northern Spotted Owls over 28 years. *The Condor* 120(3): 530-542.

Horoupian, N., C.B. Lenihan, A.S. Harestad, and I.R. Blackburn. 2004. Diet of Northern Spotted Owls in British Columbia. Report prepared for B.C. Ministry of Water, Land and Air Protection. 13 pp.

Irwin, L.L., D.F. Rock, and G.P. Millers. 2024. Stand Structures Used by Northern Spotted Owls in Managed Forests. *Journal of Raptor Research*: Vol. 34 : Iss. 3 , Article 3.  
<https://digitalcommons.usf.edu/jrr/vol34/iss3/3>

Jenkins, J. M. A., D. B. Lesmeister, E. D. Forsman, K. M. Dugger, S. H. Ackers, L. S. Andrews, S. A. Gremel, B. Hollen, C. E. McCafferty, M. S. Pruett, J. A. Reid, S. G. Sovern, and J. D. Wiens. 2021. Conspecific and congeneric interactions shape increasing rates of breeding dispersal of northern spotted owls. *Ecological Applications* 00(00):e02398. 10.1002/eap.2398.

Jenkins, J.M.A., D.B. Lesmeister, E.D. Forsman, K.M. Dugger, S.H. Ackers, L.S. Andrews, C.E. McCafferty, M.S. Pruett, J.A. Reid, S.G. Sovern, R.B. Horn, S.A. Gremel, J.D. Wiens, and Z. Yang. 2019. Social status, forest disturbance, and Barred Owls shape long-term trends in breeding dispersal distance of Northern Spotted Owls. *Condor* 121: 1–17.

Johnson, D.H. 1993. Spotted Owls, Great Horned Owls, and forest fragmentation in the central Oregon Cascades. M.S. thesis, Oregon State Univ., Corvallis, OR, U.S.A.

Kelly, E.G., E.D. Forsman, and R.G. Anthony. 2003. Are Barred Owls displacing Spotted Owls? *Condor* 105:45–53.

Kelly, E.G., and E.D. Forsman. 2004. Recent records of hybridization between barred owls (*Strix varia*) and northern spotted owls (*S. occidentalis caurina*). *Auk* 121:806–810.

Krawchuk, M.A., G.W. Meigs, J.M. Cartwright, J.D. Coop, R. Davis, A. Holz, C. Kolden, and A.J.H. Meddens. 2020. Disturbance refugia within mosaics of forest fire, drought, and insect outbreaks. *Frontiers in Ecology and the Environment* 18(5):235–244, doi:10.1002/fee.2190.

Kremsater, L. and F.L. Bunnell. 1999. Edge effects: theory, evidence and implications to management of western North American forests *in* Forest fragmentation: wildlife and management implications. J.A. Rochelle, L.A. Lehmann, J. Wisniewski (Eds.). Koninklijke Brill NV, Leiden, Netherlands. pp. 117-153.

Kroll, A.J., J.E. Jones, A.B. Stringer, and D.J Meekins. 2016. Multistate Models Reveal Long-Term Trends of Northern Spotted Owls in the Absence of a Novel Competitor. *PLoS ONE* 11(4): e0152888. doi:10.1371/journal.pone.0152888

Lamberson, R.H., B.R. Noon, C. Voss, and R. McKelvey. 1994. Reserve design for terrestrial species: The effects of patch size and spacing on the viability of the Northern Spotted Owl. *Conservation Biology* 8:185–195.

Leskiw, T. and R.J. Gutiérrez. 1998. Possible predation of a spotted owl by a barred owl. *Western Birds* 29(3):225-226.

Lesmeister, D.B., R.J. Davis, S.G. Sovern and Z. Yang. 2021. Northern spotted owl nesting forests as fire refugia: a 30-year synthesis of large wildfires. *Fire Ecology*: 17, 32.  
<https://doi.org/10.1186/s42408-021-00118-z>

Littell, J.S., E.E. Oneil, D. McKenzie, J.A. Hicke, J.A. Lutz, R.A. Norheim, and M.M. Elsner. 2010. Forest ecosystems, disturbance, and climatic change in Washington State, USA. *Climatic Change*: 102(1–2): 129–158. doi:10.1007/s10584-010-9858-x

Livezey, K.B. 2009a. Range Expansion of barred owls, Part I: Chronology and Distribution. *The American Midland Naturalist* 161(1): 49-56.

Livezey, K.B. 2009b. Range Expansion of barred owls, Part II: Facilitating Ecological Changes. *The American Midland Naturalist* 161(2): 323-349.

Long, L. L. and J.W. Wolfe. 2019. Review of the Effects of Barred Owls on Spotted Owls. *The Journal of Wildlife Management*, 83 (6), 1281–1296. <https://doi.org/10.1002/jwmg.21715>

Marcot, B.G., M.G. Raphael, N.H. Schumaker, and B. Galleher. 2013. How big and how close? Habitat patch size and spacing to conserve a threatened species. *Natural Resource Modeling* 26(2): 194-214.

McCulligh, J. 2024. Northern Spotted Owl Captive Breeding Program: Final Report for 2023/2024 – Project No. COA-F24-W-3891. Prepared for the Fish and Wildlife Compensation Program. <http://a100.gov.bc.ca/pub/siwe/details.do?id=5467>

McCulligh, J. 2019. Northern Spotted Owl Captive Breeding Program: Final Report for 2018/2019 - Project No. COA-F19-W-2676. Prepared for the Fish and Wildlife Compensation Program. <http://a100.gov.bc.ca/pub/siwe/details.do?id=5467>

NatureServe. 2021. NatureServe Explorer: An online encyclopedia of life [web application]. Version 7.1. NatureServe, Arlington, Virginia. <http://explorer.natureserve.org>.

North, M., G. Steger, R. Denton, G. Eberlein, T. Munton, and K. Johnson. 2000. Association of weather and nest-site structure with reproductive success in California Spotted Owls. *The Journal of Wildlife Management* 64:797–807.

Northern Spotted Owl Breeding Program (NSOBP). n.d.. History. <https://www.nsobreedingprogram.com/history>

Perlman, K.R. 2017. Using a Two-Species Individual-Based Model to Examine the Population Responses of Northern Spotted Owls to Experimental Removals of Barred Owls in the Pacific Northwest. Thesis, Oregon State University, Corvallis, USA [https://ir.library.oregonstate.edu/concern/graduate\\_thesis\\_or\\_dissertations/6969z5287](https://ir.library.oregonstate.edu/concern/graduate_thesis_or_dissertations/6969z5287)

Price, K. and D. Daust. 2016. Climate Change Vulnerability of BC's Fish and Wildlife: First Approximation. B.C. Ministry of Forests, Lands, and Natural Resource Operations - Competitiveness and Innovation Branch. <https://www2.gov.bc.ca/assets/gov/environment/natural-resource-stewardship/nrs-climate-change/adaptation/climate20change20vulnerability20of20bcs20fish20and20wildlife20final20june6.pdf>.

Pritchard et. al. 2021. Adapting North American forests to climate change and wildfires: 10 common questions. *Ecological Applications* 31 (8) :e02433.

Sovern, S.G., Forsman, E.D., Olson, G.S., Biswell, B.L., Taylor, M., and R.G. Anthony. 2014. Bared Owls and Landscape Attributes Influence Territory Occupancy of Northern Spotted Owls. *The Journal of Wildlife Management* 78(8): 1436-1443.

Spies, T.A., P.A. Stine, R. Gravenmier, J.W. Long, and M.J. Reilly (technical coordinators). 2018. Synthesis of science to inform land management within the Northwest Forest Plan area. General Technical Report. PNW-GTR-966. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 1020 p. 3 vol.

Spotted Owl Management Inter-Agency Team (SOMIT). 1997. Spotted Owl management plan: Strategic component. B.C. Minist. Environment, Lands and Parks and B.C. Ministry of Forests, Victoria, BC. 81 pp.

Sutherland, G.D., D. O'Brien, A. Fall, F.L. Waterhouse, A. Harestad, and J.B. Buchanan (editors). 2007. A framework to support landscape analyses of habitat supply and effects on populations of forest-dwelling species: a case study based on the Northern Spotted Owl. Technical Report 038. British Columbia Ministry of Forests and Range, Research Branch, Victoria, British Columbia. <https://www.for.gov.bc.ca/hfd/pubs/Docs/Tr/Tr038.pdf>.

Thom, D., Golivets, M., Edling, L., Meigs, G.W., Gourevitch, J.D., Sonter, L.J., Galford, G.L., Keeton, W.S. 2019. The climate sensitivity of carbon, timber, and species richness covaries with forest age in boreal–temperate North America. *Global Change Biology* 25:2446-2458 <https://doi.org/10.1111/gcb.14656>.

Thomas, J.W., E.D. Forsman, J.B. Lint, E.C. Meslow, B.R. Noon, and J. Verner. 1990. A Conservation Strategy for the Northern Spotted Owl. Rep. of Interagency Scientific Committee to Address the Conservation of the Northern Spotted Owl (Portland, OR). 427 pp + maps.

U.S. Department of the Interior (USDI). 1992. Recovery plan for the Northern Spotted Owl: Draft. U.S. Fish and Wildlife Service, Washington, DC. 662 pp + maps.

USFWS. 2024. Final Bared Owl Management Strategy. [https://www.fws.gov/sites/default/files/documents/2024-08/final-bared-owl-management-strategy-2024\\_508.pdf](https://www.fws.gov/sites/default/files/documents/2024-08/final-bared-owl-management-strategy-2024_508.pdf)

USFWS. 2020. Revised Transmittal of Guidance: Estimating the Effects of Auditory and Visual Disturbance to Northern Spotted Owls and Marbled Murrelets in Northwestern California. [https://www.fws.gov/arcata/es/birds/nso/documents/2020\\_MAMU\\_NSO\\_Disturbance\\_Guide\\_Combined\\_Final\\_signed.pdf](https://www.fws.gov/arcata/es/birds/nso/documents/2020_MAMU_NSO_Disturbance_Guide_Combined_Final_signed.pdf).

USFWS. 2011. Revised recovery plan for the northern spotted owl (*Strix occidentalis caurina*). Portland, OR, USA: U.S. Department of Interior.

USGS. 2021. Fire Refugia in Old-Growth Forests: Predicting Habitat Persistence to Support Land Management in an Era of Rapid Global Change. [https://www.usgs.gov/centers/casc-sc/science/fire-refugia-old-growth-forests-predicting-habitat-persistence-support-land?qt-science\\_center\\_objects=0#qt-science\\_center\\_objects](https://www.usgs.gov/centers/casc-sc/science/fire-refugia-old-growth-forests-predicting-habitat-persistence-support-land?qt-science_center_objects=0#qt-science_center_objects).

- Van Lanen, N. J., A. B. Franklin, K. P. Huyvaert, R. F. Reiser, and P.C. Carlson. 2011. Who hits and hoots at whom? Potential for interference competition between Barred and Northern Spotted owls. *Biological Conservation* 144:2194–2201.
- Wang, T., A. Hamann, D. Spittlehouse, and C. Carroll. 2016. Locally Downscaled and Spatially Customizable Climate Data for Historical and Future Periods for North America. *PLoS One* 11(6): e0156720.
- Wasser, S.K., K. Bevis, G. King, and E. Hanson. 1997. Noninvasive Physiological Measures of Disturbance in the Northern Spotted Owl. *Conservation Biology* 11:4, 1019-1022. <https://conbio.onlinelibrary.wiley.com/doi/epdf/10.1046/j.1523-1739.1997.96240.x>.
- Waterhouse, F.L., I.A. Manley, A.S. Harestad, and P.K. Ott. 2012. Nest structures and habitats of the northern spotted owl in three ecological subregions of British Columbia. Prov. B.C., Victoria, B.C. Tech Rep 069. [www.for.gov.bc.ca/hfd/pubs/Docs/Tr/Tr069.htm](http://www.for.gov.bc.ca/hfd/pubs/Docs/Tr/Tr069.htm).
- Waters, J.R. and C.J. Zabel. 1995. Northern flying squirrel densities in fir forests of northeastern California. *Journal of Wildlife Management*: 59, pp. 858-866.
- Wiens, J.D., R.A. Anthony, and E.D. Forsman. 2014. Competitive interactions and resource partitioning between northern spotted owls and barred owls in western Oregon. *Wildlife Monographs* 185: 1-50.
- Wiens, J.D., K. M. Dugger, D. B. Lesmeister, K. E. Dilione, and D. C. Simon. 2020. Effects of Barred Owl (*Strix varia*) Removal on Population Demography of Northern Spotted Owls (*Strix occidentalis caurina*) in Washington and Oregon—2019 Annual Report. US Department of Interior, and US Geological Survey. p.1-28.
- Wiens, J.D., J. Dugger, J. Higley, D.B. Lesmeister, A.B. Franklin, K.A. Hamm, G.C. White, K.E. Dilione, D.C. Simon, R.R. Bown, P.C. Carlson, C.B. Yackulic, J.D. Nichols, J.E. Hines, R.J. Davis, D.W. Lamphear, C. McCafferty, T.L. McDonald, and S.G. Sovern. 2021. Invader removal triggers competitive release in a threatened avian predator. *Proceedings of the National Academy of Sciences* 118 (31) e2102859118; DOI: 10.1073/pnas.2102859118.
- Wilk, R.J., D.B. Lesmeister, and E.D. Forsman. 2018. Nest trees of northern spotted owls (*Strix occidentalis caurina*) in Washington and Oregon, USA. *PLoS ONE* 13(5): e0197887. <https://doi.org/10.1371/journal.pone.0197887>.
- Wilson, T.M. and E.D. Forsman. 2013. Thinning effects on Spotted Owl prey and other forest-dwelling small mammals. In: *Density management in the 21st century: west side story* P.D. Anderson and K.L. Ronnenberg (editors). U.S. Department of Agriculture, Forest Service, Portland Oregon. General Technical Report. pnw-gtr-880, pp. 79–90. [www.fs.fed.us/pnw/pubs/pnw\\_gtr880.pdf](http://www.fs.fed.us/pnw/pubs/pnw_gtr880.pdf).
- Yackulic, C.B., L.L. Bailey, K.M. Dugger, R.J. Davis, A.B. Franklin, E.D. Forsman, S.H. Ackers, L.S. Andrews, L.V. Diller, S.A. Gremel, K.A. Hamm, D.R. Herter, J.M. Higley, R.B. Horn, C. McCafferty, J.A. Reid, J.T. Rockweit, and S.G. Sovern. 2019. The past and future roles of competition and habitat in the range-wide occupancy dynamics of Northern Spotted Owls. *Ecological Applications* 29(3):e01861. 10.1002/eap.1861

## 11. Personal Communications

Ian Blackburn. *Manager, Resource Stewardship for Ecosystems - B.C. Ministry of Forests, Lands, Natural Resource Operations and Rural Development. 2021.*

Joseph Buchanan. *Wildlife Biologist/Natural Resource Scientist – Washington Department of Fish and Wildlife. 2019.*

Joel Gillis. *Spotted Owl Biologist – B.C. Ministry of Forests, Lands, Natural Resource Operations and Rural Development. 2019.*

Joel Gillis. *Spotted Owl Biologist – B.C. Ministry of Forests, Lands, Natural Resource Operations and Rural Development. 2020.*

Joel Gillis. *Spotted Owl Biologist – B.C. Ministry of Forests, Lands, Natural Resource Operations and Rural Development. 2021.*

Government of B.C representatives, *B.C. Minister of Water, Lands and Natural Resource Operations.* to Honourable Stephen Guilbeault, Minister of Environment and Climate Change. 2023.

Joanna Hirner. *Conservation Specialist – B.C. Parks. 2020.*

## Appendix A: Effects on the Environment and Other Species

A strategic environmental assessment (SEA) is conducted on all SARA recovery planning documents, in accordance with the [Cabinet Directive on the Environmental Assessment of Policy, Plan and Program Proposals](#)<sup>15</sup>. The purpose of a SEA is to incorporate environmental considerations into the development of public policies, plans, and program proposals to support environmentally sound decision-making and to evaluate whether the outcomes of a recovery planning document could affect any component of the environment or any of the [Federal Sustainable Development Strategy](#)'s<sup>16</sup> (FSDS) goals and targets.

Recovery planning is intended to benefit species at risk and biodiversity in general. However, it is recognized that strategies may also inadvertently lead to environmental effects beyond the intended benefits. The planning process based on national guidelines directly incorporates consideration of all environmental effects, with a particular focus on possible impacts upon non-target species or habitats. The results of the SEA are incorporated directly into the strategy itself, but are also summarized below in this statement.

Conservation of habitat for the Spotted Owl will benefit a multitude of vertebrate, invertebrate, and plant species that use mature and old-growth coniferous forests. Harper and Milliken (1994) concluded there were approximately 71 species of vertebrates closely associated with late-successional and old forests within the range of the Spotted Owl in Canada (four amphibians, 34 birds, 17 mammals, and 16 fish). Examples of other species at risk whose habitats overlap with those of Spotted Owls include Marbled Murrelet (*Brachyramphus marmoratus*), Northern Goshawk (*Accipiter gentilis*), and Western Screech-Owl (*Megascops kennicottii*) *kennicottii* and *macfarlanei* subspecies. The large landscapes required to manage and conserve populations of Spotted Owls lend themselves to application of ecosystem-based approaches to forest management. The conservation of habitat for the Spotted Owl will help maintain functioning late-successional forest ecosystems, and help regulate water and nutrient cycles. Further, many of the old-growth stands conserved for the Spotted Owl are likely to function as refugia as climate change-mediated disturbances increase in frequency and extent. Protection of these areas for Spotted Owls will also improve the climate change resilience of other old-forest-dependent species.

Barred Owls are also known to compete with or prey upon a number of other native species, including species at risk. Predation by Barred Owls is one of the highest-level IUCN-CMP threats for Western Screech-Owl *kennicottii* and *macfarlanei* subspecies, the ranges of which overlap significantly with that of the Spotted Owl. Control of Barred Owls within the Spotted Owl range will therefore also support recovery of Western Screech-Owl.

---

<sup>15</sup> [www.canada.ca/en/environmental-assessment-agency/programs/strategic-environmental-assessment/cabinet-directive-environmental-assessment-policy-plan-program-proposals.html](http://www.canada.ca/en/environmental-assessment-agency/programs/strategic-environmental-assessment/cabinet-directive-environmental-assessment-policy-plan-program-proposals.html)

<sup>16</sup> [www.fsd-sfdd.ca/index.html#/en/goals/](http://www.fsd-sfdd.ca/index.html#/en/goals/)