



Guide on best practices for seaweed harvesting in Quebec



Fanny Noisette

Fucus sp. assemblage at Anse-aux-Fraises in the clear waters of Anticosti Island, QC

Fisheries and Oceans Canada
Regional Science Branch
Maurice Lamontagne Institute
Quebec region

© His Majesty the King of Canada, represented by Minister
of the Department of Fisheries and Oceans, 2025

Cat. no. Fs124-13/2025E-PDF

ISBN 978-0-660-76677-5

Authors: Stéphanie Roy, Virginie Roy, Andrew Smith and Rénaud Belley

Fisheries and Oceans Canada
Regional Science Branch
Quebec region
Maurice Lamontagne Institute
850, route de la Mer
Mont-Joli, Quebec
G5H 3Z4


This guide was developed from the following literature review (in preparation) :
Roy, S., Roy, V., Smith, A. et Belley, R. 2025. Revue de la biologie et des conditions de
récolte des macroalgues de l'Atlantique Nord : suggestions de modifications et d'ajouts
pour le Québec. Rapp. tech. can. sci. halieut. aquat. 0000 : v + x p.

Table of contents

Foreword.....	3
Importance of seaweeds.....	4
List of commercial species in Quebec.....	5
Vertical distribution - where to find seaweed species?.....	6
General practices for sustainable harvest.....	7
Brown seaweeds	
Sea colander (<i>Agarum clathratum</i>).....	8
Sugar kelp (<i>Saccharina latissima</i>).....	9
<i>Hedophyllum nigripes</i>	10
Oarweed (<i>Laminaria digitata</i>).....	11
Winged kelp (<i>Alaria esculenta</i>).....	12
<i>Saccorhiza dermatodea</i>	13
Knotted wrack (<i>Ascophyllum nodosum</i>).....	14
Fucus (<i>Fucus</i> spp.).....	15
Sea lace (<i>Chorda filum</i>).....	16
<i>Chordaria flagelliformis</i>	17
Red seaweeds	
Irish moss (<i>Chondrus crispus</i>).....	18
Dulse (<i>Palmaria palmata</i>).....	19
<i>Devaleraea ramentacea</i>	20
Nori (<i>Porphyra</i> spp.).....	21
Green seaweeds	
Sea lettuce (<i>Ulva</i> spp.).....	22
Summary table.....	23
References.....	25

Foreword

This guide was developed for seaweed harvesters and anyone wishing to deepen their knowledge on seaweed harvesting practices. It promotes sustainable seaweed management in Quebec in the maritime zone illustrated in turquoise on the map. This guide is based on a literature review addressing seaweed harvesting conditions of multiple species harvested in Quebec and in similar climatic regions (Roy et al. 2025, in preparation).



Estuary and Gulf of St. Lawrence

This guide aims to raise awareness on the importance of sustainable seaweed harvesting, with a focus on harvesting techniques respecting seaweed ecosystems. It is important to note that this guide is intended to serve as an informative resource promoting appropriate seaweed harvesting practices. However, it does not constitute an official regulation. The recommendations are designed to complement and not to replace existing regulations.




Stéphanie Roy

Kelp beds at low tide, Anse-au-Lard, Rimouski, QC

Importance of seaweeds

Seaweed ecosystems provide numerous services to humanity, also called ecosystem services, such as oxygen production, carbon dioxide fixation, as well as various marine species, using them as nurseries and refuges from predators. To preserve these services, it is crucial to maintain the integrity of these ecosystems during harvesting. Sustainable harvesting requires particular attention to harvesting techniques, which must be adapted to the biology of seaweeds. These techniques will be explained in more detail below. Prevailing conditions at harvest sites must also be considered. For example, if the harvest site is located near a coastal erosion zone, subject to a particularly high rate of herbivory or exposed to strong dislodgement following a storm, it is recommended to limit the harvest as much as possible at the site until the situation stabilizes to preserve ecosystem services.



List of commercial species in Quebec

Brown seaweeds



- **Sea colander**
(*Agarum clathratum*)
- **Sugar kelp**
(*Saccharina latissima*)
- ***Hedophyllum nigripes***
- **Oarweed**
(*Laminaria digitata*)
- **Winged kelp**
(*Alaria esculenta*)
- ***Saccorhiza dermatodea***
- **Knotted wrack**
(*Ascophyllum nodosum*)
- **Fucus**
(*Fucus* spp.)
- **Sea lace**
(*Chorda filum*)
- ***Chordaria flagelliformis***

Dominates the coast of the Estuary and Gulf of St. Lawrence

Air vesicles keeping an upright position in the water

Red seaweeds



- **Irish moss**
(*Chondrus crispus*)
- **Dulse**
(*Palmaria palmata*)
- ***Devaleraea ramentacea***
- **Nori**
(*Porphyra* spp.)

Specialized pigments allowing them to colonise the deep waters

Often attached to a living support (e.g. kelp stipe)

Non-flagellated sexual cells limiting their dispersal

Green seaweeds




- **Sea lettuce**
(*Ulva* spp.)

Can proliferate quickly

Can be a pollution indicator when in very large quantity

Vertical distribution - where to find the species?



Not all species described in this guide are shown in this figure.

General practices for sustainable harvest


For all seaweed species, harvest suggestions go as follow:

Harvest site (in order of importance)

- **Harvest a maximum of 33% of the biomass per species per site (one blade out of three)**
Conservation of the biomass and associated benefits
- **Favour harvesting during the growth phase and avoid harvesting during reproduction**
Maintains biomass through rapid regrowth following harvest and better population renewal through more efficient reproduction
- **Leave the holdfast attached to the substrate. If problematic, uprooting of holdfasts of a maximum of 10% of the harvested biomass per species may be acceptable**
Maximum loss to conserve benefits and promote effective regrowth
- **Apply fallow periods**
Allows the population to return to the pre-harvest state
- **Maintain an unharvested zone (buffer zone) between each harvested area, ensuring that each zone is of similar size (mosaic harvest)**
Allows more efficient recolonization due to proximity to an unexploited zone
- **Avoid harvesting fertile individuals**
Ensures better population renewal through efficient reproduction
- **Avoid harvesting juveniles**
More effective population regeneration

Harvested individual (in order of importance)

- **Cut above the meristem**
Leaving the meristem in place allows for faster regrowth of the blade
- **Avoid harvesting reproductive structures (leave at least an intact branch)**
Ensures better population renewal through efficient reproduction
- **Use sharp object**
Less damaging to the seaweed and less risk of accidental uprooting



Sea colander (*Agarum clathratum*)



Other names: *Agarum cibrosum*

Life expectancy: Several years

Age of maturity: n.a.

Reproduction period: Autumn, winter

Fertile individual: Presence of a dull beige area on the blade

Growth period: Spring, summer

Meristem: Basal area of the blade



One of the few species to be less grazed by the green sea urchin due to chemical compounds that decrease its edibility.



Christian Nozais

Anticosti Island, QC

Specific sustainable harvesting practices

Harvest a maximum of 33% of the biomass (one blade out of three)

Conservation of biomass and associated benefits

Cut at a minimum of 10 cm above the stipe and blade junction

Leaving the meristem in place allows for faster regrowth of the blade

Favour harvesting during growth from May to July

Maintains biomass through rapid regrowth

Avoid harvesting during reproduction from September to February

More efficient reproduction leads to better population renewal

Fallow period of 2 years

Population recovery to the pre-harvest state

Sugar kelp (*Saccharina latissima*)



Other names: Sweet kombu, *Saccharina longicuris*, *Laminaria saccharina*, *Laminaria longicuris*

Life expectancy: 2 to 5 years

Age of maturity: 8 to 15 months

Reproduction period: Autumn

Fertile individual: Presence of a dark brown area in the center of the blade

Growth period: Spring, summer

Meristem: Basal area of the blade



The blade can be more or less undulated depending on the wave exposure, where sheltered environments have more undulated blades than exposed environments.



**Brown
algae**

Anticosti Island, QC

Christian Nozais

Specific sustainable harvesting practices

Harvest a maximum of 33% of the biomass (one blade out of three)

Conservation of biomass and associated benefits

Cut at a minimum of 20 cm above the stipe and blade junction

Leaving the meristem in place allows for faster regrowth of the blade

Favour harvesting during growth from May to July

Maintains biomass through rapid regrowth

Avoid harvesting during reproduction in September and October

More efficient reproduction leads to better population renewal

Fallow period of 2 years

Population recovery to the pre-harvest state

Harvest within a 15 m diameter area and leave 15 m between harvested areas

Allows for more effective recolonization due to proximity to an unharvested area

Hedophyllum nigripes



Other names: -

Life expectancy: Several years

Age of maturity: n.a.

Reproduction period: n.a. (Possibly similar to other kelp species, like *L. digitata*)

Fertile individual: Presence of dark brown areas on the blade

Growth period: n.a. (Possibly similar to other kelp species, like *L. digitata*)

Meristem: Basal area of the blade



Can easily be misidentified as *L. digitata*, because they are morphologically similar. It is possible to differentiate the two species by the presence of mucus on the stipe compared to *L. digitata*.



Brown algae

Hansneset, Spitsbergen, Norway
Modified from Dankworth et al. 2020

Specific sustainable harvesting practices

Harvest a maximum of 33% of the biomass (one blade out of three)

Conservation of the biomass and associated benefits

Cut at a minimum of 20 cm above the stipe and blade junction

Leaving the meristem in place allows for faster blade regrowth

Favour harvesting during growth from May to July

Maintains biomass through rapid regrowth

Avoid harvesting during reproduction in September and October

More efficient reproduction leads to better population renewal

Fallow period of 2 years

Population recovery to the pre-harvest state

Harvest within a 15 m diameter area and leave 15 m between harvested areas

Allows for more effective recolonization due to proximity to an unharvested area

Oarweed (*Laminaria digitata*)



Other names: Digitate kelp, horsetail kelp, tangle

Life expectancy: 3 to 5 years

Age of maturity: 18 to 20 months

Reproduction period: Autumn

Fertile individual: Presence of dark brown areas on the blade

Growth period: Spring, summer

Meristem: Basal area of the blade



Can easily be misidentified with *H. nigripes*, because they are morphologically similar. It is possible to differentiate the species by the absence of mucus on the stipe compared to *H. nigripes*.



Fanny Noisette

Anticosti Island, QC

Specific sustainable harvesting practices

Harvest a maximum of 33% of the biomass (one blade out of three)

Conservation of the biomass and associated benefits

Cut at a minimum of 20 cm above the stipe and blade junction

Leaving the meristem in place allows for faster blade regrowth

Favour harvesting during growth from May to July

Maintains biomass through rapid regrowth

Avoid harvesting during reproduction in September and October

More efficient reproduction leads to better population renewal

Fallow period of 2 years

Population recovery to the pre-harvest state

Harvest within a 15 m diameter area and leave 15 m between harvested areas

Allows for more effective recolonization due to proximity to an unharvested area

Winged kelp (*Alaria esculenta*)



Other names: Atlantic wakame, dabberlocks

Life expectancy: 5 to 10 years

Age of maturity: 8 to 14 months

Reproduction period: Summer

Fertile individual: Presence of dark brown areas on the sporophylls

Growth period: Spring, summer

Meristem: Basal area of the blade



Reproductive structures called sporophylls are located beneath its blade, unlike other kelp species that bear them directly on their blades.



Stéphanie Roy

Saint-Fabien-sur-Mer, QC

Specific sustainable harvesting practices

Harvest a maximum of 33% of the biomass (one blade out of three)

Conservation of the biomass and associated benefits

Cut at a minimum of 20 cm above the stipe and blade junction

Leaving the meristem in place allows for faster blade regrowth

Favour harvesting during growth in May and June

Maintains biomass through rapid regrowth

Avoid harvesting during reproduction in July and August

More efficient reproduction leads to better population renewal

Fallow period of 2 years

Population recovery to the pre-harvest state

Saccorhiza dermatodea



Other names: -

Life expectancy: Approximately one year

Age of maturity: Less than a year

Reproduction period: Autumn

Fertile individual: Presence of a dark area on the blade

Growth period: Spring, summer

Meristem: Basal area of the blade



The blade can sometimes split lengthwise and look like *L. digitata* and *H. nigripes*. It can be distinguished by its flattened stipe towards the blade as opposed to the other two species that have cylindrical stipes.



Eve Tremblay-Morel

Brown algae

Anticosti Island, QC

Specific sustainable harvesting practices

Harvest a maximum of 33% of the biomass (one blade out of three)

Conservation of the biomass and associated benefits

Cut at a minimum of 20 cm above the stipe and blade junction

Leaving the meristem in place allows for faster blade regrowth

Favour harvesting during growth from May to July

Maintains biomass through rapid regrowth

Avoid harvesting during reproduction in September and October

More efficient reproduction leads to better population renewal

Knotted wrack (*Ascophyllum nodosum*)



Other names: -

Life expectancy: 3 to 15 years (blade), 40 years (holdfast)

Age of maturity: 5 years

Reproduction period: Spring, summer

Fertile individual: Presence of yellow or green receptacles at the tip of the blade

Growth period: Summer

Meristem: Holdfast and apical area of the blade



Very slow renewal rate, so it is particularly important to use fallow periods and buffer zones (see below).



Brown algae

Anticosti Island, QC

Eve Tremblay-Morel

Specific sustainable harvesting practices

Harvest a maximum of 33% of the biomass (one blade out of three)

Conservation of biomass and associated benefits

Cut at a minimum of 30 cm above the holdfast

Leaving the meristem in place allows for faster regrowth of the blade

Leave at least one branch

Preserve part of the reproductive structures to allow for faster regrowth

Favour harvesting during growth period from July to September

Maintain biomass through rapid regrowth

Avoid harvesting during the reproduction period in May and June

More effective reproduction leads to better population renewal


Fallow period of 3 years

Population recovery to pre-harvest state

Harvest within a 15 m diameter area and leave 15 m between harvested areas

Allows for more effective recolonization due to proximity to an unharvested area

Fucus (*Fucus* spp.)



Other names: Bladderwrack

Life expectancy: 2 to 5 years

Age of maturity: 1 to 2 years

Reproduction period: Spring, summer

Fertile individual: Presence of yellow or green receptacles at the tip of the blade

Growth period: Summer

Meristem: Holdfast and apical area of the blade



The species and subspecies present in Quebec are : *Fucus vesiculosus*, *Fucus distichus*, *Fucus distichus* subsp. *edentatus*, *Fucus distichus* subsp. *evanescens* and *Fucus spiralis*.



Fanny Noisette

Brown algae

Anticosti Island, QC

Specific sustainable harvesting practices

Harvest a maximum of 33% of the biomass (one blade out of three)

Conservation of biomass and associated benefits

Cut at a minimum of 15 cm above the holdfast

Leaving the meristem in place allows for faster regrowth of the blade

Leave at least one branch

Preserve part of the reproductive structures to allow for faster regrowth

Favour harvesting during growth period from July to September

Maintain biomass through rapid regrowth

Avoid harvesting during the reproduction period in May and June

More effective reproduction leads to better population renewal

Fallow period of 3 years

Population recovery to pre-harvest state

Harvest within a 15 m diameter area and leave 15 m between harvested areas

Allows for more effective recolonization due to proximity to an unharvested area

Sea lace (*Chorda filum*)



Other names: -

Life expectancy: Approximately one year

Age of maturity: Less than a year

Reproduction period: Autumn

Fertile individual: Presence of dark spots all along the blade

Growth period: Spring, summer


Meristem: Apical area of the blade (adult) and basal area of the blade (juvenile)



The meristem is located in the apical zone of the blade, making it impossible to harvest adults without also harvesting the meristem.



Small hairs are located along the entire blade, with a particularly high hair density on the meristem.



Rénauld Belley & Anne-Sara Sean

Brown algae

***C. filum* among other seaweeds such as *A. esculenta* and *Porphyra* spp.**

Baie-Sainte-Catherine, QC

Specific sustainable harvesting practices

Harvest a maximum of 33% of the biomass (one blade out of three)

Conservation of biomass and associated benefits

Favour harvesting during growth from May to July

Maintains biomass through rapid regrowth

Avoid harvesting during reproduction in September and October

More efficient reproduction leads to better population renewal

Chordaria flagelliformis



Other names: -

Life expectancy: Approximately 1 year

Age of maturity: Less than a year

Reproduction period: n.a. Possibly similar to other annual species

Fertile individual: n.a.

Growth period: Spring, summer

Meristem: n.a.



Information on the life cycle is scarce. Research should be conducted to better understand the life cycle of this commercial species.



Stephanie Roy

**Brown
algae**

Rimouski, QC

Specific sustainable harvesting practices

Harvest a maximum of 33% of the biomass (one blade out of three)

Conservation of biomass and associated benefits

Cut at a minimum of 15 cm above the holdfast

Leaving the meristem in place allows for faster regrowth of the blade

Leave at least one branch

Preserve part of the reproductive structures to allow for faster regrowth

Favour harvesting during growth from May to July

Maintains biomass through rapid regrowth

Avoid harvesting during reproduction in September and October

More efficient reproduction leads to better population renewal

Irish moss (*Chondrus crispus*)



Other names: Carrageen moss

Life expectancy: 2 to 4 years

Age of maturity: n.a.

Reproduction period: Summer, autumn

Fertile individual: Presence bulging areas on the blade

Growth period: Spring, summer

Meristem: Apical and marginal area of the blade



Can be mistaken for *Mastocarpus stellatus*, which has a similar morphology, but its stipe curls along its entire length, unlike *C. crispus*.



Eve Tremblay-Morel

Red
algae

Anticosti Island, QC

Specific sustainable harvesting practices

Harvest a maximum of 33% of the biomass (one blade out of three)

Preserve biomass and its associated benefits

Cut above the holdfast, leaving a portion of the blade

Leaving the meristem intact allows for faster blade regrowth

Leave at least one branch

Preserve part of the reproductive structures to enable quicker regrowth

Favour harvesting during the growth period in May and June

Maintain biomass through rapid regrowth

Avoid harvesting during the reproduction period from July to October

More effective reproduction leads to better population renewal

Fallow period of 3 years

Population recovery to pre-harvest state

Harvest within a 3 m diameter area and leave 3 m between harvested areas

Allows for more effective recolonization due to proximity to an unharvested area

Dulse

(*Palmaria palmata*)



Other names: -

Life expectancy: Several years

Age of maturity: 9 to 12 months (male), a few days (female)

Reproduction period: Autumn, winter

Fertile individual: Presence of a dark, almost black, area in the center of the blade

Growth period: Spring, summer

Meristem: Apical and marginal area of the blade



Due to the differences in their maturation rates, females will be fertilized by males from earlier generations.



Red
algae

Stéphanie Roy

Rimouski, QC

Specific sustainable harvesting practices

Harvest a maximum of 33% of the biomass (one blade out of three)

Conservation of biomass and associated benefits

Cut above the holdfast, leaving a portion of the blade

Leaving the meristem in place allows for faster regrowth of the blade

Leave at least one branch

Preserve part of the reproductive structures to enable quicker regrowth

Favour harvesting during growth from May to July

Maintains biomass through rapid regrowth

Avoid harvesting during reproduction from September to February

More efficient reproduction leads to better population renewal

Sallow time of 2 or 3 years

Population recovery to the pre-harvest state

Devaleraea ramentacea



Other names: -

Life expectancy: Several years

Age of maturity: n.a.

Reproduction period: n.a. (Possibly similar to species from the same family, like *P. palmata*)


Fertile individual: Presence of dark areas on the blade

Growth period: n.a. (Possibly similar to species from the same family, like *P. palmata*)

Meristem: n.a.



There is very little information available on this species. However, it belongs to the same order as *P. palmata*, so its biology and ecology are likely similar.



Red algae

Anticosti Island, QC

Christian Nozais

Specific sustainable harvesting practices

Harvest a maximum of 33% of the biomass (one blade out of three)

Conservation of biomass and associated benefits

Cut above the holdfast, leaving a portion of the blade

Leaving the meristem in place allows for faster regrowth of the blade

Leave at least one branch

Preserve part of the reproductive structures to enable quicker regrowth

Favour harvesting during growth from May to July

Maintains biomass through rapid regrowth

Avoid harvesting during reproduction from September to February

More efficient reproduction leads to better population renewal

Fallow period of 2 or 3 years

Population recovery to the pre-harvest state

Nori

(*Porphyra* spp.)



Other names: -

Other genera with a similar life cycle and morphology: *Wildemania* sp., *Pyropia* sp.

Life expectancy: Approximately one year

Age of maturity: 45 days

Reproduction period: Spring, autumn

Fertile individual: Presence of a dark or yellow border on the outer edge of the blade

Growth period: Spring, summer

Meristem: Basal area of the blade



When reproduction is over, a thin white border is visible on the outer edge of the blade (like on the photo).



Red
algae

Rénaud Belley & Anne-Sara Sean

Baie-Sainte-Catherine, QC

Specific sustainable harvesting practices

Harvest a maximum of 33% of the biomass (one blade out of three)

Conservation of biomass and associated benefits

Cut above the holdfast, leaving a portion of the blade

Leaving the meristem in place allows for faster regrowth of the blade

Favour harvesting during growth in July

Maintains biomass through rapid regrowth

Avoid harvesting during reproduction in May, June, August and September

More efficient reproduction leads to better population renewal

Sea lettuce (*Ulva* spp.)



Other names: *Enteromorpha* sp., *Ulva* sp.

Life expectancy: Approximately one year

Age of maturity: A few weeks

Reproduction period: Summer

Fertile individual: Presence of a yellow-brown border on the outer edge of the blade

Growth period: Spring, summer

Meristem: Mitotic points on the blade



Opportunist species, with rapid growth, sometimes considered a nuisance species.



Green algae

Baie-Sainte-Catherine, QC

Specific sustainable harvesting practices

Harvest a maximum of 33% of the biomass (one blade out of three)

Conservation of biomass and associated benefits

Cut above the holdfast, leaving a portion of the blade

Leaving the meristem in place allows for faster regrowth of the blade

Favour harvesting during growth in May and June

Maintains biomass through rapid regrowth

Avoid harvesting during reproduction in July and August

More efficient reproduction leads to better population renewal

Summary of specific sustainable harvesting practices for 15 commercially valuable seaweeds in the Quebec region

 Brown seaweeds
  Red seaweeds
  Green seaweeds
 n.a. : not applicable

Latin name	Harvest site				Individual harvested			
	Vernacular name	Maximum harvest	Preferred harvest period	Period to avoid	Suggested fallow	Buffer zone	Cutting height	Leave a branch
 <i>Agarum clathratum</i>	Sea colander	33%	May to July	September to February	2 years	To consider	10 cm above the stipe and blade junction	n.a.
 <i>Hedophyllum nigripes</i>	-	33%	May to July*	September and October*	2 years	Yes (15 m)	20 cm above the stipe and blade junction	n.a.
 <i>Laminaria digitata</i>	Oarweed	33%	May to July	September and October	2 years	Yes (15 m)	20 cm above the stipe and blade junction	n.a.
 <i>Saccharina latissima</i>	Sugar kelp	33%	May to July	September and October	2 years	Yes (15 m)	20 cm above the stipe and blade junction	n.a.
 <i>Alaria esculenta</i>	Winged kelp	33%	May and June	July and August	2 years	To consider	20 cm above the stipe and blade junction	n.a.
 <i>Saccorhiza dermatodea</i>	-	33%	May and July	September and October	None	To consider	20 cm above the stipe and blade junction	n.a.
 <i>Ascophyllum nodosum</i>	Knotted wrack	33%	July to September	May and June	3 years	Yes (15 m)	30 cm above the holdfast	Yes
 <i>Fucus</i> spp.	Fucus	33%	July to September	May and June	3 years	Yes (15 m)	15 cm above the holdfast	Yes
 <i>Chorda filum</i>	Sea lace	33%	May to July	September and October	None	To consider	None	n.a.
 <i>Chordaria flagelliformis</i>	-	33%	May to July	September and October	None	To consider	15 cm above the holdfast	Yes

 Brown seaweeds

 Red seaweeds

 Green seaweeds

n.a. : not applicable

Latin name	Harvest site				Individual harvested			
	Vernacular name	Maximum harvest	Preferred harvest period	Period to avoid	Suggested fallow	Buffer zone	Cutting height	Leave a branch
 <i>Chondrus crispus</i>	Irish moss	33%	May and June	July to October	3 years	Yes (3 m)	Above the holdfast with a portion of the blade	Yes
 <i>Palmaria palmata</i>	Dulse	33%	May to July	September to February	2-3 years	To consider	Above the holdfast with a portion of the blade	Yes
 <i>Devaleraea ramentacea</i>	-	33%	May to July*	September to February*	2-3 years	To consider	Above the holdfast with a portion of the blade	Yes
 <i>Porphyra</i> spp. Nori	Nori	33%	July	May, June, August and September	None	To consider	Above the holdfast with a portion of the blade	n.a.
 <i>Ulva</i> spp.	Sea lettuce	33%	May and June	July and August	None	To consider	Above the holdfast with a portion of the blade	n.a.

*Growth and reproduction periods are not available, but could be similar to the ones from similar species.

References

- Adey, W.H., and Hayek, L.A.C. 2011. Elucidating Marine Biogeography with Macrophytes: Quantitative Analysis of the North Atlantic Supports the Thermogeographic Model and Demonstrates a Distinct Subarctic Region in the Northwestern Atlantic. *Northeastern Naturalist* 18(mo8): 1–128. doi:10.1656/045.018.m801.
- Bartsch, I., Wiencke, C., Bischof, K., Buchholz, C.M., Buck, B.H., Eggert, A., Feuerpfeil, P., Hanelt, D., Jacobsen, S., Karez, R., Karsten, U., Molis, M., Roleda, M.Y., Schubert, H., Schumann, R., Valentin, K., Weinberger, F., and Wiese, J. 2008. The genus *Laminaria* sensu lato: recent insights and developments. *European J. Phycol.* 43(1): 1–86. doi:10.1080/09670260701711376.
- Baud, A. 2023. Base biologique et génétique de la domestication de l'algue rouge *Palmaria palmata*. Ph. D, Université de la Sorbonne.
- Belsher, T., Bove, J., and de Cayeux, M.B. 1986. Étude bibliographiques de quelques espèces planctoniques et benthiques littorales de la Manche. IFREMER 4.
- Birkett, D.A., Maggs, C.A., Dring, M.J., and Boaden, P.J.S. 1998. Infralittoral reef biotopes with kelp species - An overview of dynamic and sensitivity characteristics for conservation management of marine SACs. p. 174p.
- Buggeln, R. 1977. Physiological investigations on *Alaria esculenta* (Laminariales, Phaeophyceae). II. Role of translocation in blade growth. *J. Phycol.* 13: 212–218.
- Chabot, R., and Rossignol, A. 2003. Algues et faune du littoral du Saint-Laurent maritime: Guide d'identification. Institut des sciences de la mer de Rimouski, Rimouski; Pêches et Océans Canada (Institut Maurice-Lamontagne), Mont-Joli, 113 p..
- Chapman, A.R.O. 1995. Functional ecology of furoid algae: twenty-three years of progress. *Phycologia* 34(1): 1–32. doi:10.2216/i0031-8884-34-1-1.1.
- Collén, J., Cornish, M.L., Craigie, J., Ficko-Blean, E., Hervé, C., Krueger-Hadfield, S.A., Leblanc, C., Michel, G., Potin, P., Tonon, T., and Boyen, C. 2014. *Chondrus crispus* - A Present and Historical Model Organism for Red Seaweeds. In *Advances in Botanical Research*. Elsevier. pp. 53–89. doi:10.1016/B978-0-12-408062-1.00003-2.
- Côte-Laurin, M.-C., Berger, K., and Tamigneaux, É. 2016. Manuel pour la récolte commerciale des macroalgues au Québec. Mérimov: 89 p.
- Dankworth, M., Heinrich, S., Fredriksen, S., and Bartsch, I. 2020. DNA barcoding and mucilage ducts in the stipe reveal the presence of *Hedophyllum nigripes* (Laminariales, Phaeophyceae) in Kongsfjorden (Spitsbergen). *J. Phycol.* 56(5): 1245–1254. doi:10.1111/jpy.13012.
- Diehl, N., Li, H., Scheschonk, L., Burgunter-Delamare, B., Niedzwiedz, S., Forbord, S., Sæther, M., Bischof, K., and Monteiro, C. 2024. The sugar kelp *Saccharina latissima* I: recent advances in a changing climate. *Annals of Botany* 133(1): 183–212. doi:10.1093/aob/mcad173.
- Dudgeon, S., and Petraitis, P.S. 2005. First year demography of the foundation species, *Ascophyllum nodosum*, and its community implications. *Oikos* 109(2): 405–415. doi:10.1111/j.0030-1299.2005.13782.x.
- Franke, K., Liesner, D., Heesch, S., and Bartsch, I. 2021. Looks can be deceiving: contrasting temperature characteristics of two morphologically similar kelp species co-occurring in the Arctic. *Botanica Marina* 64(3): 163–175. doi:10.1515/bot-2021-0014.
- Fredersdorf, J., Müller, R., Becker, S., Wiencke, C., and Bischof, K. 2009. Interactive effects of radiation, temperature and salinity on different life history stages of the Arctic kelp *Alaria esculenta* (Phaeophyceae). *Oecologia* 160(3): 483–492. doi:10.1007/s00442-009-1326-9.
- Gagnon, P., Himmelman, J.H., and Johnson, L.E. 2004. Temporal variation in community interfaces: kelp-bed boundary dynamics adjacent to persistent urchin barrens. *Mar. Biol.* 144(6): 1191–1203. doi:10.1007/s00227-003-1270-x.
- Gagnon, P., Johnson, L.E., and Himmelman, J.H. 2005. Kelp patch dynamics in the face of intense herbivory: Stability of *Agarum clathratum* (Phaeophyta) stands and associated flora on urchin barrens. *J. Phycol.* 41(3): 498–505. doi:10.1111/j.1529-8817.2005.00078.x.
- Gantt, E., Berg, G.M., Bhattacharya, D., Blouin, N.A., Brodie, J.A., Chan, C.X., Collén, J., Cunningham, F.X., Gross, J., Grossman, A.R., Karpowicz, S., Kitade, Y., Klein, A.S., Levine, I.A., Lin, S., Lu, S., Lynch, M., Minocha, S.C., Müller, K., Neefus, C.D., De Oliveira, M.C., Rymarquis, L., Smith, A., Stiller, J.W., Wu, W.-K., Yarish, C., Zhuang, Y., and Brawley, S.H. 2010. *Porphyra*: Complex Life Histories in a Harsh Environment: *P. umbilicalis*, an Intertidal Red Alga for Genomic Analysis. In *Red Algae in the Genomic Age*. Edited by J. Seckbach and D.J. Chapman. Springer Netherlands, Dordrecht. pp. 129–148. doi:10.1007/978-90-481-3795-4_7.
- Gendron-Lemieux, I., Gilmore-Solomon, L., Tamigneaux É. et Tremblay-Gratton, A. 2021. Essais de culture de *Porphyra* sp dans une perspective d'épuration des effluents de viviers de homard. Merinov, Rapport de R-D no 21-01. 73 p.
- Graham, L.E., Graham, J.M., Cook, M.E., and Wilcox, L.W. 2016. *Algae*. In Third edition. LJLM Press, Madison.
- Grant, W.S., Lydon, A., and Bringloe, T.T. 2020. Phylogeography of split kelp *Hedophyllum nigripes*: northern ice-age refugia and trans-Arctic dispersal. *Polar Biol* 43(11): 1829–1841. doi:10.1007/s00300-020-02748-6.
- Grote, B. 2019. Recent developments in aquaculture of *Palmaria palmata* (Linnaeus) (Weber & Mohr 1805): cultivation and uses. *Reviews in Aquaculture* 11(1): 25–41. doi:10.1111/raq.12224.

References

- Guiry, M.D. 1982. *Devaleraea*, A New Genus of the Palmariaceae (Rhodophyta) in the North Atlantic and North Pacific. J. Mar. Biol. Ass. 62(1): 1–13. doi:10.1017/S0025315400020063.
- Hatchett, W.J., Coyer, J.A., Sjøtun, K., Jueterbock, A., and Hoarau, G. 2022. A review of reproduction in the seaweed genus *Fucus* (Ochrophyta, Fucales): Background for renewed consideration as a model organism. Front. Mar. Sci. 9: 1051838. doi:10.3389/fmars.2022.1051838.
- Hayden, H.S., Blomster, J., Maggs, C.A., Silva, P.C., Stanhope, M.J., and Waaland, J.R. 2003. Linnaeus was right all along: *Ulva* and *Enteromorpha* are not distinct genera. European J. Phycol. 38(3): 277–294. doi:10.1080/1364253031000136321.
- Hill. 2008. Oarweed (*Laminaria digitata*): Marine Evidence-based Sensitivity Assessment (MarESA) Review. MarLIN - Marine Life Information Network. Available from http://www.marlin.ac.uk/assets/pdf/species/marlin_species_1386_2019-03-21.pdf [accessed 15 August 2024].
- Hurd, C.L., Harrison, P.J., Bischof, K., and Lobban, Christopher.S. 2014. Seaweed Ecology and Physiology. In Second edition. Cambridge University Press.
- Keats, D., South, G., and Steele, D. 1985. Algal biomass and diversity in the upper subtidal at a pack-ice disturbed site in eastern Newfoundland. Mar. Ecol. Prog. Ser. 25: 151–158. doi:10.3354/meps025151.
- Keats, D.W. 1991. Refugial *Laminaria* Abundance and Reduction in Urchin Grazing in Communities in the North-West Atlantic. J. Mar. Biol. Ass. 71(4): 867–876. doi:10.1017/S0025315400053510.
- Knoop, J., Griffin, J.N., and Barrento, S. 2020. Cultivation of early life history stages of *Porphyra dioica* from the British Isles. J Appl Phycol 32(1): 459–471. doi:10.1007/s10811-019-01930-6.
- Kraan, S. 2020. Concise review of the genus *Alaria* Greville, 1830. J Appl Phycol 32(6): 3543–3560. doi:10.1007/s10811-020-02222-0.
- Longtin, C., and Saunders, G. 2016. The relative contribution of *Saccharina nigripes* (Phaeophyceae) to the Bay of Fundy Laminariaceae: spatial and temporal variability. Mar. Ecol. Prog. Ser. 543: 153–162. doi:10.3354/meps11566.
- Longtin, C.M., and Saunders, G.W. 2015. On the utility of mucilage ducts as a taxonomic character in *Laminaria* and *Saccharina* (Phaeophyceae) – the conundrum of *S. groenlandica*. Phycologia 54(5): 440–450. doi:10.2216/15-19.1.
- Lukeman, R.J., Beveridge, L.F., Flynn, A.D., and Garbary, D.J. 2012. A mathematical model of the commercial harvest of *Palmaria palmata* (Palmariales, Rhodophyta) on Digby Neck, Nova Scotia, Canada. ALGAE 27(1): 43–54. doi:10.4490/algae.2012.27.1.043.
- Mantri, V.A., Kazi, M.A., Balar, N.B., Gupta, V., and Gajaria, T. 2020. Concise review of green algal genus *Ulva* Linnaeus. J Appl Phycol 32(5): 2725–2741. doi:10.1007/s10811-020-02148-7.
- Mathieson, A.C., and Dawes, C.J. 2017. Seaweeds of the Northwest Atlantic. In University of Massachusetts Press.
- Mathieson, A.C., and Dawes, C.J. 2020. Biology and ecology of Northwest Atlantic seaweeds. In Handbook of Algal Science, Technology and Medicine. Elsevier. pp. 71–90. doi:10.1016/B978-0-12-818305-2.00005-X.
- MPO. 1998. Ascophylle noueuse (*Ascophyllum nodosum*). MPO – Sciences, Rapport sur l'état des stocks C3-57.
- MPO. 1999a. Coastal Zone Species Profile Series - Winged Kelp. (13).
- MPO. 1999b. Coastal Zone Species Profiles Series - Finger Kelp. (14)
- MPO. 1999c. Coastal Zone Species Profile Series - Knotted wrack. (16).
- MPO. 2009. Évaluation écologique de la mousse d'Irlande (*Chondrus crispus*) dans la zone de protection marine Bassin Head. Secr. can. de consult. sci. du MPO, Avis sci. 2009/059.
- Munda, I.M. 1992. Gradient in seaweed vegetation patterns along the North Icelandic coast, related to hydrographic conditions. Hydrobiologia 242(3): 133–147. doi:10.1007/BF00019962.
- Neiva, J., Serrão, E.A., Paulino, C., Gouveia, L., Want, A., Tamigneaux, É., Ballenghien, M., Mauger, S., Fouqueau, L., Engel-Gautier, C., Destombe, C., and Valero, M. 2020. Genetic structure of amphiatlantic *Laminaria digitata* (Laminariales, Phaeophyceae) reveals a unique range-edge gene pool and suggests post-glacial colonization of the NW Atlantic. European J. Phycol. 55(4): 517–528. doi:10.1080/09670262.2020.1750058.
- Norton, T.A. 1972. The development of *Saccorhiza dermatodea* (Phaeophyceae, Laminariales) in culture. Phycologia. 11 : 81–86.
- Novacek, I., Lubbers, G.W., and Breeman, A.M. 1990. Thermal ecotypes of amphiatlantic algae. I. Algae of Arctic to cold-temperate distribution (*Chaetomorpha melagonium*, *Devaleraea ramentacea* and *Phycodrys rubens*). Helgolander Meeresunters 44(3–4): 459–474. doi:10.1007/BF02365480.
- Parke, M. 1948. Studies on the British Laminariaceae. I. Growth in *Laminaria saccharina* (L.) Lamour. J. Mar. Biol. Ass. UK, 27, 651–709.
- Pizzolla, P.F. 2008. Sea lettuce (*Ulva lactuca*). Available from <https://www.marlin.ac.uk/species/detail/1467>.
- Schiel, D.R., and Foster, M.S. 2006. The Population Biology of Large Brown Seaweeds: Ecological Consequences of Multiphase Life Histories in Dynamic Coastal Environments. Annu. Rev. Ecol. Evol. Syst. 37(1): 343–372. doi:10.1146/annurev.ecolsys.37.091305.110251.
- South, G.R., and Burrows, E.M. 1967. Studies on marine algae of the British Isles. 5. *Chorda filum* (L.) Stackh. British Phycological Bulletin 3(2): 379–402. doi:10.1080/00071616700650261.

References

- Stévant, P., Schmedes, P.S., Le Gall, L., Wegeberg, S., Dumay, J., and Rebours, C. 2023. Concise review of the red macroalga dulse, *Palmaria palmata* (L.) Weber & Mohr. *J Appl Phycol* 35(2): 523–550. doi:10.1007/s10811-022-02899-5.
- Sutherland, J.E., Lindstrom, S.C., Nelson, W.A., Brodie, J., Lynch, M.D.J., Hwang, M.S., Choi, H., Miyata, M., Kikuchi, N., Oliveira, M.C., Farr, T., Neefus, C., Mols-Mortensen, A., Milstein, D., and Müller, K.M. 2011. A new look at an ancient order : generic revision of the Bangiales (Rhodophyta). *J. Phycol.* 47(5): 1131–1151. doi:10.1111/j.1529-8817.2011.01052.x.
- Tamigneaux, É., Rondeau, M.-H., Gauthier-Barrette, C., and Fortin, M.-G. 2022. Adaptation du processus de culture de l'alarie succulente (*Alaria esculenta*), une macroalgue arctique d'intérêt pour le secteur alimentaire et cosmétique. Merinov. Rapport de projet. no 000872, 26 p.
- Vandermeulen, H. 2013. Information to Support Assessment of Stock Status of Commercially Harvested Species of Marine Plants in Nova Scotia: Irish Moss, Rockweed and Kelp. DFO Can. Sci. Advis. Sec. Res. Doc. 2013/042. vi + 50 p.
- White, N. 2006. Sea lace or Dead man's rope (*Chorda filum*): Marine Evidence-based Sensitivity Assessment (MarESA) Review. MarLIN - Marine Life Information Network. Available from http://www.marlin.ac.uk/assets/pdf/species/marlin_species_1366_2019-03-21.pdf [accessed 15 August 2024].
- Wilding, C., Tillin, H., Stewart, E.J., Burrows, M., and Smale, D. 2021. Hand harvesting of seaweed: Evidence review to support sustainable management. NRW Evidence Report Series, NRW, Bangor.