


Health Reports

All-cause and circulatory disease-related hospitalization, by generation status: Evidence from linked data

by Edward Ng, Claudia Sanmartin, Jack V. Tu and Douglas G. Manuel

Release date: October 21, 2015



Statistics
Canada

Statistique
Canada

Canada

How to obtain more information

For information about this product or the wide range of services and data available from Statistics Canada, visit our website, www.statcan.gc.ca.

You can also contact us by

email at infostats@canada.ca

telephone, from Monday to Friday, 8:30 a.m. to 4:30 p.m., at the following toll-free numbers:

- Statistical Information Service 1-800-263-1136
- National telecommunications device for the hearing impaired 1-800-363-7629
- Fax line 1-877-287-4369

Depository Services Program

- Inquiries line 1-800-635-7943
- Fax line 1-800-565-7757

Standards of service to the public

Statistics Canada is committed to serving its clients in a prompt, reliable and courteous manner. To this end, Statistics Canada has developed standards of service that its employees observe. To obtain a copy of these service standards, please contact Statistics Canada toll-free at 1-800-263-1136. The service standards are also published on www.statcan.gc.ca under “Contact us” > “Standards of service to the public.”

Note of appreciation

Canada owes the success of its statistical system to a long-standing partnership between Statistics Canada, the citizens of Canada, its businesses, governments and other institutions. Accurate and timely statistical information could not be produced without their continued co-operation and goodwill.

Standard table symbols

The following symbols are used in Statistics Canada publications:

- . not available for any reference period
- .. not available for a specific reference period
- ... not applicable
- 0 true zero or a value rounded to zero
- 0^s value rounded to 0 (zero) where there is a meaningful distinction between true zero and the value that was rounded
- ^P preliminary
- ^r revised
- X suppressed to meet the confidentiality requirements of the *Statistics Act*
- ^E use with caution
- F too unreliable to be published
- * significantly different from reference category ($p < 0.05$)

Published by authority of the Minister responsible for Statistics Canada

© Minister of Industry, 2015

All rights reserved. Use of this publication is governed by the Statistics Canada [Open Licence Agreement](#).

An HTML version is also available.

Cette publication est aussi disponible en français.

All-cause and circulatory disease-related hospitalization, by generation status: Evidence from linked data

by Edward Ng, Claudia Sanmartin, Jack V. Tu and Douglas G. Manuel

Abstract

Background: Immigrants tend to have better health than the Canadian-born. However, the “healthy immigrant” effect diminishes over time and varies by source country. This study examines whether lower hospitalization rates persist from the first (G1) to the second generation (G2) of immigrants, compared with other Canadians (G3+). All-cause and circulatory disease-related hospitalization rates were examined by generation, with special attention to people of Chinese and South Asian descent.

Data and methods: Data from the 2006 Census-hospitalization linkage database (which excludes Quebec) were analysed using logistic regression. Age-standardized all-cause (excluding pregnancy) and circulatory disease-related hospitalization rates were derived for the urban population aged 30 or older for the 2006/2007 to 2008/2009 fiscal years.

Results: Over the generations, immigrants’ all-cause and circulatory disease-related hospitalization rates converged with those of the Canadian population overall. Compared with G3+, the age-adjusted odds of all-cause hospitalization among men were 0.49 (CI = 0.48-0.51) for recent G1 immigrants, 0.78 (CI = 0.77-0.79) for long-term G1 immigrants, and 0.95 (CI = 0.94-0.97) for G2. Adjustments for socioeconomic status reduced the difference, especially between G2 and G3+. For South Asians, rates converged for circulatory disease, notably among men. Hospitalization rates for people of Chinese descent rose across generations, but remained significantly below rates for G3+.

Interpretation: The lower circulatory disease-related hospitalization risk experienced by G1 is maintained in G2 among people of Chinese descent, but not among South Asians.

Keywords: Cardiovascular diseases, census, Chinese, data linkage, hospital records, immigrant health, medical records, socioeconomic status, South Asian

Since the 1970s, the origins of immigrants to Canada have shifted toward non-European source countries, such as India and China.¹ As a result, 8% of the total population self-identify as South Asian or Chinese, a percentage that is projected to reach about 15% by 2031.² Canada’s changing ethnic make-up and its long-run impact on subsequent generations can have implications for health care.

While immigrants tend to have better health than the Canadian-born population, variations by disease and by source country are considerable.^{3,4} For example, regardless of where they live, South Asians have one of the highest rates of heart/circulatory disease in the world.⁵⁻⁷ In Canada, circulatory disease mortality risk among South Asians tends to be similar to that of the Canadian-born population, but higher than that of other immigrant/ethnic groups.^{3,8,9} Conversely, circulatory disease risk among the Chinese population is relatively low.^{10,11} A recent review of immigrant health research found that Canadians of South Asian descent tend to experience increased risks of heart disease and hypertension as duration of residence in Canada lengthens, and noted a need to study the effects from a generational perspective.¹²

Evidence of the extent to which health advantages (or disadvantages) of first-generation immigrants are transferred to future generations is mixed.¹³⁻¹⁵ In a 1976 study of immigrant health, Marmot and Syme¹⁶ found a higher prevalence of heart disease among male Japanese immigrants in California and

Hawaii than among men in Japan. However, first-generation Japanese immigrants in Hawaii had a lower prevalence of heart disease than the second generation, whose health profiles were closer to those of non-immigrant Hawaiians. In Canada, generational analyses of health tend to focus on youth and school children, specifically risk behaviours and weight gain, with little attention to cardiac health.¹⁷⁻¹⁹ A study of immigrants in Montreal reported no health advantage in obesity or self-rated health among second-generation adults, especially those in disadvantaged neighborhoods.²⁰

Information about the use of health care services by generation status is limited. A key barrier is the lack of immigration and generation status variables in administrative health data.²¹ Area-based studies have used the percentage of immigrants in a neighbourhood as a proxy for immigrant status,²² but such ecological approaches are limited. Alternatively, linking administrative health data to sources that contain information on immigrant and generation status opens the possibility for detailed examinations of immigrant health.³

This study used data from the 2006 Census of Population (long-form respondents) linked to administrative records to determine if hospitalization patterns among first-generation immigrants persist in the second generation, and if patterns differ between South Asians and Chinese subgroups, when socioeconomic covariates are taken into account.

Data and methods

Data sources

The 2006 Census (long-form) was linked to the Canadian Institute for Health Information's Discharge Abstract Database (DAD). The long-form census data represent about 20% of the non-institutional population and provide information about the Canadian population by generation status, place of birth and period of arrival of the first generation, as well as socioeconomic characteristics.

The DAD contains annual information about inpatient hospitalizations, excluding the province of Quebec, which does not routinely submit data to the DAD. As a result, immigrants living in Quebec were excluded from this study, as were hospitalizations of residents of other provinces that occurred in Quebec.

Approximately 4.65 million census long-form respondents (excluding Quebec) were linked to the DAD for the three fiscal years from 2006/2007 to 2008/2009. A hierarchical deterministic linkage using date of birth, sex, and residential postal code was conducted. Postal code information from the Historical Tax File was used to account for changes in residence. The weighted coverage rates in the linked data were relatively high (78% to 80%). A validation study²³ concluded that the linked file is suitable for health-related research and is broadly representative of immigrants in Canada.

Additional information about the linkage process and results are reported elsewhere.²³ The linkage was approved by Statistics Canada's Policy Committee²⁴ and is governed by the Directive on Record Linkage.²⁵

Study sample

The study cohort consisted of people aged 30 or older who were living in urban communities—census metropolitan areas (CMAs) or census agglomerations (CAs)²⁶ outside the territories—at the time of the 2006 Census (May 16, 2006). The majority of immigrants, especially recent arrivals, live in these urban communities.²⁷ Non-permanent residents (people in Canada on employment or student authorizations) and refugee

claimants were excluded. The final study cohort comprised 2,212,755 respondents, among whom 548,460 hospitalizations were recorded over the three years.

Measures

Generation status was defined based on immigrant status and parental place of birth. First-generation (G1) respondents are those who were, or had been, landed immigrants in Canada. A landed immigrant is a person who is not a Canadian citizen by birth, but who has been granted the right to live in Canada permanently.¹ Immigrants were categorized by their period of arrival: long-term (before 1996) or recent (from 1996 to 2006). Second-generation (G2) respondents are non-immigrants with at least one parent born outside Canada. Third-plus-generation (G3+) respondents are non-immigrants with both parents born in Canada. Although the study focuses on immigrants from China (including Hong Kong) and South Asia (Bangladesh, Bhutan, India, Maldives, Nepal, Pakistan, and Sri Lanka), these origins are not singled out in the G3+ comparison population.

Additional covariates from the census are age group (30 to 49, 50 to 64, and 65 or older), sex, secondary school graduation (yes/no), and before-tax low-income status (yes/no/not applicable). The low-income measure is based on the low-income cut-off threshold, determined by the average spending on food, clothing and shelter of all Canadians. A person is identified as in a low-income situation if the total income of his or her economic family (or the person if not in an economic family) was below the low-income threshold, which varies by economic family size and the size of the area of residence.²⁶

The primary outcome is at least one inpatient acute-care hospitalization between April 1, 2006 and March 31, 2009. Two types of hospitalization are considered in this analysis: all-cause, excluding those related to pregnancy (ICD10 codes O00 to O99,) and circulatory disease-related (ICD10 codes I00 to I93), which accounted for 17% (91,821) of discharges in the linked database.

Statistical methods

Descriptive statistics were produced to profile the study population by generation status and by the selected origins: South Asia and China. Crude and age-standardized hospitalization rates (ASHR) were derived for all-cause and circulatory disease-related hospitalizations using the overall cohort population as the reference population. Statistical testing for differences in age-standardized rates was conducted using logarithmic transformation to adjust for the skewness in the distribution of the standardized rates for inferential purposes in analysis of rare events.²⁸ Multivariate logistic regressions were used to estimate the odds of being hospitalized at least once, by generation status, using G3+ as the reference population. Separate models were estimated by sex and by origin. Two models were estimated: 1) age-adjusted, and 2) fully adjusted, controlling for age, low-income status, and secondary school graduation status. Analysis was conducted using SAS version 9.2.²⁹

Results

First-generation immigrants (G1) made up 31% of the urban population aged 30 or older; the second generation (G2) made up 21% (Table 1). The third-plus generation (G3+) accounted for the remaining 48%.

Among G1 respondents, 23% were recent immigrants (1996 to 2006), and 25% were from South Asia or China.

Approximately 3% of G2 respondents reported that their parent(s) were born in South Asia or China. G2 respondents of South Asian and Chinese descent were relatively young—85% and 79%, respectively, were aged 30 to 49, compared with about 50% for G2 overall.

Distributions by education level were similar across generations, with about 80% having graduated from secondary school. However, more than 90% of G2 respondents of South Asian and Chinese descent were secondary school graduates.

A relatively large percentage of G1 respondents lived in low-income situa-

**All-cause and circulatory disease-related hospitalization, by generation status:
Evidence from linked data • Research Article**

tions: 18% versus 10% of G2 and G3+. The figure was particularly high—27%—for Chinese immigrants.

A rising gradient in age-standardized hospitalization rates (ASHRs) was observed across the generations (Table 2). All-cause ASHRs for G1, G2 and G3+ were 609, 792 and 839 per 10,000, respectively; the corres-

ponding ASHRs for circulatory disease were 119, 142 and 152 per 10,000. The ASHRs for G1 and G2 were significantly lower than those for G3+. Patterns were similar among those of South Asian and Chinese descent. The lowest ASHRs were among G1 from China.

Logistic regression revealed rising odds of all-cause hospitalization by

generation for both men and women (Table 3). Compared with G3+, the age-adjusted odds of at least one hospitalization among men were 0.50 for recent G1, 0.79 for long-term G1, and 0.96 for G2. The trend was similar for women. Full adjustment for education and income reduced the differences, but they remained significant. These patterns prevailed among those of South Asian and Chinese descent.

The results for circulatory disease-related hospitalizations were similar, but some differences emerged (Table 3). For men, differences between G2 and G3+ were not significant when adjusting for covariates. For women, the odds of hospitalization for G2 converged with G3+ even before adjustment.

Results varied by origin, with higher odds of circulatory disease-related hospitalization among people of South Asian descent, and lower odds among those of Chinese descent. Among South Asians, the odds of at least one circulatory disease-related hospitalization were not significantly lower for G2 men and women, or for long-term G1 men. Among the Chinese, even after adjustment, the odds of at least one circulatory disease-related hospitalization were significantly lower for G1 and G2, compared with G3+.

**Table 1
Selected characteristics of long-form 2006 Census respondents aged 30 or older in urban areas, by generation status and South Asian or Chinese descent, Canada excluding Quebec and territories, 2006**

Characteristics	Non-immigrants					Immigrants		
	Total	Third-plus generation (G3+)	Second generation (G2)			Total	First generation (G1)	
		Total	Total	South Asia	China [†]		Total	South Asia
Number	2,212,755	1,058,300	468,755	4,755	8,460	745,700	88,320	91,290
% distribution	100	47.8	21.2	0.2	0.4	33.7	4.0	4.1
			Column %					
Age group (years)								
30 to 49	49	53	49	85	79	45	59	52
50 to 64	30	32	23	9	12	31	27	28
65 or older	21	15	28	7	9	24	14	20
Male	48	48	48	49	51	47	51	46
Years since immigration								
Less than 10	23	44	37
10 or more	77	56	63
Secondary school graduation	77	77	79	93	93	77	79	78
Low-income situation	13	10	10	8	8	18	21	27

[†] includes Hong Kong
... not applicable

Source: 2006 Census of Canada, long-form.

**Table 2
All-cause and circulatory disease-related crude and age-standardized hospitalization rates per 10,000 population, by generation status and South Asian or Chinese descent, long-form 2006 Census respondents aged 30 or older in urban areas, Canada excluding Quebec and territories, 2006/2007 to 2009/2010**

Type of hospitalization	Total			South Asia		China [‡]	
	Non-immigrants		Immigrants	Non-immigrants	Immigrants	Non-immigrants	Immigrants
	Third-plus generation (G3+) [†]	Second generation (G2)	First generation (G1)	Second generation (G2)	First generation (G1)	Second generation (G2)	First generation (G1)
All-cause excluding pregnancy							
Crude rate	757	897	664	362	466	372	332
Age-standardized rate	839	792*	609*	597*	544*	578*	356*
Circulatory disease-related							
Crude rate	130	169	133	52	105	43	45
Age-standardized rate	152	142*	119*	125	129*	79*	46*

[†] reference group

[‡] includes Hong Kong

* significantly different from reference group (p < 0.05)

Notes: Total cohort population was used for age-standardization. Statistical testing was conducted only for age-standardized rates.

Source: Linked 2006 Census-2006/2007 to 2009/2010 Discharge Abstract Database.

Discussion

Convergence in hospitalization rates over immigrant generations was apparent for both men and women. Recent immigrants were the least likely to have had at least one hospitalization during the 2006/2007-to-2008/2009 period, followed by long-term first-generation, and then, second-generation Canadians.

All three groups were less likely to have been hospitalized, compared with the third-plus generation. While a trend toward convergence emerged for circulatory disease-related hospitalizations, second-generation Canadians overall were no different from the third-plus generation, especially for women. This was more pronounced among South Asians than among those of Chinese descent.

Convergence of hospitalization rates in successive generations is consistent with the theory that immigrants tend to be healthier upon arrival, but that their health deteriorates over time.^{30,31} It is argued that as immigrants and their offspring integrate into the host society, they adopt the norms, attitudes and behaviours of the new country, and experience the accompanying health consequences.³²

Table 3

Age- and fully-adjusted odds ratios relating generation status and South Asian or Chinese descent to all-cause and circulatory disease-related hospitalization, by sex, long-form 2006 Census respondents aged 30 or older in urban areas, Canada excluding Quebec and territories, 2006/2007 to 2009/2010

Sex, generation status and South Asian or Chinese descent	All-cause hospitalization (excluding pregnancy)						Circulatory disease-related hospitalization					
	Age-adjusted odds ratio	95% confidence interval		Fully adjusted [†] odds ratio	95% confidence interval		Age-adjusted odds ratio	95% confidence interval		Fully adjusted [†] odds ratio	95% confidence interval	
		from	to		from	to		from	to		from	to
Men												
Third-plus generation (G3+) [‡]	1.00	1.00	1.00	1.00
Total												
Second generation (G2)	0.95*	0.94	0.97	0.97*	0.96	0.99	0.95*	0.93	0.98	0.97	0.94	1.00
First generation (G1)												
Long-term (10 or more years)	0.78*	0.77	0.79	0.79*	0.78	0.80	0.82*	0.80	0.84	0.83*	0.81	0.85
Recent (less than 10 years)	0.49*	0.48	0.51	0.49*	0.48	0.51	0.55*	0.52	0.59	0.56*	0.52	0.60
South Asia												
Second generation (G2)	0.71*	0.60	0.84	0.74*	0.63	0.88	0.87	0.64	1.20	0.91	0.65	1.25
First generation (G1)												
Long-term (10 or more years)	0.74*	0.71	0.77	0.75*	0.72	0.78	0.97	0.91	1.04	1.01	0.94	1.08
Recent (less than 10 years)	0.50*	0.47	0.53	0.49*	0.46	0.52	0.71*	0.64	0.79	0.71*	0.64	0.79
China[§]												
Second generation (G2)	0.64*	0.56	0.72	0.67*	0.59	0.76	0.52*	0.40	0.69	0.55*	0.42	0.72
First generation (G1)												
Long-term (10 or more years)	0.44*	0.42	0.46	0.43*	0.41	0.45	0.37*	0.34	0.41	0.37*	0.33	0.40
Recent (less than 10 years)	0.31*	0.28	0.34	0.30*	0.27	0.32	0.24*	0.19	0.29	0.23*	0.19	0.28
Women												
Third-plus generation (G3+) [‡]	1.00	1.00	1.00	1.00
Total												
Second generation (G2)	0.96*	0.94	0.97	0.97*	0.96	0.99	0.97	0.94	1.00	0.98	0.95	1.01
First generation (G1)												
Long-term (10 or more years)	0.74*	0.73	0.75	0.74*	0.73	0.75	0.79*	0.77	0.82	0.78*	0.75	0.90
Recent (less than 10 years)	0.53*	0.52	0.52	0.51*	0.5	0.53	0.52*	0.48	0.56	0.49*	0.45	0.54
South Asia												
Second generation (G2)	0.74*	0.62	0.85	0.73*	0.63	0.89	0.67	0.42	1.07	0.71	0.44	1.13
First generation (G1)												
Long-term (10 or more years)	0.67*	0.64	0.70	0.66*	0.63	0.69	0.77*	0.70	0.85	0.74*	0.67	0.82
Recent (less than 10 years)	0.54*	0.51	0.57	0.51*	0.48	0.54	0.67*	0.58	0.78	0.61*	0.52	0.71
China[§]												
Second generation (G2)	0.72*	0.64	0.80	0.74*	0.66	0.83	0.63*	0.46	0.88	0.67*	0.48	0.92
First generation (G1)												
Long-term (10 or more years)	0.43*	0.41	0.45	0.41*	0.39	0.43	0.34*	0.30	0.38	0.31*	0.27	0.34
Recent (less than 10 years)	0.36*	0.34	0.39	0.34*	0.31	0.36	0.25*	0.20	0.32	0.23*	0.18	0.29

[†] adjusted for age, secondary school graduation, and low income status

[‡] reference group G3+ is for overall population

[§] includes Hong Kong

* significantly different from reference group ($p < 0.05$)

... not applicable

Source: Linked 2006 Census-2006/2007 to 2009/2010 Discharge Abstract Database.

Research on cardiovascular disease points to acculturation as a contributor to the increasing risk among immigrants and the second generation.³³ For instance, people of Mexican origin who were born in the United States had higher levels of hypertension, smoking, cholesterol, diabetes and obesity than

their Mexican immigrant counterparts^{34,35} and consumed less “heart healthy diets.”³⁶ In Canada, cardiovascular disease risks among South Asians and Chinese immigrants tend to increase with duration of residence.¹¹

The present study found higher odds of circulatory disease-related hospitalization among first- and second-generation South Asians, compared with the Chinese. Similar findings have been reported in the United States, the United Kingdom, South Africa, the Caribbean, and Singapore.^{5,37-39} For example, in the United Kingdom, an excess risk of coronary artery problems among first- and second-generation South Asians was reported.⁴⁰ A study of Indian immigrants in London, England found that they had higher body mass index, systolic blood pressure and fasting blood glucose than their siblings in India.⁴¹ South Asians in Canada have a high prevalence of risk factors such as diabetes, which may explain some of the findings in this study.^{7,11,42} By contrast, the Chinese have the most favourable circulatory disease risk factor profile of the major ethnic groups in Canada, although it worsens with duration of residence. Contrary to these findings, an Australian study reported that the risk of circulatory- and diabetes-related mortality among older South Asians decreased with duration of residence, while immigrants from the United Kingdom and Ireland experienced increasing mortality over time.⁴³

People of South Asian descent tend to have one of the highest rates of circulatory disease in the world.^{6,44} Nonetheless, results of this study revealed a lower risk of hospitalization among recent immigrants from South Asia, compared with third-plus-generation Canadians. This may reflect the selection process that involves medical screening of potential immigrants.⁴⁵ However, additional analyses (data not shown) found relatively high circulatory disease-related hospitalization rates among younger South Asians, especially G1, which is consistent with an early onset of heart problems.^{46,47} As well, long-term male immigrants

from South Asia did not differ from G3+ in the likelihood of experiencing a circulatory disease-related hospitalization. Case-control research has found that South Asians tend to be hospitalized later in the course of acute myocardial infarction,⁴⁸ which may indicate differences in care-seeking behaviour and/or cultural or language problems.

This study found that people of Chinese descent, both G1 and G2, had significantly lower odds of circulatory disease-related hospitalization than G3+, even with adjustment for education and income. Consistent with these results, an analysis of linked census and mortality data³ revealed lower circulatory disease mortality rates among recent and long-term and immigrants (G1) from East Asia (Hong Kong, Japan, Macao, Mongolia, North Korea, People’s Republic of China, Taiwan, Singapore, South Korea), compared with the Canadian-born population (largely G3+). The same study showed that South Asian male immigrants who arrived after 1971 also had significantly lower circulatory disease mortality rates, but South Asian female immigrants’ mortality rates were similar to those of Canadian-born women. The latter is not consistent with the significantly lower hospitalization rates for G1 South Asian women found in this analysis. This indicates potential under-utilization of circulatory disease-related health care by female immigrants from South Asia.

Limitations

The limitations of the linked census-hospitalization dataset include the absence of information about diet, smoking, exercise and social capital (connectedness and support). A lack of information about non-hospital care received from family doctors and specialists or about alternative care, which may be more prevalent among some ethnic groups, is also important because low hospitalization rates may reflect efficient use of and access to primary or alternative health care, not better health. As well, the G2 South Asian and Chinese

What is already known on this subject?

- Canadian studies of immigrant integration have explored economic outcomes for the second generation, but patterns of health and health care use by generation status have received little attention thus far.
- South Asian immigrants have a circulatory disease mortality risk similar to the Canadian-born population, but higher than other immigrant groups such as the Chinese.
- There is a need to determine if the relatively high circulatory disease mortality risk among South Asians is reflected in hospitalization, and if the health risks affect subsequent generations.

What does this study add?

- Using linked census-hospital data, this study examined all-cause (excluding pregnancy) and circulatory disease-related hospitalizations among first- and second-generation of immigrants, particularly, South Asians and Chinese.
- Based on age-standardized hospitalization rates, South Asian immigrants’ health advantage in circulatory disease is lost by the second generation.
- Among the Chinese, all-cause and circulatory disease-related hospitalization risk generally increased across generations but remained significantly lower than for Canadians overall.

populations are still relatively young, and their group sizes are small. For example, the unadjusted and adjusted odds of circulatory disease-related hospitalization among G2 women of South Asian descent—0.67 and 0.71—would seem to be significantly low, but the confidence intervals were wide. Further research is needed to determine if this is a sample size issue or if circulatory disease-related hospitalizations are actually less prevalent in this group. Finally, linkage of the long-form census with hospital records could differ by place of birth. In particular, linkage of 2006 Census results to tax data to obtain changes in postal code information, which relied on names, may yield lower linkage rates for certain subgroups (for example, Chinese).

This would be an issue only if the link was accomplished via the additional postal code information derived from the tax data. The majority of links for this study were established in the first iteration with an exact match of the postal code on the census record.

Conclusion

The results of this analysis suggest that immigrants' health advantage diminishes in the second generation, particularly for circulatory disease. For those of South Asian descent, the health advantage was lost in the second generation, while for those of Chinese descent, the advantage decreased, but persisted compared with the third-plus generation. These findings

have potentially important implications for public health and future needs for health care services. ■

Acknowledgements

The authors thank David Mowat and Julie Stratton of the Office of the Medical Officer of Health Region of Peel for reviewing an earlier version of this paper. Jack V. Tu is supported by a Canada Research Chair in Health Services Research and a Career Investigator Award from the Heart and Stroke Foundation-Ontario.

References

- Chui T, Tran K, Maheux H. *Immigration in Canada: A Portrait of the Foreign-born Population, 2006 Census* (Catalogue 97-557-XIE) Ottawa: Statistics Canada, 2007.
- Statistics Canada. *Projections of the Diversity of the Canadian Population* (Catalogue 91-551-X) Ottawa: Statistics Canada, 2010.
- Ng E, LHAD Research Team. *Insights into the Healthy Immigrant Effect: Mortality by Period of Immigration and Birthplace*. Health Analysis Division Working Paper No. 8 (Catalogue 82-622) Ottawa: Statistics Canada, 2011.
- Rotermann M. The impact of considering birthplace in analyses of immigrant health. *Health Reports* 2011; 22(4): 37-43.
- Bhopal R. Epidemic of cardiovascular disease in South Asians. *British Medical Journal* 2002; 324(7338): 625-6.
- Celermajer DS, Chow CK, Marijon E, et al. Cardiovascular disease in the developing world: prevalences, patterns and the potential of early disease detection. *Journal of the American College of Cardiology* 2012; 16(14): 1207-16.
- Lau D. Excess prevalence and mortality rates of diabetes and cardiovascular disease among South Asians: A call to action. *Canadian Journal of Diabetes* 2010; 34(2): 102-4.
- Sheth T, Nair C, Nargundkar N, et al. Cardiovascular and cancer mortality among Canadians of European, South Asian and Chinese origin from 1979 to 1993: an analysis of 1.2 million deaths. *Canadian Medical Association Journal* 1999; 161(2): 132-8.
- Muening P, Jia H, Khan K. Hospitalization for heart disease, stroke and diabetes mellitus among Indian-born persons: a small area analysis. *BMC Cardiovascular Disorders* 2004; 4:19. doi:10.1186/1471-2261-4-19.
- Chiu M, Austin PC, Manuel DG, et al. Comparison of cardiovascular risk profiles among ethnic groups using population health surveys between 1996 and 2007. *Canadian Medical Association Journal* 2010; 182(2): E301-10.
- Chiu M, Austin PC, Manuel DG, et al. Cardiovascular risk factor profiles of recent immigrants vs long-term residents of Ontario: a multi-ethnic study. *Canadian Journal of Cardiology* 2012; 28: 20-6.
- Hyman I. *Immigration and Health: Reviewing Evidence of the Healthy Immigrant Effect in Canada*. Joint Centre of Excellence for Research on Immigration and Settlement (CERIS) Working Paper No. 55. Toronto: CERIS, 2007.
- Leão TS, Johansson L-M, Sundquist K. Hospitalization due to alcohol and drug abuse in first- and second-generation immigrants: a follow-up study in Sweden. *Substance Use and Misuse* 2006; 41: 283-96.
- Hosper K, Nierkens V, Nicolaou M, et al. Behavioural risk factors in two generations of non-Western migrants: do trends converge towards the host population? *European Journal of Epidemiology* 2007; 22: 163-72.
- Almeida J, Johnson RM, Matsumoto A, et al. Substance use, generation and time in the United States: the modifying role of gender for immigrant urban adolescents. *Social Science & Medicine* 2012; 75: 2069-75.
- Marmot MG, Syme SL. Acculturation and coronary heart disease in Japanese-Americans. *American Journal of Epidemiology* 1976; 104(3): 225-47.
- Quon EC, McGrath JJ, Roy-Gagnon M-H. Generation of immigration and body mass index in Canadian youth. *Journal of Pediatric Psychology* 2012; 37(8): 843-53.
- Maximova K, O'Loughlin J, Gray-Donald K. Healthy weight advantage lost in one generation among immigrant elementary school children in multi-ethnic, disadvantaged, inner-city neighborhoods in Montreal, Canada. *Annals of Epidemiology* 2011; 21(4): 238-44.
- Hamilton HA, Noh S, Adlaf EM. Adolescent risk behaviours and psychological distress across immigrant generations. *Canadian Journal of Public Health* 2009; 100(3): 221-5.
- Zunzunegui MV, Forster M, Gauvin L, et al. Community unemployment and immigrants' health in Montreal. *Social Science and Medicine* 2006; 63: 485-500.
- Gagnon AJ. *Responsiveness of the Canadian Health Care System Towards Newcomers*. Commission on the Future of Health Care in Canada Discussion Paper No. 40 (Catalogue CP32-79/40-2002E-IN) Ottawa: Royal Commission on Health Care in Canada, 2002.
- Carrière G, Peters PA, Sanmartin C. Area-based methods to calculate hospitalization rates for the foreign-born population in Canada, 2005-2006. *Health Reports* 2012; 23(3): 43-51.

**All-cause and circulatory disease-related hospitalization, by generation status:
Evidence from linked data • Research Article**

23. Rotermann M, Sanmartin C, Carrière G, et al. Comparing two approaches to link census and hospital data in Canada. *Health Reports* 2014; 25(10): 3-14.
24. Statistics Canada. *Approved Record Linkages*. Linkage of the Census of Population 2006 to the Discharge Abstract Database and the Canadian Mortality Database for Purposes of the Longitudinal Health and Administrative Data Initiative. Ottawa: Statistics Canada, 2012. Available at : <http://www.statcan.gc.ca/record-enregistrement/2012-eng.htm#a15>
25. Statistics Canada. *Directive on Record Linkage*. Ottawa: Statistics Canada, 2011. Available at : <http://www.statcan.gc.ca/record-enregistrement/policy4-1-politique-4-1-eng.htm>
26. Statistics Canada. *2006 Census Dictionary* (Catalogue 92-566-X) Ottawa: Minister of Industry, 2010.
27. Statistics Canada. *Immigration and Ethno-cultural Diversity in Canada, National Household Survey, 2011* (Catalogue 99-010-X2011001) Ottawa: Statistics Canada, 2013.
28. Carriere KC, Roos LL. A method of comparison for standardized rates of low-incidence events. *Medical Care* 1997; 35(1): 57-69.
29. SAS Institute Inc. *SAS 9.2 User's Guide*. Cary, North Carolina: SAS Institute Inc., 2009.
30. Newbold KB. Health status and health care utilization of the foreign-born in Canada: A longitudinal analysis, 1994/1995-2000/01. *Journal of Health Services Research and Policy* 2005; 10(2): 77-83A.
31. Ng E, Wilkins R, Gendron F, Berthelot J-M. Dynamics of immigrants' health in Canada: evidence from the National Population Health Survey. *Healthy Today, Healthy Tomorrow? Findings from the National Population Health Survey*. Ottawa: Statistics Canada, 2005.
32. Hyman I. *Immigration and Health*. Health Policy Working Paper Series Working Paper No. 01-05. Ottawa: Health Canada, 2001.
33. Morales LS, Leng M, Escarce JJ. Risk of cardiovascular disease in first and second generation Mexican-Americans. *Journal of Immigrant and Minority Health* 2011; 13: 61-8.
34. Sundquist J, Winkleby M. Country of birth, acculturation status and abdominal obesity in a national sample of Mexican American women and men. *International Journal of Epidemiology* 2000; 29(3): 470-7.
35. Singh GK, Siapush M. All-cause and cause-specific mortality of immigrants and native born in the United States. *American Journal of Public Health* 2001; 91: 392-9.
36. Dixon LB, Sundquist J, Winkleby M. Differences in energy, nutrient and food intakes in a US sample of Mexican-American women and men: findings from the third national health and nutrition examination survey, 1988-1994. *American Journal of Epidemiology* 2000; 152(6): 548-57.
37. Kuppuswamy VC, Gupta S. Excess coronary heart disease in South Asians in the United Kingdom. *British Medical Journal* 2005; 330: 1223-4.
38. Harding S, Rosato M, Teyhan A. Trends for coronary heart disease and stroke mortality among migrants in England and Wales 1979-2003: Slow declines notable for some groups. *Heart* 2008; 94: 463-70.
39. Yusuf S, Reddy S, Ounpuu S, Anand S. Global burden of cardiovascular diseases: part II: variations in cardiovascular disease by specific ethnic groups and geographic regions and prevention strategies. *Circulation* 2001; 104: 2855-64.
40. Shaikat N, DeBono DP, Jones DR. A comparison of risk factors in south Asian and north European coronary artery patients with their young male offspring. *British Heart Journal* 1994; 71: 27.
41. Bhatnager D, Anand IS, Durrington PN, et al. Coronary risk factors in people with the Indian subcontinent living in West London and their siblings in India. *Lancet* 1995; 345: 405-9.
42. Ayesha R, de Souza RJ, Kandasamy S, et al. Cardiovascular risk among South Asians living in Canada: a systematic review and meta-analysis. *Canadian Medical Association Journal* 2014; 2(3): E183-91. Accessed October 1, 2014. doi: 10.9778/cmajo.20130064.
43. Gray, L, Harding S, Reid A. Evidence of divergence with duration of residence in circulatory disease mortality in migrants to Australia. *European Journal of Public Health* 2007; 17(6): 550-4.
44. Goyal A, Yusuf S. The burden of cardiovascular disease in the Indian subcontinent. *Indian Journal of Medical Research* 2006; 124: 235-44.
45. Gushulak BD, Williams LS. National immigrant health policy: existing policy, changing needs, and future directions. *Canadian Journal of Public Health* 2004; 95(3): 127-9.
46. Prasad DS, Kabir Z, Dash AK, Das BC. Childhood cardiovascular risk factors in South Asians: a cause of concern for adult cardiovascular disease epidemic. *Annals of Pediatric Cardiology* 2011; 4(2): 166-71. Accessed February 12, 2014. doi:10.4103/0974-2069.84663
47. McKeigue PM, Ferrie JE, Pierpoint T, et al. Association of early-onset coronary heart disease in South Asian men with glucose intolerance and hyperinsulinemia. *Circulation* 1993; 87: 152-61.
48. Gupta M, Doobay AV, Singh N, et al. Risk factors, hospital management and outcomes after acute myocardial infarction in South Asian Canadians and matched control subjects. *Canadian Medical Association Journal* 2002; 166: 717-22.