


Health Reports

The 2006 Canadian Birth-Census Cohort

by Tracey Bushnik, Seungmi Yang, Michael S. Kramer,
Jay S. Kaufman, Amanda J. Sheppard and Russell Wilkins

Release date: January 20, 2016



Statistics
Canada

Statistique
Canada

Canada

How to obtain more information

For information about this product or the wide range of services and data available from Statistics Canada, visit our website, www.statcan.gc.ca.

You can also contact us by

email at STATCAN.infostats-infostats.STATCAN@canada.ca

telephone, from Monday to Friday, 8:30 a.m. to 4:30 p.m., at the following toll-free numbers:

- | | |
|---|----------------|
| • Statistical Information Service | 1-800-263-1136 |
| • National telecommunications device for the hearing impaired | 1-800-363-7629 |
| • Fax line | 1-877-287-4369 |

Depository Services Program

- | | |
|------------------|----------------|
| • Inquiries line | 1-800-635-7943 |
| • Fax line | 1-800-565-7757 |

Standards of service to the public

Statistics Canada is committed to serving its clients in a prompt, reliable and courteous manner. To this end, Statistics Canada has developed standards of service that its employees observe. To obtain a copy of these service standards, please contact Statistics Canada toll-free at 1-800-263-1136. The service standards are also published on www.statcan.gc.ca under “Contact us” > “Standards of service to the public.”

Note of appreciation

Canada owes the success of its statistical system to a long-standing partnership between Statistics Canada, the citizens of Canada, its businesses, governments and other institutions. Accurate and timely statistical information could not be produced without their continued co-operation and goodwill.

Standard table symbols

The following symbols are used in Statistics Canada publications:

- . not available for any reference period
- .. not available for a specific reference period
- ... not applicable
- 0 true zero or a value rounded to zero
- 0^s value rounded to 0 (zero) where there is a meaningful distinction between true zero and the value that was rounded
- ^P preliminary
- ^r revised
- X suppressed to meet the confidentiality requirements of the *Statistics Act*
- ^E use with caution
- F too unreliable to be published
- * significantly different from reference category ($p < 0.05$)

Published by authority of the Minister responsible for Statistics Canada

© Minister of Industry, 2016

All rights reserved. Use of this publication is governed by the Statistics Canada [Open Licence Agreement](#).

An HTML version is also available.

Cette publication est aussi disponible en français.

The 2006 Canadian Birth-Census Cohort

by Tracey Bushnik, Seungmi Yang, Michael S. Kramer, Jay S. Kaufman,
Amanda J. Sheppard and Russell Wilkins

Abstract

Background: Evidence on socioeconomic and ethnocultural disparities in perinatal health in Canada tends to be limited to analyses by neighbourhood or for selected provinces. In 2010, the Canadian Institutes of Health Research awarded funding for a project on perinatal outcomes. This article describes the resulting 2006 Canadian Birth-Census Cohort Database.

Data and methods: From the Canadian Live Birth, Infant Death and Stillbirth Database, 687,340 records of children born in Canada from May 16, 2004 through May 15, 2006 to mothers whose usual place of residence was Canada were selected as in-scope births. Deterministic rules were applied to link each person on the birth record—child, mother, father—to 2006 Census data. The cohort was restricted to records linked to a long-form questionnaire ($n = 135,426$), and a cohort weight was developed. Cohort rates (unweighted and weighted) for five birth outcomes—preterm birth, small-for-gestational age, large-for-gestational age, stillbirth, and infant mortality—were compared with rates for all in-scope births across birth characteristics. Cohort rates for these birth outcomes were examined across selected census characteristics.

Results: Linkage rates were 91% for births surviving to age 1, 76% for stillbirths, and 80% for infant deaths matched to a birth registration. The cohort estimates were similar to those for all in-scope births, particularly after the cohort weight was applied. The cohort data produced plausible estimates of selected birth outcomes across maternal ethnocultural categories and levels of education.

Interpretation: The 2006 Canadian Birth-Census Cohort data can help inform perinatal surveillance and research in Canada.

Keywords: Birth outcomes, ethnocultural, infant mortality, record linkage, socioeconomic, stillbirths

Reducing health disparities is an ongoing population health goal in Canada and other countries.¹ A step toward achieving this goal is to exploit existing data on the nature and extent of variations in health across socioeconomic and ethnocultural groups. Evidence on disparities in perinatal health in Canada has generally been limited to analyses by neighbourhood characteristics,² or for selected provinces,³⁻⁵ owing to a lack of socioeconomic and ethnocultural information in most routinely collected perinatal data.

Perinatal health is monitored and documented by the Canadian Perinatal Surveillance System, led by the Public Health Agency of Canada in collaboration with Statistics Canada. As in previous years, the System's 2013 report⁶ presented data from existing databases, including vital statistics, for live births, stillbirths, and infant deaths. However, these national databases contain little socioeconomic or ethnocultural information.

In 2010, the Canadian Institutes of Health Research funded a project on socioeconomic position, ethnocultural background, and perinatal outcomes. The aim was to link the databases used for perinatal surveillance to long-form census data for 1996 and 2006, thereby creating *birth-census* cohorts.

Approval for this linkage was granted by Statistics Canada's Policy Committee (now known as the Executive Management Board) in 2012. The project received ethics approval from the Pediatric Research Ethics Board of the McGill University Health Centre Research Institute, and was supported by the Public

Health Agency of Canada, the Canadian Perinatal Surveillance System, and Health Canada.

This article is an overview of the creation, content, and quality of the 2006 Canadian Birth-Census Cohort Database. Similar information about the 1996 Canadian Birth-Census Cohort Database is available in its userguide.⁷

Methods

Overview

Births that met the inclusion criteria (within the study period; to women resident in Canada) were selected from a database of previously linked live birth, infant death and stillbirth records. Those records were matched to 2006 Census records from both short- and long-form questionnaires and then restricted to those from long-form questionnaires to create the 2006 Canadian Birth-Census Cohort Database. The match to *all* census records was performed to reduce the possibility of false positive matches to a long-form record, as birth records best matched to a short-form census record were flagged as ineligible for inclusion in the cohort. It also allowed evaluation of the linkage rate of births to census data across various birth characteristics. To ensure respondent privacy, Statistics Canada employees involved in the process accessed only the identifying information required for linkage, not health-related information. When linkage was completed, identifying information was removed from the final analytical file.

Authors: Tracey Bushnik (tracey.bushnik@canada.ca) is with the Health Analysis Division at Statistics Canada. Seungmi Yang and Jay S. Kaufman are with the Department of Epidemiology, Biostatistics and Occupational Health at McGill University. Michael S. Kramer is with the Departments of Pediatrics and of Epidemiology, Biostatistics and Occupational Health at McGill University and the McGill University Health Centre Research Institute. Amanda J. Sheppard is with the Dalla Lana School of Public Health at the University of Toronto. Russell Wilkins is with the Department of Epidemiology and Community Medicine at the University of Ottawa.

Data sources

Vital statistics

Provincial and territorial vital statistics registrars collect information on live births, stillbirths and deaths in their respective jurisdictions. In collaboration with the registrars, Statistics Canada compiles this information into three national databases, which have been combined to create the Canadian Live Birth, Infant Death and Stillbirth Database. That database is a census of all live births and stillbirths, in which live births have been matched to the Canadian Mortality Data Base to identify infants who died during their first year (0 to 364 days). The database identifies four types of events: live births for which a death record within the first year was not found (*births surviving to age 1*); stillbirths; deaths during the first year for which a birth registration was found (*infant deaths matched to a birth registration*); and deaths during the first year for which a birth registration was not found (*infant deaths not matched to a birth registration*). The database contains information such as birth weight and gestational age; maternal and paternal age at child's birth; and information about the death such as cause.

From the 1985 to 2008 Canadian Live Birth, Infant Death and Stillbirth Database, 687,340 records of children born in Canada from May 16, 2004 through May 15, 2006 (one day before census day) to mothers whose usual place of residence was Canada were selected. These were the in-scope births. The two-year period before census day was chosen to generate adequate sample size, while limiting the time elapsed between the birth date and census day. The latter is important for the analysis of associations between perinatal outcomes at the time of birth and time-varying characteristics captured on the census such as maternal education.

2006 Census of Population

Using short- or long-form (20% sample) questionnaires, the 2006 Census of Population collected information on individuals living in all households

that were enumerated. The short-form questionnaire collected each person's name, address and postal code, date of birth, sex, marital status, mother tongue, and relationship to "Person 1" (head of household). The long-form questionnaire collected the short-form information, plus data on characteristics such as ethnicity, Aboriginal identity, education, and income.⁸

Only people enumerated by the census could be linked to corresponding birth records. Reasons for not being enumerated included emigration, death, and census undercoverage. Net undercoverage of the 2006 Census was 2.7% for the population younger than age 5 (an estimated 47,213 children), 2% to 6% for women aged 20 to 44 (undercoverage was greater for women without a partner), and 10.6% for people living in Indian reserves and settlements that participated in the census.⁹

A total of 30,537,738 individuals were in the 2006 Census data to which the births were initially linked.

Linkage variables

Five key linkage variables were available in the Canadian Live Birth, Infant Death and Stillbirth Database: child's date of birth (DOB), child's sex, mother's DOB, father's DOB, and postal code of mother's residence at the time of the child's birth. Names were used only in conjunction with one or more of the above linkage variables. The child's DOB was complete for all records; sex of the child, postal code and mother's DOB were complete for almost all records (99%). The father's information was less complete: year of birth was complete for 95% of births surviving to age 1; 73% for stillbirths; and 87% for infant deaths matched to a birth registration. The child's DOB and sex could not be used as linkage variables for stillbirths or for infant deaths that occurred before census day, as those children would not have been enumerated in the census household. Nor was parental DOB or postal code at birth available as linkage variables for infant deaths not matched to a birth registration, because of the lack of birth registration data for those events.

Linkage process

Based on a series of linkage rules (referred to as waves) that were ordered hierarchically from most to least discriminatory, each person on the birth record—child, mother, father—was linked deterministically (by exact matches) to the census data. The strongest waves included perfect matches between records for at least two DOBs, the full postal code and child sex, and accounted for 71% of the total number of matches. Waves of lesser strength used names and allowed for links among fewer linkage variables. This was important because up to two years had passed between the day of birth and census day, and the family composition and place of residence listed on the birth record could have changed. The linkage strategy did not allow for individuals on the same birth record to be linked to *multiple* census households. When this occurred (rarely), the link based on the greater number and/or better quality of linkage variables was retained, and the other links were discarded.

Most waves involved a two-step match process. The initial match was as described above. The second match dealt with instances where not everyone from the birth record was found within a given census household during the initial match. The second step aimed to find, within such census households, the person or persons from the birth record who initially had not been matched. Not finding these other persons could reflect a situation such as a lone-parent family, but it could also have resulted from data error. Thus, the second match included permutations of the linkage variables such as month/day inversions.

Quality control

Lower linkage rates were achieved for stillbirths and infant deaths, and for births surviving to age 1 to mothers younger than 25. To improve these rates, some linkage constraints were relaxed, such as allowing a first name versus a full name match. Additional potential links to census households that had completed a long-form questionnaire (approximately 1,400) were evaluated manually. The addition of manually

approved links increased the combined number of stillbirths and infant deaths by 10 percentage points, and the number of births to young mothers by one percentage point.

When linkage work was complete, a 100% manual review and verification of all in-scope records matched to a census household that had completed a long-form questionnaire showed that the overall false positive match rate was less than 1%.

Creation of analytical cohort

In-scope birth records linked to a census household that completed a long-form questionnaire were considered part of the analytical cohort. Person-, census-family- and economic-family-level census variables were assigned to each person on the birth record (child, mother, father) who had linked to an individual in a census household. Household- and dwelling-level census variables were

added to each linked birth record as a whole because all persons on the birth record were presumed to reside in the same household.

Creation of a cohort weight

Because the analytical cohort was roughly a 20% sample of in-scope births, a cohort weight was generated to produce estimates about the characteristics of all in-scope births. The cohort weight was

Table 1
Comparison of cohort and in-scope births, by selected birth and maternal characteristics, Canada, 2004 to 2006

Birth and maternal characteristics	In-scope population		Linkage rate†	Birth-Census Cohort		Birth-Census Cohort (weighted)		Ratio (B/A)	Ratio (C/A)
	Number	% distribution (A)		Number	% distribution (B)	Number	% distribution (C)		
Total	687,340	100.0	90.3	135,426	100.0	687,340	100.0	1.00	1.00
Type of birth									
Surviving to age 1	679,431	98.8	90.5	134,002	98.9	679,399	98.8	1.00	1.00
Stillbirth	4,240	0.6	76.3	808	0.6	4,242	0.6	0.97	1.00
Infant death (matched to birth registration)	2,984	0.4	79.8	616	0.5	3,699	0.5	1.05	1.01
Infant death (not matched to birth registration)	685	0.1	0.0	0	0.0	0	0.0
Province/Territory of birth									
Newfoundland and Labrador	9,111	1.3	91.6	1,794	1.3	9,111	1.3	1.00	1.00
Prince Edward Island	2,664	0.4	91.2	457	0.3	2,664	0.4	0.87	1.00
Nova Scotia	17,345	2.5	91.0	3,307	2.4	17,345	2.5	0.97	1.00
New Brunswick	13,966	2.0	92.1	2,777	2.1	13,966	2.0	1.01	1.00
Quebec	151,307	22.0	92.8	29,250	21.6	151,307	22.0	0.98	1.00
Ontario	270,086	39.3	90.0	48,638	35.9	270,086	39.3	0.91	1.00
Manitoba	29,247	4.3	90.3	8,584	6.3	29,247	4.3	1.49	1.00
Saskatchewan	24,367	3.5	89.1	6,458	4.8	24,367	3.5	1.35	1.00
Alberta	84,367	12.3	89.7	17,424	12.9	84,367	12.3	1.05	1.00
British Columbia	81,970	11.9	86.9	15,026	11.1	81,970	11.9	0.93	1.00
Yukon	685	0.1	91.1	266	0.2	685	0.1	1.97	1.00
Northwest Territories	1,452	0.2	87.3	825	0.6	1,452	0.2	2.88	1.00
Nunavut	773	0.1	82.1	620	0.5	773	0.1	4.07	1.00
Postal code at birth									
Urban	563,406	82.0	90.4	101,496	74.9	562,915	81.9	0.91	1.00
Rural	120,245	17.5	90.6	32,970	24.3	121,907	17.7	1.39	1.01
Sex of child									
Male	352,849	51.3	90.2	69,325	51.2	352,779	51.3	1.00	1.00
Female	334,432	48.7	90.4	66,089	48.8	334,501	48.7	1.00	1.00
Year of birth									
2004	214,647	31.2	89.8	42,222	31.2	214,647	31.2	1.00	1.00
2005	344,654	50.1	90.3	67,670	50.0	344,654	50.1	1.00	1.00
2006	128,039	18.6	91.0	25,534	18.9	128,039	18.6	1.01	1.00
Maternal age group (years)									
Younger than 20	28,203	4.1	77.3	6,736	5.0	28,218	4.1	1.21	1.00
20 to 24	110,969	16.1	85.1	22,460	16.6	111,082	16.2	1.03	1.00
25 to 29	211,655	30.8	91.0	41,381	30.6	211,889	30.8	0.99	1.00
30 to 34	215,677	31.4	92.8	41,603	30.7	215,913	31.4	0.98	1.00
35 to 39	99,497	14.5	93.0	19,346	14.3	99,633	14.5	0.99	1.00
40 to 44	19,446	2.8	92.4	3,691	2.7	19,443	2.8	0.96	1.00
45 or older	859	0.1	91.4	165	0.1	870	0.1	0.97	1.01
Maternal place of birth									
In Canada	502,649	73.1	92.0	104,535	77.2	514,031	74.8	1.06	1.02
Outside Canada	172,551	25.1	86.3	29,048	21.4	163,489	23.8	0.85	0.95
Single or multiple birth									
Single	665,362	96.8	90.3	131,285	96.9	666,026	96.9	1.00	1.00
Twin	20,452	3.0	91.3	4,007	3.0	20,472	3.0	0.99	1.00
Triplet or more	841	0.1	91.0	134	0.1	842	0.1	0.81	1.00

... not applicable

† percentage of in-scope birth records linked to a census record from a long- or short-form questionnaire

Sources: Canadian Live Birth, Infant Death and Stillbirth Database 1985 to 2008; 2006 Canadian Birth-Census Cohort Database.

developed from the census household weight, calibrated to marginal totals for the in-scope births, to adjust for missed linkages.¹⁰ Those marginal totals were based on known characteristics for which the linkage rates varied, including type of birth event, year of birth, province of birth, and maternal age group. A set of bootstrap weights that captured both sampling and stochastic variability was also developed to allow users to calculate the corresponding variance of an estimate.¹⁰

Analysis

All estimates were produced using SUDAAN 11.0.1. The SMCOUNT and SMCONF options were used to produce small proportion confidence intervals (CIs) proposed by Korn and Graubard¹¹ (also known as “exact” CIs or CIs based

on the binomial distribution) for in-scope population and unweighted cohort estimates. Weighted cohort estimates with corresponding logit CIs were produced using the cohort weight and the bootstrap weights, respectively.¹⁰

Results

The initial match from in-scope birth records to all census records resulted in matches for 90% of the in-scope births. The linkage rates were 91% for births surviving to age 1, 76% for stillbirths, 80% for infant deaths matched to a birth registration, and 0% for infant deaths not matched to a birth registration (Table 1).

Stillbirths and infant deaths were less likely to be linked (fewer linkage variables available), as were births in British Columbia, the Northwest Territories and

Nunavut, and births to mothers younger than age 25 or to mothers not born in Canada. Birth records missing a postal code, sex of child, age of mother, or maternal place of birth were also less likely to be linked to a census household (data not shown).

Representativeness of cohort

The 2006 Birth-Census Cohort consists of 135,426 linked records. Cohort membership was dependent on overall linkage rates and on the sampling strategy for the 2006 Census. In 2006, one in five occupied private dwellings in self-enumeration areas (householders completed questionnaires) received a long-form questionnaire. All dwellings in areas enumerated by canvassers (generally, remote and northern areas and most Indian reserves, Indian settle-

Table 2
Rates of non-fatal perinatal outcomes for cohort and all in-scope births, by province/territory of birth, maternal age group, and maternal place of birth, Canada, 2004 to 2006

Province/Territory of birth, maternal age group, maternal place of birth	Preterm birth (less than 37 weeks)						Small-for-gestational age (below 10th percentile)						Large-for-gestational age (above 90th percentile)								
	In-scope		Birth-Census Cohort				In-scope		Birth-Census Cohort				In-scope		Birth-Census Cohort						
	Rate per 100	Rate per 100	95% confidence interval		95% confidence interval		Rate per 100	Rate per 100	95% confidence interval		95% confidence interval		Rate per 100	Rate per 100	95% confidence interval		95% confidence interval				
			from	to	from	to			from	to	from	to			from	to					
Total	7.9	7.9	7.7	8.0	7.9	7.8	8.1	8.3	7.8	7.6	7.9 [†]	8.3	8.2	8.5	11.2	12.2	12.1	12.4 [†]	11.1	11.0	11.3
Province/Territory of birth																					
Newfoundland and Labrador	8.4	8.5	7.3	9.9	9.3	7.7	11.1	6.5	6.2	5.1	7.4	7.1	5.8	8.5	15.1	16.7	15.0	18.6	14.6	13.0	16.4
Prince Edward Island	5.4	6.1	4.1	8.7	6.8	4.5	10.2	7.2	5.8	3.8	8.3	6.2	4.1	9.2	15.0	15.1	11.9	18.7	13.7	10.8	17.2
Nova Scotia	8.1	7.5	6.6	8.5	8.1	7.0	9.3	8.0	7.5	6.6	8.5	8.3	7.2	9.5	13.2	15.0	13.8	16.3 [†]	13.9	12.6	15.2
New Brunswick	8.1	7.6	6.6	8.6	7.5	6.4	8.7	7.7	7.8	6.8	8.9	8.9	7.7	10.3	13.2	13.5	12.3	14.9	12.5	11.3	13.9
Quebec	7.7	7.8	7.5	8.1	7.8	7.4	8.1	8.1	7.6	7.3	8.0 [†]	8.2	7.9	8.6	9.8	11.1	10.7	11.4 [†]	10.0	9.6	10.4
Ontario	7.7	7.5	7.2	7.7	7.8	7.5	8.1	8.7	8.3	8.0	8.5 [†]	8.7	8.4	8.9	11.0	11.5	11.2	11.8 [†]	11.1	10.8	11.4
Manitoba	8.5	9.0	8.4	9.6	8.6	7.8	9.5	7.8	7.8	7.2	8.4	8.8	8.0	9.6 [†]	14.6	16.1	15.3	16.9 [†]	13.4	12.6	14.3 [†]
Saskatchewan	7.7	7.8	7.1	8.5	7.7	6.9	8.7	7.3	7.2	6.6	7.9	8.0	7.2	8.9	13.9	15.7	14.8	16.6 [†]	12.4	11.5	13.4 [†]
Alberta	9.3	9.1	8.7	9.6	9.1	8.6	9.6	8.6	8.2	7.8	8.6	8.6	8.1	9.1	10.5	11.1	10.6	11.6 [†]	10.2	9.7	10.7
British Columbia	7.5	7.3	6.8	7.7	7.1	6.7	7.6	7.7	6.7	6.3	7.2 [†]	7.2	6.8	7.7	11.5	12.6	12.0	13.1 [†]	11.8	11.3	12.4
Yukon	6.0	4.2	2.1	7.3	4.1	1.7	9.3	6.7	5.0	2.7	8.4	6.4	3.5	11.7	13.0	13.1	9.3	17.9	10.4	6.8	15.5
Northwest Territories	6.1	6.3	4.7	8.2	6.6	4.5	9.6	6.0	4.3	3.0	5.9 [†]	5.1	3.2	8.2	18.5	19.5	16.8	22.5	19.8	16.3	24.0
Nunavut	12.9	12.9	10.3	15.8	13.3	11.0	15.9	5.0	5.2	3.6	7.3	5.3	3.9	7.1	17.7	17.6	14.7	20.9	17.1	14.5	20.0
Maternal age group (years)																					
Younger than 20	8.5	8.3	7.6	8.9	9.3	8.4	10.2	9.9	8.1	7.4	8.7 [†]	10.8	9.8	11.8	9.9	13.5	12.7	14.3 [†]	9.8	8.9	10.7
20 to 24	7.7	7.7	7.3	8.0	7.6	7.2	8.1	9.8	8.7	8.3	9.1 [†]	9.7	9.2	10.1	9.7	11.5	11.1	11.9 [†]	9.6	9.2	10.0
25 to 29	7.5	7.3	7.1	7.6	7.3	7.0	7.6	8.3	7.8	7.6	8.1 [†]	8.2	8.0	8.5	10.5	11.6	11.3	11.9 [†]	10.5	10.2	10.8
30 to 34	7.7	7.6	7.4	7.9	7.7	7.5	8.0	7.4	7.2	7.0	7.5	7.5	7.2	7.8	11.9	12.4	12.1	12.8 [†]	11.8	11.4	12.1
35 to 39	8.8	9.0	8.6	9.5	8.9	8.4	9.4	7.7	7.7	7.3	8.1	8.1	7.7	8.5	12.8	13.5	13.0	14.0 [†]	13.1	12.5	13.6
40 to 44	10.7	10.7	9.7	11.7	11.2	10.0	12.5	8.7	7.8	7.0	8.8	8.3	7.3	9.4	12.3	13.0	11.9	14.1	12.4	11.3	13.7
45 or older	17.6	15.3	10.2	21.8	18.3	11.9	27.1	9.0	11.4	6.8	17.6	12.1	7.4	19.1	14.0	14.1	8.9	20.7	12.4	7.7	19.4
Maternal place of birth																					
In Canada	8.1	8.0	7.9	8.2	8.1	7.9	8.2	7.3	6.9	6.8	7.1 [†]	7.4	7.2	7.6	12.4	13.4	13.2	13.6 [†]	12.2	12.0	12.4
Outside Canada	7.4	7.3	7.0	7.6	7.6	7.2	7.9	11.0	10.8	10.4	11.1	11.2	10.8	11.5	8.0	8.1	7.8	8.4	7.9	7.6	8.2

[†] confidence interval does not contain point estimate for in-scope births

Sources: Canadian Live Birth, Infant Death and Stillbirth Database 1985 to 2008; 2006 Canadian Birth-Census Cohort Database.

ments, Indian government districts and “terres reserves”), and most persons in non-institutional collective dwellings (excluding children in orphanages and children’s homes) received a long-form questionnaire.¹² This resulted in over-representation of certain groups in the long-form questionnaire sample.

Table 1 shows the number of in-scope births and the number of births (unweighted and weighted) in the cohort. Two sets of ratios are presented: the ratio of the percentage of the cohort to the percentage of in-scope records across selected birth characteristics, and the ratio of the percentage of the *weighted* cohort to the percentage of in-scope records. Categories with a ratio greater than 1 were more likely to be in the cohort. The ratios comparing the unweighted cohort to all in-scope births reflect the sampling strategy of the census:

cohort members were more likely to be from Manitoba, Saskatchewan, Yukon, the Northwest Territories and Nunavut, from rural areas, or to be born to mothers younger than age 20; they were less likely to be from Prince Edward Island, to have mothers who were born outside of Canada, or to be from triplets or higher-order births. When the cohort weight was applied, the ratios comparing the weighted cohort to all in-scope births almost always rounded to 1.0.

Perinatal outcomes

Rates of five birth outcomes for the cohort and for all in-scope births were compared: preterm birth (less than 37 weeks), small-for-gestational age (SGA, sex-specific birth weight below 10th percentile for gestational age), large-for-gestational age (LGA, sex-

specific birth weight above 90th percentile for gestational age), fetal mortality (gestational age of 20 or more weeks or birth weight of at least 500 grams), and infant mortality (death 0 to 364 days after birth). All outcomes were derived as described in the *Perinatal Health Report 2008*,¹³ with SGA and LGA based on Canadian reference values.¹⁴ Tables 2 and 3 show the rates (with 95% confidence intervals (CIs) for the cohort estimates generated using the bootstrap weights) of those outcomes across province of birth, maternal age at child’s birth, and maternal place of birth. (Because a large percentage of stillbirth and infant death records lacked maternal place of birth, this variable was excluded from Table 3.) Rates for the in-scope population that fell outside the 95% CIs for the cohort are noted in the tables. All estimates met the minimum

Table 3
Rates of fatal perinatal outcomes for cohort and all in-scope births, by province/territory of birth and maternal age group, Canada, 2004 to 2006

Province/Territory of birth and maternal age group	Fetal mortality (crude)							Infant mortality (0 to 364 days)								
	In-scope	Birth-Census Cohort				Birth-Census Cohort (weighted)			In-scope	Birth-Census Cohort				Birth-Census Cohort (weighted)		
		Rate per 1,000	Rate per 1,000	95% confidence interval		Rate per 1,000	95% confidence interval			Rate per 1,000	Rate per 1,000	95% confidence interval		Rate per 1,000	95% confidence interval	
				from	to		from	to				from	to		from	to
Total	6.0	5.9	5.5	6.3	6.1	5.9	6.3	5.4	4.6	4.2	5.0 [†]	5.4	5.2	5.6		
Province/Territory of birth																
Newfoundland and Labrador	5.4	5.0	2.3	9.5	3.5	1.8	6.9	5.4	4.5	1.9	8.8	3.2	1.4	7.5		
Prince Edward Island	3.4	0.0	0.0	1.9	0.0	0.0		
Nova Scotia	9.3	9.7	6.6	13.6	9.8	6.8	14.3	4.2	3.4	1.7	6.0	3.9	2.1	7.1		
New Brunswick	4.9	4.3	2.2	7.5	4.7	2.8	8.0	3.7	2.9	1.2	5.7	4.0	1.7	9.5		
Quebec	4.0	3.4	2.8	4.2	4.1	3.3	5.0	4.7	5.2	4.4	6.1	4.7	4.3	5.1		
Ontario	6.4	5.7	5.0	6.4	6.2	5.6	6.8	5.5	2.7	2.2	3.2 [†]	5.5	5.2	5.9		
Manitoba	8.2	10.5	8.5	12.9 [†]	9.7	7.4	12.6	7.1	8.5	6.6	10.7	7.8	5.8	10.3		
Saskatchewan	6.4	8.2	6.2	10.7	8.1	5.6	11.7	6.8	7.8	5.8	10.3	7.0	4.9	10.0		
Alberta	7.0	6.5	5.3	7.8	6.6	5.4	8.1	6.7	6.6	5.4	7.9	6.7	6.1	7.4		
British Columbia	6.6	6.4	5.2	7.8	6.9	5.6	8.5	4.3	4.0	3.1	5.2	4.5	3.9	5.0		
Yukon	F	F	F	F	F	F		
Northwest Territories	8.6	8.8	3.6	18.1	7.1	3.6	14.1	4.9	7.3	2.7	15.9	13.9	4.5	42.1		
Nunavut	F	F	F	11.7	9.7	3.6	21.0	13.4	6.7	26.8		
Maternal age group (years)																
Younger than 20	8.1	8.6	6.6	11.1	8.8	6.4	12.1	8.8	10.6	8.4	13.4	10.2	7.8	13.5		
20 to 24	6.0	5.2	4.3	6.2	5.3	4.3	6.4	5.3	5.9	4.9	7.0	5.9	4.9	7.3		
25 to 29	5.2	5.0	4.4	5.8	5.2	4.5	5.9	3.9	3.5	2.9	4.1	4.3	3.6	5.0		
30 to 34	5.5	5.3	4.6	6.0	5.7	5.0	6.5	3.6	3.8	3.3	4.5	5.0	4.3	5.8		
35 to 39	7.3	7.5	6.3	8.8	7.5	6.4	8.7	4.4	4.4	3.5	5.4	5.9	4.6	7.4		
40 or older [†]	11.1	12.2	9.0	16.2	13.5	9.6	18.8	6.2	7.1	4.7	10.3	10.2	6.9	15.2		


... not applicable

F too unreliable to be published

[†] because of small sample sizes, 40 to 44 and 45 or older age groups were combined

[‡] confidence interval does not contain point estimate for in-scope births

Sources: Canadian Live Birth, Infant Death and Stillbirth Database 1985 to 2008; 2006 Canadian Birth-Census Cohort Database.

Figure 1
Infant mortality rates, including and excluding Ontario, Canada, 2004 to 2006

I = 95% confidence interval

Sources: Canadian Live Birth, Infant Death and Stillbirth Database 1985 to 2008; 2006 Canadian Birth-Census Cohort Database.

sample size requirement of 5 in both the numerator and the denominator.

The rates of preterm birth in the cohort were generally consistent with those for all in-scope births, in that the 95% CIs contained the rates for all in-scope births

(Table 2). However, rates of SGA and LGA for the cohort and in-scope births differed across the three characteristics. Applying the cohort weight eliminated the differences by maternal age group and place of birth, but not for certain provinces.

Fetal mortality rates in the cohort differed slightly from those of all in-scope births for Prince Edward Island (no fetal deaths in the cohort) and Manitoba (Table 3). Applying the cohort weight did not adjust for the lack of cohort fetal deaths in Prince Edward Island.

The overall infant mortality rate was lower for the cohort, primarily because no census record could be found for infant deaths not matched to a birth registration. Applying the weight adjusted the overall estimated infant mortality rate to match that of the in-scope population, because the cohort weight was calibrated to infant death totals, including the 685 not matched to a birth registration. However, the weight did not adjust for the lack of cohort infant deaths in Prince Edward Island.

Almost all (97%) infant deaths not matched to a birth registration occurred in Ontario. Because of concerns about the quality of Ontario's birth registration data,¹³ the province tends to be excluded from most national estimates published by the Canadian Perinatal Surveillance System. Ontario births were excluded entirely from the 1996 Canadian Birth-Census Cohort Database because critical linkage variables were missing from the birth records, and because of the

Table 4
Rates of non-fatal perinatal outcomes for cohort births, by maternal ethnocultural category and highest level of education, 2006 Canadian Birth-Census Cohort

Maternal ethnocultural category and highest level of education	Preterm birth (less than 37 weeks)						Small for gestational age (below 10th percentile)						Large for gestational age (above 90th percentile)					
	Birth-Census Cohort			Birth-Census Cohort (weighted)			Birth-Census Cohort			Birth-Census Cohort (weighted)			Birth-Census Cohort			Birth-Census Cohort (weighted)		
	Rate per 100	95% confidence interval		Rate per 100	95% confidence interval		Rate per 100	95% confidence interval		Rate per 100	95% confidence interval		Rate per 100	95% confidence interval		Rate per 100	95% confidence interval	
	from	to		from	to		from	to		from	to		from	to		from	to	
Total	7.9	7.7	8.0	7.9	7.8	8.1	7.8	7.6	7.9	8.3	8.2	8.5	12.2	12.1	12.4	11.1	11.0	11.3
Ethnocultural category																		
Aboriginal identity	9.2	8.8	9.6	9.1	8.4	9.8	5.5	5.1	5.8	6.3	5.8	6.9	21.5	20.9	22.2	18.9	18.1	19.7
Visible minority	8.1	7.8	8.5	8.4	8.0	8.8	12.3	11.9	12.8	12.6	12.2	13.0	6.9	6.5	7.2	6.7	6.4	7.0
Neither Aboriginal nor visible minority	7.5	7.3	7.7	7.7	7.5	7.8	6.9	6.8	7.1	7.2	7.1	7.4	11.8	11.6	12.1	11.8	11.6	12.0
Highest level of education																		
Less than secondary graduation	8.9	8.5	9.2	8.7	8.3	9.2	8.9	8.5	9.3	10.5	10.0	11.0	13.8	13.4	14.3	11.0	10.5	11.5
Secondary graduation	7.8	7.5	8.1	7.9	7.5	8.3	8.0	7.7	8.3	8.7	8.4	9.1	12.7	12.3	13.1	11.5	11.1	11.9
Postsecondary diploma or certificate	8.0	7.8	8.3	8.2	8.0	8.6	7.5	7.2	7.7	7.9	7.6	8.2	12.0	11.7	12.3	11.2	10.9	11.5
University degree	7.0	6.8	7.3	7.1	6.8	7.4	7.2	6.9	7.4	7.4	7.2	7.7	11.1	10.8	11.4	10.9	10.5	11.2

Note: Estimates based on 133,858 mothers in census households. Maternal ethnocultural category and maternal education were derived from census responses.

Source: 2006 Canadian Birth-Census Cohort Database.

Table 5
Rates of fatal perinatal outcomes for cohort births, by maternal ethnocultural category and highest level of education, 2006 Canadian Birth-Census Cohort

Maternal ethnocultural category and highest level of education	Fetal mortality						Infant mortality					
	Birth-Census Cohort			Birth-Census Cohort (weighted)			Birth-Census Cohort			Birth-Census Cohort (weighted)		
	Rate per 1,000	95% confidence interval		Rate per 1,000	95% confidence interval		Rate per 1,000	95% confidence interval		Rate per 1,000	95% confidence interval	
		from	to		from	to		from	to		from	to
Total	5.9	5.5	6.3	6.1	5.9	6.3	4.6	4.2	5.0	5.4	5.2	5.6
Ethnocultural category												
Aboriginal identity	8.5	7.2	10.0	9.0	7.2	11.2	9.3	7.9	10.8	9.7	8.0	11.8
Visible minority	5.8	4.9	6.9	6.3	5.4	7.3	3.9	3.2	4.8	6.2	5.1	7.5
Neither Aboriginal nor visible minority	5.2	4.7	5.6	5.7	5.3	6.1	3.7	3.3	4.1	4.7	4.4	5.1
Highest level of education												
Less than secondary graduation	7.5	6.5	8.7	6.8	5.7	8.1	8.7	7.5	9.9	8.0	6.7	9.5
Secondary graduation	6.4	5.5	7.4	7.1	6.1	8.2	3.5	2.9	4.3	5.1	4.1	6.4
Postsecondary diploma or certificate	5.4	4.7	6.1	6.0	5.3	6.8	3.8	3.2	4.4	5.0	4.3	5.8
University degree	4.5	3.8	5.2	4.8	4.2	5.6	3.5	2.9	4.1	4.7	3.9	5.6

Note: Estimates based on 133,858 mothers in census households. Maternal ethnocultural category and maternal education were derived from census responses.

Source: 2006 Canadian Birth-Census Cohort Database.

documented data quality concerns for this period. Excluding Ontario raises the cohort's overall infant mortality rate (Figure 1). However, the exclusion makes little difference to the weighted cohort estimates because the cohort weight adjusts for the missing 685 infant deaths.

Perinatal outcomes by maternal census characteristics


Rates for all five birth outcomes were calculated by the mother's ethnocultural background and her highest level of education. Ethnocultural background was grouped into three categories: Aboriginal identity; visible minority (Chinese, South Asian, Black, Filipino, Latin American, Southeast Asian, Arab, West Asian, Korean, Japanese, other visible minority, multiple visible minority); and neither Aboriginal nor visible minority. Highest level of maternal education, based on most advanced certificate, diploma or degree, was grouped into four categories: less than secondary graduation, secondary graduation, postsecondary certificate or diploma (short of a bachelor's degree, including trades certificate), and university degree (bachelor's degree or higher).

Cohort rates of preterm and LGA birth and fetal and infant mortality were higher, and the rate of SGA was lower, among mothers who reported Aboriginal identity than among non-Aboriginal

mothers (Tables 4 and 5). Applying the cohort weight produced similar results, although for LGA and infant mortality, the differences between Aboriginal and non-Aboriginal mothers were somewhat attenuated. Births to mothers from a visible minority had higher rates of SGA and lower rates of LGA than births to other mothers. Rates of fetal and infant mortality were similar among mothers from a visible minority and mothers who were neither Aboriginal nor visible minority. Excluding Ontario had little effect on the patterns in the unweighted and weighted rates of infant mortality across maternal ethnocultural categories (Figure 2 presents weighted rates).

Cohort rates of preterm, SGA and LGA birth, fetal mortality and infant mortality were lower at higher levels of maternal education. Applying the cohort weight yielded a similar pattern for SGA, preterm birth and infant mortality, but flattened the gradient for LGA and lessened the gradient for fetal mortality. Excluding Ontario had little effect on the patterns in rates across maternal levels of education (data not shown).

Figure 2
Weighted infant mortality rates, including and excluding Ontario, by maternal ethnocultural category, 2006 Canadian Birth-Census Cohort



I = 95% confidence interval
Sources: 2006 Canadian Birth-Census Cohort Database.

Discussion

The purpose of the 2006 Canadian Birth-Census Cohort Database is to provide information on the nature and extent of variations in perinatal health across socioeconomic and ethnocultural groups. The results of this analysis suggest that the cohort can help accomplish this goal.

Cohort eligibility was dependent on linkage rates to the census, and on the census sampling of households for the long-form questionnaire. Despite an overall linkage rate of 90%, variations emerged across certain characteristics. A substantial number of those differences were reduced or eliminated by applying the cohort weight; the resulting weighted cohort estimates were consistent with those of all in-scope births.

What is already known on this subject?

- The analysis of perinatal health disparities in Canada has been limited by a lack of socioeconomic and ethnocultural information in most routinely collected perinatal data.
- Various studies suggest that disparities exist and that disadvantaged socioeconomic position is associated with higher risk of adverse birth outcomes.

What does this study add?

- Existing data can be used to uncover the nature and extent of variations in perinatal health across socioeconomic and ethnocultural groups.
- With the 2006 Canadian Birth-Census Cohort Database, it is possible to examine, for all of Canada, perinatal outcomes such as preterm birth, small and large for gestational age, and fetal and infant mortality across a wide range of parental socioeconomic and ethnocultural characteristics.

Differences in birth outcomes across the selected socioeconomic and ethnocultural characteristics were similar to those based on other data sources. The higher rates of preterm birth, LGA, and fetal and infant mortality for births to Aboriginal mothers are consistent with other studies,¹⁵⁻¹⁷ as are the higher rates of preterm birth and infant mortality among mothers with lower educational attainment.¹⁸⁻²⁰ Although these general patterns held whether or not the cohort weight was applied, the importance of applying the weight was apparent for estimates directly affected by census long-form oversampling of remote northern areas and most Indian reserves.

Strengths and limitations

The 2006 Canadian Birth-Census Cohort has several important strengths. It is population-based with a large sample and a cohort weight that permits inference about the *population* of births that the cohort represents. This allows for detailed analyses of perinatal outcomes by characteristics including education, income, ethnicity and Aboriginal identity. Contextual effects (such as neighbourhood) and the health effects of environmental exposures^{21,22} can also be examined. Analyzing results for the 2006 cohort together with the 1996 cohort will reveal the extent to which differences in perinatal outcomes across socioeconomic and ethnocultural groups changed over that 10-year period.

Analysis of the cohort involves a number of limitations. Among the linked records, not all individuals on the birth record were found in the same census household, resulting in missing information for some children, mothers or fathers. Furthermore, the validity of the cohort estimates across the socioeconomic and ethnocultural characteristics could not be evaluated as easily as the cohort estimates of perinatal outcomes across birth characteristics. The latter could be compared directly with the rates for all in-scope births, whereas the former relies on comparisons with the findings of other studies.

To further assess the cohort's face validity, a subsample analysis was undertaken. Quebec is the only province to report information about maternal education to the national birth database. Consequently, it was possible to examine birth outcomes by maternal education for cohort members born in Quebec, and to compare them with rates for all in-scope births in Quebec. Across levels of maternal education, rates of preterm birth, SGA and LGA for the cohort were comparable to those calculated for all in-scope births in Quebec; applying the cohort weight further reduced differences (data not shown). However, small sample sizes resulted in wide confidence intervals for estimates of fetal and infant mortality. Thus, potential bias for fatal outcomes across socioeconomic measures remains a concern because of the relatively low linkage rates for stillbirths and infant deaths.

Conclusion

With two years of birth data and a broad range of socioeconomic and ethnocultural characteristics now linked at the individual level, the 2006 Canadian Birth-Census Cohort offers information that can help inform perinatal surveillance and research in Canada, particularly with respect to non-fatal outcomes. ■

Acknowledgements

The authors are grateful to Statistics Canada employees Martin Lessard, James Brennan and Patrick Gallifa for performing the data linkage; Wei Qian for developing a cohort weight; Lauren Pinault, Jessica Pembroke, Raymond Reaume, and Zimei Zhang for manual review and verification; and Michael Tjepkema and Julie Bernier for managerial oversight and support. Funding for this study was provided by the Canadian Institutes of Health Research (MOP-111122), with support from Statistics Canada and Health Canada (Air Health Effects Research, Population Studies Division).

References

1. Commission on Social Determinants of Health. *Closing the Gap in a Generation: Health Equity through Action on the Social Determinants of Health. Final Report of the Commission on Social Determinants of Health*. Geneva: World Health Organization, 2008.
2. Gilbert NL, Auger N, Wilkins R, Kramer MS. Neighbourhood income and neonatal, postneonatal and sudden infant death syndrome (SIDS) mortality in Canada 1991-2005. *Canadian Journal of Public Health* 2013; 104(3): 187-92.
3. Auger N, Park AL, Zoungrana H, et al. Rates of stillbirth by gestational age and cause in Inuit and First Nations populations in Quebec. *Canadian Medical Association Journal* 2013; 185(6): E256-62.
4. Joseph KS, Liston RM, Dodds L, et al. Socioeconomic status and perinatal outcomes in a setting with universal access to essential health care services. *Canadian Medical Association Journal* 2007; 177(6): 583-90.
5. Luo Z-C, Wilkins R, Heaman M, et al. Birth outcomes and infant mortality among First Nations, Inuit, and non-Indigenous women by northern versus southern residence, Quebec. *Journal of Epidemiology and Community Health* 2012; 66: 328-33.
6. Public Health Agency of Canada. *Perinatal Health Indicators for Canada 2013: A Report of the Canadian Perinatal Surveillance System*. Available at: http://publications.gc.ca/collections/collection_2014/aspc-phac/HP7-1-2013-eng.pdf: 2013
7. Statistics Canada. *1996 Canadian Birth-Census Cohort Userguide*. Available on request from Health Analysis Division.
8. Statistics Canada. *2006 Census 2B Questionnaire*. Available at: http://www23.statcan.gc.ca/imdb-bmdi/pub/instrument/3901_Q2_V3-eng.pdf.
9. Statistics Canada. *2006 Census Technical Report: Coverage* (Catalogue 92-567-X2006001). Available at: <http://www12statcan.gc.ca/census-recensement/2006/ref/rp-guides/rp/coverage-couverture/pdf/92-567-X2006001-eng.pdf>.
10. Statistics Canada. *2006 Canadian Birth-Census Cohort User Guide*. Available on request from Health Analysis Division.
11. Korn EL, Graubard BI. Confidence intervals for proportions with small expected number of positive counts estimated from survey data. *Survey Methodology* 1998; 24(2): 193-201.
12. Statistics Canada. *2006 Census Technical Report: Sampling and Weighting* (Catalogue 92-568-X2006001). Available at: <http://www12statcan.gc.ca/census-recensement/2006/ref/rp-guides/rp/sw-ep/pdf/92-568-X2006001-eng.pdf>.
13. Public Health Agency of Canada. *Perinatal Health Report 2008*. Available at: <http://www.phac-aspc.gc.ca/publicat/2008/cphr-rspsc/pdf/cphr-rspsc08-eng.pdf>: 2008.
14. Kramer MS, Platt RW, Wen SW, et al. A new and improved population-based Canadian reference for birth weight for gestational age. *Pediatrics* 2001; 108(2): E35.
15. Smylie J, Fell D, Ohlsson A, Joint Working Group on First Nations, Indian, Inuit and Métis Infant Mortality of the Canadian Perinatal Surveillance System. A review of aboriginal infant mortality rates in Canada: Striking and persistent aboriginal/non-aboriginal inequities. *Canadian Journal of Public Health* 2010; 101(2): 143-8.
16. Shah PS, Zao J, Al-Wassia H, Shah V. Pregnancy and neonatal outcomes of aboriginal women: A systematic review and meta-analysis. *Women's Health Issues* 2011; 21(1): 28-39.
17. Kierans W, Luo Z-C, Taylor-Clapp S, et al. *Infant Macrosomia among First Nations in British Columbia: Prevalence, Trends and Characteristics*. Victoria, British Columbia: British Columbia Vital Statistics Agency, Ministry of Health, 2007.
18. Arntzen A, Samuelsen SO, Bakketeig LS, Stoltenberg C. Socioeconomic status and risk of infant death. A population-based study of trends in Norway, 1967-1998. *International Journal of Epidemiology* 2004; 33: 279-88.
19. Blumenshine P, Egerter S, Barclay CJ, et al. Socioeconomic disparities in adverse birth outcomes. *American Journal of Preventive Medicine* 2010; 39(3): 263-72.
20. Thompson JM, Irgens IM, Rasmussen S, Daltveit AK. Secular trends in socio-economic status and the implications for preterm birth. *Paediatric and Perinatal Epidemiology* 2006; 20(3): 182-7.
21. Health-Environment Expert Panel. *Health Environment Linkages Expert Panel Report to Statistics Canada. Environment Accounts and Statistics Division, December 19, 2008*. Ottawa: Statistics Canada, 2008.
22. Statistics Canada. *Canadian Environmental Sustainability Indicators: Air Quality Indicators: Data Sources and Methods, 2007* (Catalogue 16-254-x2008000). Available at: <http://www.statcan.gc.ca/pub/16-254-x/16-254-x2008000-eng.pdf>: 2008.