


Health Reports

Development of a population-based microsimulation model of body mass index

by Deirdre Hennessy, Rochelle Garner, William M. Flanagan,
Ron Wall and Claude Nadeau

Release date: June 21, 2017



Statistics
Canada

Statistique
Canada

Canada

How to obtain more information

For information about this product or the wide range of services and data available from Statistics Canada, visit our website, www.statcan.gc.ca.

You can also contact us by

email at STATCAN.infostats-infostats.STATCAN@canada.ca

telephone, from Monday to Friday, 8:30 a.m. to 4:30 p.m., at the following toll-free numbers:

- Statistical Information Service 1-800-263-1136
- National telecommunications device for the hearing impaired 1-800-363-7629
- Fax line 1-877-287-4369

Depository Services Program

- Inquiries line 1-800-635-7943
- Fax line 1-800-565-7757

Standards of service to the public

Statistics Canada is committed to serving its clients in a prompt, reliable and courteous manner. To this end, Statistics Canada has developed standards of service that its employees observe. To obtain a copy of these service standards, please contact Statistics Canada toll-free at 1-800-263-1136. The service standards are also published on www.statcan.gc.ca under “Contact us” > “Standards of service to the public.”

Note of appreciation

Canada owes the success of its statistical system to a long-standing partnership between Statistics Canada, the citizens of Canada, its businesses, governments and other institutions. Accurate and timely statistical information could not be produced without their continued co-operation and goodwill.

Standard Table symbols

The following symbols are used in Statistics Canada publications:

- . not available for any reference period
- .. not available for a specific reference period
- ... not applicable
- 0 true zero or a value rounded to zero
- 0^s value rounded to 0 (zero) where there is a meaningful distinction between true zero and the value that was rounded
- ^P preliminary
- ^r revised
- X suppressed to meet the confidentiality requirements of the *Statistics Act*
- ^E use with caution
- F too unreliable to be published
- * significantly different from reference category ($p < 0.05$)

Published by authority of the Minister responsible for Statistics Canada

© Minister of Industry, 2017

All rights reserved. Use of this publication is governed by the Statistics Canada [Open Licence Agreement](#).

An HTML version is also available.

Cette publication est aussi disponible en français.

Development of a population-based microsimulation model of body mass index

by Deirdre Hennessy, Rochelle Garner, William M. Flanagan, Ron Wall and Claude Nadeau

Abstract

Background: The increasing prevalence of overweight and obesity has necessitated the development of body mass index (BMI) projection models such as the POpulation HEalth Model (POHEM). This study describes the POHEM-BMI model, a microsimulation tool that can be used to support evidence-based health policy making for obesity reduction.

Methods: The National Population Health Survey, the Canadian Community Health Survey (CCHS), and the Canadian Health Measures Survey (CHMS) were used to develop and validate a predictive model of BMI for adults and childhood BMI history. Models were incorporated into POHEM and used to transition BMI over time in a fully dynamic simulated Canadian population.

Results: POHEM-BMI projections of self-reported and measured adult BMI and childhood BMI history agreed well with CCHS and CHMS validation estimates. Among men and women, average BMI is projected to increase by more than one BMI unit between 2001 and 2030. Projections of self-reported BMI show that 59% of the adult population will be overweight or obese by 2030; projections of measured BMI show that the percentage will be 66%.

Interpretation: Using empirically developed BMI prediction models for adults and childhood BMI history integrated into the POHEM framework, validated projections of BMI for the Canadian population can be produced. Projections of BMI trends could have important applications in tracking the prevalence of related diseases and in planning and comparing intervention strategies.

Keywords: Health projections, obesity, overweight, physical activity, POHEM, population surveillance

In both the developed and the developing world, obesity has increased, with global prevalence more than doubling since 1980 to 13%.¹⁻⁵ This dramatic change has popularly been labelled the “obesity epidemic.”

Obesity is a risk factor for several diseases, including diabetes, osteoarthritis, cardiovascular disease, kidney diseases, and certain cancers. As the prevalence of obesity rises, so will the prevalence of these conditions,⁶⁻⁹ which are associated with substantial morbidity, increased risk of mortality, and greater economic costs to individuals and society.¹⁰

As a consequence, obesity surveillance has become an international priority.¹¹ Halting the rising prevalence of obesity has been established as a global goal by the World Health Organization.¹¹ To understand the implications for population health, some countries have constructed models to project body mass index (BMI) and obesity trends over time.^{12,13}

Microsimulation modelling is particularly useful for studying BMI trends, as it can simultaneously account for population dynamics such as aging, migration, and mortality. As well, the longitudinal framework of such models allows BMI to evolve over the life course of simulated individuals, to interact with factors such as physical activity, and to contribute to the risk of multiple diseases.¹⁴ While typically used to evaluate programs such as taxation and pension policy,¹⁵ microsimulation modelling is increasingly used in the health arena.¹⁶⁻¹⁹ It offers a method of comparing the consequences of health policy changes in advance of implementation.

The POpulation HEalth Model (POHEM), developed at Statistics Canada,^{14,20,21} has been used to quantify the health and economic burden of osteoarthritis,²²⁻²⁴ to investigate trends in risk factors for cardiovascular diseases,²⁵ and to project trends in

BMI and physical activity.^{21,26} POHEM-BMI was a collaboration between the Public Health Agency of Canada and Statistics Canada.

This article describes the development of BMI prediction models for adults and of childhood BMI history, and compares projected BMI estimates with estimates from nationally representative survey data to establish validity. Longer-term projections of adult BMI are presented to demonstrate the utility of POHEM-BMI for tracking future risk factor and disease trends to support evidence-based policy making.

POpulation HEalth Model: Overview

POHEM is a continuous-time, Monte Carlo microsimulation tool in which the basic unit of analysis is the individual. POHEM integrates data distributions and equations derived from nationally representative cross-sectional and longitudinal surveys, vital statistics, and disease registries.

Producing projections from POHEM entails several steps.²⁰ The initial population, comprised of adults aged 20 or older, comes from the cross-sectional 2001 Canadian Community Health Survey (CCHS). Initial values for BMI and other characteristics are updated periodically, based on predictive algorithms and transition matrices. As individuals age and their risk factors are updated, their risk of disease changes. Simultaneously, as individuals age and die, the population demographics change, based on observed and projected demographic data.²⁷ The modelled estimates of risk factor or disease prevalence are validated,²⁸ and if necessary, calibrated. Finally, baseline and counterfactual projections for risk factor or disease prevalence are produced.

POpulation HEalth Model: BMI modules

In the adult model, the initial BMI values come from the 2001 CCHS. Subsequent values are determined by multivariate BMI prediction models constructed using the longitudinal National Population Health Survey (NPHS).²⁹ Both self-reported and measured BMI can be projected. Measured BMI is derived from self-reported BMI using a conversion algorithm. Within POHEM-BMI, self-reported BMI is a predictor of multiple diseases, health status, and mortality. As well, it is involved in estimating other risk factors, including physical activity, cholesterol, and hypertension. Figure 1 displays the predictors in the adult BMI model and the outcomes and other risk factors for which BMI is used as a predictor. The BMI and physical activity modules have an interacting relationship—physical activity is used as a predictor of BMI, which, in turn, predicts physical activity. The POHEM-Physical Activity model has been described elsewhere.³⁰

POHEM-BMI does not simulate children. However, for actors born into the simulation at age 20, a “BMI history” that shows what their BMI would have been at ages 6 to 19 is created. This BMI history module is stand-alone, in that it allows for static reporting of childhood and adolescent BMI history among 20-year-olds created in any year, but does not interact with other parts of the model. The childhood BMI history model was developed and validated using the 2004 CCHS, which collected height and weight measurements for children aged 2 or older.

Development of a predictive BMI model for adults

Data source


The longitudinal component of the NPHS followed a group of respondents randomly selected in 1994/1995 to be representative of the Canadian household

population living in the 10 provinces. Residents of Indian Reserves and Crown lands, health care institutions and some remote areas, and full-time members of the Canadian Forces were excluded. The initial sample of 17,276 respondents was re-interviewed every second year. The design, sample, and interview procedures of the NPHS are described elsewhere.²⁹⁻³¹

Data from the first seven NPHS cycles (1994/1995 through 2006/2007) were used to derive the predictive model of self-reported BMI change for adults. The longitudinal data showed how sociodemographic and health behaviour characteristics contributed to the evolution of self-reported BMI over time.

Weight, and therefore BMI, tends to be under-reported in surveys.³² To address this limitation, an algorithm was developed to convert self-reported to measured BMI, using data from the 2004 CCHS, which collected both self-reported and measured height and weight for a subsample of approximately 5,000 individuals.

Figure 1
Adult body mass index (BMI) predictors, outcomes and other risk factors with BMI as a predictor in POpulation HEalth Model (POHEM) framework



Source: POpulation HEalth Model.

Validation data sources

Validation involved comparisons of POHEM-BMI projections with estimates from four CCHS cycles between 2001 and 2014. The CCHS,^{33,34} a nationally representative cross-sectional survey, has collected self-reported information on BMI, other lifestyle risk factors and chronic diseases at regular intervals since 2001. It covers the non-institutionalized household population aged 12 or older in all provinces and territories, except

residents of Indian Reserves or Crown lands, institutions and certain remote areas, and full-time members of the Canadian Forces.

Projections of measured BMI were validated against estimates from the Canadian Health Measures Survey (CHMS),³⁵⁻³⁹ a direct health measures survey. Between March 2007 and December 2013, three CHMS cycles were completed at 49 sites across Canada. The survey provides nationally representative estimates from a sample of Canadians aged 6 to 79 living in private households. The CHMS excludes residents of Indian Reserves or Crown lands, institutions and certain remote regions, and full-time members of the Canadian Forces.

tion of self-reported BMI, education (less than secondary graduation, secondary graduation, some postsecondary, postsecondary graduation), and age group.

Adjustment for calendar time: When new individuals are born into the simulation at age 20, they “inherit” an initial BMI value from an individual in the starting population (CCHS 2001). Because it may be unrealistic for 20-year-olds in future years to have the same BMI distribution as 20-year-olds in 2001, the ability to alter BMI over time is programmed. Additional analysis of the CCHS from 2001 to 2009 was undertaken to construct an algorithm that adjusts BMI for calendar time separately for males and females. This feature can be switched on or off in POHEM-BMI as needed. The current results were generated without adjusting for calendar time.

What is already known on this subject?

- The global prevalence of obesity has more than doubled since 1980.
- Obesity is an important risk factor for several diseases, including diabetes, osteoarthritis, cardiovascular disease, kidney disease, and certain cancers.
- As a consequence, obesity surveillance has become an international priority.
- To understand implications for population health, some countries have constructed models to project body mass index (BMI) and obesity trends over time.

What does this study add?

- The POPELation HEalth Model (POHEM) is a microsimulation tool developed at Statistics Canada; POHEM-BMI projects trends in body mass index.
- POHEM-BMI projections of self-reported and measured BMI agree well with the survey estimates, especially for the percentage obese.
- By 2030, approximately 59% of Canadian adults will be overweight/obese according to POHEM projections of self-reported BMI; according to POHEM projections of measured BMI, 66% of adults will be overweight/obese.

Modelling techniques and model selection

BMI predictive model: Sex-specific, auto-regressive models were constructed to predict adult BMI. Previous BMI values were the main explanatory variables; respondents could have up to four previous BMI measurements from the longitudinal data (NPHS). Depending on the autoregressive order of the model (from 1 to 4), a different set of coefficients for the covariates was used. For illustrative purposes, Appendix Table A shows the coefficients for the first order model. Other covariates were age, physical activity (leisure-time physical activity; biking and walking for transportation and errands; usual-day physical activity), and smoking. As well as smoking and physical activity, variables that captured the individual’s longitudinal trend (downward or upward) in these behaviours were included to more precisely reflect exposure to these risks over time. The BMI predictive model initially included income, education, and geography as potential covariates, but they did not confer additional predictive power, and so were dropped during model selection.

Conversion model: Self-reported BMI was converted to measured BMI using the equation and coefficients in Appendix Table B. In this algorithm, measured BMI is modelled as a func-

Development of a predictive model of childhood BMI history

The predictive model of childhood BMI history is based on measured data from the 2004 CCHS.^{33,34} The same data source was used to validate the POHEM-BMI output.

To predict childhood BMI history, sex-specific growth curve modelling techniques were used. The model of BMI distribution covers 6- to 19-year-olds and does not include any covariates besides 4 cubic spline functions of age (details about this model are available on request). Although the model was built using cross-sectional data, a sequence of plausible BMI values by age that exhibited a similar correlation structure to that in longitudinal data was sought. Additional analysis using the NPHS revealed that the auto-correlation structure of children’s BMI over time resembled that of an autoregressive moving average (ARMA) model, which meant it exhibited slow exponential decay after a sharper initial decline. Using the ARMA model, correlated quantiles of BMI for children were created.

Integrating predictive models into the POHEM environment

The BMI prediction models were integrated into the POHEM framework by implementing the algorithms in Modgen code.⁴⁰ Pieces of code that control the population dynamics, risk factors and disease status updates are written as separate modules and compiled into a Microsoft Visual Studio solution that generates an executable file. The POHEM executable uses the parameters, including population counts, mortality rates and equation coefficients, to run the simulation and generate output.

The outputs are a set of pre-programmed tables that show average projected BMI, projected distribution of BMI categories (underweight, normal weight, overweight, and obese), and the distribution of adult weight categories given child weight category. The tables can be customized for any outcome. Because of the modular format, newly developed pieces of code can be integrated with the existing program. Once modules are implemented and validated,

additional functionality can be added. For instance, POHEM can perform counterfactual analyses by changing model inputs and then projecting alternative distributions of the outcome of interest.

A set of intervention parameters allows for changes (reductions or increases) to the BMI of the population. These interventions can be targeted by various population characteristics, including year, age group, sex, BMI category, and disease risk. With this facility, it is possible to conduct comparative analysis of different interventions.

Validation framework

Kopec et al. have detailed recommendations for validating disease simulation models,²⁸ the most pertinent of which for this study relate to parameter quality, computer implementation of the model, and evidence of model performance. The BMI prediction model parameters were derived from statistical analyses of population-based surveys. These data sources (NPHS, CCHS and CHMS) and the methods used to analyse them are well-documented.^{29-31,33-39} POHEM-BMI


was implemented in Modgen code,⁴⁰ a computer language developed for microsimulation, which has been used to program a number of other models at Statistics Canada and internationally. Evidence for model performance was gathered by comparing the projected outputs to input data (used to develop the models) and to external data (not used to develop the models).

Statistical analysis

The POHEM-BMI projections and validation estimates from the CCHS (various cycles) and CHMS (cycles 1 through 3) were age-standardized to the 2001 adult (20 or older) population in Figure 2 to allow comparisons over time. The data in Figures 3, 4 and 5 are unstandardized. World Health Organization cut-offs for BMI were used to classify adults and children into BMI categories.^{41,42} Variance estimates that account for design effects were calculated using the bootstrap technique for survey-based estimates. Analyses were conducted using SAS 9.3 and Stata 11.

Figure 2
Prevalence of self-reported and measured overweight and obesity based on POpulation HHealth Model (POHEM) and survey data, population aged 20 or older, Canada, 2001 to 2014

percent in BMI category



I = 95% confidence interval

Sources: POHEM projections for 2001 through 2014; Canadian Community Health Survey (self-reported); Canadian Health Measures Survey (measured).

Results

Adult BMI


The predictors (and coefficients) used to model adult BMI are described in Figure 1 and Appendix Table A.

After integration of the models into the POHEM framework, projections of self-reported and measured BMI (Figure 2) were validated by comparing simulation outputs to estimates from the CCHS and CHMS, respectively. The

projections and estimates agree well, especially for the percentage obese.


POHEM-BMI projections of average self-reported BMI and the distribution of self-reported BMI categories from 2001 to 2030 are shown in Figure 3 for men

Figure 3
Population Health Model (POHEM) projections of self-reported body mass index (BMI) category and average BMI, men aged 20 or older, Canada, 2001 to 2030



Source: POHEM projections for 2001 through 2030.


Figure 4
Population Health Model (POHEM) projections of self-reported body mass index (BMI) category and average BMI, women aged 20 or older, Canada, 2001 to 2030



Source: POHEM projections for 2001 through 2030.

Figure 5
POHEM Health Model (POHEM) projected distribution of childhood body mass index (BMI) categories of 20-year-olds in 2015, compared with distribution of BMI categories of 9-year-olds in 2004 Canadian Community Health Survey (CCHS)

percent in BMI category



I = 95% confidence interval

Sources: POHEM projections for 2015; 2004 Canadian Community Health Survey.

and in Figure 4 for women. For both sexes, average self-reported BMI is projected to rise by more than one BMI unit between 2001 and 2030. The percentage overweight is projected to remain relatively stable, while the percentage obese increases among both sexes. According to POHEM projections of *self-reported* BMI, about 59% of the adult population will be overweight/obese by 2030. Men have a higher initial average self-reported BMI than do women, and a greater percentage are projected to become overweight/obese. Also, the increase in obesity among men is projected to be steeper than that among women. Trends are similar in projections of *measured* BMI, but the total projected prevalence of overweight/obesity is higher, reaching 66% of the adult population by 2030 (data not shown).

History of childhood BMI

Modelled childhood BMI history was validated by comparing POHEM-BMI outputs with data from the 2004 CCHS. The projected childhood BMI history of 20-year-olds in 2015 (who had been 9 years old in 2004) was compared with that of 9-year-olds surveyed by the CCHS in 2004. Figure 5 shows close agreement

between the POHEM-BMI projections and CCHS estimates of the percentage of 9-year-olds who were normal weight, overweight or obese.

Discussion

Based on empirically developed BMI prediction models, by 2030, approximately 59% of Canadian adults will be overweight/obese according to POHEM projections of self-reported BMI (66% according to POHEM projections of measured BMI). These projected increases are comparable to those of the Foresight microsimulation model, which has been used for the United Kingdom (UK), the United States, and other countries,^{10,12,43} and to statistical projections (non-microsimulation) for Australia.¹³ A greater increase in the prevalence of obesity among men than women was also noted for Russia and Poland by the Foresight researchers.^{44,45} However, the increase in the prevalence of overweight/obesity projected by POHEM-BMI is not as steep as the Foresight models, which predict levels as high as 72% for the UK population by 2035,⁹ and 80% for the Irish population by 2030.⁴⁶ Similar results were reported by Walls et al.¹³

for Australia, where 72% of the adult population is projected to be overweight/obese by 2025.

The slower rise in overweight/obesity in Canada has also been shown in analyses by the Organization for Economic Co-operation and Development.⁴⁷ However, it is important to consider how POHEM-BMI differs from other models. POHEM-BMI was constructed with longitudinal self-reported BMI data, whereas Foresight was originally developed using cross-sectional measured BMI data from the UK Health Survey for England.^{10,12,43} Self-reported data consistently underestimate measured BMI.³² Although measured BMI data would be the ideal input for POHEM, direct measures are costly to collect. To address this limitation, POHEM-BMI contains a conversion algorithm that allows projections of both self-reported and measured BMI.

The models also differ in their use of covariates: POHEM-BMI includes age, sex, smoking and physical activity as predictors, while Foresight generated cross-sectional BMI parameters for the UK by age, sex, ethnicity, social class, and geographical region.¹² As well, BMI and physical activity interact

in POHEM-BMI. Although Foresight projects BMI by covariates, it is not a multivariate predictive model in the same way as POHEM-BMI. POHEM-BMI predicts BMI using the covariates, and the simulated value of BMI is then used as a predictor of other risk factors and diseases (Figure 1). This method of modelling allows for considerable heterogeneity in risk profiles among individuals,⁴⁸ so that the life course trajectories of individuals vary in ways that plausibly represent observed data. Also, the multivariate nature of POHEM-BMI offers many intervention points. For instance, co-occurring interventions that target both BMI and physical activity could be implemented, as well as interventions that tackle individual risk factors or diseases.

Notwithstanding the differences between them, the models have similar applications. In population health, policy and program evaluation has generally been undertaken after implementation. Microsimulation projections offer a way to measure the future burden of a risk factor or disease, accounting for aging and other dynamic population changes. Such analysis can determine if risk factor or disease prevalence is increasing, decreasing, or stabilizing. As well, quan-

tification of the extent of a problem (in this case, the prevalence of overweight and obesity) is, itself, worthwhile.

Information about the future incidence and prevalence of risk factors and disease by age, sex and sociodemographic characteristics is helpful for deploying public health and prevention resources. Using POHEM-BMI or similar models to project an intervention's potential impacts on the prevalence of obesity and obesity-related diseases can provide evidence that the benefits outweigh the costs.

Limitations

POHEM-BMI should be considered in the context of certain limitations.

First, the estimates and projections produced by POHEM-BMI are only as good as the input data. Survey data are subject to potential bias resulting from sampling error, incomplete coverage, non-response, and measurement error. Despite Statistics Canada's efforts to identify and minimize such biases, they may still exist.

Second, as a measure of overweight and obesity, BMI itself is a limitation. BMI cannot account for differences in body composition. The standard cut-offs

used to define overweight and obesity may not be appropriate for all ethnicities.⁴⁹ In addition, POHEM's BMI predictive models do not consider certain important causal determinants of weight gain, such as diet quality and neighbourhood walkability.

Conclusions

Using POHEM-BMI, it is possible to produce validated projections of overweight and obesity for the Canadian population. Such projections could have important applications for surveillance of risk factor and disease prevalence, as well as for planning and comparative analysis of intervention strategies.

Acknowledgements

The Science Integration Division of the Public Health Agency of Canada provided financial support for the development of the body mass index module of the Population Health Model (POHEM). The authors thank all researchers who have used POHEM in their work and have helped refine the model over time. ■

References

- World Health Organization. *Fact Sheet on Obesity and Overweight*. Available at: <http://www.who.int/topics/obesity/en>
- Finucane MM, Stevens GA, Cowan MJ, et al. National, regional, and global trends in body-mass index since 1980: Systematic analysis of health examination surveys and epidemiological studies with 960 country-years and 9.1 million participants. *Lancet* 2011; 377: 557-67.
- NCD Risk Factor Collaboration. Trends in adult body-mass index in 200 countries from 1975 to 2014: A pooled analysis of 1698 population-based measurement studies with 19.2 million participants. *Lancet* 2011; 387: 1377-96.
- Ng M, Fleming T, Robinson M, et al. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980–2013: A systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2014; 384: 766-81.
- Stevens GA, Singh GM, Lu Y, et al. National, regional, and global trends in adult overweight and obesity prevalences. *Population Health Metrics* 2012; 10: 22.
- The Emerging Risk Factors Collaboration. Separate and combined associations of body-mass index and abdominal adiposity with cardiovascular disease: Collaborative analysis of 58 prospective studies. *Lancet* 2011; 377: 1085-95.
- The Global Burden of Metabolic Risk Factors for Chronic Diseases Collaboration (BMI Mediated Effects). Metabolic mediators of the effects of body-mass index, overweight, and obesity on coronary heart disease and stroke: A pooled analysis of 97 prospective cohorts with 1.8 million participants. *Lancet* 2014; 383: 970-83.
- Renahan AG, Tyson M, Egger M, et al. Body-mass index and incidence of cancer: A systematic review and meta-analysis of prospective observational studies. *Lancet* 2008; 371: 569-78.
- Bhimjiyani A, Knuchel-Takano A, Jaccard A, et al. Tipping the scales: Why preventing obesity makes economic sense. 2015. Available at: http://www.cancerresearchuk.org/sites/default/files/tipping_the_scales_-_cruk_full_report11.pdf
- Wang YC, McPherson K, Marsh T, et al. Health and economic burden of the projected obesity trends in the USA and the UK. *Lancet* 2011; 378: 815-25.
- World Health Organization. *Global Action Plan for the Prevention and Control of Noncommunicable Diseases 2013–2020*. Geneva, Switzerland: World Health Organization, 2013.
- Government of the United Kingdom. *Foresight Tackling Obesities: Future Choices*. Government Office for Science, 2007. Available at: <https://www.gov.uk/government/collections/tackling-obesities-future-choices>. Accessed July 30, 2016.
- Walls HL, Magliano DJ, Stevenson CE, et al. Projected progression of the prevalence of obesity in Australia. *Obesity* 2012; 20, 872-8.
- Wolfson MC. POHEM—A framework for understanding and modelling the health of human populations. *World Health Statistics Quarterly* 1994; 47(3-4): 157-76.
- Spielauer M. What is dynamic social science microsimulation? Ottawa: Modelling Division, Statistics Canada, 2007.
- Zucchelli E, Jones A, Rice N. The evaluation of health policies through dynamic microsimulation methods. *International Journal of Microsimulation* 2012; 5: 2-20.
- Fone D, Hollinghurst S, Temple M, et al. Systematic review of the use and value of computer simulation modelling in population health and health care delivery. *Journal of Public Health Medicine* 2003; 25(4): 325-35.
- Rutter CM, Zaslavsky AM, Feuer EJ. Dynamic microsimulation models for health outcomes: A review. *Medical Decision Making* 2011; 31: 10-8.
- Glied S, Tilipman N. Simulation modeling of health care policy. *Annual Review of Public Health* 2010; 31: 439-55.
- Hennessy DA, Flanagan WM, Tanuseputro P, et al. The Population Health Model (POHEM): An overview of rationale, methods and applications. *BMC Population Health Metrics* 2015; 13: 24.
- Bancej C, Jayabalasingham B, Wall RW, et al. Trends and projections of obesity among Canadians. *Health Promotion and Chronic Disease Prevention in Canada: Research, Policy and Practice* 2015; 35(7): 109-12.
- Kopec JA, Sayre EC, Flanagan WM, et al. Development of a population-based microsimulation model of osteoarthritis in Canada. *Osteoarthritis and Cartilage* 2010; 18: 303-11.
- Sharif B, Kopec J, Bansback N, et al. Projecting the direct cost burden of osteoarthritis in Canada using a microsimulation model. *Osteoarthritis and Cartilage* 2015; 23(10): 1654-63.
- Sharif B, Garner R, Hennessy D, et al. Productivity costs of work loss associated with osteoarthritis in Canada from 2010 to 2031. *Osteoarthritis and Cartilage* 2017; 25(2): 249-58. doi: 10.1016/j.joca.2016.09.011.
- Manuel DG, Tuna M, Hennessy D, et al. Modelled projections of preventable risks for cardiovascular disease in Canada to 2021. *Canadian Medical Association Journal Open* 2014; 2: 2.
- Nadeau C, Wong SL, Flanagan W, et al. Development of a population-based microsimulation model of physical activity in Canada. *Health Reports* 2013; 24(10): 11-9.
- Bohnert N, Chagnon J, Coulombe S, et al. *Population Projections for Canada (2013 to 2063), Provinces and Territories (2013 to 2038): Technical Report on Methodology and Assumptions* (Catalogue 91-620-X) Ottawa: Statistics Canada, 2015.
- Kopec JA, Finès P, Manuel DG, et al. Validation of population-based disease simulation models: A review of concepts and methods, *BMC Public Health* 2010; 10: 710.
- Statistics Canada. *National Population Health Survey*. Available at: <http://www.statcan.gc.ca/nphs>. Accessed March 24, 2011.
- Swain L, Catlin G, Beaudet MP. The National Population Health Survey—Its longitudinal nature. *Health Reports* 1999; 10(4): 69-82.
- Tambay J-L, Catlin G. Sample design of the National Population Health Survey. *Health Reports* 1995; 7(1): 29-38.
- Connor Gorber S, Tremblay M, Moher D, Gorber B. A comparison of direct vs. self-report measures for assessing height, weight and body mass index: A systematic review. *Obesity Reviews* 2007; 8: 307-26.
- Statistics Canada. *Canadian Community Health Survey*. Available at: <http://www.statcan.gc.ca/cchs>. Accessed March 24, 2011.
- Béland Y. Canadian Community Health Survey—Methodological overview. *Health Reports* 2002; 13(3): 9-14.
- Tremblay MS, Connor Gorber S. Canadian health measures survey: Brief overview. *Canadian Journal of Public Health* 2007; 98(6): 453-6.
- Day B, Langlois R, Tremblay M, Knoppers BM. Canadian Health Measures Survey: ethical, legal and social issues. *Health Reports* 2007; 18(Suppl): 37-51.
- Tremblay M, Langlois R, Bryan S, et al. Canadian Health Measures Survey pre-test: Design, methods, results. *Health Reports* 2007; 18(Suppl): 21-30.
- Tremblay M, Wolfson M, Connor Gorber S. Canadian Health Measures Survey: Rationale, background and overview. *Health Reports* 2007; 18(Suppl): 7-20.

39. Giroux S, Labrecque F, Quigley A. *Sampling Documentation for Cycle 2 of the Canadian Health Measures Survey*. Methodology Branch Working Paper 002. Ottawa: Statistics Canada, 2013.
40. Statistics Canada. *Introduction to Modgen*. Available at: <http://www.statcan.gc.ca/eng/microsimulation/modgen/new/introduction/introduction>
41. WHO Expert Committee. *Physical Status: The Use and Interpretation of Anthropometry* (WHO Technical Report Series No. 854) Geneva: World Health Organization, 1995.
42. WHO Multicentre Growth Reference Study Group. *WHO Child Growth Standards: Length/Height-for-Age, Weight-for-Age, Weight-for-Length, Weight-for-Height and Body Mass Index-for-Age: Methods and Development*. Geneva: World Health Organization, 2006.
43. Webber L, Divajeva D, Marsh T, et al. The future burden of obesity related disease in the 53 WHO European-Region countries and the impact of effect interventions: A modelling study. *British Medical Journal Open* 2014; 4: e004787.
44. Rtveldzde K, Marsh T, Webber L, et al. Obesity trends in Russia. The impact on health and healthcare costs. *Science Research: Health* 2012; 4(12a): 1471-84.
45. Webber L, Halicka E, Marsh T, et al. Projected incidence of overweight and obesity and related disease incidence across Poland. *Central European Journal of Public Health* 2014; 22(1): 17-23.
46. Keaver L, Webber L, Dee A, et al. Application of the UK Foresight Obesity Model in Ireland: The health and economic consequences of projected obesity trends in Ireland. *PLoS One* 2013; 8(11): e79827.
47. Organization for Economic Cooperation and Development. *Obesity Update 2014*. Available at: <http://www.oecd.org/els/health-systems/Obesity-Update-2014.pdf>
48. Manuel DG, Rosella L, Hennessy D, et al. Predictive risk algorithms in a population setting: An overview. *Journal of Epidemiology and Community Health* 2012; 66(10): 859-65.
49. Chiu M, Austin PC, Manuel DG, et al. Deriving ethnic-specific BMI cutoff points for assessing diabetes risk. *Diabetes Care* 2011; 34(8): 1741-8.

Appendix

Table A
Coefficients for model of self-reported adult (age 20 or older) body mass index value, by sex and for auto-regressive model order 1, derived from data from the National Population Health Survey 1994/1995 through 2006/2007

Terms	Men	Women
Intercept	0.198	0.169
Age group		
20 to 29	0.032	0.027
30 to 39	0.027	0.025
40 to 49	0.026	0.026
50 to 59	0.024	0.026
60 to 69	0.018	0.024
70 to 79	0.010	0.011
80 or older	0	0
Previous BMI value(s), log		
At time t-2 years	0.933	0.940
At time t-4 years	0	0
At time t-6 years	0	0
At time t-8 years	0	0
Leisure-time physical activity (minutes per day)		
None	0.003	0.005
Less than 30	0.004	0.008
30 to less than 60	0.003	0.004
60 or more	0	0
Cycling for transportation or errands		
No	0	0
Yes	-0.005	-0.001
Longitudinal trend in walking for transportation or errands		
None	0	0
Downward	0.003	0.0001
Upward	0.0008	0.0009
Longitudinal trend in usual daily physical activity		
None	0	0
Downward	0.003	0.003
Upward	0.0004	-0.0006
Smoking status		
Never smoker	0	0
Former smoker	0.003	0.003
Current smoker	-0.002	-0.004
Longitudinal trend in smoking		
None	0	0
Downward	0.007	0.021
Upward	-0.003	-0.004

Note: The BMI prediction model for adult men and women is: $\log(\text{BMI}) = \alpha + \beta_A \text{Age} + \beta_H \log(\text{Previous BMI values}) + \beta_{PA} \text{Physical Activity} + \beta_S \text{Smoking} + \sigma \epsilon$

Sources: 1994/1995 through 2006/2007 National Population Health Survey.

Table B
Coefficients for model to convert self-reported adult (20 or older) body mass index (BMI) value to measured BMI value, by sex, derived from subsample data from the 2005/2006 Canadian Community Health Survey

Terms (x)	Mean (μ)		Variance (τ)	
	Men	Women	Men	Women
BMI based on self-reported height and weight	1.038	1.080	0.037	0.090
BMI category based on self-reported height and weight				
Underweight	0.276	0.174
Normal weight	-0.638	-0.097
Overweight	-0.322	-0.203
Obese	-0.608	-0.142
Age group				
20 to 24	-0.634	-0.756
25 to 34	-0.303	-0.544
35 to 44	-0.538	-0.482
45 to 54	-0.370	-0.234
55 to 64	0.067	-0.176
65 to 74	0.475	-0.024
75 or older	0	0
Educational attainment				
Less than secondary graduation	0.569	-0.016	0.122	-0.822
Secondary graduation	0.616	-0.662	-0.227	-1.405
Some postsecondary	0.644	-0.533	0.467	-1.353
Postsecondary graduation	0.778	-0.357	-0.023	-1.390

... not applicable

Note: The BMI conversion model for adults is: $BMI^{\text{measured}} | x, z \sim \text{logistic}(\mu = \beta'x(\text{Self-reported BMI, Age group, Education level}), \tau)$, where $\text{logistic}(\mu, \tau)$ refers to logistic distribution with mean μ and variance τ .

Source: 2005/2006 Canadian Community Health Survey.