

2710
c.1

DFO - Library / MPO - Bibliothèque




12063322

Central Coast Juvenile Herring Survey, August 2003

S. Henderson, M. Thompson, and T.W. Therriault

Fisheries and Oceans Canada
Science Branch, Pacific Region
Pacific Biological Station
Nanaimo, British Columbia
V9T 6N7



2005

Canadian Manuscript Report of Fisheries and Aquatic Sciences 2710

SH
223
F55
no. 2710
c.1



Fisheries
and Oceans

Pêches
et Océans

Canada

Canadian Manuscript Report of Fisheries and Aquatic Sciences

Manuscript reports contain scientific and technical information that contributes to existing knowledge but which deals with national or regional problems. Distribution is restricted to institutions or individuals located in particular regions of Canada. However, no restriction is placed on subject matter, and the series reflects the broad interests and policies of the Department of Fisheries and Oceans, namely, fisheries and aquatic sciences.

Manuscript reports may be cited as full publications. The correct citation appears above the abstract of each report. Each report is abstracted in *Aquatic Sciences and Fisheries Abstracts* and indexed in the Department's annual index to scientific and technical publications.

Numbers 1-900 in this series were issued as Manuscript Reports (Biological Series) of the Biological Board of Canada, and subsequent to 1937 when the name of the Board was changed by Act of Parliament, as Manuscript Reports (Biological Series) of the Fisheries Research Board of Canada. Numbers 1426 - 1550 were issued as Department of Fisheries and the Environment, Fisheries and Marine Service Manuscript Reports. The current series name was changed with report number 1551.

Manuscript reports are produced regionally but are numbered nationally. Requests for individual reports will be filled by the issuing establishment listed on the front cover and title page. Out-of-stock reports will be supplied for a fee by commercial agents.

Rapport manuscrit canadien des sciences halieutiques et aquatiques

Les rapports manuscrits contiennent des renseignements scientifiques et techniques qui constituent une contribution aux connaissances actuelles, mais qui traitent de problèmes nationaux ou régionaux. La distribution en est limitée aux organismes et aux personnes de régions particulières du Canada. Il n'y a aucune restriction quant au sujet; de fait, la série reflète la vaste gamme des intérêts et des politiques du ministère des Pêches et des Océans, c'est-à-dire les sciences halieutiques et aquatiques.

Les rapports manuscrits peuvent être cités comme des publications complètes. Le titre exact paraît au-dessus du résumé de chaque rapport. Les rapports manuscrits sont résumés dans la revue *Résumés des sciences aquatiques et halieutiques*, et ils sont classés dans l'index annuel des publications scientifiques et techniques du Ministère.

Les numéros 1 à 900 de cette série ont été publiés à titre de manuscrits (série biologique) de l'Office de biologie du Canada, et après le changement de la désignation de cet organisme par décret du Parlement, en 1937, ont été classés comme manuscrits (série biologique) de l'Office des recherches sur les pêcheries du Canada. Les numéros 901 à 1425 ont été publiés à titre de rapports manuscrits de l'Office des recherches sur les pêcheries du Canada. Les numéros 1426 à 1550 sont parus à titre de rapports manuscrits du Service des pêches et de la mer, ministère des Pêches et de l'Environnement. Le nom actuel de la série a été établi lors de la parution du numéro 1551.

Les rapports manuscrits sont produits à l'échelon régional, mais numérotés à l'échelon national. Les demandes de rapports seront satisfaites par l'établissement auteur dont le nom figure sur la couverture et la page du titre. Les rapports épuisés seront fournis contre rétribution par des agents commerciaux.

Canadian Manuscript Report of
Fisheries and Aquatic Sciences 2710

2005

CENTRAL COAST JUVENILE HERRING SURVEY, AUGUST 2003

by

S. Henderson, M. Thompson, and T.W. Therriault

Fisheries and Oceans Canada
Science Branch, Pacific Region
Pacific Biological Station
Nanaimo, British Columbia
V9T 6N7

© Her Majesty the Queen in Right of Canada, 2005
Cat No. Fs 97-4/2710E ISSN -0706-6473

Correct citation for this publication:

Henderson, S., Thompson, M., and Therriault, T.W. 2005. Central Coast juvenile herring survey, August 2003. Can. Manusc. Rep. Fish. Aquat. Sci. 2710: v + 36 p.

TABLE OF CONTENTS

LIST OF FIGURES.....	iv
LIST OF TABLES.....	iv
ABSTRACT.....	v
RÉSUMÉ.....	v
INTRODUCTION.....	1
METHODS.....	1
Fish Sampling.....	1
Plankton Sampling.....	2
Stomach Contents.....	3
RESULTS.....	3
Herring.....	4
Plankton.....	5
Herring Stomachs.....	5
Other Species Stomachs.....	6
CONCLUSION.....	6
ACKNOWLEDGEMENTS.....	6
REFERENCES.....	7

LIST OF FIGURES

Figure 1. The five major British Columbia herring stock assessment regions.....	7
Figure 2. Northern Central Coast stock assessment area with set locations and section numbers.....	8
Figure 3. Southern Central Coast stock assessment area with set locations and section numbers.....	9
Figure 4. Length frequency distribution of all sampled herring.....	10
Figure 5. Length frequency distributions of sampled herring by set location.....	11
Figure 6. Length-weight relationship of all sampled herring.....	15
Figure 7. Proportion of herring relative to total catch by location.....	16

LIST OF TABLES

Table 1. Details of the 39 sets made during the 2003 Central Coast juvenile herring survey.....	17
Table 2. Species quantity and weights by set.....	18
Table 3. Percent species occurrence for all 2003 Central Coast sets.....	23
Table 4. Mean lengths, weights and total catch weights of age-0+ and age-1+ herring by set location.....	24
Table 5. Categories of organisms, by phylum, identified in plankton and stomach samples.....	25
Table 6. Copepods identified in plankton and stomach samples.....	26
Table 7. Zooplankton density by tow location.....	27
Table 8. Herring stomach fullness and state of content	31
Table 9. Occurrence of identifiable food items in age-0+ herring stomachs, by set location.....	32
Table 10. Occurrence of identifiable food organisms in age-1+ herring stomachs, by set locations.....	33
Table 11. Stomach contents of other sampled species.....	35

ABSTRACT

Henderson, S., Thompson, M., and Therriault, T.W. 2005. Central Coast juvenile herring survey, August 2003. Can. Manuscr. Rep. Fish. Aquat. Sci. 2710: v + 36 p.

This Central Coast survey was conducted from August 5-17, 2003. Thirty-nine purse seine sets were made in Statistical Management Areas 6, 7, 8 and 9, from Meyers Passage in the north to Fish Egg Inlet in the south. The survey serves to address information gaps on the distribution, abundance, size and feeding habits of juvenile herring in these areas.

Twenty-two species of fish were recorded in the purse seine catches with herring being the most frequently captured. A total of 7225 juvenile herring were measured and weighed resulting in a distinctly bimodal length frequency distribution representing age-0+ and age-1+ fish. Age-0+, age-1+ and age-2+ or older herring occurred in 87.2%, 76.9% and 35.9% of the sets respectively. A total of 224 herring stomachs and 46 possible herring predator stomachs were analysed for stomach content. Twenty oblique plankton tows were performed during the survey. Juvenile calanoid copepods and *Pseudocalanus minutus* were the most prominent organisms encountered.

RÉSUMÉ

Henderson, S., Thompson, M., and Therriault, T.W. 2005. Central Coast juvenile herring survey, August 2003. Can. Manuscr. Rep. Fish. Aquat. Sci. 2710: v + 36 p.

Ce relevé a été réalisé sur la Côte centrale entre le 5 et le 17 août 2003. Nous avons effectué 39 pêches à la senne à poche dans les secteurs statistiques 6, 7, 8 et 9, du passage Meyers au nord au bras Fish Egg au sud. Ce relevé vise à combler les lacunes en ce qui concerne la répartition, l'abondance, la taille et les habitudes alimentaires des jeunes harengs dans ces secteurs.

Vingt-deux espèces ont été observées dans les prises, le hareng venant en première place. Au total, nous avons mesuré et pesé 7 225 harengs juvéniles, et nous avons obtenu une distribution des fréquences de longueur nettement bimodale, représentant les poissons d'âge 0+ et d'âge 1+. Des harengs d'âge 0+, d'âge 1+ et d'âge 2+ ou davantage étaient présents respectivement dans 87,2 %, 76,9 % et 35,9 % des coups de senne. Au total, nous avons analysé le contenu stomacal de 224 harengs et de 46 prédateurs potentiels du hareng. Pendant la mission, nous avons aussi effectué 20 traits en oblique de filet à plancton. Les organismes les plus souvent observés dans ces traits étaient des copépodes calanoïdes juvéniles et le *Pseudocalanus minutus*.

INTRODUCTION

Pacific herring (*Clupea pallasii*) are an important commercial species and a vital forage species for many marine mammals, birds and other fish in British Columbia's coastal waters. Herring spawn principally on marine vegetation in the subtidal and upper intertidal zone between February and June, with peak spawning between March and April (Humphreys and Hourston 1978). Larvae hatch in two to three weeks, and disperse with surface currents, metamorphosing into juvenile herring at a length of ~25mm (Hourston and Haegele 1980). Herring are considered juvenile until they are three years of age and have joined the sexually mature spawning population (Hay and McCarter 1999). During daylight hours, juvenile herring congregate in schools, occasionally forming mixed aggregates with other pelagic species, close to shore near the bottom (Haegele 1997). At dusk, these fish migrate into surface waters to feed on plankton. During this time they are vulnerable to purse seine operations.

Relatively little is known about the distribution, abundance, size and feeding habits of juvenile herring in the Central Coast of British Columbia (Figure 1). In 2002, a juvenile herring survey was designed to address information gaps and learn about the general biology of herring in this area. The survey used an ecosystem based approach to biological sampling. Therefore, in addition to juvenile herring, all other fish species, plankton samples and predator stomach samples were retained for analysis. Juvenile herring stomach contents also were examined from six set locations. This approach will potentially provide a better understanding of the role and relationships juvenile herring have in Central Coast waters, and may provide an empirical forecast of recruitment to the herring roe fishery based on relative juvenile abundance.

METHODS

In 2003, the Central Coast juvenile herring survey was conducted from August 5-17. Thirty-nine sets were made at 12 locations within Statistical Management Areas 6, 7, 8 and 9 (Table 1). The study area extended from Meyers Passage in the north to Fish Egg Inlet in the south (Figures 2 and 3). The 2003 set locations were nearly identical to set locations used in 2002, the first year of this survey (Henderson et al. 2004). The sampling sites originally were chosen based on known historical herring spawning sites, and represent both nearshore and open water habitats (Haegele and Armstrong 2003).

Fish Sampling

The 12m, aluminium-hulled Fisheries Research Vessel *Walker Rock* was used for all fishing events. A 183m long and 27m deep seine net of knotless web, resulting in an area fished of ~2665m², was used for all fishing events. The body of the net had 46m of 22.2mm mesh at the tow end followed by 91m of 19.0mm mesh, and the bunt end was 46m of 9.5mm mesh. The net fished to a depth of 10m, and was able to retain fish greater than 30mm in length. All sets were made after dusk when herring are feeding near the surface. All sets were made "blind" at predetermined set locations. Up to four sets were completed per night, depending on location. No proposed sets were lost due to adverse weather or oceanic (tide) conditions. For most sets, it was possible to land the entire catch for biological sampling. When possible, a minimum of 200

herring from each juvenile age class (age-0+ and age-1) were retained in addition to any older herring. On occasion, it was not practical to land a large set in its entirety, so sub-sampling was necessary. When sub-sampling was necessary, a 40kg tote was filled with randomly selected fish and retained for biological sampling. The remainder of the set was released over the corkline, its size estimated as the number of totes released. The fish were sorted to species immediately after each set, and herring were further sorted to year-class based on visual observation. The number of juvenile herring caught in each set was determined by dividing the total catch weight by the age-specific mean weight of sub-sampled herring from each age class. The number of other species caught was determined in the same manner, with juvenile and adult salmon treated separately to minimize errors when calculating total catch (Table 2). All sorted fish were weighed, bagged and preserved in a 3.7% seawater formalin solution, with the exception of large predator species (e.g. adult salmon, dogfish and mackerel). These fish were individually weighed and measured in the field, and their stomachs were removed and preserved for later analysis at the Pacific Biological Station.

To minimize the effects of fish preservation, all fish samples were processed within eight weeks of collection. From each set, up to 200 herring in each age-class and all other fish species caught were identified, weighed and measured. Herring were measured to standard length; salmon to fork length; dogfish, hake and pollock to total length; and squid to distal mantle length. All other fish species were measured to standard length. Seventy herring, 50 age-0+ and 20 age-1+, from each of six selected sets were retained for stomach content analysis.

Plankton Sampling

Twenty stepped oblique plankton tows were performed during the survey, generally one tow every second seine set. The tows were always completed after dusk and immediately before or after a fishing event. Dual 19cm diameter bongo nets with 350µm mesh were used for sampling, resulting in 'left' and 'right' bongo plankton samples (only left samples were processed). The bongos were lowered to 20m (10m in shallow areas) and raised by an electric winch at a rate of 1m every 15sec (or 1m every 30sec for shallow areas). A General Oceanics® 2030R model flowmeter was attached to the left bongo to determine the volume of seawater filtered. Volume filtered was calculated using the following equation (McCarter and Hay 2002):

$$V = (A \cdot F \cdot K) / 999,999 \quad \text{Equation 1}$$

Where:

V = volume of water filtered through the plankton net (m³)

A = area of net opening (0.02835m²)

F = number of revolutions recorded by the flow meter (end reading – start reading)

K = standard speed rotor constant for 7cm rotor (26,873)

Upon retrieval, the bongo nets were washed with a high pressure deck hose, and the samples preserved in 3.7% seawater formalin.

In the laboratory, a volumetric splitter was used to reduce the sample size to a point where organisms could be conveniently counted and identified in a counting tray using a stereo

microscope under 30X magnification. Sample splitting continued until a target size of roughly 300 organisms was reached (Thompson et al. 2003).

Generally, plankters were not identified to species but classified into readily recognizable categories. Copepods were however identified to species when possible. Densities for all plankters were determined and expressed as plankters·m⁻³.

STOMACH CONTENTS

Herring

Juvenile herring stomachs were retained from six set locations: Set 1 (Fish Egg Inlet), Set 5 (Kwakshua Channel), Set 22 (Kitasu Bay), Set 28 (Powell Anchorage), Set 31 (Spiller Channel) and Set 36 (Hunter Channel) for further analysis. From each of these sets, 20 age-0+ and 20 age-1+ herring stomach samples were extracted in the laboratory and preserved in ~3.7% formalin in individually labelled vials. Each stomach was rated for fullness (empty, trace, half full and full), state of digestion (fresh, partly, mostly and totally) and, when possible, its contents identified and enumerated. Stomach contents were categorized into 28 groups of organisms representing 10 phyla. Copepods were further classified to species.

Other Species

Stomachs of possible herring predators; pink, coho, chum and sockeye salmon, chub mackerel, spiny dogfish shark, and starry flounder were sampled (fish >25cm). Fish were weighed and measured in the field, and their stomachs were removed and preserved in ~3.7% seawater formalin. Stomachs were analyzed for prey items, fullness and level of digestion. Identifiable contents were weighed and measured.

RESULTS

Thirty-nine sets were made during the 2003 survey; four in section 091 (Fish Egg Inlet), three in section 085 (Kwakshua Channel), six in section 076 (Kildidt Sound), four in section 074 (Thompson Bay), six in section 067 (Laredo Inlet), four in section 077 (Milbanke Sound), six in section 072 (Powell Anchorage), four in section 073 (Bella Bella) and two in section 084 (Burke Channel). Section numbers and names correspond to the classification by Midgley (2003), and should not be confused with the individual set location names (Table 1).

All 2003 set locations correspond to set locations used in the 2002 Central Coast survey, with the following exceptions. Two new sets (38 and 39) were made further inland (NW) in Burke Channel, replacing a 2002 set near the entrance (omitted due to strong tide). Also, one new set was made in Spider Anchorage (set 13), which was aborted in 2002 due to adverse weather conditions (Figures 2 and 3, Table 1).

Twenty-two species of fish and invertebrates from 4 phyla were identified in the seine catches. Fish species most frequently encountered (>15% occurrence) included: Pacific herring, pink salmon, chinook salmon, walleye pollock, coho salmon, shiner perch and capelin. Opal squid also were encountered in 43.5% of the sets (Tables 2 and 3).

Herring

A total of 7225 herring were measured resulting in a length frequency distribution that was distinctly bimodal. Based on this length frequency distribution, the length designations for the two juvenile herring age-classes are (Figure 4):

0+ = herring less than or equal to 85mm standard length

1+ = herring between 86mm and 129mm standard length

2+ or older = herring greater than or equal to 130mm standard length

Age-0+ herring occurred in 87.2% of the sets (Table 3). The largest catches of age-0+ herring by weight occurred in Kitasu Bay (section 067), Fish Egg Inlet (section 091) and Hunter Channel (section 073). Table 4 shows the average length and weight, and the total herring catch weight for age-0+ herring at each set location. The mean length of all sampled age-0+ herring (n=3933) was 61.70mm.

Age-1+ herring occurred in 76.9% of the sets (Table 3). The largest catches of age-1+ herring occurred in Fish Egg Inlet (section 091), Kitasu Bay (section 067) and Kildidt Sound (section 076). Table 4 shows the average length and weight, and the total herring catch weight for age-1+ herring at each set location. The mean length of all sampled age-1+ herring (n=3129) was 103.29mm.

Age-2+ herring occurred in 35.9% of the sets (Table 3). The largest catches occurred in Kildidt Sound (section 076) and Kwakshua Channel (section 085). No age-2+ herring were caught in Spider Anchorage, East Higgins Passage or Powell Anchorage. The mean length of all sampled age-2+ herring (n=163) was 145.60mm.

Herring length frequency distributions varied by location. The histograms display the length frequency for sampled herring only (Figure 5) but provide a representation of juvenile herring length distribution during the survey. The relationship between length and weight for all sampled herring was determined by fitting a logistic function to the length-weight data (Figure 6).

Plankton

There were 26 categories of organisms identified in 20 plankton samples (Table 5 and 6). Most samples contained a mild to heavy amount of *Coscinodiscus* diatoms, but were not quantified. An average of 12.31m³ of water was filtered per plankton tow. Copepods occurred in all samples. Unidentifiable juvenile calanoid copepods and *Pseudocalanus minutus* were the only organisms to occur in all samples. Amphipods, barnacle larvae, coelenterates, crab megalopea and zoea, larval euphausiids, larvaceans, calanoid copepods (*Acartia longiremis*, *Centropages abdominalis*, *Metridia pseudopacifica*, *Pseudocalanus minutus*), and unidentifiable juvenile calanoid copepods made up >75% of samples. Ctenophores, ectoprocts, prosobranch gastropods, polychaetes, miscellaneous eggs (mainly euphausiid) and the calanoid copepod *Calanus marshallae* occurred in 50 – 74% of samples. Chaetognaths, adult euphausiids, teleost larvae,

siphonophores, calanoid copepods (*Calanus pacificus*, *Eucalanus elongates*, *Paracalanus parvus*, *Tortanus discaudatus*), and the cyclopoid copepod *Oithona* sp. occurred in 25 – 49% of samples. Barnacle nauplii, isopods (cumaceans and mysids), echinoderms, euphausiids, ostracods, pelecypods, calanoid copepods (*Acartia* sp., *Calanus* sp., *Eucalanus bungii*, *Gaidius* sp., *Oncaea borealis*) the cyclopoid copepod *Corcaeus anglicus*, and unidentified copepod nauplii, cyclopoid copepods and harpacticoid copepods occurred in <25% of samples (Table 7).

Herring Stomachs

A total of 121 age-0+ herring stomachs were examined (Table 8). Of these, 117 contained identifiable stomach contents. On average, unidentifiable calanoid copepods occurred most frequently followed by barnacle nauplii, *Podon* sp. and *Evadne* sp. cladocerans and coscinodiscus diatoms (Table 9).

A total of 103 age-1+ herring stomachs were examined (Table 8). Of these, 92 contained identifiable stomach contents. On average, crab zoea occurred most frequently, followed by unidentifiable calanoid copepods, coscinodiscus diatoms and turbellarians (Table 10).

Other Species

Pink, coho, chum and sockeye salmon, chub mackerel, spiny dogfish shark, and starry flounder were sampled. A total of 46 possible herring predator stomachs were sampled, with only seven stomachs containing identifiable herring (Table 11). Several stomachs had material that was unidentifiable, so only the stomach content weight was recorded. Three salmon species (pink, coho and chum) were found to contain juvenile herring (age-0+ and age-1+). Chub mackerel stomachs contained euphausiids but no herring.

CONCLUSION

Thirty-nine stations were sampled resulting in 22 different fish species recorded from the purse seine sets. A total of 7225 herring were measured and weighed creating a distinct bimodal histogram representing two juvenile herring age groups. A total of 224 herring stomachs were examined for contents. The dominant prey item was calanoid copepods for age-0+ herring and crab zoea for age-1+ herring. Forty-six possible herring predator stomachs were examined, with only seven stomachs containing identifiable juvenile herring. Twenty plankton tows were performed resulting in unidentifiable juvenile calanoid copepods and *Pseudocalanus minutus* being the most prominent organisms.

ACKNOWLEDGMENTS

This study was funded by the Herring Conservation and Research Society (HCRS), and we are grateful for their continuing support. All plankton and herring stomach samples were processed by Carol Cooper of Zotec Services. Bob Armstrong of Armstrong Biotech Services and Dennis Chalmers conducted the field work and sampling. Fisheries and Oceans provided the research vessel and laboratory facilities. Kristen Daniel provided editorial comments regarding ArcView© maps.

REFERENCES

- Haegle, C.W. 1997. The occurrence, abundance and food of juvenile herring and salmon in the Strait of Georgia, British Columbia in 1990 to 1994. *Can. Manuscr. Rep. Fish. Aquat. Sci.* 2390: 124 p.
- Haegle, C.W., and Armstrong, R.W. 2003. Juvenile herring survey of the Central Coast of British Columbia, August 11-24, 2002. Contractors report for the Herring Conservation and Research Society: 34 p.
- Hay, D.E., and McCarter, P.B. 1999. Age of sexual maturation and recruitment in Pacific herring. *Can. Stock Assessment Secretariat Research Document* 99/175: 42 p.
- Henderson, S., Therriault, T.W., Haegle, C.W., Schweigert, J., and Thompson, M. 2004. Central Coast Juvenile Herring Survey, August 2002. *Can. Manuscr. Rep. Fish. Aquat. Sci.* 2687: 33 p.
- Hourston, A.S., and Haegle, C.W. 1980. Herring on Canada's Pacific coast. *Can. Spec. Publ. Fish. Aquat. Sci.* 48: 23 p.
- Humphreys, R.D., and Hourston, A.S. 1978. British Columbia herring spawn deposition survey manual. *Fish. Mar. Serv. Misc. Spec. Publ.* 38: 40 p.
- McCarter, P.B., and Hay, D.E. 2002. Eulachon embryonic egg and larval outdrift sampling manual for ocean and river surveys. *Can. Tech. Rep. Fish. Aquat. Sci.* 2451: 33 p.
- Midgley, P. 2003. Definitions and codings of localities, herring sections, and stock assessment regions for British Columbia herring data. *Can. Manuscr. Rep. Fish. Aquat. Sci.* 2634: 113 p.
- Thompson, M., Hrabok, C., Hay, D.E., Schweigert, J., Haegle, C., and Armstrong, B. 2003. Juvenile herring surveys: methods and data base. *Can. Manuscr. Rep. Fish. Aquat. Sci.* 2651: 31 p.




Figure 1. The five major British Columbia herring stock assessment regions: Prince Rupert District (PRD), Queen Charlotte Islands (QCI), Central Coast (CC), west coast Vancouver Island (WCVI) and the Strait of Georgia (SOG).




Figure 2. Northern Central Coast stock assessment area with set locations (triangles) and section numbers (large 3-digit numbers).




Figure 3. Southern Central Coast stock assessment area with set locations (triangles) and section numbers (large 3-digit numbers).




Figure 4. Length frequency distribution of all sampled herring.




Figure 5. Length frequency distributions of sampled herring by set location.

Figure 5 continued...




Spider AnchorageThompson BayMeyers Passage

Figure 5 continued...





Figure 5 continued...






Figure 6. Length-weight relationship for all sampled herring ($W = 7.62 \times 10^{-6} L^{3.085}$, $R^2 = 0.9904$, $n = 7225$).




Figure 7. Proportion of herring relative to catch by location.

Table 1. Details of the 39 sets made during the 2003 Central Coast juvenile herring survey.

Set	Month	Day	Set Code	Station	Section	Set Location Name	DD Lat (N)	DD Long (W)	Tow Time
1	8	5	1	1	091	Fish Egg Inlet	51.6210	127.7450	2245
2	8	5	1	2	091	Fish Egg Inlet	51.6040	127.7730	2420
3	8	5	1	3	091	Fish Egg Inlet	51.6170	127.8320	130
4	8	5	1	4	091	Fish Egg Inlet	51.6470	127.8280	230
5	8	6	2	2	085	Kwakshua Channel	51.6550	128.0530	2245
6	8	6	2	3	085	Kwakshua Channel	51.6480	128.0910	2340
7	8	6	2	4	085	Kwakshua Channel	51.6550	128.1150	2430
8	8	7	4	2	076	Kildidt Sound	51.8790	128.1590	2230
9	8	7	4	1	076	Kildidt Sound	51.8620	128.1460	2320
10	8	7	4	3	076	Kildidt Sound	51.8370	128.1540	2415
11	8	7	4	4	076	Kildidt Sound	51.8497	128.1155	120
12	8	8	3	1	076	Spider Anchorage	51.8630	128.2380	2220
13	8	8	3	2	076	Spider Anchorage	51.8530	128.2352	2350
14	8	9	5	1	074	Thompson Bay	52.1210	128.4070	2230
15	8	9	5	2	074	Thompson Bay	52.1340	128.3780	2315
16	8	9	5	3	074	Thompson Bay	52.1480	128.3980	2400
17	8	9	5	4	074	Thompson Bay	52.1590	128.3540	110
18	8	11	6	2	067	Meyers Passage	52.6840	128.5770	2215
19	8	11	6	1	067	Meyers Passage	52.6730	128.5720	2300
20	8	12	7	1	067	Kitasu Bay	52.5500	128.7990	2220
21	8	12	7	2	067	Kitasu Bay	52.5450	128.7710	2340
22	8	12	7	3	067	Kitasu Bay	52.5410	128.7420	2455
23	8	12	7	4	067	Kitasu Bay	52.5250	128.7510	210
24	8	13	8	1	077	East Higgins Pass	52.4020	128.5580	2210
25	8	13	8	2	077	East Higgins Pass	52.4360	128.5950	2325
26	8	13	8	3	077	East Higgins Pass	52.4710	128.6380	2450
27	8	13	8	4	077	East Higgins Pass	52.4920	128.6930	145
28	8	14	9	1	072	Powell Anchorage	52.2910	128.3760	2230
29	8	14	9	2	072	Powell Anchorage	52.2840	128.3550	2320
30	8	15	10	1	072	Spiller Channel	52.2500	128.2740	2210
31	8	15	10	2	072	Spiller Channel	52.2680	128.2710	2255
32	8	15	10	3	072	Spiller Channel	52.2770	128.2350	2335
33	8	15	10	4	072	Spiller Channel	52.2900	128.2590	2430
34	8	16	11	4	073	Hunter Channel	52.0780	128.1530	2205
35	8	16	11	2	073	Hunter Channel	52.0570	128.1410	2245
36	8	16	11	3	073	Hunter Channel	52.0730	128.1350	2330
37	8	16	11	1	073	Hunter Channel	52.0570	128.1150	2430
38	8	17	12	2	084	Burke Channel	52.1239	127.4380	2135
39	8	17	12	3	084	Burke Channel	52.1693	127.4946	2335

Section	Official herring management section names
067	Laredo Inlet
072	Powell Anchorage
073	Bella Bella
077	Milbanke Sound

Table 2. Species* quantity and weights by set.

Set	Location Name	Species	Quantity	Weight (g)
1	Fish Egg Inlet	0+ Herring	34692	15410
		1+ Herring	7920	18790
		Capelin	701	182
		Juvenile Pollock	16	26
		Three-spine		
		Stickleback	4	3
2	Fish Egg Inlet	0+ Herring	20460	35600
		1+ Herring	60	600
		Shiner Perch	5	45
3	Fish Egg Inlet	0+ Herring	20588	21000
		1+ Herring	545	3600
		2+ Herring	12	227
		Juvenile Chinook		
		salmon	2	52
		Juvenile Pollock	12	472
		Shiner Perch	2	25
4	Fish Egg Inlet	0+ Herring	4715	5800
		1+ Herring	9	80
5	Kwakshua Channel	0+ Herring	2486	4600
		1+ Herring	620	9800
		2+ Herring	31	1800
		Juvenile Chinook		
		salmon	2	222
6	Kwakshua Channel	Juvenile Pink salmon	1	41
		0+ Herring	8	21
		1+ Herring	135	2055
7	Kwakshua Channel	Squid		4000
		0+ Herring	1	4
		1+ Herring	164	2260
		2+ Herring	3	94
		Adult Pink salmon	2	3200
		Shiner Perch	2	-
		Three-spine		
8	Kildidt Sound	Stickleback	2	-
		0+ Herring	26	90
		1+ Herring	439	6000
		Poacher	1	0.5
		Squid	-	2100

Table 2 continued...

Set	Location Name	Species	Number	Weight (g)
9	Kildidt Sound	0+ Herring	12	40
		1+ Herring	914	13800
		2+ Herring	70	2700
		Juvenile Hake	30	449
		Squid	-	1500
10	Kildidt Sound	0+ Herring	10	42
		1+ Herring	427	5400
		2+ Herring	11	401
		Juvenile Chinook salmon	3	915
		Juvenile Hake	1	16
		Squid	-	7900
11	Kildidt Sound	Pacific Sardine	6	1330
12	Spider Anchorage	0+ Herring	14	38
		Juvenile Coho salmon	1	17
		Squid	-	2000
13	Spider Anchorage	0+ Herring	37	112
		Juvenile Coho salmon	1	108
		Juvenile Pink salmon	5	510
		Squid	-	300
14	Thompson Bay	1+ Herring	349	5100
		2+ Herring	2	60
		Squid	-	1300
15	Thompson Bay	0+ Herring	2	6
		1+ Herring	130	1800
		Spiny Dogfish	1	86
		Squid	-	3000
16	Thompson Bay	0+ Herring	8	39
		1+ Herring	4	45
		Shiner Perch	1	15
		Squid	-	20
17	Thompson Bay	0+ Herring	2	4
		Juvenile Hake	1	4
		Squid	-	1000
18	Meyers Passage	0+ Herring	9862	28600
		1+ Herring	42	574
		2+ Herring	1	27
		Adult Pink salmon	1	1600

Table 2 continued...

Set	Location Name	Species	Number	Weight (g)
19	Meyers Passage	0+ Herring	91	272
		1+ Herring	513	6700
		2+ Herring	2	50
		Juvenile Pollock	1	141
20	Kitasu Bay	0+ Herring	54917	49700
		1+ Herring	24	33
		Adult Pink salmon	3	6200
		Capelin	84	109
		Juvenile Rockfish	6	3
21	Kitasu Bay	0+ Herring	5900	15380
		1+ Herring	4827	27610
		2+ Herring	5	77
		Juvenile Coho salmon	2	129
		Juvenile Pink salmon	3	53
		Squid	5	-
		Whitebait Smelt	3	15
22	Kitasu Bay	0+ Herring	4802	15800
		1+ Herring	1401	19100
		2+ Herring	5	127
		Adult Coho salmon	2	7700
		Adult Pink salmon	1	2000
		Juvenile Chinook salmon	6	942
		Juvenile Coho salmon	11	1570
		Juvenile Pink salmon	1	19
		Squid	-	1300
23	Kitasu Bay	0+ Herring	879	3500
		1+ Herring	4	38
		Juvenile Chinook salmon	4	455
		Shiner Perch	1	16
		Sockeye	1	1500
		Squid	8	-
24	East Higgins Pass	Adult Chum salmon	1	4000
		Adult Pink salmon	1	2100
		Juvenile Coho salmon	4	652
		Juvenile Pink salmon	26	711
		Mackeral	26	53100

Table 2 continued...

Set	Location Name	Species	Number	Weight (g)
25	East Higgins Pass	Adult Chum salmon	2	8100
		Adult Pink salmon	2	3300
		American Shad	1	827
		Juvenile Chinook salmon	3	622
		Juvenile Pink salmon	8	202
		Mackeral	24	50100
26	East Higgins Pass	Adult Chum salmon	1	2400
		Juvenile Pink salmon	1	2100
27	East Higgins Pass	0+ Herring	36	51
		1+ Herring	5	48
		Spiny Dogfish	3	4300
28	Powell Anchorage	0+ Herring	5608	8300
		Adult Chum salmon	1	6000
		Juvenile Pollock	5	374
		Shiner Perch	1	3
		Squid	17	500
		Starry Flounder	1	1800
29	Powell Anchorage	0+ Herring	3289	5000
		1+ Herring	13	147
		Juvenile Pollock	10	573
		Shiner Perch	14	55
30	Spiller Channel	0+ Herring	6	19
		1+ Herring	1	12
		2+ Herring	1	58
		Adult Coho salmon	1	2500
		Juvenile Chinook salmon	1	154
		Squid	2	-
31	Spiller Channel	0+ Herring	200	502
32	Spiller Channel	0+ Herring	42	129
		1+ Herring	5	61
33	Spiller Channel	0+ Herring	17	47
		1+ Herring	2	24
		Juvenile Chinook salmon	1	108

Table 2 continued...

Set	Location Name	Species	Number	Weight (g)
34	Hunter Channel	0+ Herring	1148	3100
		1+ Herring	535	6000
		2+ Herring	8	309
		Capelin	19	32
		Juvenile Pollock	22	1290
35	Hunter Channel	0+ Herring	12755	41452
		1+ Herring	532	6347
		Capelin	2	11
		Juvenile Pollock	19	1478
36	Hunter Channel	0+ Herring	13723	41032
		1+ Herring	420	5166
		Capelin	16	24
		Juvenile Chinook salmon	1	36
		Juvenile Pollock	2	175
37	Hunter Channel	0+ Herring	9026	19044
		1+ Herring	64	555
		Capelin	72	49
		Squid	2	-
38	Burke Channel	0+ Herring	7643	36992
		1+ Herring	518	6029
		2+ Herring	8	283
		Adult Pink salmon	2	3800
		Juvenile Smelt	3	30
		Spiny Dogfish	2	3300
		Squid	1	-
		Three-spine Stickleback	7	20
39	Burke Channel	0+ Herring	8	40
		1+ Herring	139	1863
		2+ Herring	4	149
		Juvenile Coho salmon	4	43
		Juvenile Pink salmon	1	24
		Three-spine Stickleback	8	16

* Invertebrates such as euphausiids (Phylum Arthropoda), comb jellies (phylum Ctenophora) and jellyfish (Phylum Cnidaria) were frequently encountered but not included in this table due to quantifying difficulties.

Table 3. Percent species* occurrence in all 2003 Central Coast sets.

Common Name	Scientific Name	Set Numbers	Percent Occurrence (%)
0+ Herring	<i>Clupea pallasii</i> in year of birth	1,2,3,4,5,6,7,8,9,10,12,13,15,16,17,18,19,20,21,22,23,27,28,29,30,31,32,33,34,35,36,37,38,39	87.2
1+ Herring	<i>Clupea pallasii</i> in year after birth	1,2,3,4,5,6,7,8,9,10,14,15,16,18,19,20,21,22,23,27,29,30,32,33,34,35,36,37,38,39	76.9
2+ Herring	<i>Clupea pallasii</i> in 2nd year after birth	3,5,7,9,10,14,18,19,21,22,30,34,38,39	35.9
No Herring caught		11,24,25,26	10.3
Opal squid	<i>Loligo opalescens</i>	6,7,8,9,10,12,13,14,15,16,17,21,22,23,28,29,37	43.5
Pink salmon	<i>Oncorhynchus gorbuscha</i>	5,7,13,18,20,21,22,24,25,26,38,39	30.8
Chinook salmon	<i>Oncorhynchus tshawytscha</i>	3,5,10,22,23,25,30,33,36	23.1
Walleye pollock	<i>Theragra chalcogramma</i>	1,3,19,28,29,34,35,36	20.5
Coho salmon	<i>Oncorhynchus kisutch</i>	12,13,21,22,24,30,39	18.0
Shiner perch	<i>Cymatogaster aggregata</i>	2,3,7,16,23,28,29	18.0
Capelin	<i>Mallotus villosus</i>	1,20,34,35,36,37	15.4
Chum salmon	<i>Oncorhynchus keta</i>	24,25,26,28	10.3
Three-spine stickleback	<i>Gasterosteus aculeatus</i>	1,7,38,39	10.3
Pacific hake	<i>Merluccius productus</i>	9,10,17	7.7
Spiny dogfish	<i>Squalus acanthias</i>	15,27,38	7.7
Red squid	<i>Berryteuthis magister</i>	30,38	5.1
Chub mackerel	<i>Scomber japonicus</i>	24,25	5.1
Sockeye salmon	<i>Oncorhynchus nerka</i>	23	2.6
Whitebait smelt	<i>Allosmerus elongatus</i>	21	2.6
Surf smelt	<i>Hypomesus pretiosus</i>	38	2.6
Northern spearnose poacher	<i>Agonopsis vulsa</i>	8	2.6
Pacific sardine	<i>Sardinops sagax</i>	11	2.6
Rockfish	<i>Sebastes sp.</i>	20	2.6
American shad	<i>Alosa sapidissima</i>	25	2.6
Starry flounder	<i>Platichthys stellatus</i>	28	2.6

*Invertebrates such as euphausiids (Phylum Arthropoda), comb jellies (Phylum Ctenophora) and jellyfish (Phylum Cnidaria) were not quantified.

Table 4. Mean lengths, weights and total catch weights of age-0+ and age-1+ herring by set location. Herring management sections in brackets.

Set Location	Mean 0+ Length (mm)	Mean 0+ Weight (g)	Total Catch Weight 0+ (Kg)	Mean 1+ Length (mm)	Mean 1+ Weight (g)	Total Catch Weight 1+ (Kg)
Fish Egg Inlet (091)	55.1	1.78	145.040	100.5	11.37	83.040
Kwakshua Channel (085)	56.3	1.94	4.625	109.8	14.92	14.115
Kildidt Sound (076)	71.3	4.00	0.172	106.9	13.88	25.200
Spider Anchorage (076)	64.8	2.93	0.149	0	0	0
Thompson Bay (074)	69.9	4.10	0.049	107.0	14.30	6.945
Meyers Passage (067)	64.2	3.03	28.872	105.8	13.42	7.274
Kitasu Bay (067)	70.9	4.16	340.570	106.6	14.05	60.751
East Higgins Passage (077)	50.5	1.40	0.051	95.0	9.61	0.048
Powell Anchorage (072)	51.1	1.53	13.300	101.1	12.12	0.147
Spiller Channel (072)	62.6	2.63	0.696	102.1	12.29	0.098
Hunter Channel (073)	62.3	2.77	104.628	100.9	11.58	17.968
Burke Channel (084)	74.9	4.87	37.032	100.5	12.60	7.892
All Locations	61.9	2.80	675.185	103.3	12.74	223.478

Table 5. Categories of organisms, by phylum, identified in plankton and stomach samples. The code (capital letters) is used to identify organisms in Tables 7, 9 and 10.

Coelenterata

COEL Medusae
SIPH Siphonophores

Ctenophora

CTEN Ctenophores

Turbellaria

TURB Turbellarians – flatworms

Annelida

POLY Polychaetes – segmented worms

Mollusca

GAST Prosobranch gastropods
PELE Pelecypods
PTER Pteropods (Ophistobranch gastropods), mostly *Clione sp.*

Arthropoda

AMPH Amphipods, mostly gammarid and hyperiid, some caprellid
BARN Barnacle, unknown stage
BARNc Barnacle cirriped
BARNn Barnacle nauplius
CLAD Cladocerans; *Podon sp.* and *Evadne sp.*
COPE Copepods (See Table 6 for a list of species)
CRAM Crab megalopea, including porcellinadea
CRAZ Crab zoea, including porcellinadea
EUPA Euphausiid adults
EUPH Euphausiid, unknown stage
EUPL Euphausiid larvae (nauplii, protozoa and zoea)
INSE Insects
ISOP Isopods, cumaceans and mysids
OSTR Ostracods
SHRI Shrimp (natant decapod) zoea

Ectoprocta

ECTO Ectoprocts, mostly *Membranipora sp.* larvae (cyphonautes)

Echinodermata

ECHI Echinoderm larvae

Chaetognatha

CHAE Chaetognaths, mostly *Sagitta sp.*

Chordata

LARV Larvaceans, mostly *Oikopleura sp.*, and tunicate larvae
TELA Teleost larvae

Miscellaneous

EGGS Unidentified eggs, polychaete or teleost

Table 6. Copepods identified in plankton and stomach samples. The code (capital letters) is used to identify organisms in Tables 7, 9 and 10.

CNAU Unidentified copepod nauplii

DIAT Diatoms, mainly *Coscinodiscus*

Calanoid copepods

ACAR	<i>Acartia sp.</i>
ALON	<i>Acartia longiremis</i>
CABD	<i>Centropages abdominales</i>
CALA	<i>Calanus sp.</i>
CCOL	<i>Cadacia columbiae</i>
CGRA	<i>Chiridius gracilis</i>
CMAR	<i>Calanus marshallae</i>
CPAC	<i>Calanus pacificus</i>
EBUN	<i>Eucalanus bungii</i>
EELO	<i>Eucalanus elongatus</i>
ELON	<i>Epilabidocera longipedata</i>
GAID	<i>Gaidius sp.</i>
JCAL	Juvenile calanoids, unknown species
MPAC	<i>Metridia pacifica</i>
MPSE	<i>Metridia pseudopacifica</i>
OBOR	<i>Oncaea borealis</i>
PMIN	<i>Pseudocalanus minitus</i>
PPAR	<i>Paracalanus parvus</i>
TDIS	<i>Tortanus discaudatus</i>
UCAL	Unidentified calanoid

Cyclopoid copepods

CANG	<i>Corcaeus anglicus</i>
CYCL	Cyclopoid
OITH	<i>Oithona sp.</i>
UCYC	Unidentified cyclopoids

Harpacticoid copepods

UHAR Unidentified harpacticoids

Table 7. Zooplankton density per set in number of organisms per m³ of water sampled. Species codes can be found in Table 5 and 6.

Set	Location	Water Volume (m ³)	ACAR	ALON	AMPH	BARN	BARNn	CABD	CALA	CANG	CCOL	CHAE	CLAD
1	Fish Egg Inlet	14.78142	-	22.7	15.7	10.3	-	0.5	-	-	-	4.3	7.6
5	Kwakshua Channel	9.08964	-	169.0	169.0	70.4	-	14.1	-	-	56.3	-	42.2
7	Kwakshua Channel	11.86810	-	17.5	1.0	76.2	-	-	-	-	-	-	6.7
8	Kildidt	11.70812	-	62.9	6.1	153.7	-	19.1	-	-	-	-	32.8
10	Kildidt	13.09545	-	29.3	2.7	26.9	-	88.0	-	-	-	0.3	66.0
13	Spider Anchorage	13.72473	-	135.2	-	33.8	-	16.3	5.8	-	-	-	46.6
14	Thompson Bay	17.04792	-	337.9	0.9	22.5	-	322.9	-	-	-	0.9	52.6
16	Thompson Bay	10.80532	-	204.3	3.7	20.7	-	266.5	-	3.0	-	-	20.7
19	Meyers Passage	18.78570	-	30.7	0.1	316.9	-	6.8	-	-	-	-	3.4
21	Kitasu Bay	8.45425	-	847.9	-	-	45.4	1150.7	-	-	-	-	-
23	Kitasu Bay	8.19370	-	1046.7	-	7.8	-	179.7	-	15.6	-	-	7.8
25	East Higgins Pass	17.23458	-	126.3	-	-	-	14.9	-	-	-	-	-
27	East Higgins Pass	15.55470	-	203.7	0.5	-	47.3	6.2	-	-	-	-	61.7
28	Powell Anchorage	18.50458	-	21.6	1.0	13.8	-	3.9	-	-	-	-	9.1
30	Spiller Channel	15.40766	-	155.8	3.1	8.3	-	20.8	-	-	-	-	-
32	Spiller Channel	19.53003	-	147.5	1.6	9.8	-	32.8	-	-	3.3	-	13.1
34	Hunter Channel	6.55039	-	161.2	15.7	444.9	-	-	-	-	-	0.3	97.7
36	Hunter Channel	8.36512	-	84.2	13.4	639.6	-	30.6	-	-	-	4.1	110.9
38	Burke Channel	3.16016	101.3	-	13.3	45.6	-	15.2	-	-	-	-	75.9
39	Burke Channel	4.28998	126.8	-	-	82.1	-	14.9	-	-	1.9	-	29.8
% Occurrence			10.0	90.0	75.0	85.0	10.0	90.0	5.0	10.0	15.0	25.0	85.0

Table 7 continued...

Set	CMAR	CNAU	COEL	CPAC	CRAM	CRAZ	CTEN	EBUN	ECHI	ECTO	EELO	EGGS	EUPA	EUPH
1	4.3	-	2.2	0.5	0.5	4.3	1.6	-	-	0.5	-	6.5	2.7	-
5	183.1	-	-	-	-	28.2	-	-	-	-	-	70.4	-	-
7	0.3	-	-	-	0.3	23.3	32.7	-	-	5.4	-	-	-	-
8	0.3	-	9.2	0.3	-	389.5	8.9	-	-	-	-	21.9	-	-
10	0.3	-	9.5	0.6	4.0	61.7	5.2	-	-	4.9	0.9	100.2	0.9	-
13	-	-	0.3	-	6.1	61.8	2.0	-	-	1.2	0.3	-	-	23.3
14	0.9	-	0.5	-	8.9	86.3	-	-	-	-	-	-	-	11.3
16	-	-	2.2	-	19.2	144.4	3.0	-	-	8.9	-	-	-	-
19	0.2	-	1.0	-	0.9	15.9	2.4	-	-	40.9	0.2	54.5	-	-
21	-	-	0.9	-	18.0	3.3	-	-	-	-	-	302.8	121.1	-
23	-	-	9.3	-	22.5	87.9	-	-	-	-	-	70.3	-	-
25	0.9	-	0.2	0.2	0.5	1.4	-	-	-	-	-	-	2.8	-
27	-	-	17.0	-	9.8	34.5	-	-	-	6.2	-	4.1	-	-
28	-	-	0.8	-	1.1	13.0	2.8	-	-	1.3	-	-	-	-
30	7.0	-	-	-	2.3	0.8	-	-	-	-	-	105.9	1.1	-
32	1.6	-	0.4	-	0.8	7.4	-	-	-	-	-	85.2	1.6	-
34	-	-	1.4	0.2	0.9	53.9	15.3	-	-	48.9	-	102.6	-	-
36	-	-	0.7	-	0.7	22.7	19.8	-	-	45.9	0.2	65.0	0.5	-
38	-	-	7.0	-	3.2	-	69.0	1.3	35.4	10.1	0.6	-	1.9	45.6
39	-	387.9	12.1	-	-	14.9	270.4	0.9	208.9	14.9	-	-	-	-
% Occurrence	50.0	5.0	85.0	25.0	85.0	95.0	60.0	10.0	10.0	60.0	25.0	60.0	40.0	15.0

Table 7 continued...

Set	EUPL	GAID	GAST	ISOP	JCAL	LARV	MPSE	OBOR	OITH	OSTR	PELE	PMIN	POLY	PPAR	SHRI	SIPH
1	5.4	-	-	-	28.7	5.4	76.9	-	-	-	-	7.0	0.5	1.1	-	0.5
5	197.1	-	98.6	-	535.1	98.6	5477.9	-	42.2	-	14.1	112.7	14.1	56.3	-	-
7	7.1	-	6.7	0.3	7.1	5.4	48.5	-	-	-	-	8.1	5.4	-	3.7	-
8	27.3	-	10.9	-	16.7	8.2	3.4	-	-	2.7	-	21.9	-	-	22.5	0.3
10	119.7	-	-	-	77.6	58.6	0.3	-	12.2	-	-	19.5	-	-	15.6	5.5
13	28.0	-	12.8	1.7	62.4	31.5	-	-	14.0	-	-	5.8	-	12.8	23.9	37.0
14	30.5	-	-	-	145.0	22.5	-	-	7.5	-	-	67.6	-	-	12.2	-
16	19.2	-	11.8	-	74.0	32.6	-	-	-	-	-	14.8	-	3.0	27.4	31.1
19	264.2	-	6.8	-	15.1	6.8	0.4	-	-	-	-	37.5	0.5	-	0.5	-
21	30.3	-	-	-	171.7	181.7	-	-	15.1	-	-	863.0	-	-	33.6	0.5
23	7.8	-	31.2	-	71.8	93.7	-	-	-	-	-	156.2	-	-	4.9	-
25	3.5	-	-	-	193.3	-	37.6	-	3.7	-	-	304.5	-	-	0.2	-
27	2.6	-	2.1	-	49.4	18.5	1.0	-	-	-	-	129.6	0.5	-	26.7	-
28	0.4	-	-	1.8	1.4	0.4	0.4	-	0.4	0.1	-	3.0	0.4	0.4	2.8	-
30	0.5	-	-	-	56.9	-	18.7	-	-	-	-	16.6	3.6	-	1.0	-
32	-	-	4.1	-	168.8	-	43.0	-	-	-	-	19.7	7.0	-	1.2	-
34	68.9	-	4.9	-	86.4	219.8	1.4	-	-	-	-	4.9	4.9	4.9	2.1	-
36	58.6	-	19.6	0.2	173.8	65.0	25.6	-	-	-	11.5	30.6	3.8	-	11.2	-
38	66.5	1.9	45.6	-	1093.6	344.3	22.8	1.9	0.6	81.0	172.1	182.3	15.8	35.4	6.3	-
39	105.4	-	15.9	-	1150.6	238.7	2.8	-	-	59.7	104.4	104.4	52.2	67.1	-	-
% Occurrence	95.0	5.0	65.0	20.0	100.0	85.0	75.0	5.0	40.0	20.0	20.0	100.0	60.0	40.0	85.0	30.0

Table 7 continued...

Set	TDIS	TELA	UCYC	UHAR	Total
1	0.5	-	-	-	213.5
5	-	-	-	-	7464.4
7	-	1.3	-	-	278.2
8	-	0.3	-	-	843.3
10	-	0.3	-	-	740.8
13	1.2	-	-	-	602.9
14	-	0.5	-	-	1173.4
16	-	-	-	-	958.7
19	-	-	-	-	862.8
21	-	0.5	-	-	3849.5
23	-	-	-	-	1882.1
25	-	-	-	-	765.0
27	65.8	-	-	-	768.1
28	1.7	0.1	-	-	165.8
30	2.1	-	-	-	494.5
32	-	-	-	-	644.9
34	-	-	-	-	1443.0
36	-	4.3	-	-	1550.7
38	-	-	6.3	0.6	2616.4
39	-	-	-	-	3183.7
% Occurrence	25.0	35.0	5.0	5.0	

Table 8. Herring stomach fullness (empty, trace, half full, or full) and state of digestion (fresh, partly digested, mostly digested or totally digested).

Fullness	State of digestion				Total
	1 - fresh	2 - partly	3 - mostly	4 - totally	
0 +					
herring					
0 - empty	9	53	17	7	86
1 - trace	-	31	4	-	35
2 - half	-	-	-	-	-
3 - full	-	-	-	-	-
Total	9	84	21	7	121
1 +					
herring					
0 - empty	9	37	10	15	71
1 - trace	5	17	2	-	24
2 - half	3	2	-	-	5
3 - full	-	3	-	-	3
Total	17	59	12	15	103

Table 9. Quantity and percent occurrence of identifiable prey items in 0+ herring stomachs (n) by location. Tables 5 and 6 describe organism abbreviations.

N	Fish Egg Inlet		Kwakshua Channel		Kitasu Bay		Powell Anchorage		Spiller Channel		Hunter Channel	
	20		20		20		19		19		19	
	Quantity	Percent Occurrence	Quantity	Percent Occurrence	Quantity	Percent Occurrence	Quantity	Percent Occurrence	Quantity	Percent Occurrence	Quantity	Percent Occurrence
ALON	2	5	1	5	77	60	30	47.37	228	84.21	-	-
AMPH	-	-	5	25	-	-	-	-	-	-	1	5.26
BARNc	-	-	8	25	-	-	1	5.26	6	26.32	32	73.68
BARNn	10	30	58	90	-	-	228	89.47	58	73.68	65	63.16
CABD	-	-	3	15	135	70	2	5.26	12	21.05	2	5.26
CANG	3	15	-	-	1	5	101	52.63	18	52.63	3	15.79
CLAD	33	55	209	95	-	-	467	89.47	382	89.47	2	10.53
COPE	1	5	-	-	-	-	82	94.74	12	47.37	-	-
CRAZ	-	-	4	20	-	-	2	10.53	1	5.26	1	5.26
CYCL	-	-	-	-	-	-	3	5.26	-	-	-	-
DIAT	9650	100	1385	75	-	-	157	68.42	115	57.89	14	31.58
ECTO	12	35	16	40	-	-	116	78.95	24	52.63	7	21.05
EGGS	26	30	67	60	608	75	1	5.26	391	84.21	23	47.37
EUPH	-	-	10	5	-	-	-	-	-	-	-	-
EUPL	-	-	5	25	1	5	1	5.26	-	-	-	-
GAST	-	-	-	-	-	-	1	5.26	-	-	-	-
INSE	-	-	-	-	-	-	-	-	-	-	1	5.26
LARV	-	-	132	75	4	5	-	-	-	-	25	52.63
OITH	-	-	-	-	-	-	8	26.32	-	-	-	-
PELE	1	5	45	80	-	-	55	78.95	38	63.16	26	57.89
PPAR	1	5	2	10	-	-	32	15.79	6	21.05	-	-
SHRI	-	-	1	5	-	-	-	-	-	-	-	-
TURB	13	45	2	10	138	95	13	63.16	19	57.89	16	52.63
UCAL	20	30	354	90	311	90	604	100	592	89.47	42	47.37
UHAR	1	5	1	5	-	-	-	-	-	-	-	-

Table 10. Quantity and percent occurrence of identifiable prey items in 1+ herring (n) by location. Tables 5 and 6 describe organism abbreviations.

N	Fish Egg Inlet		Kwakshua Channel		Kitasu Bay		Powell Anchorage		Spiller Channel		Hunter Channel	
	20	Percent	20	Percent	18	Percent	13	Percent	4	Percent	17	Percent
	Quantity	Occurrence	Quantity	Occurrence	Quantity	Occurrence	Quantity	Occurrence	Quantity	Occurrence	Quantity	Occurrence
ALON	-	-	93	70	121	38.89	2	15.38	2	25	1	5.88
AMPH	1	5	27	65	-	-	1	7.69	-	-	1	5.88
BARNc	3	10	8	30	1	5.56	1	7.69	-	-	24	35.29
BARNn	-	-	30	55	-	-	1	7.69	-	-	8	23.53
CABD	-	-	18	35	81	44.44	2	7.69	-	-	11	11.76
CANG	-	-	-	-	-	-	19	38.46	-	-	-	-
CGRA	13	35	-	-	-	-	-	-	-	-	-	-
CHAE	-	-	1	5	-	-	-	-	-	-	-	-
CLAD	-	-	20	40	-	-	-	-	-	-	-	-
CMAR	-	-	5	10	-	-	-	-	-	-	-	-
COPE	-	-	-	-	-	-	1	7.69	-	-	-	-
CRAM	-	-	2	5	9	27.78	25	46.15	2	25	10	29.41
CRAZ	-	-	41	70	-	-	82	84.62	4	50	20	47.06
DIAT	145.3	60	1080	85	-	-	2	15.38	-	-	-	-
EGGS	-	-	89	45	65	33.33	-	-	9	25	-	-
ELON	5	15	-	-	-	-	-	-	-	-	-	-
EUPA	5	15	3	10	-	-	-	-	10	50	1	5.88
EUPH	-	-	-	-	2	5.56	-	-	1	25	-	-
EUPL	-	-	20	35	-	-	-	-	-	-	-	-
GAST	-	-	2	5	-	-	-	-	-	-	-	-
INSE	-	-	1	5	-	-	-	-	-	-	-	-
ISOP	1	5	1	5	-	-	-	-	-	-	-	-
LARV	-	-	354	80	31	16.67	-	-	-	-	24	5.88
MPAC	46	5	1281	90	-	-	-	-	-	-	-	-

Table 10 continued...

N	Fish Egg Inlet		Kwakshua Channel		Kitasu Bay		Powell Anchorage		Spiller Channel		Hunter Channel	
	20		20		18		13		4		7	
	Quantity	Percent Occurrence	Quantity	Percent Occurrence	Quantity	Percent Occurrence	Quantity	Percent Occurrence	Quantity	Percent Occurrence	Quantity	Percent Occurrence
OITH	-	-	2	5	-	-	-	-	-	-	-	-
PELE	-	-	5	20	-	-	-	-	-	-	-	-
PMIN	-	-	2	10	-	-	-	-	-	-	-	-
POLY	-	-	17	15	-	-	-	-	-	-	-	-
PPAR	-	-	8	20	-	-	-	-	-	-	-	-
PTER	1	5	16	20	-	-	-	-	-	-	-	-
SHRI	-	-	5	10	6	11.11	9	30.77	1	25	1	5.88
TELA	-	-	3	5	-	-	-	-	-	-	-	-
TURB	21	45	25	55	39	66.67	21	61.54	4	50	7	23.53
UCAL	603	95	173	100	160	55.56	6	15.38	-	-	110	41.18
UHAR	-	-	1	5	-	-	1	7.69	-	-	-	-

Table 11. Herring predator stomach contents. When possible, gut species were identified to species and total length (mm) and weight (g) were recorded.

Set	Species	Length (mm)	Weight (g)	Stomach Weight (g)	Gut Species	Quantity	Gut Species Length (mm)	Gut Species Weight (g)
7	Pink	495	1500	13.39	-	empty	-	-
7	Pink	524	1700	11.75	-	empty	-	-
18	Pink	508	1600	10.91	-	empty	-	-
20	Pink	572	2600	20.26	-	empty	-	-
20	Pink	540	2000	19	-	empty	-	-
20	Pink	533	1600	19	Herring	1	72	4.72
22	Pink	540	2000	31.61	Herring	1	112	15.29
22	Coho	648	3400	48.11	Herring	1	127	11.91
22	Coho	679	4300	43.17	-	-	-	3.27
					-	-	-	3.86
23	Sockeye	495	1500	6.47	-	empty	-	-
24	Chum	660	4000	70.5	Euphausiids	-	-	15.25
24	Pink	559	2100	28.63	Euphausiids	-	-	13.77
24	Chub Mackerel	556	2000	16.96	Euphausiids	-	-	12.27
24	Chub Mackerel	552	2000	10.81	Euphausiids	-	-	6.91
24	Chub Mackerel	552	2000	17.87	Euphausiids	-	-	12.7
24	Chub Mackerel	559	2000	20.05	Euphausiids	-	-	16.63
24	Chub Mackerel	546	2000	17.24	Euphausiids	-	-	10.07
24	Chub Mackerel	552	2000	17.47	Euphausiids	-	-	25.97
24	Chub Mackerel	594	2000	23.46	Euphausiids	-	-	0.62
24	Chub Mackerel	581	2000	15.36	Euphausiids	-	-	26.74
24	Chub Mackerel	559	2000	14.47	Euphausiids	-	-	15.83
24	Chub Mackerel	568	2000	16.49	Euphausiids	-	-	9.51
24	Chub Mackerel	540	2000	15.86	Euphausiids	-	-	30.99
24	Chub Mackerel	540	2000	15.08	Euphausiids	-	-	36.37
25	Chub Mackerel	533	2100	19.75	Euphausiids	-	-	7.22
25	Pink	483	1500	8.06	-	-	-	2.44
					Herring	3	-	2.37
25	Chum	686	4400	61.84	Euphausiids	-	-	0.92
25	Chub Mackerel	556	2100	23.15	Euphausiids	-	-	11.09
25	Pink	518	1800	12.68	Euphausiids	-	-	3.19
25	Chum	635	3700	32.15	-	-	-	2.47
25	Chub Mackerel	549	2100	18.14	Euphausiids	-	-	4.35
25	Chub Mackerel	524	2100	17.52	Euphausiids	-	-	21.12
25	Chub Mackerel	591	2100	18.75	Euphausiids	-	-	20.94
25	Chub Mackerel	559	2100	15.81	Euphausiids	-	-	1.96
26	Pink	552	2100	18.29	Herring	-	-	15.38
					Euphausiids	-	-	77.49
26	Chum	597	2400	3.73	Euphausiids	-	-	12.94
27	Spiny Dogfish	597	900	35.08	Euphausiids	-	-	0.79
27	Spiny Dogfish	648	1000	38.62	Euphausiids	-	-	2.12

Table 11 continued...

Set	Species	Length (mm)	Weight (g)	Stomach Weight (g)	Gut Species	Quantity	Gut Species Length (mm)	Gut Species Weight (g)
27	Spiny Dogfish	787	2400	164.91	Pollock	1	-	58.59
					-	-	-	50.89
27	Spiny Dogfish	787	2400	164.91	-	-	-	2.68
28	Chum	737	6000	40.96	-	-	-	0.96
					Herring	-	54	1.76
					Herring	-	51	1.59
28	Starry Flounder	489	1800	16.85	-	-	-	7.22
30	Coho	635	2500	22.09	Euphausiids	9	-	0.68
38	Pink	546	2000	8.7	-	empty	-	-
38	Spiny Dogfish	667	2300	51.88	-	4	-	12.57
					Herring	1	-	3.79
38	Pink	552	1800	7.28	-	empty	-	-
38	Spiny Dogfish	584	1000	36.35	-	-	-	1.24

SH 223 F55 no.2710 c.1

Henderson, S.

Central coast juvenile

herring survey, August 2003

287007_ 12063322 _ _ c.1

