



Natural Resources  
Canada

Ressources naturelles  
Canada

## CANADIAN GEOSCIENCE MAP 466

SURFICIAL GEOLOGY

# McCANN LAKE

Northwest Territories

NTS 75-G



Map Information  
Document

Geological Survey of Canada  
Canadian Geoscience Maps

2025

Canada 



## **MAP NUMBER**

Natural Resources Canada, Geological Survey of Canada  
Canadian Geoscience Map 466

## **TITLE**

Surficial geology, McCann Lake, Northwest Territories, NTS 75-G

## **SCALE**

1:100 000

## **CATALOGUE INFORMATION**

Catalogue No. M183-1/466-2025E-PDF  
ISBN 978-0-660-77376-6  
<https://doi.org/10.4095/pjfeux56fu>

## **COPYRIGHT**

© His Majesty the King in Right of Canada, as represented by the Minister of Natural Resources, 2025

Information contained in this publication or product may be reproduced, in part or in whole, and by any means, for personal or public non-commercial purposes, without charge or further permission, unless otherwise specified.

You are asked to:

- exercise due diligence in ensuring the accuracy of the materials reproduced;
- indicate the complete title of the materials reproduced, and the name of the author organization; and
- indicate that the reproduction is a copy of an official work that is published by Natural Resources Canada (NRCan) and that the reproduction has not been produced in affiliation with, or with the endorsement of, NRCan.

Commercial reproduction and distribution is prohibited except with written permission from NRCan. For more information, contact NRCan at [copyright-droitdauteur@nrcan-rncan.gc.ca](mailto:copyright-droitdauteur@nrcan-rncan.gc.ca).

## **RECOMMENDED CITATION**

Dyke, A.S. and Campbell, J.E., 2025. Surficial geology, McCann Lake, Northwest Territories, NTS 75-G; Geological Survey of Canada, Canadian Geoscience Map 466, scale 1:100 000. <https://doi.org/10.4095/pjfeux56fu>

## **ABSTRACT**

The McCann Lake map area lies in the Rae Province of the Canadian Shield. Quaternary deposits are extensive and thick, with bedrock outcrops scattered throughout. Till is

extensively streamlined to the west-southwest by late deglacial ice flow from the Keewatin Ice Divide and is incised by west-southwest-flowing subglacial meltwater corridors that extent across the map area, spaced 5 to 15 km apart. The ice-flow chronology is recorded both in the landforms and the striae record. Rare measurements record an old flowset oriented northwest-southeast. Well-defined indicators in cross-cutting relationships reveal a regional clockwise rotation in ice-flow directions evolving from an old southward to a late deglacial westward flow. Short-lived proglacial lakes, as evidenced by wave-cut scarps, beaches, terraced outwash and ice-contact deltas, and patches of glacial lake sediments, occupied paleodrainage areas to Hudson Bay and possibly headwaters of the Mackenzie River through the Taltson River system.

## **RÉSUMÉ**

La région cartographique de McCann Lake se situe dans la Province de Rae du Bouclier canadien. Les dépôts quaternaires y sont étendus et épais, et sont percés çà et là par des affleurements du substratum rocheux. Le till a largement été fuselé vers l'ouest-sud-ouest par un écoulement glaciaire de la fin de la déglaciation en provenance de la ligne de partage glaciaire du Keewatin et il est entaillé par des corridors d'eau de fonte sous-glaciaire de direction ouest-sud-ouest, espacés de 5 à 15 km, qui s'étendent à l'ensemble de la région cartographique. La chronologie des écoulements glaciaires est révélée à la fois par les reliefs glaciaires et le registre des stries. De rares mesures rendent compte d'un écoulement ancien d'orientation nord-ouest-sud-est. Des indicateurs d'écoulement glaciaire bien définis qui affichent des relations de recoupement révèlent une rotation régionale de la direction de l'écoulement glaciaire dans le sens des aiguilles d'une montre, depuis un écoulement ancien dirigé vers le sud jusqu'à un écoulement dirigé vers l'ouest à la fin de la déglaciation. Des lacs proglaciaires de courte durée, dont témoignent des escarpements sculptés par les vagues, des plages, des terrasses de délavage, des deltas de contact glaciaire, ainsi que des amas de sédiments glaciolacustres, occupaient le paléobassin versant de la baie d'Hudson, mais peut-être aussi le cours supérieur du fleuve Mackenzie, par le biais du réseau hydrographique de la rivière Taltson.

## **LICENCE AGREEMENT**

View the licence agreement at

<https://open.canada.ca/en/open-government-licence-canada>

## **ACCORD DE LICENCE**

Voir l'accord de licence à

<https://ouvert.canada.ca/fr/licence-du-gouvernement-ouvert-canada>

## **SHEET 1 OF 1, SURFICIAL GEOLOGY**

### **GENERAL INFORMATION**

Authors: A.S. Dyke and J.E. Campbell

Geology based on aerial photograph interpretation by A.S. Dyke, 2014 and fieldwork by J.E. Campbell (2014–2016).

Revisions by J.E. Campbell (2024)

Geological data conforms to Surficial Data Model v. 2.5.1 (Deblonde et al., 2024).

Accompanying field site, till composition and ice-flow datasets for NTS mapsheet 75-G are available in GSC Open File 8714 (Campbell et al., 2020).

Geomatics by L. Robertson

Cartography by D. Viner and A. Page

Scientific editing by A. Weatherston

Initiative of the Geological Survey of Canada, conducted under the auspices of the South Rae Bedrock and Surficial Mapping Activity, Rae Project as part of Natural Resources Canada's Geo-mapping for Energy and Minerals (GEM-2) program

Map projection Universal Transverse Mercator, zone 13  
North American Datum 1983

Base map at the scale of 1:50 000 from Natural Resources Canada, with modifications  
Elevations in metres above mean sea level

Mean magnetic declination 2025, 10°47'E, decreasing 7.7' annually  
Readings vary from 9°58'E in the SE corner to 11°35'E in the NW corner of the map.

This map is not to be used for navigational purposes.

Title photograph: Subglacial meltwater corridors composed of hummocks and ridges (foreground) are common in the area covered by the McCann NTS map sheet. Here, north of McArthur Lake, a corridor dissects well defined streamlined terrain composed of drumlinoid ridges (background). Photograph by J.E. Campbell. NRCan photo 2023-420

The Geological Survey of Canada welcomes corrections or additional information from users ([gscpublications-cgcpublishings@nrcan-rncan.gc.ca](mailto:gscpublications-cgcpublishings@nrcan-rncan.gc.ca)).

Data may include additional observations not portrayed on this map. See map info document accompanying the downloaded data for more information about this publication.

This publication is available for free download through the NRCan Open Science and Technology Repository (<https://ostrnrcan-dostrncan.canada.ca>).

### **MAP VIEWING FILES**

The published map is distributed as a Portable Document File (PDF), and may contain a subset of the overall geological data for legibility reasons at the publication scale.

### ***CARTOGRAPHIC REPRESENTATIONS USED ON MAP***

This map utilizes ESRI Cartographic Representations in order to customize the display of standard GSC symbols for visual clarity on the PDF of the map only. The digital data still contains the original symbol from the standard GSC symbol set. The following legend features have Cartographic Representations applied:

- Roche moutonnée, Striation, Geomorphology lines

### ***DEFINITION QUERIES USED ON MAP***

This map utilizes definition queries in order to customize the display for visualization on the PDF of the map only and does not affect the digital data. The following features have a definition query applied:

- Field stations

### ***DESCRIPTIVE NOTES***

The McCann Lake map area lies in the south Rae Craton of the Canadian Shield, in an area where the rocks are chiefly granite gneiss with substantial belts of metamorphosed sediments and basic intrusive rocks. This region of the Shield has extensive, but broken, drift cover characterized by streamlined glacial bedforms and parallel esker systems related to ice and meltwater flow during deglaciation from an ice divide located over Keewatin to the east-northeast (Prest et al., 1968; Aylsworth and Shilts, 1989; McMartin and Henderson, 2004).

Within the map area, bedrock outcrop constitutes about 15 to 20% of the surface. Exposures are distributed throughout the area, but are somewhat more common in the northwest quarter than elsewhere. Thin drift (unit Tv), where additional small exposures are common, occurs extensively in a 20 km wide belt of high ground trending south-southwest from the easternmost reach of Talston River. Elsewhere, the till is moderately thick and is extensively streamlined. Drumlins and flutings are characteristic of areas where till is thick (unit Tb). They form numerous densely packed fields, with features typically 200 to 400 m long, with some extending to lengths of 4 km. Crag-and-tail forms are characteristic of areas where rock outcrop and shallow subcrop is more common; individual tails are rarely longer than 1 km. The till is typically bouldery silty sand to sandy.

About 10 parallel subglacial meltwater corridors that are incised in the till cover trend southwest-west across the map area. They are spaced about 5 to 15 km apart, and are typically 2 to 4 km wide, several of which extend across the width (100 km) of the map area. The signature deposits of the corridors are large eskers, trains of either hummocky glaciofluvial deposits or terraced glaciofluvial deposits, and equally or more common trains of hummocky and/or ridged till. Each of these deposits repeatedly alternate along individual corridors. With the exception of the terraced proglacial deposits, the others formed subglacially. The hummocky and ridged till deposits within these corridors are a result of the subglacial meltwater flow, either as erosional till remnants of the surrounding till or a depositional diamicton features (Campbell et al., 2020). In this interpretation, much of the hummocky till formed by meltwater erosion of previously deposited streamlined till. In places (as also in NTS 75-B; Dyke et al., 2022) drumlins were not completely eroded. The hummocky till associated with the large esker nearest the northeast corner of the map area is interfingered with irregular ridges extending from the adjacent till blanket. The ridges there appear to be residuals of meltwater erosion and they retain flutings on flat tops. In form, they are transitional to Rogen moraines. Note that numerous areas of Rogen moraine mapped in this area by Aylsworth and Shilts (1989) are all interpreted

instead here as hummocky till, because these deposits, with the exception noted, lack organized ridge systems of Rogen moraines. In other instances, the hummocky till within the corridors is composed of a sandy diamicton whose morphology and composition suggest a depositional genesis (Haiblen, 2017; Vérité et al., 2023).

The main deglacial ice-flow pattern is well known. Across the breadth of the map area, the flow splayed (diverged) gradually from southwestward in the southwest corner to nearly westward at the north end. The ubiquity of streamlining of the till and copious reworking by meltwater indicates that during deglaciation, at least, the ice was warm-based everywhere. In several places within and near the subglacial corridors, the main (strong) streamline pattern is overprinted by weak flutings of local extent, including, in places, fluting of glaciofluvial and till hummocks (after hummock formation). Where they occur outside the corridor they can be seen as areas of late flow convergence toward adjacent subglacial cavities, probably close to the receding ice margins.

Systematic mapping of predominantly small erosional ice-flow indicators (e.g. striations) reveal at least five phases of ice flow (Campbell et al., 2020). Old flow(s) of unknown sense, to the southeast to south-southeast and/or the northwest to north-northwest were recorded at a few sites. This ice-flow event likely predates-Late Wisconsin glacial maximum and may represent flow from an ancestral Keewatin ice centre located to the northwest (e.g. Dyke and Prest, 1987). Well defined indicators (both landforms and striations) in crosscutting relationships indicate a regional clockwise rotation in ice-flow directions evolving from an old southward to a late deglacial westward flow (Fig. 1). These ice-flow indicators are grouped into four events based on the dominant recorded orientation. Ice-flow indicators of each event vary by 15 to 20 degrees. The south flow ( $185^{\circ}$  to  $200^{\circ}$ ; phase 1 in Fig. 1) is interpreted as an early flow, likely prior and during the Last Glacial Maximum (LGM). The dominant regional flow directions were to the southwest, possibly at LGM or early deglacial ( $\sim 212^{\circ}$  to  $230^{\circ}$ ; phase 2 in Fig. 1) shifting to west-southwest ( $\sim 239^{\circ}$  to  $254^{\circ}$ ; phase 3 in Fig. 1) followed by a late, overprinting westward ( $260^{\circ}$  to  $280^{\circ}$ ; phase 4 in Fig. 1) flow. The southwest-to-west regional flowsets (phases 2, 3, and 4) are attributed to the deglaciation.

In the landform record, the locally older southwest-oriented flowsets in this area are recorded by large bedforms of 2 to 4 km length, overprinted by distinctly smaller bedforms. In places, the reworking of the older bedforms has resulted in an echelon sets of younger drumlins (Ely et al., 2022). The widespread preservation of older southwest bedforms indicates that reworking of the drift sheet during the formation of the youngest flowset was far from complete. Hence, much of the drift dispersion is related to the older event(s), including the Phase 1 flow, which is preserved in the striae record (Campbell et al., 2020). The major topographic grain of the map area is aligned northeast-southwest, suggesting that these event(s) may have been more erosive.

Evidence of glacial lakes occurs in both major drainage systems. Most of the map area is in the Mackenzie River drainage basin. The Taltson River traverses the area from the southeast to the northwest, discharging to Great Slave Lake, which, during deglaciation, was occupied by glacial Lake McConnell (Craig, 1965; Smith, 1994). The northeast corner of the map area, about 10% of it, is in the headwaters of the Thelon River drainage, which flows towards Hudson Bay. Prior to the present work, there were no glacial lakes indicated in this region (Prest et al., 1968; Aylsworth and Shilts, 1989; Stokes and Clark, 2004). Nonetheless, there is widespread evidence of an ever-changing sequence of proglacial lakes in the form of shoreline features, esker deltas, lake spillways, and deltas and perched outwash terraces at the mouths of spillways and meltwater channels. Multiple

strandline levels (highest ~445 m a.s.l.) and spillway sequences indicate complexities of lake evolution that are not currently resolved.

In the Thelon headwaters, fragmentary shoreline features, generally only a few kilometres in length, occur 10 to 40 m above modern lake levels. This is high enough to divert drainage into the Taltson River. No distinct spillway is evident at the low point between the two drainages. This may indicate that water levels were contiguous across the drainage divide, which seems possible from shoreline levels between McArthur and Penylan lakes. Within the Taltson River drainage, evidence of glacial lakes at many levels is seen in shoreline features and more spectacularly in the form of spillways that opened up with progressive ice-marginal recession, the largest sequence of which is located south of McArthur Lake. Most, perhaps all, of these lakes occurred simply because the major tributaries of the Taltson River drain northeastward, toward the receding ice margin. The spillways terminate close to or just above modern lake/river levels, and glaciofluvial terraces graded to baselines slightly above modern lake levels, as well as low shorelines, indicate that sections of the river's gradient may have been isostatically reversed. An unusual set of 19 parallel northwest-trending meltwater channels occurs about 10 km north of McArthur Lake. These channels are interpreted as ice marginal or possibly lake spillways although there are no shorelines associated with their onsets.

Extensive bogs occupy nearly all topographic basins. They record paludification of about 20% of the surface. Some are fringed by shoreline features, which suggests that they may be underlain by lake sediment. If lake sediment is absent, then postglacial paludification has been a very effective geomorphic agent.

In the absence of end moraines, ice-marginal recession is assumed to have occurred in a roughly parallel manner up the deglacial flowlines. In this map area, that assumption is confirmed by numerous ice-contact escarpments that trend across meltwater corridors on the ice-proximal sides of esker nodes, esker deltas, or outwash terraces; by heads of outwash trains; by points of egress of proglacial meltwater channels; by trends of lake spillways that opened along the receding margin; and here and there by sets of crevasse fillings. Furthermore, some short "eskers" that join larger systems at high angles may be interpreted as features that formed along crevasses parallel to ice margins. Taking these features together, numerous short segments of ice margins can be recognized. Along the 100 km lengths of two most continuous eskers near the north and south ends of the map area, there are about 30 such features. The spacing is highly irregular and the average spacing of 3.3 km is several times the long-term average rate of recession across the area behind the Cree Lake Moraine, as approximated by Dyke (2004). Nevertheless, where features are closely spaced, 0.5 to 1 km apart, they provide a reasonable approximation of an annual recession rate in an area devoid of end moraines.

### **ACKNOWLEDGMENTS**

We would like to express our deepest appreciation to Martin Roy (supervisor from UQAM), Louis Robertson, Sally Pehrsson, Sean Eagles, and Guy Buller (GSC) as well as field assistants Gabriel Lauzon, Anna Haiblen, and Benjamin McNeil. Logistical support was provided by Helicopter Transport Services in La Ronge, Saskatchewan, Discovery Mining Service, Yellowknife, and Adam and Julie Duff of Scott's General Store, Stony Rapids.

## REFERENCES

- Aylsworth, J.M. and Shilts, W.W., 1989. Glacial features around the Keewatin Ice Divide: Districts of Mackenzie and Keewatin; Geological Survey of Canada, Paper 88-24. <https://doi.org/10.4095/127320>
- Campbell, J.E., McCurdy, M.W., Lauzon, G., Regis, D., and Wygergangs, M., 2020. Field data, till composition and ice-flow history, south Rae Craton, Northwest Territories: results from the GEM-2 South Rae project – Surficial Mapping activity; Geological Survey of Canada, Open File 8714, 1 .zip file. <https://doi.org/10.4095/327218>
- Craig, B.G., 1965. Glacial Lake McConnell and the surficial geology of parts of Slave River and Redstone River map areas, District of Mackenzie; Geological Survey of Canada, Bulletin 122. <https://doi.org/10.4095/100639>
- Deblonde, C., Campbell, J.E., Chow, W., Cocking, R.B., Huntley, D.H., Parent, M.P., Rice, J.M., Robertson, L., Smith, I.R., Weatherston, A.J., and Zawadzka, K., 2024. Surficial Data Model: the science language of the integrated Geological Survey of Canada data model for surficial geology maps; Geological Survey of Canada, Open File 8236, ver. 2.5.1, 1 .zip file. <https://doi.org/10.4095/332530>
- Dyke, A.S., 2004. An outline of North American deglaciation with emphasis on central and northern Canada; *in* Developments in Quaternary Sciences, (ed.) J. Ehlers and P.L. Gibbard; Elsevier, p. 373–424. [https://doi.org/10.1016/S1571-0866\(04\)80209-4](https://doi.org/10.1016/S1571-0866(04)80209-4)
- Dyke, A.S. and Prest, V.K., 1987. Late Wisconsinan and Holocene history of the Laurentide Ice Sheet; *Géographie physique et Quaternaire*, v. 41, p. 237–263. <https://doi.org/10.7202/032681ar>
- Dyke, A.S., Campbell, J.E., and Lauzon, G., 2022. Surficial geology, Abitau Lake, Northwest Territories, NTS 75-B; Geological Survey of Canada, Canadian Geoscience Map 350, scale 1:100 000. <https://doi.org/10.4095/330072>
- Ely, J.C., Stevens, D., Clark, C.D., and Butcher, F.E.G., 2022. Numerical modelling of subglacial ribs, drumlins, herringbones, and mega-scale glacial lineations reveals their developmental trajectories and transitions; *Earth Surface Processes and Landforms*, p. 1–23. <https://doi.org/10.1002/esp.5529>
- Haiblen, A.M., 2017. Glacial history and landform genesis in the Lac de Gras area, Northwest Territories. M.Sc. dissertation, Department of Earth Sciences, Simon Fraser University, Burnaby, British Columbia.
- McMartin, I. and Henderson, P., 2004. Evidence from Keewatin (Central Nunavut) for Paleo-Ice Divide Migration; *Géographie physique et Quaternaire*, v. 58, no. 2-3, p. 163–186. <https://doi.org/10.7202/013137ar>
- Prest, V.K., Grant, D.R., and Rampton, V., 1968. Glacial Map of Canada; Geological Survey of Canada, Map 1253A, scale 1:5 000 000. <https://doi.org/10.4095/108979>

Smith, D.G., 1994. Glacial Lake McConnell: paleogeography, age, duration, and associated river deltas, Mackenzie River basin, western Canada; Quaternary Science Reviews, v. 13, no. 9, p. 829–843.  
[https://doi.org/10.1016/0277-3791\(94\)90004-3](https://doi.org/10.1016/0277-3791(94)90004-3)

Stokes, C.R. and Clark C.D., 2004. Evolution of late glacial ice-marginal lakes on the northwestern Canadian Shield and their influence on the location of the Dubawnt Lake palaeo-ice stream; Palaeogeography, Palaeoclimatology, Palaeoecology v. 215, no. 1-2, p. 155–171. <https://doi.org/10.1016/j.palaeo.2004.09.006>

Vérité, J., Livingstone, S.J., Ravier, E., McMartin, I., Campbell, J.E., Lewington, E.L.M., Dewald, N., Clark, C.D., Sole, A.J., and Storrar, R.D., 2023. Conceptual model for the formation of bedforms along subglacial meltwater corridors (SMCs) by variable ice-water-bed interactions; Earth Surface Processes and Landforms.  
<https://doi.org/10.1002/esp.5725>

#### **ADDITIONAL INFORMATION**

The Additional Information folder of this product's digital download contains figures and tables that appear in the map surround as well as additional geological information not depicted on the map, nor this document, nor the geodatabase.

- Figure 1 PDF

#### **AUTHOR CONTACT**

Questions, suggestions, and comments regarding the geological information contained in the data sets should be addressed to:

J.E. Campbell  
Geological Survey of Canada  
601 Booth Street  
Ottawa ON  
K1A 0E8  
[Janet.Campbell@nrcan-rncan.gc.ca](mailto:Janet.Campbell@nrcan-rncan.gc.ca)

#### **COORDINATE SYSTEM**

Projection: Universal Transverse Mercator  
Units: metres  
Zone: 13  
Horizontal Datum: NAD83  
Vertical Datum: mean sea level

#### **BOUNDING COORDINATES**

Western longitude: 108°00'00"W  
Eastern longitude: 106°00'00"W  
Northern latitude: 62°00'00"N  
Southern latitude: 61°00'00"N

### **SOFTWARE VERSION**

Data has been originally compiled and formatted for use with ArcGIS™ desktop version 10.8.2 developed by ESRI®.

### **DATA MODEL INFORMATION**

#### **Surficial**

The Geological Survey of Canada (GSC) through the Geo-mapping for Energy and Minerals Program (GEM) has undertaken the Geological Map Flow to develop protocols for the collection, management (compilation, interpretation), and dissemination of surficial and bedrock geology data and map information. To this end, a data model has been created.

The Surficial Data Model (SDM) was designed using ESRI geodatabase architecture. The XML workspace document provided can be imported into a geodatabase, and the geodatabase will then be populated with the feature datasets, feature classes, tables, relationship classes, subtypes, and domains.

Shapefile and table (.dbf) versions of the data are included within the data. Column names have been simplified and the text values have been maintained within the shapefile attributes. The direction columns are numerical, to display rotation for points, and the symbol fields will hold the correct values to be matched to the appropriate style file.

For a more in depth description of the data model please refer to the official publication:

Deblonde, C., Campbell, J.E., Chow, W., Cocking, R.B., Huntley, D.H., Parent, M.P., Rice, J.M., Robertson, L., Smith, I.R., Weatherston, A.J., and Zawadzka, K., 2024. Surficial Data Model: the science language of the integrated Geological Survey of Canada data model for surficial geology maps; Geological Survey of Canada, Open File 8236, ver. 2.5.1, 1 .zip file. <https://doi.org/10.4095/332530>